1 **Endemic Chikungunya Fever in Kenyan Children**

- Doris K. Nyamwaya^{1, a}, Mark Otiende^{1,a}, Donwilliams O. Omuoyo¹, George Githinji¹, Henry K. 2
- 3 Karanja¹, John N. Gitonga¹, Zaydah de Laurent¹, James R. Otieno¹, Rosemary Sang², Everlyn
- Kamau¹, Stanley Cheruiyot¹, Edward Otieno¹, Charles N. Agoti¹, Philip Bejon^{1,3}, Samuel M. 4
- Thumbi^{4,5,6}, George M. Warimwe^{1,3*} 5

6

7	¹ KEMRI-Wellcome Trust Research Programme, P.O. Box 230-80108, Kilifi, Kenya; ² KEMRI-Centre
8	for Virus Research, Nairobi, Kenya; ³ Centre for Tropical Medicine and Global Health, University of

- Oxford, Old Road Campus, NDM Research Building, Oxford OX3 7FZ, UK; ⁴Paul G Allen School 9
- for Global Animal Health, Washington State University, Pullman, WA 99164-7090, USA; ⁵Centre for 10
- 11 Global Health Research, Kenya Medical Research Institute, P.O. Box 1578-4100, Kisumu, Kenya;⁶
- 12 Institute of Tropical and Infectious Diseases, University of Nairobi, P.O Box 19676 - 00202

13 Nairobi.

- 14
- 15 ^acontributed equally
- 16 *Correspondence:
- 17 George Warimwe, KEMRI-Wellcome Trust Research Programme, P.O. Box 230-80108, Kilifi, 18 Kenya
- 19 Email: gwarimwe@kemri-wellcome.org; Phone: +254(0)709 983 000
- 20

21 ABSTRACT

22	Background. Chikungunya virus (CHIKV) was first identified in Tanzania in 1952. Several epidemics
23	including East Africa are described, but there are no descriptions of longitudinal surveillance of
24	endemic disease. Here, we estimate the incidence of CHIKV and describe viral phylogeny in coastal
25	Kenya.
26	Methods. Over a 5-year period (2014-2018), 11,708 febrile illnesses in 5,569 children visiting two
27	primary healthcare facilities linked to a demographic surveillance system in coastal Kenya were
28	recorded and blood samples obtained. Reverse-transcriptase PCR was used to identify CHIKF cases
29	in 3,500 children randomly selected from the 5,569 children.
30	Results. We found CHIKF to be endemic in this setting, associated with 12.7% (95% CI 11.60, 13.80)
31	of all febrile presentations to primary healthcare. The prevalence of CHIKV infections among
32	asymptomatic children in a community survey was 0.7% (95% CI 0.22, 2.12). CHIKF incidence
33	among children <1 year of age was 1703 cases/100,000-person years and 46 cases/100,000-person years
34	among children aged ≥ 10 years. Recurrent CHIKF episodes, associated with fever and viraemia, were
35	observed among 19 of 170 children with multiple febrile episodes during the study period and
36	confirmed by genome sequencing. All sequenced viral genomes mapped to the ECSA genotype albeit
37	distinct from CHIKV strains associated with the 2004 East African epidemic.
38	Conclusions. CHIKF may be a substantial public health burden in primary healthcare on the East
39	African coast outside epidemic years, and recurrent infections are common.
40	

41 BACKGROUND

- 42 Chikungunya fever (CHIKF) is a zoonotic mosquito-borne febrile illness characterised by acute, often
- 43 chronic, debilitating polyarthralgia and polyarthritis that can last for months to years [1-3]. The
- 44 disease is caused by chikungunya virus (CHIKV), a positive sense RNA virus in the family
- 45 Togaviridae that was first isolated from a febrile patient in Tanzania in 1952 [4]. CHIKV transmission

46	between humans is mainly mediated by the geographically widespread Aedes aegypti and Ae.
47	albopictus mosquitoes, with geographic spread and spillover from sylvatic transmission cycles in
48	monkeys thought to account for the periodic re-emergence of human disease outbreaks [5-8].
49	The CHIKV genome is approximately 11.8kb in length and encodes four non-structural proteins
50	(nsP1 to nsP4) required for viral replication, three major structural proteins (the capsid protein, and
51	envelope proteins E1 and E2), and two small polypeptides (6K/TF and E3) [9]. Exposure to CHIKV
52	results in acquisition of protective virus neutralising antibodies that target the E2 protein, and is the
53	basis for ongoing efforts to develop CHIKF vaccines [10, 11]. Phylogenetic studies have defined three
54	CHIKV genotypes, namely East, Central and Southern Africa (ECSA), West Africa, and the Asian
55	genotype [12]. In 2004, CHIKV re-emerged in coastal Kenya causing one of the largest epidemics on
56	record, affecting millions of people as it spread along the Indian Ocean islands, India, southeast Asia
57	and Europe [8, 13, 14]. This epidemic was associated with emergence of the Indian Ocean Lineage
58	(IOL), which mapped within the ECSA genotype and included viruses with adaptive mutations in the
59	E1 protein that increased their transmissibility by Aedes albopictus mosquitoes [15, 16].
60	CHIKF epidemics are now frequently reported globally [8, 17], but despite its original discovery and
61	later re-emergence in East Africa, very little is known regarding inter-epidemic CHIKV exposure in
62	the region. CHIKF cases have previously been detected in children in Kenya and Tanzania during
63	inter-epidemic periods [18-20], which together with the high anti-CHIKV antibody seroprevalence
64	observed in these settings suggest endemic CHIKV transmission [21, 22]. To address these
65	knowledge gaps, we conducted a primary healthcare-based study linked to demographic surveillance
66	and community survey to estimate the prevalence and incidence of CHIKV infections among children
67	in Kilifi, coastal Kenya. Our study period, 2014 to 2018, included the most recent CHIKF epidemic
68	year, 2016 [23], allowing an unprecedented assessment of exposure risk and disease burden before,
69	during and after a CHIKF epidemic in this setting.

71 **METHODS**

72 **Study population**

73	We conducted a population-based observational cohort study between March 2014 and October 2018
74	at two dispensaries, Ngerenya and Pingilikani, based within the Kilifi Health Demographic
75	Surveillance System (KHDSS) [24]. The KHDSS covers an area of 891km ² with approximately
76	290,000 residents that are enumerated every four months at the household level [24]. Ngerenya and
77	Pingilikani have different rates of mosquito-borne pathogen exposure as inferred from previous
78	malaria studies; Ngerenya and environs experiences low intensity malaria transmission, whereas
79	malaria transmission around Pingilikani is moderate [25]. However, no studies on arbovirus exposure
80	had been conducted previously. Our approach was to screen children presenting with fever (axillary
81	temperature of \geq 37.5°C) at the dispensaries for CHIKV infection by reverse-transcriptase polymerase
82	chain reaction (RT-PCR). Children presenting at the dispensaries were linked to the KHDSS data
83	using a unique identifier, allowing us to estimate incidence in the population, and to determine the
84	influence of various demographic variables on the incidence estimates. Ethical approval was provided
85	by the Kenya Medical Research Institute Scientific and Ethics Review Unit (SSC Nos. 3296, 2617
86	and 3149).

87 **CHIKV RT-PCR assays**

88 CHIKV infection was defined as detection of CHIKV viral RNA by RT-PCR using a published

89 primer-probe set targeting the CHIKV non-structural protein 1 (nsP1) region[26], namely CHIKV 874

90 (5'-AAAGGGCAAACTCAGCTTCAC-3'), CHIKV 961 (5'-GCCTGGGCTCATCGTTATTC-3')

91 and CHIKV 899-FAM (5'-FAM-CGCTGTGATACAGTGGTTTCGTGTG-TAMRA-3'). Briefly,

92 100μ l of finger-prick blood samples collected at presentation to the dispensaries, and stored at -80°C,

- 93 was used for total RNA isolation using TRIzolTM Reagent (ThermoFisher) as per manufacturer
- 94 instructions. Presence of CHIKV viral RNA was then determined using the QuantiFast RT-PCR Kit
- 95 (Qiagen) in a 25µl reaction with primers and probe at a final concentration of 100nM and 20nM,
- 96 respectively, and 5µl total RNA template. RT-PCR assays were done on a 7500 Real-Time PCR

97	System (Applied Biosystems) with cycling conditions as follows: 50°C for 20 minutes, 95°C for 15
98	minutes, followed by 45 cycles of 94°C for 15 seconds and 60°C for 1 minute. A second confirmatory
99	screen was performed on some of the RT-PCR positive samples using an assay targeting the CHIKV
100	nsP4 region as previously described[27]. For both assays, a positive result was defined as a cycle
101	threshold (Ct) value of <40. Viral RNA from a cultured CHIKV isolate obtained from a febrile
102	Kenyan patient was used as a positive control, while RT-PCR mastermix without template was used
103	as a negative control.
104	CHIKV genome sequencing and analysis
105	RT-PCR positive CHIKV samples were sequenced using the Nanopore MinION technology following
106	PCR amplification using methods described in the PrimalSeq approach ¹⁷ . Full details of our
107	sequencing and bioinformatics workflow can be found in the Supplementary Material. In brief,
108	CHIKV-specific multiplex primers were designed using the Primal Scheme software and used for pre-
109	enrichment of CHIKV viral RNA in the samples using a multiplex PCR method. Barcodes and
110	sequencing adapters were ligated into each sample, after which all samples were pooled into a single
111	reaction tube and the library loaded on to a MinION sequencing device for sequencing. The sequences
112	generated in this study (N=10) were deposited in GenBank, accession numbers: MT526796-
113	MT526807. MUSCLE was used to align the Kilifi CHIKV genome sequences with those available in
114	GenBank from other geographical locations (see Supplementary Material). Maximum Likelihood
115	phylogenies were reconstructed from the alignment using RAxML[28] using the GTR substitution
116	model with 4 gamma categories (GTR+G4) and visualized in FigTree v1.4.4.

117 Sampling and statistical analyses

118 We selected a random sample of 3500 out of 5,669 children who visited the two dispensaries with

119 febrile illness. Associations between available demographic variables and CHIKV RT-PCR positivity

120 were assessed using odds ratios estimated using logistic regression models. CHIKF incidence within

121 the dispensary catchment area, defined as radius of 5km from each dispensary, was expressed per

122 100,000 person-years observed and incidence rate ratios (IRRs) comparing various demographic

123	variables determined by Poisson regression. Univariate analyses were performed using Chi2 test
124	(categorical variables) or the Mann-Whitney U test (continuous variables). All statistical analyses
125	were carried out in Stata TM version 15 with a two-sided p value <0.05 used for statistical significance.

126

127 **RESULTS**

128 CHIKV infections are common among children in coastal Kenya

- 129 Between March 2014 and October 2018, there were 29,819 visits by children aged up to 15 years to
- 130 the two dispensaries (6,746 in Ngerenya, 23,073 in Pingilikani), of which 13,696 were febrile (Figure
- 131 1). Of the 13,696 fevers, 1,256 lacked a unique patient identifier and 732 samples were missing. After
- these exclusions, there were 11,708 febrile visits from a total of 5,569 children (median 1 febrile visit
- 133 per child during the study period, IQR 1 2) eligible for CHIKV RT-PCR analysis (Figure 1).
- 134 A sample of 3,500 febrile children out of the 5,569 across the two dispensaries were randomly
- 135 selected and samples from their first (index) febrile episode during the study period screened for
- 136 CHIKV infection (Figure 1). The median age of these 3,500 children was 3.1 years (IQR 1.3 6.4),
- 137 with 1,701 being resident in Ngerenya and 1,799 in Pingilikani, respectively (Figure 1). Our study
- 138 period, 2014 to 2018, was at least 10 years since the 2004 CHIKF epidemic occurred in Kenya, with
- 139 ~90% our study population having been born after that epidemic.
- 140 Of the 3,500 children 443 were RT-PCR positive (12.7%, 95% CI 11.60, 13.80; Table 1), with RT-
- 141 PCR positivity being twice as frequent in Ngerenya (16.7%, 95% CI 15.00, 18.54) than in Pingilikani
- 142 (8.8%, 95% CI 7.61, 10.24). Most of the 443 CHIKF cases had a clinical diagnosis of upper
- 143 respiratory tract infection (URTI) but in no instance was a diagnosis of CHIKF assigned (Table 1).
- 144 The treatments prescribed for the management of CHIKF cases were comparable to those for other
- 145 causes of fever at the dispensaries, but fewer antimalarials were prescribed for CHIKF cases (Table
- 146 1). CHIKF case prevalence was highest in 2016 (23.1%), when an epidemic was reported in Kenya
- 147 [23], and was significantly lower in the pre- and post-epidemic years where it ranged between 5% and

- 148 8% (Table 1). In contrast, the prevalence of asymptomatic CHIKV infections estimated by cross-
- sectional sampling of a different group of 435 children in the same study location in 2016 was 0.7%
- 150 (95% CI 0.22, 2.12); this yielded a clinical-to-asymptomatic CHIKV infection ratio of 18:1 during the
- 151 epidemic year.

152 Incidence of CHIKF in the community

153 We calculated the incidence of CHIKF restricting our analysis to children aged up to 15 years whose

154 residence was within the dispensary catchment area. This represented a total of 139,840 person-years

- 155 of observation (pyo) over the study period at both locations. For the numerator (CHIKF cases), we
- applied the observed CHIKV RT-PCR positivity rates (Table 1) to the total number of febrile children
- 157 presenting to the respective dispensary on the assumption that the case positive rate among the 3,500
- 158 we randomly selected would hold true for the remaining 2,069 samples.
- 159 The incidence of CHIKF during the study period was 340 cases/100,000 pyo (95% CI 310, 373; Table

160 2). However, an inverse relationship was evident between CHIKF incidence and age, consistent with

- 161 acquisition of protective immunity against disease due to ongoing CHIKV exposure in this setting
- 162 (Table 2). CHIKF incidence was significantly higher in Ngerenya as compared to Pingilikani;
- 163 furthermore, while an increase in incidence was observed in Ngerenya during the 2016 CHIKF
- 164 epidemic, such an increase was not observed in Pingilikani where CHIKF incidence had been
- 165 gradually declining since 2014 (Table 2). Marked seasonality in CHIKF incidence was observed in
- 166 Pingilikani but not Ngerenya (Table 2) suggesting potential differences in ecological risk factors of
- 167 disease. A negative correlation was observed between CHIKF incidence and residential distance from
- 168 the dispensary at both locations, which likely reflected access to care (Table 2) as has been previously
- 169 reported for malaria incidence [29].

170 Recurrent CHIKF episodes

We determined whether recurrent CHIKF episodes occurred in our study population. To do this weidentified all 443 children with first or only episodes of CHIKF and screened all subsequent samples

173	collected during febrile episodes during the period 2014-2018. We defined recurrent episodes as
174	CHIKV RT-PCR positivity during these subsequent febrile dispensary visits. Of the 443 index cases,
175	170 presented to the dispensary with fever on at least one other occasion during the study period,
176	contributing a total of 320 subsequent febrile dispensary visits with varying durations since the
177	corresponding index CHIKF case (Figure 1). Of the 170 children, 19 (11.2%, 95% CI 6.86, 16.90)
178	were CHIKV RT-PCR positive on at least one subsequent febrile episode (Figures 1 and 2). The
179	duration between the index and the subsequent RT-PCR positive febrile episodes ranged from 2 to 43
180	months (Figure 2, and Supplementary Material Table S2). Eighteen of the 19 CHIKV RT-PCR
181	positive cases were detected during or after the 2016 CHIKF epidemic year. Neither age, sex nor
182	geographic location were statistically associated with recurrent episodes of CHIKF (Chi2 test p>0.05
183	for all) and the RT-PCR Ct values and clinical diagnoses were comparable between the index case
184	and corresponding subsequent timepoints (Supplementary Material Figure S2 and Table S2).

185 Viral phylogeny

186 We attempted to sequence all index and subsequent RT-PCR positive samples from the 19 children 187 with recurrent CHIKF episodes (Figure 2). Complete and partial genome coding sequences were 188 generated from 10 samples, including an additional index sample with a low Ct value (Ct=19) from a 189 child who did not experience recurrent CHIKF episodes during the study period. All CHIKV genomes 190 generated in this study mapped to the ECSA genotype, albeit distinct from the clade containing the 191 IOL strains that emerged during the 2004 epidemic and the clade including genomes from CHIKF 192 patients sampled in Mandera, northern Kenya, during the 2016 epidemic (Figure 3). The E1 A226V 193 mutation, associated with adaptation and increased transmissibility by Aedes albopictus [15], was 194 absent from all sequences sampled in Kenya [23]. Time-resolved phylogenies showed strong temporal 195 clustering of CHIKV sequences from coastal Kenya that may suggest antigenic changes over time 196 driven by acquisition of herd immunity (Figure 3). Index-recurrent episode case genome sequence 197 pairs were available for four children with respective time intervals of 2.9, 3.5, 22.3 and 28.3 months 198 between the index and recurrent episodes (Figure 3). The degree of relatedness of the sequence pairs

was similarly time-dependent, with the sequence pair with the longest time interval (28.3 months)being most divergent (Figure 3).

201

202 **DISCUSSION**

203 In summary, we find that CHIKF is endemic in coastal Kenya with the case burden being particularly 204 high in young children aged <1 year. Our population-based cohort approach in primary healthcare 205 facilities could detect an increase in CHIKF cases during the 2016 epidemic, supporting the utility of 206 such a surveillance framework in the early detection of CHIKF epidemics. During the epidemic year 207 an increase in CHIKF incidence was only noted in Ngerenya while that in Pingilikani (located 208 approximately 40km south of Ngerenya) had been declining [25]; in our dataset, clinical malaria 209 accounted for 57% of all fevers in Pingilikani but only 5% in Ngerenya, while URTI accounted for 210 70% of all fevers in Ngerenya but only 10% in Pingilikani. We observed significant CHIKF 211 incidence in both locations, and an inverse relationship between CHIKF incidence and age was 212 evident at both locations. This suggests that immunity is acquired in older children, and that CHIKF 213 transmission is significant in both locations despite the varying malaria ecology. 214 Consistent with previous reports of CHIKF in children in East Africa [19, 30], none of the patients in 215 this study had a clinical diagnosis of CHIKF or reported symptoms of joint pathology. Nineteen 216 children were CHIKV RT-PCR positive during subsequent febrile episodes, with viral genome 217 sequences from these recurrent episodes not showing high degrees of relatedness to the index case. 218 Chronic relapsing and remitting CHIKF symptoms have been observed in adult patients during 219 previous epidemics, as an inflammatory post-infectious arthropathy [31-33] with viral RNA detected 220 in the joint tissues in one study [34]. However, we are aware of no reports of recurrent discrete 221 CHIKF episodes associated with reinfection. This may simply be due to the rarity of longitudinal 222 surveillance studies for CHIKF, particularly in children. In our study, the relatedness of virus 223 genomes from first and second infections in the same child was similar to the relatedness of virus 224 genomes from different children, suggesting that these were reinfections rather than relapses.

225 Neutralising antibodies that develop following CHIKV infection have been shown to correlate with 226 decreased risk of clinical illness in a longitudinal study in the Phillipines [35, 36], though CHIKF 227 cases in that study were all due to the Asian CHIKV genotype. The possibility that the ECSA 228 genotype viruses, which tend to be more diverse [12] and associated with a higher clinical disease 229 burden than Asian genotypes [37], results in a variable immune response in children in East Africa 230 needs to be ruled out. The inverse relationship between age and incidence of CHIKF suggests that 231 there is an acquisition of protective immunity, although the recurrent infections suggest that this 232 immunity is not rapidly acquired in some children. 233 Only finger-prick whole blood samples for RT-PCR analysis were available for this study, thus 234 precluding any immunological work. Future longitudinal studies with serial blood sampling for 235 immunological assays will help determine the natural course of infection, including duration of 236 viraemia, and identify host and viral factors that may underlie the occurrence of CHIKF relapses in 237 coastal Kenya. 238 Consistent with other reports of CHIKV strains circulating in East Africa [12, 23], the viral genomes 239 generated in this study all mapped to the ECSA genotype. Further, in contrast to the 2016 CHIKF 240 epidemic viruses from Mandera (northern Kenya) that formed a single well-supported clade, 241 sequences from coastal Kenya were highly diverse and fell within multiple well-supported clades that 242 whose divergence increased with time. Together, these data suggest ongoing endemic CHIKV 243 transmission at the Kenyan coast possibly sustained through frequent emergence of diverse CHIKV 244 strains from local sylvatic transmission cycles [6].

245 Our data and that from others in the region [30] further reinforce the need for inclusion of CHIKF

high on the list of differential diagnoses to consider when faced with a febrile child at primary

247 healthcare facilities in coastal Kenya. Access to rapid diagnostics would be of value in antibiotic

stewardship, and control of transmission in the community would substantially reduce the burden on

249 healthcare facilities. Further studies on the associated clinical outcomes, including the incidence of

- 250 severe ('atypical') manifestations of CHIKF, should help determine the wider public health
- 251 significance of CHIKF in coastal Kenya.

252

253 FUNDING

- 254 This work was commissioned by the National Institute for Health Research (NIHR) Global Health
- 255 Research programme [16/136/33] using UK aid from the UK Government. The views expressed in
- this publication are those of the authors and not necessarily those of the NIHR or the Department of
- 257 Health and Social Care. GMW is supported by an Oak foundation fellowship and a Wellcome Trust
- 258 grant [grant number 203077_Z_16_Z].
- 259

260 CONFLICT OF INTEREST

- 261 All authors declare no conflicts of interest.
- 262

263 ACKNOWLEDGEMENTS

- 264 This manuscript was submitted for publication with permission from the Director of the Kenya
- 265 Medical Research Institute.

266

267 **REFERENCES**

- 268 1. Robinson MC. An epidemic of virus disease in Southern Province, Tanganyika Territory, in
- 269 1952-53. I. Clinical features. Trans R Soc Trop Med Hyg **1955**; 49(1): 28-32.

- 270 2. Lumsden WH. An epidemic of virus disease in Southern Province, Tanganyika Territory, in
- 271 1952-53. II. General description and epidemiology. Trans R Soc Trop Med Hyg **1955**; 49(1):
- 272 33-57.
- 273 3. Tritsch SR, Encinales L, Pacheco N, et al. Chronic joint pain 3 years after chikungunya virus
- 274 infection largely characterized by relapsing-remitting symptoms. J Rheumatol **2019**.
- 4. Ross RW. The Newala epidemic. III. The virus: isolation, pathogenic properties and
- 276 relationship to the epidemic. J Hyg (Lond) **1956**; 54(2): 177-91.
- 277 5. Althouse BM, Guerbois M, Cummings DAT, et al. Role of monkeys in the sylvatic cycle of
- chikungunya virus in Senegal. Nat Commun **2018**; 9(1): 1046.
- 279 6. Eastwood G, Sang RC, Guerbois M, Taracha ELN, Weaver SC. Enzootic Circulation of
- 280 Chikungunya Virus in East Africa: Serological Evidence in Non-human Kenyan Primates.
- 281 Am J Trop Med Hyg **2017**; 97(5): 1399-404.
- 282 7. Weaver SC, Lecuit M. Chikungunya virus and the global spread of a mosquito-borne disease.
- 283 N Engl J Med **2015**; 372(13): 1231-9.
- 8. Wahid B, Ali A, Rafique S, Idrees M. Global expansion of chikungunya virus: mapping the
 64-year history. Int J Infect Dis 2017; 58: 69-76.
- 9. Voss JE, Vaney MC, Duquerroy S, et al. Glycoprotein organization of Chikungunya virus
 particles revealed by X-ray crystallography. Nature 2010; 468(7324): 709-12.
- 288 10. Rezza G, Weaver SC. Chikungunya as a paradigm for emerging viral diseases: Evaluating
- disease impact and hurdles to vaccine development. PLoS Negl Trop Dis 2019; 13(1):
 e0006919.
- 291 11. Gouglas D, Thanh Le T, Henderson K, et al. Estimating the cost of vaccine development
 292 against epidemic infectious diseases: a cost minimisation study. Lancet Glob Health 2018;
 293 6(12): e1386-e96.
- Volk SM, Chen R, Tsetsarkin KA, et al. Genome-scale phylogenetic analyses of chikungunya
 virus reveal independent emergences of recent epidemics and various evolutionary rates. J
 Virol 2010; 84(13): 6497-504.

- 297 13. Chretien JP, Anyamba A, Bedno SA, et al. Drought-associated chikungunya emergence along
- 298 coastal East Africa. Am J Trop Med Hyg **2007**; 76(3): 405-7.
- 299 14. Sergon K, Njuguna C, Kalani R, et al. Seroprevalence of Chikungunya virus (CHIKV)
- 300 infection on Lamu Island, Kenya, October 2004. Am J Trop Med Hyg **2008**; 78(2): 333-7.
- 30115.Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in chikungunya
- 302 virus affects vector specificity and epidemic potential. PLoS Pathog **2007**; 3(12): e201.
- 303 16. Weaver SC, Forrester NL. Chikungunya: Evolutionary history and recent epidemic spread.
- 304 Antiviral Res **2015**; 120: 32-9.
- 305 17. Nsoesie EO, Kraemer MU, Golding N, et al. Global distribution and environmental suitability
 306 for chikungunya virus, 1952 to 2015. Euro Surveill 2016; 21(20).
- 307 18. Kajeguka DC, Kaaya RD, Mwakalinga S, et al. Prevalence of dengue and chikungunya virus
- infections in north-eastern Tanzania: a cross sectional study among participants presenting
 with malaria-like symptoms. BMC Infect Dis **2016**; 16: 183.
- 310 19. Chipwaza B, Mugasa JP, Selemani M, et al. Dengue and Chikungunya fever among viral
- 311 diseases in outpatient febrile children in Kilosa district hospital, Tanzania. PLoS Negl Trop
- 312 Dis **2014**; 8(11): e3335.
- 313 20. Hertz JT, Munishi OM, Ooi EE, et al. Chikungunya and dengue fever among hospitalized
 314 febrile patients in northern Tanzania. Am J Trop Med Hyg **2012**; 86(1): 171-7.
- 315 21. Weller N, Clowes P, Dobler G, et al. Seroprevalence of alphavirus antibodies in a cross-
- 316 sectional study in southwestern Tanzania suggests endemic circulation of chikungunya. PLoS
 317 Negl Trop Dis 2014; 8(7): e2979.
- LaBeaud AD, Banda T, Brichard J, et al. High rates of o'nyong nyong and Chikungunya virus
 transmission in coastal Kenya. PLoS Negl Trop Dis 2015; 9(2): e0003436.
- 320 23. Maljkovic Berry I, Eyase F, Pollett S, et al. Global Outbreaks and Origins of a Chikungunya
- 321 Virus Variant Carrying Mutations Which May Increase Fitness for Aedes aegypti:
- 322 Revelations from the 2016 Mandera, Kenya Outbreak. Am J Trop Med Hyg **2019**; 100(5):
- 323 1249-57.

- 324 24. Scott JA, Bauni E, Moisi JC, et al. Profile: The Kilifi Health and Demographic Surveillance
- 325 System (KHDSS). Int J Epidemiol **2012**; 41(3): 650-7.
- 326 25. Mogeni P, Williams TN, Fegan G, et al. Age, Spatial, and Temporal Variations in Hospital
- 327 Admissions with Malaria in Kilifi County, Kenya: A 25-Year Longitudinal Observational
- 328 Study. PLoS Med **2016**; 13(6): e1002047.
- Lanciotti RS, Kosoy OL, Laven JJ, et al. Chikungunya virus in US travelers returning from
 India, 2006. Emerg Infect Dis 2007; 13(5): 764-7.
- 27. Pabbaraju K, Wong S, Gill K, Fonseca K, Tipples GA, Tellier R. Simultaneous detection of
- Zika, Chikungunya and Dengue viruses by a multiplex real-time RT-PCR assay. J Clin Virol
 2016: 83: 66-71.
- Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and
 user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics 2019;
 35(21): 4453-5.
- Bejon P, Williams TN, Nyundo C, et al. A micro-epidemiological analysis of febrile malaria
 in Coastal Kenya showing hotspots within hotspots. Elife **2014**; 3: e02130.

339 30. Waggoner J, Brichard J, Mutuku F, et al. Malaria and Chikungunya Detected Using

- Molecular Diagnostics Among Febrile Kenyan Children. Open Forum Infect Dis 2017; 4(3):
 ofx110.
- 342 31. Couturier E, Guillemin F, Mura M, et al. Impaired quality of life after chikungunya virus
 343 infection: a 2-year follow-up study. Rheumatology (Oxford) 2012; 51(7): 1315-22.
- 344 32. Staikowsky F, Le Roux K, Schuffenecker I, et al. Retrospective survey of Chikungunya
 345 disease in Reunion Island hospital staff. Epidemiol Infect 2008; 136(2): 196-206.
- 346 33. Chang AY, Encinales L, Porras A, et al. Frequency of Chronic Joint Pain Following
- Chikungunya Virus Infection: A Colombian Cohort Study. Arthritis Rheumatol 2018; 70(4):
 578-84.
- 349 34. Hoarau JJ, Jaffar Bandjee MC, Krejbich Trotot P, et al. Persistent chronic inflammation and
 infection by Chikungunya arthritogenic alphavirus in spite of a robust host immune response.
 J Immunol 2010; 184(10): 5914-27.

352	35.	Yoon IK, Alera MT, Lago CB, et al. High rate of subclinical chikungunya virus infection and
353		association of neutralizing antibody with protection in a prospective cohort in the Philippines.
354		PLoS Negl Trop Dis 2015 ; 9(5): e0003764.
355	36.	Yoon IK, Srikiatkhachorn A, Alera MT, Fernandez S, Cummings DAT, Salje H. Pre-existing
356		chikungunya virus neutralizing antibodies correlate with risk of symptomatic infection and
357		subclinical seroconversion in a Philippine cohort. Int J Infect Dis 2020.
358	37.	Bustos Carrillo F, Collado D, Sanchez N, et al. Epidemiological Evidence for Lineage-
359		Specific Differences in the Risk of Inapparent Chikungunya Virus Infection. J Virol 2019;
360		93(4).
361		
362		
363		
364		
365		
366		
367		
368		
369		
370		
371		
372		

373 TABLES

Table 1. Demographic characteristics of study participants

	CHIKF cases /N	Crude OR (95% CI)	P value
	(%)		
Dispensary			
Ngerenya (Kilifi North)	284/1701 (16.7)	Ref	
Pingilikani (Kilifi South)	159/1799 (8.8)	0.5 (0.39, 0.59)	< 0.001
Gender			
Female	221/1647 (13.4)	Ref	
Male	222/1853 (12.0)	0.9 (0.72, 1.07)	0.2
Age (years)			
< 1	79/601 (13.1)	Ref	
1 to <5	220/1700 (12.9)	1.0 (0.75, 1.29)	0.9
5 to <10	109/899 (12.1)	0.9 (0.67, 1.24)	0.6
10 to 15	35/300 (11.7)	0.9 (0.57, 1.33)	0.5
Year			
2014	43/509 (8.4)	Ref	
2015	48/869 (5.5)	0.6 (0.41, 0.97)	0.04
2016	208/902 (23.1)	3.2 (2.29, 4.60)	< 0.001
2017	108/674 (16.0)	2.1 (1.42, 3.01)	< 0.001
2018	36/546 (6.6)	0.8 (0.48, 1.21)	0.3
Season			
Jan – Mar	142/855 (16.6)	Ref	
Apr – Jun	134/1070 (12.5)	0.7 (0.56, 0.93)	0.01
Jul – Sep	97/939 (10.3)	0.6 (0.44, 0.76)	< 0.001
Oct – Dec	70/636 (11.0)	0.6 (0.46, 0.84)	0.002

Clinical diagnosis					
Other	61/661 (9.2)	Ref			
Malaria	83/690 (12.0)	1.3 (0.95, 1.91)	0.1		
Pneumonia	27/300 (9.0)	1.0 (0.60, 1.56)	0.9		
URTI	218/1428 (15.3)	1.8 (1.31, 2.39)	< 0.001		
Gastroenteritis	14/95 (14.7)	1.7 (0.91, 3.18)	0.1		
Undifferentiated fever	40/326 (12.3)	1.4 (0.90, 2.10)	0.1		
Treatment					
Antibiotic	313/2393 (13.1)	1.1 (0.90,1.40)	0.3		
Antimalarial	117/1104 (10.6)	0.8 (0.60, 0.94)	0.01		
Other	41/291 (14.1)	1.1 (0.81, 1.62)	0.4		

375

376

Table 2. Incidence of CHIKF in the study population

	Ngerenya N=74,799 person-years		Pingilikani N=65,041 person-years	
	Incidence	IRRs	Incidence	IRRs
	(95% CI)	(95% CI)	(95% CI)	(95% CI)
Overall incidence	408 (363, 457)	-	267 (228, 311)	-
Year				
2014	-	-	643 (513, 794)	Ref
2015	67 (32, 123)	Ref	289 (205, 397)	0.43 (0.30, 0.64)
2016	1205 (1035, 1395)	18.08 (9.56, 34.18)	187 (120, 279)	0.28 (0.18, 0.44)
2017	611 (493, 750)	9.17 (4.77, 17.61)	62 (27, 122)	0.09 (0.05, 0.19)

2018	104 (56, 178)	1.57 (0.69, 3.57)	93 (44, 170)	0.14 (0.07, 0.27)
Gender				
Female	382 (321, 452)	Ref	328 (268, 398)	Ref
Male	432 (368, 505)	1.13 (0.90, 1.42)	207 (160, 264)	0.63 (0.46, 0.86)
Age (years)				
< 1	2209 (1776, 2715)	Ref	1198 (856, 1631)	Ref
1 to <5	791 (670, 927)	0.36 (0.28, 0.46)	587 (476, 717)	0.49 (0.34, 0.71)
5 to <10	198 (146, 263)	0.09 (0.06, 0.13)	76 (44, 124)	0.06 (0.04, 0.11)
10 to 15	24 (9, 52)	0.01 (0.005, 0.02)	68 (38, 112)	0.06 (0.03, 0.10)
Distance from				
dispensary (Km)				
<1	800 (531, 1156)	Ref	560 (306, 939)	Ref
1 to <2	930 (742, 1152)	1.16 (0.76, 1.79)	1038 (760, 1384)	1.85 (1.02, 3.37)
2 to <3	657 (531, 804)	0.82 (0.54, 1.25)	837 (658, 1049)	1.50 (0.85, 2.65)
3 to <4	513 (396, 654)	0.64 (0.41, 1.00)	224 (128, 363)	0.40 (0.20, 0.82)
4 to <5	118 (70, 186)	0.15 (0.08, 0.27)	118 (59, 211)	0.21 (0.10, 0.46)
5 to <6	34 (13, 74)	0.04 (0.02, 0.10)	20 (7, 43)	0.04 (0.01, 0.09)
Season				
Jan - Mar	463 (371, 572)	Ref	227 (160, 312)	Ref
Apr - Jun	348 (269, 444)	0.75 (0.54, 1.04)	528 (422, 651)	2.33 (1.58, 3.42)
Jul - Sep	423 (335, 527)	0.91 (0.67, 1.24)	234 (166, 321)	1.03 (0.66, 1.62)
Oct - Dec	394 (304, 503)	0.85 (0.62, 1.17)	50 (20, 103)	0.22 (0.09, 0.49)

381 FIGURE LEGENDS

- Figure 1: Study population. The flow of study participants, reasons for exclusion and results from
 CHIK RT-PCR screening are shown.
- 384 Figure 2: Recurrent CHIKF episodes. The distribution of febrile episodes for each of 19 children
- 385 who had at least two CHIKV RT-PCR positive febrile episodes is shown. The timing of the CHIKV
- 386 RT-PCR positive results is shown, with the index episode shown as the first data point in the follow
- 387 up timeline. Children with prefix 'NGE' were resident in Ngerenya and those with 'PIN' from
- 388 Pingilikani, respectively. Further details of these children can be found in Supplementary Material
- 389 Table S2.

390 Figure 3: Phylogenetic analysis of CHIKV genomes from Kenya. BEAST MCC tree of 123 ECSA

391 genomes collected across the world, including 24 from Kenya is shown in the left panel. On the right

392 panel is an extraction highlighting the CHIKV sequences generated in this study. The tips are

- 393 coloured according to geographic location or child identity (as in Figure 2, for those with paired
- index-recurrent episode sequence pairs), with posterior probability support shown for select nodes
- 395 with probability >0.8. The same and more detailed MCC tree can be found in the Supplementary
- 396 Material Figure S1.
- 397
- 398
- 399
- 400
- 401
- 402
- 403

- 405
- 406
- 407
- 408 FIGURES
- 409 Figure 1



411

412 **Figure 2**



Figure 3

