## Ischemic Stroke Risk in Patients on Direct Oral Anticoagulants with

#### Levetiracetam: A Pharmacovigilance Study

Mohammed Abou Kaoud, Ran Nissan, Amitai Segev, Avi Sabbag, David Orion, and Elad Maor

#### Running title: Stroke in DOACS and Levetiracetam

(a) Chaim Sheba Medical Center, Tel Hashomer, Israel(b) Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel

#### **Corresponding author:**

Elad Maor, M.D., Ph.D. Sheba Medical Center and Tel-Aviv University Phone: 972-35302646 Fax: +972-35305789 Email: elad.maor@sheba.health.gov.il Twitter: @maor\_elad Postal address: Dr. Elad Maor, The Olga & Lev Leviev Heart Center Sheba Medical Center Hospital- Tel Hashomer 52621 Ramat Gan, ISRAEL

Funding/Support: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Manuscript total word count: 5632

#### Abstract

Background: Levetiracetam is widely used in post stroke epilepsy. However, it is suspected to possess P-glycoprotein induction properties and therefore a potential significant interaction with DOACs . Our aim was to search for ischemic stroke signals with levetiracetam and the DOACs.

Methods: In this retrospective, pharmacovigilance study, we used the Food and Drug Administration adverse event reporting system to identify ischemic stroke events associated with DOACs and concomitant use of levetiracetam. We evaluated disproportionate reporting by the reporting odds ratio adjusted to age and sex (adj.ROR) and the lower bound of the shrinkage 95% confidence interval ( $\Omega_{025} > 0$  is deemed significant for an interaction).

Results: We identified 1,841 (1.5%), 3,731 (5.3%), 338 (4.9%), and 1,723 (1.3%), ischemic stroke reports with apixaban, dabigatran, edoxaban, and rivaroxaban respectively. When heparin/enoxaparin was used as the comparator the adjusted ROR of the interaction effect was 3.57 (95%CI, 2.81–4.58) between DOACs and

levetiracetam. The shrinkage analysis detected an interaction between each of the DOACs and levetiracetam resulting in higher reports of ischemic stroke with the combination compared to each drug alone. The logistic model and shrinkage analysis failed to detect an interaction when queried for hemorrhagic stroke. Conclusions: We show a strong signal for the levetiracetam interaction with apixaban, dabigatran, edoxaban, and rivaroxaban leading to a 3-5 folds increased reporting risk of ischemic stroke. Our findings suggest the need for pharmacodynamic monitoring, while concomitantly prescribing levetiracetam with the DOACs.

#### INTRODUCTION

The introduction of direct oral anticoagulants (DOACs) has significantly altered the management of atrial fibrillation (AF) due to their improved safety profile and ease of use compared with warfarin. Contemporary guidelines prefer DOACs over warfarin for stroke prevention among patients with AF, excluding patients with mechanical valve replacement or significant mitral stenosis (1). While there is no preference for one specific DOAC over another, in the clinic DOACs are personalized according to patient age, gastrointestinal bleeding risk, chronic kidney disease, and drug interactions (2).

Since stroke is one of the most common structural etiologies of epilepsy, the concomitant use of DOACs and an anticonvulsant is not uncommon (3). Post-stroke seizures account for 11% of all epilepsy, 22% of all cases of status epilepticus, and 55% of newly diagnosed seizures amongst older people (4, 5, 6). Since AF is a common cause of ischemic stroke, the rate of AF-stroke-epilepsy

triplet, and thus DOAC-anticonvulsant treatment, is not uncommon among older adults (7).

To date, the combination of DOAC and some anticonvulsants remains controversial. The European Heart Rhythm Association (EHRA) 2018 guide did not support using anticonvulsants (carbamazepine, levetiracetam, phenobarbital, phenytoin, topiramate, and valproic acid) in patients concurrently taking DOACs (8). The 2021 EHRA guide states that after inquiry with the drug manufacturer there is unfortunately no study which reliably investigated the effect of levetiracetam on DOAC plasma levels and clinical events in a sufficiently large 'real world' cohort of concomitantly treated patients (33). However, in practice, levetiracetam is widely used for post-stroke epilepsy due to its favorable safety profile and fewer drug interactions.

Therefore, our study aim was to investigate reports to the Federal Drug Agency (FDA) regarding ischemic and non-ischemic stroke with and without levetiracetam combination in AF patients treated with DOACs.

#### **METHODS**

An observational, retrospective pharmacovigilance study was carried out using the FAERS database, a global repository of voluntary reports by healthcare professionals and consumers, and mandatory reports from manufacturers (9). The database was screened for reports containing the following terms in their brand or generic names: 'rivaroxaban,' 'apixaban,' 'dabigatran,' 'dabigatran etexilate mesylate,' dabigatran etexilate,' 'edoxaban,' 'edoxaban mesylate,' 'edoxaban tosylate,' 'enoxaparin,' 'enoxaparin sodium,' 'heparin,' 'heparin sodium,' 'heparin calcium,' 'levetiracetam,' 'carbamazepine' and 'omeprazole.' Omeprazole and carbamazepine are, respectively, negative and positive controls in this study.

The study included patients who were reported as the primary suspects for a given adverse event (AE) when novel anticoagulants accessed the market (2012-2023). In case multiple reports of the same event were detected, only the latest

case version of every event was retained, as recommended by the FDA. We further applied a population-linkage program to detect suspected duplicate reports of the same drug-event pair with different case numbers by screening for identical values in six key fields: age, sex, event date, country of occurrence, concomitant medications, and the same reasons for use. These suspected duplicate reports were excluded.

The database was then searched for the following adverse events in their Medical Dictionary for Regulatory Activities (MedDRA) preferred terms: 'ischemic stroke,' 'hemorrhagic stroke,' 'cerebral hemorrhage,' 'intracranial hemorrhage,' 'cerebrovascular accident,' 'lacunar stroke,' 'cerebellar stroke,' 'cerebral artery occlusion,' 'basal ganglia stroke', 'vertebrobasilar stroke', 'brain stem stroke', 'thrombotic stroke' and 'embolic stroke.' Those adverse events either denote a lack of effectiveness or toxicity. Detailed lists of terms included in our database guery are available at http://bioportal.bioontology.org/. The data was then filtered by DOAC's reason for use: embolic stroke in AF patients' treatment and prophylaxis. The dataset was then queried for cases with concomitant anticonvulsants used. Cases involving anticonvulsants were identified by a predefined list of anticonvulsant medications constructed using the FDA National Drug Code (NDC) file (10). Only patients above age 12 and cases with one reported anticonvulsant were included.

# Study End Points

The predefined primary endpoint was any MedDRA-preferred term describing ischemic stroke. Additional information for each report in the database, including demographic information (country, reporter occupation, reporting year, age, and sex), concomitant anticonvulsant uses, and date of AE occurrence and its outcomes, were collected for the analysis. Cases were defined as serious medical events if one or more of the following outcomes were reported: death, lifethreatening event, hospitalization, disability, or another serious medical event.

# Statistical Analysis

Descriptive statistics were calculated for patient demographics. Means and Standard deviations were generated for continuous variables. Frequencies and proportions were stated for categorical variables. A validated case-non-case method in drug safety research assessed whether ischemic and non-ischemic strokes were reported more frequently with the DOACs: apixaban, edoxaban, rivaroxaban, or dabigatran, compared with enoxaparin or heparin. We further analyzed each of the DOACs compared to the other three for reported ischemic stroke.

A logistic regression model obtained an adjusted reporting odds ratio (adj. ROR). The regression model included age, sex, and concomitant levetiracetam use. The ROR is a statistical model surrogate to the odds ratio and is an acceptable method to detect signals of adverse events and drug interactions (11, 12). Missing sex data were coded as 'Not Specified,' and the median value imputed missing age data in each group of drug recipients. R-squared and adjusted R-squared were used to assess model fitness. The ROR for concomitant levetiracetam was used to measure the strength of the association of the drug interaction.

Additionally, a refined model called the  $\Omega$  shrinkage measure was used to calculate the observed-to-expected for detecting signals of potential drug-drug interactions (DDIs). Omega  $\Omega$  is a robust observed-to-expected triplet measure of disproportionate reporting developed by the Uppsala Monitoring Centre (12, 13). When  $\Omega$  is positive, and two drugs are used together, an increased risk of a specific adverse event occurrence is emphasized over the sum of the individual risks when these same drugs are used separately (13). Thus,  $\Omega_{025} > 0$ , a positive lower bound of 95% CI, is used as a threshold for detecting the signals of the concomitant use of drug  $D_1$  and drug  $D_2$  (12). In other words, it indicates the frequency of reporting specific Drug-Drug-Event triplets in the dataset compared to what is expected based on the relative reporting for each drug alone (see eq. 1).

$$\Omega(0.25) = \Omega - \frac{\phi(0.975)}{\log 2\sqrt{n111}}$$
 (eq. 1)

Where  $\emptyset(0.975)$  is the standard normal distribution, and n111 is the number of

cases with the Drug-Drug-Event triplet.

The logistic regression analyses were performed using SPSS version 27. Two-sided

P < .05 indicated significance. Shrinkage calculation steps were inputted in Microsoft Excel version 2022. Noguchi Y. et al. 2019 described the complete shrinkage calculation (12).

## Sensitivity Analysis

First, we examined whether the drug interaction also manifests as hemorrhagic stroke. A model for hemorrhagic stroke comparing each of the DOACs; apixaban, rivaroxaban and dabigatran against the other two was performed. Edoxaban was not included in this analysis since there were no reports of hemorrhagic events with the levetiracetam-edoxaban combination. A sensitivity analysis for the levetiracetam drug interaction, included two drug combinations: apixaban-carbamazepine and apixaban-omeprazole. The former drug interaction is well established. Carbamazepine is known to induce Pglycoprotein and CYP3A4, thus reducing apixaban effectiveness (FDA and European Medical Agency physician guidance; 14,15). In contrast to carbamazepine, the combination of apixaban and omeprazole is widespread and considered safe without any known or potential drug interaction (16).

We further stratified DOACs by publication date in an additional analysis up to 2020, aiming to mitigate reporting bias. We hypothesized that from 2020 onwards, increased reports regarding DOACs, and ischemic stroke may be expected since the publication by Giustazzi et al. showed decreased effectiveness with concomitant DOAC-anticonvulsant use. Rivaroxaban was selected for further analysis since most patients in the Giustazzi et al. study were treated with rivaroxaban and more case reports were published regarding rivaroxaban (17). Further, we re-analyzed our results for the period 2015-2023, this is to account for the fact that during 2010-2014 a trial had been ongoing against Boehringer Ingelheim, dabigatran's manufacturer (18). Thus, to mitigate any additional negative reporting bias, we analyzed our data excluding the years 2012-2014.

# Standard Protocol Approvals, Registrations, and Patient Consents

The FAERS Public Dashboard is a publicly available web-based tool containing mandatory data reports from drug manufacturers and voluntary ADR reports from consumers and healthcare professionals. Hence an informed consent or ethical statement of approval by an ethical standards committee on human experimentation is not required for analysis.

## RESULTS

## **Demographics and Patient Characteristics**

The FAERS database included 19,609,956 unique safety reports from January 2012 to January 2023. Out of these reports, 125,799 for Apixaban, 69,993 for

Dabigatran, 6,965 for Edoxaban, and 136,710 for Rivaroxaban. Among eligible patients in the FAERS database that reported any of the MedDRA preferred terms for ischemic stroke are 1,841 (1.5%), 3,731 (5.3%), 338 (4.9%), and 1,723 (1.3%), treated with apixaban, dabigatran, edoxaban, and rivaroxaban respectively. In contrast, reports that described hemorrhagic stroke included 3,016 (2.2%), 1,346 (1.1%), and 1,256 (1.8%) for rivaroxaban, apixaban, and dabigatran. An additional 383 (6.9%) ischemic stroke cases were identified among enoxaparin/heparin treated patients.

The proportion on man significantly higher in both adverse events. The mean age for apixaban-treated patients was 77±10, significantly older than the other three DOACs. Enoxaparin/heparin patients were younger with a mean age of 72±13 and did not differ by sex from the DOACs. Hemorrhagic stroke resulted in poorer outcomes and more mortality compared to ischemic stroke (Table 1).

Concomitant levetiracetam reports and percentages of the total in each anticoagulant were as follows: 122 (0.1%) in apixaban, 142 (0.2%) in dabigatran,

20 (0.3%) in edoxaban, 168 (0.1%) in rivaroxaban, and 45 (0.8%) in enoxaparin/heparin. The number of reports with the DOAC-levetiracetam-adverse event triplet are listed in table 1.

## **Ischemic Stroke**

The DOACs did not differ from enoxaparin or heparin in the disproportionality analysis for ischemic stroke, (adj. ROR; 1.14, 95%Cl, 1.01-1.28). The adjusted ROR of the interaction effect was 3.57 (95%Cl, 2.81–4.58) between DOACs and levetiracetam, supporting the existence of a significant interaction.

When comparing the overall database to each DOAC separately, apixaban and rivaroxaban demonstrated the least disproportionality for ischemic stroke, (adj.ROR; 0.44, 95%CI, 0.41-0.47), and (adj.ROR; 0.35, 95%CI, 0.33-0.37) respectively. A significant disproportionality signal was identified for ischemic stroke in dabigatran (adj. ROR 3.69; 95%CI, 3.50-3.88) and in edoxaban (adj. ROR 1.85; 95%CI, 1.58-2.16) AF users (Table 2).

# DOAC-Levetiracetam Drug-Drug Interactions

Apixaban, dabigatran, edoxaban, and rivaroxaban illustrated a significant signal for the drug interaction in the regression model. The strength of association seems higher in edoxaban (adj.ROR 5.59; 95%CI, 4.23-7.39). Whereas the strength of association was slightly lower with rivaroxaban, (adj. ROR 3.69; 95%CI, 2.92-4.67).

When queried for ischemic stroke  $\Omega_{0.25}$  was greater than zero for apixaban, dabigatran, and rivaroxaban, demonstrating an interaction with levetiracetam (figure 1). The magnitude of  $\Omega_{0.25}$  was similar for the three DOACs. An interaction between levetiracetam and enoxaparin/heparin was not detected, with  $\Omega_{0.25}$ <0.

# Non-Ischemic Stroke- Hemorrhage

A significant disproportionality signal was identified for hemorrhagic stroke in rivaroxaban AF patients compared to apixaban and dabigatran (adj. ROR 1.46; 95%CI, 1.38-1.55) (Table 3). Dabigatran demonstrated a significantly lower adj. ROR for hemorrhagic stroke than the other DOACs, (adj. ROR 0.75; 95%CI, 0.70-0.78). Apixaban did not differ significantly from rivaroxaban and dabigatran.

The regression model queried for hemorrhagic stroke did not detect an interaction with levetiracetam. Likewise, the shrinkage analysis was negative for apixaban, and rivaroxaban, detecting no interaction. The concomitant use of levetiracetam and the two DOACs does not seem to increase the risk of intracerebral bleeding reports. The shrinkage analysis was tested positive for dabigatran (figure 1). Contrary to the shrinkage analysis, the regression model did not detect signals for dabigatran-levetiracetam-hemorrhagic stroke triplet (adj. ROR 1.07; 95%CI, 0.90-1.28).

# Sensitivity Analysis

In sensitivity analysis, we could not detect changes in disproportionality by publication date. The model detected signals in the positive control carbamazepine but not in the negative control, omeprazole (Table 4).

## DISCUSSION

The main finding of the current analysis is a significant signal for interaction between levetiracetam and DOACs, as demonstrated by the increased risk of ischemic stroke in the FAERS database. The interaction was confirmed using two methods accepted by regulators worldwide suggesting a robust and significant finding.

The mechanism of the DDI remains to be elucidated. However, it is hypothesized that the mechanism involves P-glycoprotein induction, an efflux protein that lowers the bioavailability of DOACs (8,19). Moreover, levetiracetam may bear a weak enzyme-inducing properties and may stimulate the activity of CYP3A4 and/or some UDP-glucuronosyltransferase (UGT) isoenzymes (19). Apixaban, dabigatran, edoxaban, and rivaroxaban are all substrates of the efflux P-gp transporter. Our analysis successfully demonstrated an interaction with all four DOACs, with 3-5 folds increased risk of stroke 3-5 and some variability between the DOACs. Rivaroxaban showed the weakest association, probably because of its high baseline bioavailability compared with the other DOACs.

Recently, there has been extensive concern regarding the interaction between levetriacetam and the DOACs. The 2021 European Heart Rhythm Association (EHRA) guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation recommends caution in the use of the antiepileptic drug, levetiracetam, due to potential P-glycoprotein-mediated drug–drug interaction (8,33). To date, data on the potential drug interaction between levetiracetam and anticoagulants from the DOAC family is scarce.

Paciullo et al. reported a case in 2020 of a 69-year-old man with AF who received rivaroxaban (20 mg/d) and developed a transient ischemic attack a few months after initiating levetiracetam for the indication of focal seizures prevention (20).

Specific anti-Xa activity for Rivaroxaban found a zero-trough level immediately before rivaroxaban administration. In 2020, Giustozzi et al. illustrated in a small prospective study (n=91) that patients with non-valvular AF treated with DOACs and anticonvulsants appear to have a relatively high rate of thromboembolic events (17). A similar conclusion was drawn in a nested case-control study supporting a diminished anticoagulant effect when combining DOACs with levetiracetam (21).

On the other hand, recent pharmacokinetic studies suggest there is no such potential drug interaction. In a case study published by Menichelli et al. (22) a 54year-old male with AF, cirrhosis, and seizures showed no significant reduction in dabigatran plasma concentration when used alongside levetiracetam. Another recent small PK study by Mavri et al. analyzed 21 patients concurrently receiving levetiracetam and DOACs, with 19 having atrial fibrillation and 2 having venous thromboembolism (23). Blood samples were collected to measure trough concentrations of DOACs and levetiracetam. The results showed that none of the patients experienced thromboembolic events during the observation period of 1,388 ± 994 days. Furthermore, there was no reduction in DOAC plasma levels during levetiracetam treatment, suggesting that levetiracetam may not significantly affect DOAC concentrations.

However, this study has two important limitations: first, because DOAC trough levels are highly variable, there is a need to analyze the area under the curve to rule out P-gp induction effect on plasma levels with and without levetiracetam. Second, the reported average trough levels of levetiracetam in that study  $(31.0 \pm 34.5 \text{ mg/L})$  although the average concentration is higher than expected, the standard deviation may not ensure that all DOAC patients were adequately exposed to levetiracetam. In the literature the lower bound confidence interval of levetiracetam concentrations is about 15 mcg/mL (34).

In another retrospective cohort study published by Ip et al., the risk of thromboembolism was evaluated in patients taking direct oral anticoagulants concurrently with antiseizure medications that modulate the cytochrome P450 or P-glycoprotein systems including levetiracetam. While CYP/P-gp-modulating antiseizure medications were associated with an increased risk of ischemic stroke in the overall analysis, no difference in thromboembolism risk or death was observed in the epilepsy subgroup using levetiracetam (24).

Earlier studies on levetiracetam showed no alterations in plasma concentrations of other P-glycoprotein substrates, such as digoxin (25, 26). Thus, it may also be reasonable to investigate pharmacodynamic interaction. Piracetam, a molecule similar in structure to levetiracetam, has been shown to have anticoagulant properties (27). However, our analysis did not show an increase in the reporting ratio of hemorrhagic stroke, defined as major bleeding, not supporting any levetiracetam anticoagulant properties or increased bleeding.

It is worthwhile to note that dabigatran demonstrated a positive shrinkage signal for hemorrhagic stroke with concomitant use of levetiracetam, however the logistic regression failed to support any drug-drug interaction. A retrospective cohort of patients from Taiwan on DOACs and 11 different anticonvulsants reported increased association of bleeding with concomitant prescription of phenytoin, valproic acid or levetiracetam, the results were explained by an increased risk of renal failure with levetiracetam (35). Further, our shrinkage analysis did not detect an interaction with heparin/enoxaparin that resulted in bleeding.

Although an interaction may exist between DOACs and levetiracetam this does not mean physicians should stop prescribing the combination for post stroke epilepsy. Levetiracetam is considered a newer and safer anticonvulsant (36). We recommend a drug monitoring strategy to be implemented as a solution for this drug interaction.

Whether the interaction is pharmacokinetic or pharmacodynamic, also has clinical implications for therapeutic drug monitoring. Naturally, a pharmacokinetic interaction is best monitored by plasma levels of DOACs. Whereas anti-Xa for apixaban and rivaroxaban and the plasma diluted thrombin time for dabigatran would be more suitable for a pharmacodynamic interaction (28, 29).

Regarding monitoring methods, Goldstein et al. examined the impact of cytochrome P450- and P-glycoprotein-inducing antiseizure medications on the pharmacokinetics of direct oral anticoagulants compared to rifampicin (a very strong inducer of DOACs elimination) (30). They suggested monitoring DOAC plasma concentrations as a helpful strategy to guide dosing and identify patients at risk for low DOAC concentrations and treatment failure when taking enzyme-inducing antiseizure medications. In the event our findings are confirmed in further studies and the mechanism for the levetiracetam interaction will be fully understood, pharmacodynamic monitoring should be preferred.

Dabigatran demonstrated a higher adjROR for ischemic stroke than apixaban, edoxaban and rivaroxaban, which could be explained by dabigatran's negative publicity a year after it accessed the market in 2010-2011 (31). The FAERS database shows a peak with the highest rates of reports between 2011-2014 (18). To account for this, we re-analyzed the data excluding the years 2010-2014, but the signal for ischemic events and the levetiracetam drug interaction did not significantly change. Our results also show that dabigatran has the lowest adj. ROR for hemorrhagic stroke, a reciprocal image of the high reporting on ischemic stroke. In the literature, dabigatran does not seem to differ in effectiveness from the other DOACs but is associated with lower risk of major bleeding (32).

# **Strengths and Limitations**

The major strengths of our study are utilizing a worldwide database, an analysis using two well validated methods for signal detection, performing an extensive sensitivity analysis to challenge our model and mitigate reporting bias, and finally, suggesting pharmacological mechanism that can explain the findings.

A limitation of our study is that some adverse events are likely not reported to national authorities for inclusion in the FAERS. This was mitigated by collecting data from all countries from 2012 to 2023. Although concomitant drug use was accounted for in our analysis, it is challenging to determine with certainty the sequence of medication use owing to missing data and temporality being associated with an adverse event rather than a drug. Another caveat is that reports lack information on dose.

Although lamotrigine is an alternative medication to post-stroke epilepsy, its use for that indication is seemingly low. There were no ischemic stroke reports in DOACs combined with lamotrigine, therefore we used omeprazole as our negative control, which is usually given to prevent gastrointestinal bleeding with DOACs. The analysis could not detect a drug interaction with DOACs and omeprazole, strengthening our results.

There were no reports of intracranial bleeding with the concomitant use of levetiracetam and edoxaban, limiting the direction analysis for edoxabanlevetiracetam interaction. Although, the regression model and shrinkage analysis detected an interaction with edoxaban, a discrepancy in their magnitudes were noted. This could be explained by a high proportion of ischemic stroke reports and a low number of non-cases reporting concomitant levetiracetam use, which emphasized this variable in the regression model. Finally, limitations to spontaneous reporting (e.g., underreporting) exist as well. However, despite its flaws, the analysis of pharmacovigilance databases remains a cornerstone for the study of adverse drug reactions and drug interactions by regulators and contributes majorly to drug labels. Data science in medicine is growing fast and is especially sensitive for detecting variables that may interfere with drug therapy.

# **Conclusions and Clinical Implications**

We show a strong signal for the levetiracetam interaction with apixaban, dabigatran, edoxaban, and rivaroxaban. The interaction is demonstrated by 3-5 folds increased reporting risk of ischemic stroke. Our findings suggest the need for pharmacodynamic monitoring, either anti-Xa for apixaban and rivaroxaban or plasma diluted thrombin time for dabigatran while concomitantly prescribing levetiracetam with the DOACs.

# Data Availability Statement

The data underlying this article are available in the FDA adverse event reporting

system (FAERS) at https://www.fda.gov/drugs/questions-and-answers-fdas-

adverse-event-reporting-system-faers/fda-adverse-event-reporting-system-faers-

public-dashboard.

#### **References:**

- Gerhard Hindricks, Tatjana Potpara, Nikolaos Dagres, Elena Arbelo, Jeroen J Bax, Carina Blomström-Lundqvist, Giuseppe Boriani, Manuel Castella, Gheorghe-Andrei Dan, Polychronis E Dilaveris, Laurent Fauchier, Gerasimos Filippatos, Jonathan M Kalman, Mark La Meir, Deirdre A Lane, Jean-Pierre Lebeau, Maddalena Lettino, Gregory Y H Lip, Fausto J Pinto, G Neil Thomas, Marco Valgimigli, Isabelle C Van Gelder, Bart P Van Putte, Caroline L Watkins, ESC Scientific Document Group, 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC, European Heart Journal, Volume 42, Issue 5, 1 February 2021, Pages 373–498, <u>https://doi.org/10.1093/eurheartj/ehaa612</u>
- 2. Chan N, Sobieraj-Teague M, Eikelboom JW. Direct oral anticoagulants: evidence and unresolved issues. Lancet. 2020 Nov 28;396(10264):1767-1776. doi: 10.1016/S0140-6736(20)32439-9.
- 3. Chang RS, Leung WC, Vassallo M, Sykes L, Battersby Wood E, Kwan J. Antiepileptic drugs for the primary and secondary prevention of seizures after stroke. Cochrane Database Syst Rev. 2022 Feb 7;2(2):CD005398. doi: 10.1002/14651858.
- 4. Camilo O, Goldstein LB. Seizures and epilepsy after ischemic stroke. Stroke. 2004 Jul;35(7):1769-75. doi: 10.1161/01.STR.0000130989.17100.96. Epub 2004 May 27.
- DeLorenzo RJ, Hauser WA, Towne AR, Boggs JG, Pellock JM, Penberthy L, Garnett L, Fortner CA, Ko D. A prospective, population-based epidemiologic study of status epilepticus in Richmond, Virginia. Neurology. 1996 Apr;46(4):1029-35.
- 6. Herman ST. Epilepsy after brain insult: targeting epileptogenesis. Neurology. 2002 Nov 12;59(9 Suppl 5):S21-6. doi: 10.1212/wnl.59.9\_suppl\_5.s21.
- Ruff CT, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD, Camm AJ, Weitz JI, Lewis BS, Parkhomenko A, Yamashita T, Antman EM. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. Lancet. 2014;383(9921):955–62. https://doi.org/10.1016/S0140-6736(13)62343-0.
- Steffel J, Verhamme P, Potpara TS, Albaladejo P, Antz M, Desteghe L, Haeusler KG, Oldgren J, Reinecke H, Roldan-Schilling V, Rowell N, Sinnaeve P, Collins R, Camm AJ, Heidbüchel H; ESC Scientific Document Group. The 2018 European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation. Eur Heart J. 2018;39(16):1330–1393.
- 9. FAERS database. https://www.fda.gov/drugs/questions-and-answers-fdas-adverse-event-reporting-system-faers/fda-adverse-event-reporting-system-faers-public-dashboard. Accessed Feb. 2023.
- Center for Drug Evaluation and Research. National Drug Code Directory. U.S. Food and Drug Administration. 2022. https://www.fda.gov/drugs/drug-approvals-and-databases/national-drugcode-directory. Accessed Feb. 2023.
- 11. van Puijenbroek EP, Bate A, Leufkens HG, Lindquist M, Orre R, Egberts AC. A comparison of measures of disproportionality for signal detection in spontaneous reporting systems for

adverse drug reactions. Pharmacoepidemiol Drug Saf. 2002 Jan-Feb;11(1):3-10. doi: 10.1002/pds.668.

- 12. Noguchi Y, Tachi T, Teramachi H. Review of Statistical Methodologies for Detecting Drug-Drug Interactions Using Spontaneous Reporting Systems. Front Pharmacol. 2019 Nov 8;10:1319. doi: 10.3389/fphar.2019.01319.
- Norén GN, Sundberg R, Bate A, Edwards IR. A statistical methodology for drug-drug interaction surveillance. Stat Med. 2008 Jul 20;27(16):3057-70. doi: 10.1002/sim.3247. Erratum in: Stat Med. 2008 Dec 20;27(29):6271-2.
- 14. Apixaban Physician Guidance. <u>https://www.medicines.org.uk/emc/product/2878/smpc#gref</u>. Accessed Feb. 2023.
- Apixaban Physician Guidance. FDA. https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=095a08ac-cf0e-497e-a682ddef38d6b29c#Section\_5.3. Accessed, Feb. 2023.
- Bolek T, Samoš M, Škorňová I, et al. Does proton pump inhibition change the on-treatment anti-Xa activity in xabans-treated patients with atrial fibrillation? A pilot study. J Thromb Thrombolysis 2019;47:140–5.doi:10.1007/s11239-018-1748-5.
- Giustozzi, M., Mazzetti, M., Paciaroni, M. et al. Concomitant Use of Direct Oral Anticoagulants and Antiepileptic Drugs: A Prospective Cohort Study in Patients with Atrial Fibrillation. Clin Drug Investig 41, 43–51 (2021). https://doi.org/10.1007/s40261-020-00982-8.
- 18. Kaba RA, Ara F, Ward DE, Emanuel S. The Dabigatran debate. Glob Cardiol Sci Pract. 2014 Oct 16;2014(3):254-6. doi: 10.5339/gcsp.2014.41.
- Galgani A, Palleria C, Iannone LF, De Sarro G, Giorgi FS, Maschio M, Russo E. Pharmacokinetic Interactions of Clinical Interest Between Direct Oral Anticoagulants and Antiepileptic Drugs. Front Neurol. 2018 Dec 7;9:1067. doi: 10.3389/fneur.2018.01067. Erratum in: Front Neurol. 2020 Jan 29;10:1381.
- 20. Paciullo F, Costa C, Gresele P. Rivaroxaban Plasma Levels and Levetiracetam. Ann Intern Med. 2020 Nov 3;173(9):772. doi: 10.7326/L20-1066.
- Gronich N, Stein N, Muszkat M. Association Between Use of Pharmacokinetic-Interacting Drugs and Effectiveness and Safety of Direct Acting Oral Anticoagulants: Nested Case-Control Study. Clin Pharmacol Ther. 2021 Dec;110(6):1526-1536. doi: 10.1002/cpt.2369.
- Menichelli D, Pastori D, Pignatelli P, Pani A. Minimizing drug-drug interactions between dabigatran and levetiracetam through clinical management: a case report. Eur Heart J Case Rep. 2023 Jan 6;7(1):ytad006. doi: 10.1093/ehjcr/ytad006.
- 23. Mavri A, Ilc S. The efficacy of direct oral anticoagulants in patients on concomitant treatment with levetiracetam. Sci Rep. 2023 Jun 7;13(1):9257. doi: 10.1038/s41598-023-33876-8.
- Ip BY, Ko H, Wong GL, Yip TC, Lau LH, Lau AY, Leng X, Leung H, Chan HH, Chan HY, Mok VC, Soo YO, Leung TW. Thromboembolic Risks with Concurrent Direct Oral Anticoagulants and Antiseizure Medications: A Population-Based Analysis. CNS Drugs. 2022 Dec;36(12):1313-1324. doi: 10.1007/s40263-022-00971-9.
- Mathy FX, Dohin E, Bonfitto F, Pelgrims B. Drug-drug interaction between levetiracetam and non-vitamin K antagonist anticoagulants. Eur Heart J. 2019 May 14;40(19):1571. doi: 10.1093/eurheartj/ehy780.

- 26. Levy RH, Ragueneau-Majlessi I, Baltes E. Repeated administration of the novel antiepileptic agent levetiracetam does not alter digoxin pharmacokinetics and pharmacodynamics in healthy volunteers. Epilepsy Res. 2001 Aug;46(2):93-9. doi: 10.1016/s0920-1211(01)00253-4.
- 27. Teng CM, Yeh HI, Lee LG. Anticoagulant and antiplatelet properties of piracetam. Taiwan Yi Xue Hui Za Zhi. 1983 Nov;82(11):1104-14.
- Sarode R. Direct oral anticoagulant monitoring: what laboratory tests are available to guide us? Hematology Am Soc Hematol Educ Program. 2019 Dec 6;2019(1):194-197. doi: 10.1182/hematology.2019000027.
- 29. Kaba RA, Ara F, Ward DE, Emanuel S. The Dabigatran debate. Glob Cardiol Sci Pract. 2014 Oct 16;2014(3):254-6. doi: 10.5339/gcsp.2014.41.
- Goldstein R, Jacobs AR, Zighan L, Gronich N, Bialer M, Muszkat M. Interactions Between Direct Oral Anticoagulants (DOACs) and Antiseizure Medications: Potential Implications on DOAC Treatment. CNS Drugs. 2023 Mar;37(3):203-214. doi: 10.1007/s40263-023-00990-0.
- 31. Pradaxa lawsuit. https://www.drugwatch.com/pradaxa/lawsuits/. Accessed Feb. 2023.
- Lau WCY, Torre CO, Man KKC, Stewart HM, Seager S, Van Zandt M, Reich C, Li J, Brewster J, Lip GYH, Hingorani AD, Wei L, Wong ICK. Comparative Effectiveness and Safety Between Apixaban, Dabigatran, Edoxaban, and Rivaroxaban Among Patients With Atrial Fibrillation : A Multinational Population-Based Cohort Study. Ann Intern Med. 2022 Nov;175(11):1515-1524. doi: 10.7326/M22-0511.
- Jan Steffel and others, 2021 European Heart Rhythm Association Practical Guide on the Use of Non-Vitamin K Antagonist Oral Anticoagulants in Patients with Atrial Fibrillation, *EP Europace*, Volume 23, Issue 10, October 2021, Pages 1612– 1676, https://doi.org/10.1093/europace/euab065.
- 34. Mendoza Aguilera M, Bellés Medall MD, Álvarez Martín T, Pascual Marmaneu Ó, Liñana Granell C, Ferrando Piqueres R. Therapeutic drug monitoring of levetiracetam in daily clinical practice: high-performance liquid chromatography versus immunoassay. Eur J Hosp Pharm. 2020 Mar;27(e1):e2-e6. doi: 10.1136/ejhpharm-2018-001616.
- 35. Wang CL, Wu VC, Chang KH, Tu HT, Kuo CF, Huang YT, Chu PH, Kuo CC, Chang SH. Assessing major bleeding risk in atrial fibrillation patients concurrently taking non-vitamin K antagonist oral anticoagulants and antiepileptic drugs. Eur Heart J Cardiovasc Pharmacother. 2020 Jul 1;6(3):147-154. doi: 10.1093/ehjcvp/pvz035.
- French JA, Gazzola DM. New generation antiepileptic drugs: what do they offer in terms of improved tolerability and safety? Ther Adv Drug Saf. 2011 Aug;2(4):141-58. doi: 10.1177/2042098611411127.

#### Figures:

	Rivaroxaban	Apixaban	Dabigatran	Edoxaban	P-
(No. of total AEs)	(136,710)	(125,799)	(69,993)	(6,965)	Value
Total AERs					
Ischemic stroke	1723 (1.3)	1841 (1.5)	3731 (5.3)	338 (4.9)	-
Hemorrhagic stroke	3016 (2.2)	1346 (1.1)	1256 (1.79)	-	
Age	73.6±13.2	76.9±9.9	73.7±10.6	74.5±11.5	<0.001,
					0.94
Sex					
Female	794 (46.1)	772 (46.5)	1478 (40.1)	46 (13.6)	
	1287 (42.7)	553 (41.1)	518 (41.2)	-	<0.001
Male	823 (47.8)	791 (47.7)	1881 (51.1)	51 (15.0)	
	1641 (54.5)	737 (54.8)	683 (54.4)	-	
Not Specified	106 (6.2)	97 (5.8)	325 (8.8)	241 (71.3)	
	88 (2.9)	28 (4.2)	55 (4.4)	-	
No. concomitant Drug-					
AE					-
Levetiracetam	34 (2.0)	32 (1.7)	33 (0.9)	4 (1.2)	
Ischemic	14 (0.5)	10 (0.7)	13 (1.0)	N/R	
Levetiracetam					
Hemorrhagic					
Mortality outcome <sup>b</sup>					
Ischemic stroke					
DOAC alone	305 (11.8)	111 (6.0)	451 (7.7)	23 (6.8)	
DOAC-levetiracetam	1 (0.04)	4 (0.2)	12 (0.2)	0 (0)	<0.001
Hemorrhagic stroke					
DOACs alone	1552 (56.9)	488 (36.3)	588 (46.8)	-	
DOAC-levetiracetam	20 (0.7)	4 (0.3)	7 (0.6)	N/R	
Publication date					
2012-2013	141 (8.2)	16 (0.87)	971 (26.0)	446 (6.4)	
	0 (0)	7 (0.5)	305 (24.3)		
2014-2015	355 (20.6)	201 (10.9)	749 (20.1)	1526 (21.9)	<0.001
	339 (11.2)	8 (0.6)	276 (21.0)		
2016-2017	571 (33.1)	408 (22.2)	582 (15.6)	1832 (26.3)	
	1072 (35.5)	313 (23.3)	215 (17.1)		
2018-2019	383 (22.2)	493 (26.8)	725 (19.4)	1651 (23.7)	
	689 (22.8)	584 (43.4)	249 (19.8)		
2020-2021	244 (14.2)	386 (21.0)	458 (12.2)	1468 (21.1)	

	854 (28.3)	313 (23.2)	115 (9.2)		
2022-2023	29 (1.68)	157 (8.5)	99 (2.7)	72 (1.0)	
	62 (2.1)	120 (8.9)	11 (0.9)		

Table 1: Characteristics and Demographics of Levetiracetam and Direct Anticoagulant Users with Associated Ischemic and non-Ischemic Stroke Adverse Events in Atrial Fibrillation Patients.

AEs-adverse events. Top figures for ischemic stroke, bottom figures for hemorrhagic stroke. Numbers in brackets are percentages of ischemic stroke events.

Table 2: Signals of Ischemic Stroke and Anticoagulants- Levetiracetam Drug Interaction in Atrial Fibrillation Patients.

	ROR 95%CI	Adj. ROR 95%Cl	ROR 95%CI for DDI*		
DOACs vs. Enoxaparin					
All DOACs	1.05 (0.94-1.18)	1.14 (1.01-1.28)	3.57 (2.81-4.58)		
DOACs Head-to-Head					
Apixaban	0.44 (0.41-0.46)	0.44 (0.41-0.47)	4.61 (3.62-5.89)		
Dabigatran	3.54 (3.38-3.70)	3.69 (3.50-3.88)	4.34 (3.41-5.52)		
Edoxaban	1.77 (1.58-1.98)	1.85 (1.58-2.16)	5.59 (4.23-7.39)		
Rivaroxaban	0.36 (0.35-0.39)	0.35 (0.33-0.37)	3.69 (2.92-4.67)		

The first row illustrates the four DOACs compared with enoxaparin. The rows below compared each of the DOACs to the other three head-to-head. \*DDI= drug-drug interaction.

	Adj. ROR 95%Cl	ROR 95%CI for DDI
Rivaroxaban	1.46 (1.38-1.55)	0.98 (0.82-1.18)
Dabigatran	0.75 (0.70-0.78)	1.07 (0.90-1.28)
Apixaban	0.95 (0.88-1.01)	1.10 (0.93-1.32)

Table 3: Secondary Outcome Signals of Hemorrhagic Stroke and the DOACs- Levetiracetam Drug Interaction in Atrial Fibrillation Patients.

#### Figure 1: Shrinkage Analysis for the Anticoagulants-Levetiracetam Interaction



 $\Omega_{025}$  > 0, a positive lower bound of 95% Cl, is used as a threshold for detecting the signals of the concomitant use of the DOACs and levetiracetam. A positive shrinkage was noticed for ischemic in the concomitant use of either apixaban, dabigatran, and rivaroxaban with levetiracetam. A negative shrinkage was noticed for hemorrhagic stroke and the concomitant use of apixaban or rivaroxaban with levetiracetam.

ROR 95%CI for DDI Shrinkage Analysis Rationale (Ω<sub>0.25</sub>) Apixaban-Carbamazepine 8.47 (5.37-13.36) +1.97 Positive control: the concomitant use of apixaban with strong CYP3A4 and P-glycoprotein inducers such as carbamazepine, may lead to a ~50% reduction in apixaban exposure. In a clinical study of atrial fibrillation patients, diminished efficacy was noted (14). The lower bound CI >1 and  $\Omega_{0.25}$ >0 for ischemic stroke detecting a signal. Apixaban-Omeprazole 1.13 (0.98-1.33) -0.45 Negative control: no interaction is documented to date; hemorrhagic stroke ROR is nonsignificant. The lower bound CI <1 and  $\Omega_{0.25}$ <0. Publication date First, in 2020, Guistozzi et al. illustrated that patients Rivaroxaban-levetiracetam with non-valvular AF 2012-2020 3.51 (2.27-5.43) +2.30 treated with rivaroxaban and levetiracetam appear to have a relatively high Dabigatran-levetiracetam rate of thromboembolic 2015-2023 3.14 (2.28-3.42) +1.31 events (17). Thus, the data was re-analyzed for 2012-2020. Second, a lawsuit occurred over Pradaxa, dabigatran's trade name, from 2010 to 2014 (18, 31). An additional analysis excluding that period did not significantly alter the results.

Table 4: Sensitivity Analysis for DOAC DDI for Ischemic or non-Ischemic Stroke in Atrial Fibrillation Patients.