

1 **Determinants of Healthcare Insurance Coverage among**
2 **Women of Reproductive Age in Tanzania: A Multilevel Mixed**
3 **Effect Analysis.**

4 Sanun Ally Kessy ^{1†}

5 Jovin R. Tibenderana ^{1†}

6 Jomo Gimonge²

7 Fabiola V Moshi²

8 ¹ Department of Epidemiology and Biostatistics, Kilimanjaro Christian Medical University
9 College, P. O. Box 2240 Moshi, Tanzania

10 ² Department of Clinical Nursing, The University of Dodoma, Dodoma, Tanzania

11 *Corresponding author: Sanun A Kessy, E-mail: sahnunally@outlook.com

12 † Authors contributed equally

13 **Abstract**

14 **Background:** Health insurance has been one of the fundamental approaches of financing
15 healthcare for sustainable Development goals (SDGs). There is a paucity of evidence on the
16 determinants of healthcare insurance coverage among women of reproductive age in Tanzania,
17 therefore this study aimed at assessing factors associated with access to health care insurance
18 among women of reproductive age in Tanzania using national Demographic and health survey
19 dataset.

20 **Methodology:** This study was an analytical cross-sectional study that utilized nationally
21 representative secondary data sourced from the Tanzania Demographic and Health Surveys

22 (TDHS) of 2022. A weighted sample of 15,254 women of reproductive age (15-49) was included
23 in the study. A multilevel regression analysis was used to examine factors associated with
24 health care insurance coverage. These results were presented using adjusted odds ratio (AOR)
25 with a 95% confidence interval.

26 **Results:** In Tanzania the proportion of health insurance coverage among women of
27 reproductive age was 5.8% whereby majority of women subscribing to employer's insurance
28 (3.9%). Older women (35-49) were more likely to be covered with health insurance compared
29 those aged 15-24 (AOR =2.75, 95% CI: 2.19, 3.45). Similarly, married women had higher
30 likelihood of health insurance coverage compared to unmarried (AOR=1.60, 95% CI: 1.33,
31 1.92). Furthermore, educated women were more likely to be covered with health insurance than
32 uneducated women (AOR=6.96, 95% CI: 4.74, 10.22). Similarly, women who were exposed to
33 mass media were more likely to be covered with health care insurance compared to unexposed
34 (AOR=1.43, 95% CI: 1.14, 1.78).

35 **Conclusion:** Less than ten percent of Tanzanian women of reproductive age (WRA) are
36 registered in health insurance, which is an unacceptably low coverage. This highlights the
37 urgent need to eliminate barriers preventing this population's access to healthcare in order to
38 achieve Standard Development Goals (SDGs) by 2030. The interplay of socioeconomic factors
39 and access to information underscores the need for multifaceted approaches to enhance
40 healthcare coverage among women of reproductive age in Tanzania.

41 **Keywords:** Health insurance, Women of reproductive age, determinants, universal health
42 coverage, Tanzania.

43 **Background**

44 Globally, health insurance has been one of the fundamental approaches of financing healthcare
45 for sustainable Development goals (SDGs) by 2030 (1). About 2 billion individuals are struggling

46 with financial difficulties, with 1 billion encountering severe out of pocket (1). Additionally, 344
47 million people globally are being pushed further into the burden of the health care costs (1,2).
48 Globally the significance of Universal Health Coverage (UHC) has been emphasized by its
49 incorporation into the Sustainable Development Goals (SDGs) stating that, all individuals should
50 have equal access to health care as right rather than a privilege with its target to be achieved by
51 2030 (3,4).

52 Universal Health Coverage (UHC) means that all individuals and communities receive health
53 services they need without suffering financial hardship. It includes the full spectrum of essential,
54 quality health services, from health promotion to prevention, treatment, rehabilitation, and
55 palliative care (2). The global trajectory is deviating from the path to achieve substantial
56 advancements in universal health coverage, a goal outlined in the Sustainable Development
57 Goals (SDGs) target 3.8 set for 2030 which aim at protecting the vulnerable population from
58 financial risks associated with severe health-related expenses (5).

59 UHC is important as it ensures provision of essential health services based on need, equitable
60 access for the entire population, and the presence of financial-risk protection mechanisms to
61 prevent economic hardship from healthcare expenses are key components of universal health
62 coverage (14). Women face considerable challenges in accessing medical services, primarily
63 stemming from limited opportunities for economic empowerment and addressing their distinct
64 healthcare requirements (5).

65 In Sub Saharan Countries, the prevalence of universal health coverage of health insurance
66 coverage to be less than 10% (23,25). The outcomes of Health Insurance (HI) implementation
67 have been varying. Findings from research in Ghana (17) and Rwanda (18) indicate that HI has
68 contributed to greater equity in access to care for the overall population, alleviating the financial
69 burden of healthcare expenses among the impoverished. Conversely, study findings from
70 Ghana suggests Health Insurance (HI) did not provide uniform protection to all its members.

71 Instead, it favored individuals with higher education, greater wealth, and those living in proximity
72 to health facilities more than others (19).

73 Furthermore, among Sub-Saharan countries there is a notable correlation between formal
74 sector employment and access to health insurance compared to informal employment sector
75 (20). Additionally, other factors such as such as income, education, household wealth status,
76 marital status, age, and place of residence, as predictors of health insurance ownership (21,23).

77 Despite notable accomplishments, the healthcare sector grapples with substantial hurdles in
78 striving to deliver high-quality care to all individuals, regardless of their financial capacity to
79 afford services. Women encounter more pronounced obstacles in accessing critical healthcare
80 services, including routine antenatal visits and childbirth services (5,26, 27).

81 In Tanzania, over 80% of the population continues to rely on out-of-pocket payments for
82 accessing healthcare services, often resulting in significant financial burdens (6). Tanzania
83 implemented health insurance initiatives in the mid-1990s, introducing the National Health
84 Insurance Fund (NHIF) in 1996 and the Community Health Fund (CHF) in 2001. The CHF,
85 designed for the informal and rural communities, operates on a voluntary basis, offering
86 coverage for up to six family members through a fixed annual payment, other health insurances
87 such as Social Health Insurance Benefit (SHIB) established as a benefit under the National
88 Social Security Fund (NSSF) and the Community Health Fund (CHF) and Tiba Kwa Kadi (TIKA)
89 (6,7).

90 The capacity of women in the reproductive age group to attain affordable health coverage and
91 care carries significant societal implications and hinges on the economic well-being of both
92 women and their families. Nevertheless, the accessibility of healthcare services for women in
93 the reproductive age group remains limited in Low- and Middle-Income Countries (LMIC),
94 primarily due to financial.

95 In Tanzania, universal health coverage is in line with government policy of improving maternal
96 health among reproductive age in order to improve maternal morbidities and mortality, more
97 than 90% of women of reproductive do not have any health insurance (10). Despite several
98 studies that have been done in Tanzania which were small scale, thus there is still a paucity of
99 national wide information regarding health insurance utilizations. Additionally, there is a notable
100 absence of published literature addressing health insurance coverage in the country. Hence to
101 address this gap, the present study aimed to evaluate the health insurance coverage and its
102 associated factors among women in the reproductive age group in Tanzania utilizing data from
103 the Demographic and Health Survey (DHS). The findings of this study offer valuable insights for
104 health planners, program developers, and healthcare professionals, providing a foundation for
105 enhancing financial protection and service utilization among women of reproductive age.

106 **Methodology**

107 **Study setting and period.**

108 The study made use of the 2022 Tanzania Demographic and Health Survey (TDHS) data, a
109 countrywide cross-sectional survey carried out every five years(1). Tanzania, the largest country in
110 East Africa, spans an area of 940,000 square kilometers, including 60,000 square kilometers of
111 inland water. The study incorporated data from all regions of the United Republic of Tanzania As of
112 2022, the estimated population of Tanzania stood at 61,741,120, with an annual population growth
113 rate of 3.2% (2,3).

114 **Study design and data source.**

115 This study was an analytical cross-sectional study that utilized nationally representative secondary
116 data sourced from the Tanzania Demographic and Health Surveys (TDHS) of 2022.The TDHS, is
117 funded by the U.S Agency for International Development and implementation is carried out by the

118 Ministry of Health (MoH) in Tanzania Mainland and Zanzibar, as well as the National Bureau of
119 Statistics (NBS), the Office of the Chief Government Statistician (OCGS), with technical support
120 provided by ICF International (2–5).

121 Data collection procedure.

122 The data collection method of the surveys involves using a standardized questionnaire that is
123 uniform across countries to gather information from women aged 15–49. The questionnaire is often
124 translated into the primary local languages of the participating nations. According to the DHS, these
125 translated questionnaires, along with the original English version, undergo pretesting in both
126 English and the local dialect to ensure their accuracy. Subsequently, during the pretest phase, field
127 workers engage in an interactive discussion of the questions, providing suggestions for
128 improvements across all versions. After the field practice, a debriefing session is conducted with the
129 pretest field personnel, and adjustments are made to the questionnaires based on the insights
130 gained from this process(1).

131 Sampling procedure and size

132 The survey employed face-to-face questionnaire interviews and utilized a stratified design with
133 multistage cluster sampling to gather information on various aspects such as population health
134 status, neonatal mortality, health behaviors, nutritional status, family planning, and demographics.

135 In the initial stage, 629 clusters were identified, from which households were then selected. Among
136 these clusters, 26 households were systematically chosen to be representative from each cluster,
137 resulting in a total of 16,354 households included in the survey. Eligibility for inclusion was based on
138 the presence of all women aged 15–49 years in the selected household on the night prior to the
139 interview. Further details on the sampling procedure and design have been previously
140 documented(1). A total of 15,254 women who had information on all the variables of interest were

141 included in the study (Table 1). We adhered to the guidelines provided by the Strengthening the
142 Reporting of Observational Studies in Epidemiology (STROBE) statement when writing the
143 manuscript(6).The DHS data is freely available for download at
144 <https://dhsprogram.com/data/available-datasets.cfm>

145 Study variables.

146 The outcome variable of this study was health insurance coverage. This was from the question “are
147 you covered by any health insurance?”. Response is coded as 0 = “No” and 1 = “Yes”. The
148 explanatory variables were age of the woman, marital status, education, employment, residence,
149 wealth index, head of household sex, head of household age, media exposure, visited by
150 healthcare worker for past12 months and visited health facility for the past 12 months. Age was
151 recorded as 15-24, 25-34 and 35-49. Marital status was categorized as married and unmarried.
152 Wealth status was categorized as poor, middle, and rich. Education was classified into three
153 categories: no education, primary education, secondary education, or higher education. A new
154 variable of media exposure was generated from household has either tv or radio. Our generated
155 study variables and coding were based on previous literature (7–10).

156 Statistical analysis

157 We used Stata software, version 18.0 (Stata Corporation, College Station, TX, USA) for all
158 statistical analyses. Initial presentation of health insurance coverage was done using percentages
159 to show the proportion of women covered. Subsequently, through cross-tabulation, we explored the
160 distribution of health insurance coverage among various background characteristics. Before
161 conducting the regression analysis, we assessed potential collinearity among the variables using
162 the variance inflation factor (VIF). Results indicated minimal collinearity, with VIF values ranging
163 from 1.44 to 1.45 and a mean of 1.61, suggesting no significant multicollinearity concerns. To
164 investigate the factors linked with health insurance coverage, we employed multilevel binary logistic

165 analysis. This approach was chosen due to the complex survey design of the DHS, which involved
166 a two-stage cluster sampling. Four models were utilized for analysis: Model 0: An empty model
167 without any explanatory variables, Model I: Including individual-level factors, Model II: Incorporating
168 community level factors and Model III: Considering all explanatory variables together. Crude and
169 adjusted odds ratios (AOR) with corresponding 95% confidence intervals (CIs) were fit for each
170 model. Both fixed and random effects were included in all models. Random effects represented the
171 variation in health insurance coverage based on primary sampling unit (PSU), measured by the
172 Intra-Cluster Correlation Coefficient (ICC). Fixed effects illustrated the relationship between
173 explanatory variables and the outcome. Model fitness was evaluated using the Akaike Information
174 Criterion (AIC), allowing us to determine how well each model fit the data. A significance level of $p <$
175 0.05 was used for all statistical tests. Furthermore, we weighed all analyses to account for
176 disproportionate sampling and non-response.

177 Ethical clearance

178 Permission to download and utilize the data was obtained from the DHS Program/ICF International
179 via <http://www.dhsprogram.com>. The data was strictly used for the current study's objectives.
180 Extensive details regarding the methodology and ethical considerations of the DHS program can be
181 found in the published reports of the Tanzania Demographic and Health survey(1,5).

182 RESULTS

183 Participants background characteristics.

184 Among 15,254 participants of this study, the mean (SD) age was 29.3(9.8). The overall health
185 insurance coverage among women of reproductive age was about 6%. Health insurance coverage
186 was prevalent (39.6%) among women aged 35+ years. Around half (51.0%) women who owned
187 health insurance were married. The majority of women (68.8%) of study participants who were

188 covered with health insurance had secondary or higher education. A significant percent (68.8%) of
189 participants who were insured were working. Slightly more than half (53.7%) of the study
190 participants who owned health insurance were from rural areas (Table 1).

191 Most (80.8%) of women of reproductive age who owned health insurance were from rich
192 household. The majority (67.1%) of women of reproductive age who were insured were living in
193 household whose head was male. Almost half (47.7%) of the women who were insured were living
194 in household whose head was aged 35-49 years of age. The majority (78.5%) of women who were
195 insured had been exposed to media. Almost all (95.1%) of the insured women had been visited by
196 field worker in the last 12 months. The majority (62.9%) of the study participants who are insured
197 visited a health facility for the past 12 months (Table 1).

198 **Table 1:** Background characteristics and coverage of health insurance (Weighted)

Variables	Covered with Health Insurance		p-value
	Total (Percentage)	Yes (Percentage)	
N	15,254 (100.0%)	888 (5.8%)	
Age (years)			
15-24	5,810 (38.1%)	263 (29.6%)	<0.001
25-34	4,609 (30.2%)	273 (30.8%)	
35-49	4,835 (31.7%)	352 (39.6%)	
Marital status			
unmarried	8,624 (56.5%)	435 (49.0%)	<0.001
married	6,630 (43.5%)	452 (51.0%)	
Education			
No education	2,450 (16.1%)	39 (4.3%)	<0.001
Primary	8,123 (53.3%)	239 (26.9%)	
Sec/higher	4,681 (30.7%)	610 (68.8%)	
Occupation			
Not working	5,452 (35.7%)	277 (31.2%)	0.032
Working	9,802 (64.3%)	610 (68.8%)	
Residence			
Urban	5,446 (35.7%)	476 (53.7%)	<0.001
Rural	9,808 (64.3%)	411 (46.3%)	
Wealth index			
Poor	5,044 (33.1%)	102 (11.4%)	<0.001
Middle	2,880 (18.9%)	69 (7.7%)	
Rich	7,330 (48.1%)	717 (80.8%)	
Sex of household head			
Male	10,918 (71.6%)	595 (67.1%)	0.017
Female	4,336 (28.4%)	292 (32.9%)	
Household head age			
15-24	539 (3.5%)	20 (2.2%)	0.089

25-34	3,245 (21.3%)	172 (19.4%)	
35-49	6,516 (42.7%)	423 (47.7%)	
50+	4,954 (32.5%)	273 (30.7%)	
Media Exposure			
No	6,653 (43.6%)	191 (21.5%)	<0.001
Yes	8,601 (56.4%)	696 (78.5%)	
Visited by fieldworker in last 12 months			
No	14,781 (96.9%)	844 (95.1%)	0.017
Yes	473 (3.1%)	43 (4.9%)	
Visited health facility last 12 months			
No	7,167 (47.0%)	329 (37.1%)	<0.001
Yes	8,087 (53.0%)	558 (62.9%)	

199

200

201

202

203

204

205 Proportional of health insurance coverage.

206 In Tanzania the proportion of health insurance coverage among women of reproductive age is
207 5.8%. The rest of the group are not covered with any health insurance (figure 1)

208 Specific types of health insurance coverage.

209 Figure 2 shows percentage of women with specific types of health insurance. Most of them (5.8%)
210 have any insurance followed by employers' insurance (3.9%) and the lowest (0.2%) being social
211 security insurance.

212 Determinants of health insurance coverage among Tanzanian 213 reproductive age women.

214 Fixed effect

215 Table 2, Model III presents determinants of health insurance coverage among Tanzanian
216 reproductive age women. Older women (35-49) were more likely to be covered with health
217 insurance compared to those aged 15-24 (AOR =2.75, 95% CI: 2.19, 3.45). Similarly, married
218 women had higher likelihood of health insurance coverage compared to unmarried (AOR=1.60,
219 95% CI: 1.33, 1.92). The odds of health insurance coverage were higher among educated women
220 (secondary/higher) compared to uneducated women (AOR=6.96, 95% CI: 4.74, 10.22). Women
221 who were exposed to media had higher odds of health insurance coverage compared to unexposed
222 (AOR=1.43, 95% CI: 1.14, 1.78). Women of reproductive age who visited health facility for the past
223 12 months had higher likelihood of health insurance coverage compared to those who didn't
224 (AOR=1.28, 95% CI: 1.09, 1.51). Additionally, participants from rich households were more likely to
225 be covered with health insurance compared to their counterparts (AOR=2.78, 95% CI: 1.99, 3.87).
226 We found that women who are living in households lead by female.

227 Random effect

228 Table 2 presents results for random effect. We observed that the Intra-Cluster Correlation (ICC)
229 value in Model III was 0.27 indicating that approximately 27% of the variability in health insurance
230 coverage is attributed to differences between the clusters. The cluster-level variation, as indicated
231 by the ICC values were 29%, 26%, 24% and 27% for Model 0, Model I, Model II and Model III
232 respectively. We opted to focus on interpreting the results of Model III for discussion, as it had the
233 lowest AIC value (5603.60) among the models, indicating a goodness of fit.

234 **Table 2:** Multilevel analysis of determinants of health insurance coverage among reproductive age women in Tanzania (N= 15,254)

Variable	Model 0	Model I	Model II	Model III
		AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Age (years)				
15-24		1		1
25-34		1.41*(1.14, 1.74)		1.30* (1.05,1.61)
35-49		3.12*** (2.49, 3.90)		2.75*** (2.19,3.45)
Marital status				
unmarried		1		1
married		1.41*** (1.19, 1.68)		1.60*** (1.33,1.92)
Education				
No education		1		1
Primary		1.91**(1.32, 2.74)		1.57**(1.08,2.27)
Sec/higher		9.91*** (6.8, 14.39)		6.96*** (4.74,10.22)
Occupation				
Not working		1		1
Working		1.02(0.86, 1.21)		0.98(0.82,1.17)
Media Exposure				
No		1		1
Yes		2.02*** (1.67, 2.46)		1.43**(1.14,1.78)
Visited by fieldworker in last months				
No		1		1
Yes		1.29(0.89, 1.86)		1.27(0.88,1.84)
Visited health facility last months				
No		1		1
Yes		1.28*(1.09, 1.51)		1.28**(1.09,1.51)
Residence				
Urban			1	1
Rural			0.97(0.75, 1.25)	1.00(0.76,1.33)
Wealth index				
Poor			1	1
Middle			1.20(0.86, 1.69)	0.83(0.58,1.17)

Rich			6.19*** (4.67, 8.21)	2.78*** (1.99,3.87)
Sex of household head				
Male			1	1
Female			1.24*** (1.06, 1.45)	1.42*** (1.19,1.70)
Random effect				
ICC (%)	29%	26%	24%	27%
Wald Chi-square		524.55	260.83	573.75
Mean VIF		1.45	1.44	1.61
Model fitness				
Log-likelihood	-3144.76	-2837.71	-2991.92	-2786.80
AIC	6293.52	5697.43	5995.83	5603.60
N	15,254	15,254	15,254	15,254
Number of clusters	628	628	628	628

235 AOR = adjusted odds ratios; CI = Confidence Interval; *p < 0.05, **p < 0.01, ***p < 0.001; 1 = Reference category; ICC = Intra-Class

236 Correlation coefficient ; AIC = Akaike Information Criterion; N= Total sample size

237 **Discussion**

238 This study aimed to analyze the determinants affecting the possession of health insurance among
239 Tanzanian women as a key approach to achieving universal health coverage. It found that
240 insignificant majority of women of about 5.8 % own any form of health insurance. The research
241 identified several factors associated with health insurance coverage, including age, economic
242 status, educational attainment, marital status, and mass media exposure, Visited health facility last
243 12 months.

244 Our study revealed that only 5.8% of Tanzanian women possess health insurance, figure is
245 contrary with Ghana having more than half percent accessing health healthcare (34). These
246 findings are in line with less than one tenth of the women of the reproductive age in Mauritania,
247 Ethiopia and Nigeria. (25,33,35). This disparity can be explained that in Ghana there is a
248 difference in health policy agendas, like livelihood empowerment against Poverty Program
249 (LEAP) in Ghana (34) A pro-poor intervention policy, use of Participatory welfare ranking (PWR)
250 and geographic targeting (GT) for poor groups helped in improving efficiency and equity needs
251 under NHIS program in Ghana. It has been mandatory and compulsory for all Ghanaians to join
252 health insurance scheme in Ghana (34).

253 This study found that compared to being unmarried, being married was associated with having
254 health insurance coverage. This suggests that having a partner or spouse is beneficial,
255 potentially because of the financial support from being in a dual-income household, which offers
256 more avenues for acquiring health insurance. Similar findings have been reported in other
257 studies conducted in East Africa, Zambia and Kenya (13,23,36) but inconsistency with the study
258 done in Malawi (4). Additionally, unmarried women may struggle to afford insurance premiums
259 and deductibles and may perceive their risk of illness as lower (23).

260 The study suggests that women 35 years of age and older are more likely to subscribe to
261 healthcare insurance, which is in line with findings from several other African nations, such as
262 South Africa, and Nigeria and the one done in sub-Saharan Africa (5,20,37). There are several
263 reasons for this trend. Firstly, people in this age group typically find employment, which makes it
264 possible for them to pay for health insurance. Furthermore, older women may have more
265 children who are more likely to become sick and frequently take on caregiving responsibilities,
266 which may lead them to look into insurance to reduce their out-of-pocket costs.

267 Women's education level had a substantial impact on whether or not they had health insurance.
268 Individuals with more educational attainment were shown to be more likely than those without
269 formal education to have health insurance. This finding adheres to other studies that show
270 education is a significant factor in predicting one's ability to obtain health insurance (34,38–40).
271 Education may help people obtain information more easily and increase their chances of
272 entering high-paying professions, which are more likely to offer insurance coverage, which
273 explains the positive association.

274 In comparison to women residing in the poor household, the current study showed that women
275 residing in the rich household had greater odds of having health insurance coverage. The
276 findings of the national survey carried out in Ethiopia were consistent with these results (41),
277 indicating a relationship between income levels and the probability of acquiring health
278 insurance. In contrast, women from wealthier homes had lower probabilities of having health
279 insurance coverage, according to different research conducted in Sudan (42).

280 The observed phenomenon of women with a history of visiting health facilities in the last 12
281 months exhibiting higher odds of having insurance coverage compared to their counterparts can
282 be attributed to several factors. It is plausible that frequent visits to health facilities provide these
283 women with increased exposure to healthcare services, thereby fostering a deeper
284 understanding of the importance of insurance coverage and potentially reducing the barrier of

285 out-of-pocket expenses. This finding aligns with existing research conducted in East African
286 countries (23).

287 Furthermore, compared to women who were not exposed to the media, those who were
288 exposed showed a higher likelihood to use health insurance. This finding is consistent with
289 research from Ethiopia, Kenya, and Sub-Saharan Africa (34,43,44). this can be explained by
290 high coverage of internet and other media channel in the country which can act as an
291 instrument for knowledge dissemination and raises awareness of the importance of health
292 insurance.

293 **Strength and limitation**

294 The most current nationwide data set was used in the present study. In addition, Furthermore,
295 the study adds to the body of knowledge regarding health insurance ownership among
296 reproductive-age women. In this regard, demographic and health surveys serve as an essential
297 resource for bridging the data gap and enabling evidence-based population health planning and
298 intervention in the recipient nation. Furthermore, the datasets are nationally representative and
299 had a sizable sample size. As such, the results can be generalized to the total female
300 population. It is important to admit limitations.

301 Due to the secondary source nature of the data analysis, it is possible that additional variables
302 or factors that could have affected health insurance coverage but were not included in the study
303 because they were not included in the DHS dataset existed. Furthermore, drawing conclusions
304 about causality is limited by the DHS's cross-sectional design. Self-reports were also used for
305 collecting data. As a result, biases in social desirability and recall may affect the results.

306 **Conclusion**

307 Less than ten percent of Tanzanian women of reproductive age (WRA) are registered in health
308 insurance, which is an unacceptably low coverage. This highlights the urgent need to eliminate
309 barriers preventing this population's access to healthcare. The following stand out as important
310 variables influencing health insurance coverage: marital status, wealth, level of education,
311 media exposure, and recent medical facility visits.

312 The complicated interactions between socioeconomic and informational factors when making
313 healthcare decisions among WRA in Tanzania are highlighted by these findings. Multifaceted
314 approaches are essential to effectively increase coverage. Initiatives aimed at empowering
315 women economically, such microfinance and income-generating businesses, can increase their
316 ability to pay insurance costs.

317 In a similar way, encouraging educational attainment especially in the areas of health and
318 finance through comprehensive education initiatives and chances for vocational training will
319 enable WRA to make knowledgeable decisions on health insurance. Interventions should also
320 prioritize the empowerment of widows and single women by ensuring them access to resources
321 and the freedom to make their own decisions. Moreover, using mass media platforms to spread
322 knowledge about health insurance advantages may also be crucial in raising WRA awareness
323 and adoption. Policymakers and other stakeholders should take note of these observations as
324 Tanzania works to achieve universal health coverage. These interventions should be focused
325 on addressing the root causes of the problem and ensuring that all population segments have
326 fair access to basic healthcare services.

327 **Acknowledgments**

328 The authors are grateful to MEASURE DHS for providing them with the data set.

329 **Funding**

330 This research did not receive any specific grant from funding agencies in the public,
331 commercial, or not-for-profit sectors.

332 **Disclosure**

333 The authors report no conflicts of interest in this work.

334 **Abbreviations**

335 AIC; Akaike Information Criteria; BIC, Bayesian Information Criteria CI, Confidence Interval;
336 CHF, community health fund; DHS; Demographic and Health Survey; EA; Enumeration Area;
337 HI, Health Insurance; LMIC, Low and Middle Income Country; NBS, National Bureau of
338 Statistics; NSSF, National Social Security Fund; PR, Prevalence Risk; PSU, Primary Sampling
339 Unit; SDGs, Sustainable development goals; UHC, Universal health coverage; VIF, Variance
340 Inflation Factor; TIKA, Tiba Kwa Kadi; SHIB, Social Health Insurance Benefit; WRA, Women of
341 Reproductive Age; TDHS, Tanzania Demographic and Health Survey.

342 **Authorship contributions**

343 Sanun Ally Kessy and Jovin R Tibenderana: Conceptualization, Data curation, Formal analysis,
344 Methodology, Visualization, Writing – original draft, Writing – review & editing. Jomo Gimonge:
345 Writing – original draft, Writing – review & editing, Visualization. Fabiola Moshi: Writing – review
346 & editing, Supervision.

347 **REFERENCE**

- 348 1. WHO. Universal health coverage (UHC) [Internet]. 2023 [cited 2024 Feb 5]. Available from:
349 [https://www.who.int/news-room/fact-sheets/detail/universal-health-coverage-\(uhc\)](https://www.who.int/news-room/fact-sheets/detail/universal-health-coverage-(uhc))
- 350 2. Sambo LG, Kirigia JM. Investing in health systems for universal health coverage in Africa.

- 351 2014;1–22.
- 352 3. Burden G, Health D, Collaborator F. Trends in future health financing and coverage : future
353 health spending and universal health coverage in 188 countries ., 2016;
- 354 4. Chauluka M, Uzochukwu BSC, Chinkhumba J. Factors Associated With Coverage of Health
355 Insurance Among Women in. 2022;2(May):1–11.
- 356 5. Shao L, Wang Y, Wang X, Id LJ, Huang R. Factors associated with health insurance
357 ownership among women of reproductive age : A multicountry study in sub-Saharan Africa.
358 2022;1–12. Available from: <http://dx.doi.org/10.1371/journal.pone.0264377>
- 359 6. Kessy SA, Tibenderana JR. Reimagining of health care : Tanzania ' s quest and bold step
360 towards universal health coverage deploying universal health insurance as a catalyst. Int J
361 Surg Glob Heal. 2023;6(e0256):1–2.
- 362 7. Amani PJ, Hurtig A karin, Frumence G, Kiwara AD, Goicolea I, Sebastian MS. Health
363 insurance and health system (un) responsiveness : a qualitative study with elderly in rural
364 Tanzania. 2021;2:1–11.
- 365 8. Kapologwe NA, Meara JG, Kengia JT, Sonda Y, Gwajima D, Alidina S, et al. Development
366 and upgrading of public primary healthcare facilities with essential surgical services
367 infrastructure: A strategy towards achieving universal health coverage in Tanzania. BMC
368 Health Serv Res. 2020;20(1):1–14.
- 369 9. MoHSW. United Republic of Tanzania, Health Sector Strategic Plan (HSSP IV): Reaching all
370 Households with Quality Health Care. July 2015 – June 2020. 2015;2020(June). Available
371 from: https://www.childrenandaids.org/Tanzania_Health-Sector-Strategic-Plan_2015
- 372 10. Demographic and Health Surveys. Demographic and Health Survey and Malaria Indicator
373 Survey (TDHS-MIS). 2022.

- 374 11. Finkelstein A, Taubman S, Wright B, Bernstein M, Gruber J, Newhouse JP, et al. The Oregon
375 Health Insurance Experiment: . Q J Econ [Internet]. 2012;127(3):1057–106. Available from:
376 [http://qje.oxfordjournals.org/content/127/3/1057.short%5Cnpapers3://publication/uuid/E87C5](http://qje.oxfordjournals.org/content/127/3/1057.short%5Cnpapers3://publication/uuid/E87C59AE-C898-4AEB-B56C-53B1FD72EE63)
377 [9AE-C898-4AEB-B56C-53B1FD72EE63](http://qje.oxfordjournals.org/content/127/3/1057.short%5Cnpapers3://publication/uuid/E87C59AE-C898-4AEB-B56C-53B1FD72EE63)
- 378 12. Shudura E, Yoseph A, Tamiso A. Utilization and Predictors of Maternal Health Care Services
379 among Women of Reproductive Age in Hawassa University Health and Demographic
380 Surveillance System Site, South Ethiopia: A Cross-Sectional Study. Adv Public Heal.
381 2020;2020.
- 382 13. Kimani JK, Ettarh R, Warren C, Bellows B. Determinants of health insurance ownership
383 among women in Kenya : evidence from the 2008 – 09 Kenya demographic and health
384 survey. Int J equity Heal Heal. 2014;13(27):1–8.
- 385 14. Saksena P, Hsu J, Evans DB. Financial Risk Protection and Universal Health Coverage:
386 Evidence and Measurement Challenges. PLoS Med. 2014;11(9).
- 387 15. Kutzin J, Sparkes SP. Health systems strengthening , universal health coverage , health
388 security and resilience. 2016;385(9980):165050.
- 389 16. Id HA, Dickson KS, Setorwu K, Id A, Kissah- K, Kofuor E, et al. Prevalence and factors
390 associated with health insurance coverage in urban sub-Saharan Africa : Multilevel analyses
391 of demographic and health survey data. PLoS One [Internet]. 2022;1–14. Available from:
392 <http://dx.doi.org/10.1371/journal.pone.0264162>
- 393 17. Blanchet NJ, Fink G, Osei-Akoto I. The effect of Ghana’s National Health Insurance Scheme
394 on health care utilisation. Ghana Med J. 2012;46(2):76–84.
- 395 18. Diop F, Yazbeck A, Bitran R. The impact of alternative cost recovery schemes on access and
396 equity in niger. Health Policy Plan. 1995;10(3):223–40.

- 397 19. Fiestas Navarrete L, Ghislandi S, Stuckler D, Tediosi F. Inequalities in the benefits of national
398 health insurance on financial protection from out-of-pocket payments and access to health
399 services: Cross-sectional evidence from Ghana. *Health Policy Plan.* 2019;34(9):694–705.
- 400 20. Kirigia JM, Sambo LG, Nganda B, Mwabu GM, Chatora R, Mwase T. Determinants of health
401 insurance ownership among South African women. *BMC Health Serv Res.* 2005;5:1–10.
- 402 21. Kumi-Kyereme A, Amo-Adjei J. Effects of spatial location and household wealth on health
403 insurance subscription among women in Ghana. *BMC Health Serv Res.* 2013;13(1).
- 404 22. Liu TC, Chen CS. An analysis of private health insurance purchasing decisions with national
405 health insurance in Taiwan. *Soc Sci Med [Internet].* 2002 [cited 2024 Jan 3];55(5):755–74.
406 Available from: <https://pubmed.ncbi.nlm.nih.gov/12190269/>
- 407 23. Weldesenbet AB, Kebede SA, Ayele BH, Tusa BS. Health insurance coverage and its
408 associated factors among reproductive-age women in east africa: A multilevel mixed-effects
409 generalized linear model. *Clin Outcomes Res.* 2021;13:693–701.
- 410 24. Dowou RK, Atanuriba GA, Adzigbli LA, Balame SK, Tahidu I, Aggrey-korsah J, et al.
411 Determinants of health insurance coverage among women in Mauritania : a multilevel
412 analysis. 2024;1–11.
- 413 25. Okedo-alex IN, Uneke CJ. A review of the incidence and determinants of catastrophic health
414 expenditure in Nigeria : Implications for universal health coverage. 2019;(May):1–18.
- 415 26. Raymond VG, Ndalla MM, Tibenderana JR. Antenatal care access in Tanzania: progress,
416 challenges, and recommendations for a healthier maternal and child well-being. *Int J Surg*
417 *Glob Heal.* 2023;6(6):1–2.
- 418 27. Masi S De, Bucagu M, Lawrie T, Oladapo OT. Integrated Person-Centered Health Care for
419 All Women During Pregnancy Implementing WHO Recommendations on Antenatal Care for

- 420 a Positive Pregnancy Experience. Global journal of health science. GhspjournalOrg
421 [Internet]. 2017;5(2):197–201. Available from:
422 <https://www.ghspjournal.org/content/5/2/197.short>
- 423 28. NBS Tanzania. Manufacturing Index. 2023 [cited 2023 Dec 16]. p. 4–7 National Bureau of
424 Statistics. Available from:
425 <https://www.nbs.go.tz/index.php/en/%0Ahttp://www.nigerianstat.gov.ng/>
- 426 29. Ministry of Health. Tanzania Ministry of Health [Internet]. 2023 [cited 2023 Dec 16]. Available
427 from: <https://www.moh.go.tz/>
- 428 30. Ministry of Health Zanzibar. Ministry of Health Zanzibar [Internet]. 2023 [cited 2023 Dec 16].
429 p. 1–184. Available from: [http://coastalforests.tfcg.org/pubs/Jozani_biodiversity_inventory](http://coastalforests.tfcg.org/pubs/Jozani_biodiversity_inventory_report_2002.pdf)
430 [report_2002.pdf](http://coastalforests.tfcg.org/pubs/Jozani_biodiversity_inventory_report_2002.pdf)
- 431 31. Macro International Inc. Measure DHS. DHS. 2021 [cited 2023 Dec 16]. The DHS Program -
432 Protecting the Privacy of DHS Survey Respondents. Available from:
433 [https://dhsprogram.com/methodology/Protecting-the-Privacy-of-DHS-Survey-](https://dhsprogram.com/methodology/Protecting-the-Privacy-of-DHS-Survey-Respondents.cfm)
434 [Respondents.cfm](https://dhsprogram.com/methodology/Protecting-the-Privacy-of-DHS-Survey-Respondents.cfm)
- 435 32. Elm E von, Douglas G Altman, Matthias Egger, Stuart J Pocock PCG& JPV for the SI,
436 Abstract. The Strengthening the Reporting of Observational Studies in Epidemiology
437 (STROBE) Statement: guidelines for reporting observational studies. Policy Pract.
438 2014;045120(November):126–40.
- 439 33. Dowou RK, Atanuriba GA, Adzigbli LA, Balame SK, Tahidu I, Aggrey-Korsah J, et al.
440 Determinants of health insurance coverage among women in Mauritania: a multilevel
441 analysis. BMC Public Health. 2024;24(1):1–10.
- 442 34. Id HA, Id A aziz S, Agbaglo E, Dowou RK, Kwabena E, Id A, et al. Mixed effects analysis of
443 factors associated with health insurance coverage among women in sub-Saharan Africa.

- 444 2021;1–15. Available from: <http://dx.doi.org/10.1371/journal.pone.0248411>
- 445 35. Tusa BS, Weldesenbet AB, Kebede SA. Spatial distribution and associated factors of
446 underweight in Ethiopia : An analysis of Ethiopian demographic and health survey , 2016.
447 2020; Available from: <http://dx.doi.org/10.1371/journal.pone.0242744>
- 448 36. Mulenga J, Mulenga MC, Musonda KMC, Phiri C. Examining gender differentials and
449 determinants of private health insurance coverage in Zambia. 2021;1–11.
- 450 37. Aregbeshola BS, Khan SM. Original Article Predictors of Enrolment in the National Health
451 Insurance Scheme Among Women of Reproductive Age in Nigeria. 2018;7(11):1015–23.
- 452 38. Kwarteng A, Akazili J, Welaga P, Dalinjong PA, Asante KP, Sarpong D, et al. The state of
453 enrollment on the National Health Insurance Scheme in rural Ghana after eight years of
454 implementation. 2020;0:1–14.
- 455 39. Rosas ER, Winkler V, Brenner S, Allegri M De. Identifying the determinants of health
456 insurance coverage among Peruvian women of reproductive age : an assessment based on
457 the national Peruvian demographic survey of 2017. 2020;4:1–9.
- 458 40. Salari P, Akweongo P, Aikins M, Tediosi F. Determinants of health insurance enrolment in
459 Ghana : evidence from three national household surveys. 2019;00(August):582–94.
- 460 41. Kebede SA, Liyew AM, Tesema GA, Agegnehu CD, Teshale AB, Alem AZ, et al. Spatial
461 distribution and associated factors of health insurance coverage in Ethiopia : further analysis
462 of Ethiopia demographic and health survey , 2016. 2020;1–10.
- 463 42. Baloul I, Dahlui M. Determinants of health insurance enrolment in Sudan : evidence from
464 Health Utilisation and Expenditure Household Survey 2009. 2014;14(July):6963.
- 465 43. Kebede DB, Belachew YB, Selbana DW, Gizaw AB. Maternal Satisfaction with Antenatal
466 Care and Associated Factors among Pregnant Women in Hossana Town. *Int J Reprod Med.*

467 2020;2020:1–8.

468 44. Kazungu JS, Barasa EW. Examining levels , distribution and correlates of health insurance
469 coverage in Kenya. 2017;22(9):1175–85.

470




Figure1: Proportion of Health insurance coverage among women of reproductive age




Figure2: Specific types of health insurance coverage