Molecular profiling of neonatal dried blood spots reveals changes in innate and adaptive immunity following fetal inflammatory response

Daniel Costa[1,2], Núria Bonet[3], Amanda Solé[2,4,5], José Manuel González de Aledo-Castillo[6], Eduard Sabidó[2,4,5], Ferran Casals[3], Carlota Rovira[7], Alfons Nadal[8], Jose Luis Marin[6], Teresa Cobo[6,*], Robert Castelo[2,9,*]

[1]Department of Pediatrics, Hospital de Figueres, Figueres, Spain; [2]Department of Experimental and Health Sciences, Universitat Pompeu Fabra (UPF), Barcelona, Spain; [3]Genomics Core Facility, Department of Experimental and Health Sciences, Universitat Pompeu Fabra (UPF), Barcelona, Spain; [4]Proteomics Unit, Centre de Regulació Genòmica (CRG), Barcelona, Spain; [5]Barcelona Institute of Science and Technology (BIST), Barcelona, Spain; [6]Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Centre for Biomedical Research on Rare Diseases (CIBER-ER), Barcelona, Spain;[7]Hospital Sant Joan de Déu, Barcelona, Spain; [8]Department of Pathology, Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Barcelona, Spain; [9]Research Programme on Biomedical Informatics, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Barcelona, Spain; *Correspondence: Robert Castelo (robert.castelo@upf.edu), Teresa Cobo (tcobo@clinic.cat)

Word count (excluding Abstract, Methods, References and Figure and Table Legends): 3,638

Abstract

The fetal inflammatory response (FIR) increases the risk of perinatal brain injury, particularly in extremely low gestational age newborns (ELGANs, < 28 weeks of gestation). One of the mechanisms contributing to such a risk is a postnatal intermittent or sustained systemic inflammation (ISSI) following FIR. The link between prenatal and postnatal systemic inflammation is supported by the presence of well-established inflammatory biomarkers in the umbilical cord and peripheral blood. However, the extent of molecular changes contributing to this association is unknown. Using RNA sequencing and mass spectrometry proteomics, we profiled the transcriptome and proteome of archived neonatal dried blood spot (DBS) specimens from 21 ELGANs. Comparing FIR-affected and unaffected ELGANs, we identified 783 gene and 27 protein expression changes of 50% magnitude or more, and an experiment-wide significance level below 5% false discovery rate. These expression changes confirm the robust postnatal activation of the innate immune system in FIR-affected ELGANs and reveal an impairment of their adaptive immunity. In turn, the altered pathways provide clues about the molecular mechanisms triggering ISSI after FIR, and the onset of perinatal brain injury.

Keywords

Premature birth; fetal inflammatory response; perinatal brain injury; transcriptomics; proteomics

Introduction

Intraamniotic infection (IAI), one of the main causes of preterm birth, can lead to a maternal (MIR) and a fetal (FIR) inflammatory response, which can be detected in placental tissues[1] (Supp. Table S1). FIR can cause damage to fetal organs presumably by triggering a systemic inflammation, leading to distant organ injury and increasing the risk of multiple adverse neonatal outcomes, such as perinatal brain damage[2,3]. Understanding the molecular dynamics of FIR and its downstream alterations before and after birth is thus crucial to developing diagnostic and therapeutic strategies that improve the clinical outcome of extreme preterm birth.

In a previous work, we found that FIR triggers a broad and complex transcriptional response in umbilical cord (UC) tissue[4], which others have shown to correlate with cognitive impairment at 10 years of age[5]. The molecular and cellular alterations induced by FIR can persist after birth[6] and interact with postnatal inflammation-initiating illnesses such as bacteremia, mechanical ventilation or necrotizing enterocolitis[7]. This interaction ultimately leads to what is known as an

intermittent or sustained systemic inflammation (ISSI)[8,9], which is in turn strongly associated

with perinatal brain injury and developmental disorders in ELGANs[9] (Extremely Low Gestational

Age Newborn, i.e. birth before the 28 week of gestation). The joint presence of placental

inflammation and ISSI appears to increase the risk of perinatal brain injury, highlighting the

important contribution of postnatal systemic inflammation to that risk[10,11]. However, little is

known about the extent of postnatal molecular changes associated with FIR, which can

contribute to understanding the link between FIR and ISSI and consequently provide an array of

candidate biomarkers and therapeutic targets for perinatal brain injury.

Here, we analyze the transcriptome and proteome of archived neonatal DBSs from ELGANs

and identify RNA and protein expression changes in whole blood that are significantly

associated with FIR. These molecular changes provide, with an unprecedented level of

resolution, a snapshot of the pathways participating in ISSI as a result of the exposure to FIR.

15   Results

16   Clinical characteristics of ELGANs and molecular profiling of DBS samples

The samples analyzed in this study were from 21 ELGANs (8 females and 13 males), who met

the eligibility criteria (see Methods). Histological acute chorioamnionitis was diagnosed in 15

cases. MIR was in stage I in two cases, stage II in one case, and stage III in 12 cases. Ten of

the 12 cases with stage III MIR showed FIR: four in stage I, five stage II, and one stage III

(Supp. Fig. S1). Other clinical characteristics and outcomes are presented in Table 1,

separately for those with and without FIR. For instance, FIR-affected ELGANs presented a

higher frequency of microbial invasion of the amniotic cavity (89% vs 29%, two-tailed Fisher's

exact test P=0.035) than unaffected ones, and had significantly higher maternal blood C-

reactive protein (CRP) levels at admission (t-test P=0.041), as well as significantly higher (t-test

P=0.034) and lower (t-test P=0.018) amniotic fluid (AF) IL-6 and glucose levels, respectively.

Our data are compatible with no differences in neonatal morbidities between FIR-affected and

unaffected ELGANs, although a higher frequency of neonatal sepsis and intraventricular

hemorrhage was observed in FIR cases. Mean gestational age (GA) and birth weight were

similar between the two groups, averaging 26 weeks and 860 g, respectively. Such an

extremely low birth weight, with a lowest value of 580 g and up to 8 neonates under 750 g,

precludes a standard blood extraction after birth for the purpose of molecular profiling in

research. On the other hand, UC blood is not an alternative either, because is not

representative of postnatal immunity[12]. For this reason, neonatal DBS samples archived at the

newborn screening program constitute a minimally invasive option to profile the transcriptome

and proteome in peripheral blood from ELGANs.

Table 1 | Clinical characteristics of ELGANs

Clinical characteristics	No FIR (n=11)	FIR (n=10)	*P* value
Baseline characteristics of pregnancy and delivery			
Cesarean section: n, (%)	8 (72.7)	3 (30.0)	0.086
Nulliparity: n, (%)	9 (81.8)	5 (50.0)	0.183
Preterm labor: n, (%)	3 (27.3)	3 (30.0)	1.000
Preterm, pre-labor rupture of membranes: n, (%)	1 (9.1)	4 (40.0)	0.149
Cervical insufficiency: n, (%)	2 (18.2)	3 (30.0)	0.635
Placenta abruption: n, (%)	4 (36.4)	0 (0.0)	0.090
Preeclampsia: n, (%)	1 (9.1)	0 (0.0)	1.000
Complete course (at least 2 doses) of antenatal steroids: n, (%)	11 (100)	7 (70.0)	0.090
Maternal CRP concentration at admission: (mg/dL), mean±SD	2.3 ± 3.4	5.8 ± 3.8	0.041
Biomarkers of intra-amniotic infection and inflammation			
MIAC: n, (%)	2 (28.6) 4 missing obs.	8 (88.9) 1 missing obs.	0.035
Clinical chorioamnionitis: n, (%)	0 (0.0)	3 (30.0)	0.090
Histological chorioamnionitis: n, (%)	5 (45.5)	10 (100.0)	0.012
Concentration of IL-6 in AF: (pg/mL), (Q1, Q3)	21,153.5 (6,695.50, 35,975.75) 5 missing obs.	112,271.5 (34,968.07, 162,866.25) 2 missing obs.	0.034
Concentration of glucose: (mg/dL), (Q1, Q3)	23.6 (10, 37) 4 missing obs.	2.4 (0, 2) 1 missing obs.	0.018
Baseline characteristics of infants			
Male: n, (%)	7 (63.6)	6 (60.0)	1.000
Birth weight: (g), mean±SD	841 ± 138	880 ± 159	0.552
Respiratory distress syndrome: n, (%)	5 (45.5)	5 (50.0)	1.000
Patent ductus arteriosus: n, (%)	6 (54.5)	6 (60.0)	1.000
Intraventricular hemorrhage: n, (%)	1 (9.1)	5 (50.0)	0.063
White matter disease: n, (%)	1 (9.1)	2 (20.0)	0.586
Retinopathy of prematurity: n, (%)	5 (45.5)	3 (37.5)	1.000

		2 missing obs.	
Necrotizing enterocolitis: n, (%)	1 (9.1)	0 (0.0)	1.000
Sepsis: n, (%)	1 (9.1)	5 (50.0)	0.063
Positive blood culture: n, (%)	1 (9.1)	2 (20.0)	0.586
GA 23-24 completed weeks: n, (%)	2 (18.2)	1 (10.0)	1.000
GA 25-26 completed weeks: n, (%)	1 (9.1)	5 (50.0)	0.063
GA 27 completed weeks: n, (%)	8 (72.7)	4 (40.0)	0.198
DBS heel sampling days > 3: n, (%)	4 (36.4)	5 (50.0)	0.678
Maximum CRP within first 8 postnatal days: (md/dL), mean±SD	1.0 ± 2.2	3.5 ± 5.3	0.187
WBC count at birth: (10^9/L), mean±SD	8.6 ± 3.5	23.0 ± 15.6	0.017
Postnatal WBC count within first 8 postnatal days: (10^9/L), mean±SD	13.2 ± 5.2	44.1 ± 13.4	< 0.001
Maximum ANC within first 8 postnatal days: (10^9/L), mean±SD	6.2 ± 4.3	29.0 ± 10.8	< 0.001
ALC at birth: (10^9/L), mean±SD	4.3 ± 1.2	6.7 ± 3.1	0.039
Lymphocyte percentage at birth: (%), mean±SD	54.3 ± 14.5	35.4 ± 16.3	0.012
Maximum ALC within first 8 postnatal days: (10^9/L), mean±SD	6.0 ± 3.2	10.8 ± 4.1	0.009
Maximum lymphocyte percentage within first 8 postnatal days: (%), mean±SD	46.9 ± 17.1	25.4 ± 10.4	0.003

CRP: C-reactive protein. AF: amniotic fluid. MIAC: microbial invasion of the amniotic cavity defined by a positive AF culture. WBC:
white blood cell. ANC: absolute neutrophil count. ALC: absolute lymphocyte count. Intraventricular hemorrhage: any grade between
I to IV according Papile et al (1978)[13] grading system. Sepsis: confirmed or suspected bacterial blood infection, which is classified
as early if within 72 hours after birth and classified as late if after the third postnatal day. Categorical characteristics were tested with
respect to FIR status using a two-tailed Fisher's exact test. Continuous characteristics were tested with respect to FIR status using a
t-test.

Figure 1: Transcriptomics and proteomics profiling of archived DBSs. (a) Sequencing depth per sample, broken by origin of sequence reads. (b) Probability of a C>T deamination due to damage, as function of DBS age in years. (c) Sample differences in terms of fold-changes between transcriptomic samples in base-2 logarithmic scale (logFC), projected along the *x* and *y*-axes. Distance between points is proportional to the dissimilarity between samples. (d) Same as (c) but calculated from proteomic samples.

Total RNA sequencing of the 21 archived DBSs produced more than 500 million paired-end reads, which we transformed into a table of 25,221 gene-level summarized count expression profiles. Nearly half of these genes (11,279) showed reliable levels of expression in at least 6 of the 21 samples and were those selected for further analysis (see Methods). Figure 1a shows the distribution of sequencing depth across different genomic elements for each sample, where only about ⅓ of the total sequencing depth derives from exonic reads. DBS samples in this cohort were stored at room temperature for one to seven years until RNA extraction and sequencing. Over this period of time, degradation and fragmentation processes affect RNA integrity and we thus investigated whether, as observed in ancient DNA samples[14], there is a

detectable damage to the RNA due to cytosine deamination on such a short timescale (see

Methods). Indeed, we found a significant linear relationship (r=0.8, P < 0.001) between the

probability of a C>T substitution due to damage and the elapsed time between DBS extraction

and RNA sequencing (Fig. 1b). Additionally, mass-spectrometry proteomics on DBSs produced

649 quantified protein expression profiles (Supplementary Data File 1), from which we selected

245 as being reliably expressed in at least 6 of the 21 samples (see Methods).

Sample-level changes of transcriptomics data, projected in two dimensions (Fig. 1c), show that

gene-level expression profiles derived from exonic reads capture the differences in placental

inflammation throughout the different stages of MIR and FIR. Analogously, DBS protein

expression profiles also show a clear separation between infants with and without MIR and FIR

(Fig. 1d). In summary, transcriptomics and proteomics of archived DBS samples can produce a

large catalog of molecular measurements that capture clinical features of the perinatal period.

Archived neonatal DBSs contain gene and protein expression changes associated with
FIR

The differential expression analysis of RNA-seq transcriptomic profiles between FIR and non-

FIR exposed ELGANs identified 783 differentially expressed (DE) genes under an experiment-

wide significance threshold of 5% false discovery rate (FDR) and a minimum 50%-fold change

(Fig. 2a, Supp. Table S2). We replicated similar magnitude changes from DE genes on a

logarithmic scale (r=0.75, R^2=0.56, P < 0.001) in the analogous differential expression analysis

of a DBS gene expression dataset[15] from the US ELGAN cohort (Fig. 2b, Supp. Table S3 and

Supp. Fig. S2 to S5), despite differences in sample preparation and gene expression profiling

technology (see Methods). The hierarchical clustering of the 21 transcriptomic samples, using

the 783 DE genes, leads to a perfect separation of the neonates who were affected by FIR, from

those that were unaffected (Fig. 3).

Figure 2: Differential expression analysis between FIR-affected and unaffected ELGANs, of transcriptomic and proteomic profiles from archived DBSs. (a) Volcano plot of transcriptomics data. Raw *P* values in negative logarithmic scale on the *y*-axis as function of the log2-fold change on the *x*-axis. Orange dots highlight genes from a NLRC4-MAS signature[16], red dots highlight genes from a G-MDSCs signature[17] and blue dots highlight transcription factor coding genes involved in regulating adaptive immunity. (b) Fold changes of this study on the *y*-axis as function of fold changes in data from the US ELGAN cohort in the *x*-axis. (c) Volcano plot of proteomics data. (d) Fold changes from proteomics data on the *y*-axis as function of fold changes from transcriptomics data on the *x*-axis.

A further question is whether there are RNA changes in peripheral blood across the different stages of MIR and FIR. Indeed, we found 58 genes with FDR < 5%, for which a combination of MIR and FIR stages, reflecting an increasing exposure to prenatal inflammation, led to a minimum increase of 50% in gene expression levels for each additional combined inflammatory stage (Supplementary Figure S6 and Table S4).

The differential expression analysis of proteomic profiles between FIR-exposed neonates and those who were unexposed identified 27 DE proteins at 5% FDR with a minimum 50%-fold change (Fig. 2c, Supp. Table S5). Figure 2d shows a comparison between log2-fold changes of the transcriptomics and proteomics analyses. Six of the 27 DE proteins are encoded by genes that are also DE by RNA-seq, while for 14 of them, their corresponding genes do not show fold-changes above 50% in the transcriptomics data. The remaining seven DE proteins do not have minimum levels of expression in the RNA-seq assay. Among them, F2, FGA, FGB and A1BG are secreted from the liver into the blood, and this may also be the case for SERPINA3 and APOE, which have a biased expression towards liver. Overall, these results show that in peripheral blood from ELGANs there are sizeable numbers of transcriptomic and proteomic changes associated with FIR that can be detected in archived DBS.

Figure 3: FIR gene expression signature and functional enrichment analysis. Heatmap of expression values for 783 DE genes with FDR < 5% and minimum 1.5-fold change between FIR-affected and unaffected ELGANs, obtained after removing FIR-unrelated variability. Dendrograms on the *x* and *y*-axes represent the hierarchical clustering of samples and genes, respectively. Leaves on the gene dendrogram are color coded according to whether genes encode for transcription factors (orange) or RNA-binding proteins (green), while those on the sample dendrogram are color coded according to FIR status, where red indicates samples derived from FIR-affected infants and blue unaffected. The right dot-matrix represents DE genes (*y*-axis) belonging to GO terms (*x*-axis) significantly enriched (FDR < 10% and

odds-ratio -OR > 1.5) by upregulated (top) and downregulated (bottom) DE genes. The bottom-left dot-

matrix represents the presence or absence of phenotypes, with grey indicating missing data.

FIR is associated with a persistent postnatal activation of the innate immune system

Protein biomarkers have helped in the discovery of a postnatal systemic inflammatory hit

following FIR[6]. However, the extent of such molecular changes was previously unknown. Our

transcriptomic data from DBS samples shows the upregulation in FIR of many genes involved in

the innate immune response (Fig. 2a; Supp. Table S2). Such genes include those coding for

pattern recognition receptors (PRRs, e.g., *NLRC4, CLEC4E, FPR1*), inflammatory transcription

factors (e.g., *CEBPD, KLF5*), acute phase response proteins (*HP*, *ORM1*), as well as multiple

inflammatory mediators including cytokines (e.g., *IL18, IL1R1, RETN*), calgranulins (S100A8,

S100A9,S100A12), chemokines (e.g., *CXCR1, CXCL1*), complement (e.g., *C5AR2, C3AR1*),

lipid mediators (e.g., *LTA4H, ALOX5AP, CYP4F3*), proteolytic enzymes (e.g., *MMP8, MMP9*)

and adhesion molecules (e.g., *ICAM3, ITGAM*). Our proteomic data from DBS samples (Fig. 2c,

Supp. Table S5) also shows protein overexpression in FIR of diverse inflammatory mediators of

the innate immune system, such as the S100A8, S100A9, LTF, MPO and DEFA3 proteins.

A functional enrichment analysis using the Gene Ontology (GO) database (Fig. 3) shows GO

terms associated with the innate immune system, such as *innate immune response in mucosa*

(OR=17.7) and *defense response to fungus* (OR=14.3), which were significantly enriched (FDR

< 10%) in upregulated genes (Supp. Table S6; see Methods). Moreover, we also observed a

significant enrichment of DE genes in curated gene sets of the innate immune response, such

as the InnateDB[18] (one-tailed Fisher's exact $P < 2.2 \times 10^{-16}$, OR=3.2). All these gene and

protein expression changes constitute the most comprehensive description to date of the

persistent postnatal activation of the innate immune system in ELGANs affected by FIR.

FIR is associated with NLRC4 inflammasome activation

The activation of the innate immune system involves the assembly of different multiprotein

complexes known as inflammasomes, which are categorized according to their constituent

PRRs. After detection of microbial and endogenous danger signals by PRRs, inflammasomes

induce, via activation of caspase-1 (CASP1), the maturation and release of IL-1β and IL-18

cytokines, bacterial clearance and pyroptosis. The release of interleukin-1 family cytokines such

as IL-1β and IL-18 has potent effects on neutrophil activation and recruitment[19]. The NLRC4

inflammasome belongs to the intracellular cytosolic PRR family of NOD-like receptors and

detects the cytosolic presence of bacterial components.

We found that FIR upregulates the expression of the NLRC4 inflammasome and IL-1 family-

related protein-coding genes (e.g., *NLRC4, PYCARD, IL18, IL1R1*). We also found a significant

overlap (Fig. 2a, one-tailed Fisher's exact P < 2.2x10^{-16}, OR=10.4) of DE genes with a

transcriptional signature of the NLRC4-infantile-onset macrophage activation syndrome

(MAS)[16]. Consistent with these results, we observed the upregulation of numerous genes

involved in neutrophil biology and function (Supp. Table S7), significantly enriching (FDR <

10%, OR > 1.5) GO terms related to neutrophil activation[20], such as *neutrophil migration*

(OR=6.3) or *positive regulation of neutrophil chemotaxis* (OR=7.5); see Fig. 3 and

Supplementary Table S6. These findings suggest that FIR leads to an initial proinflammatory-

driven response (*IL-6*, *IL-1b*, *CXCL8*), followed by a posterior neonatal robust activation of the

IL-18/IL-1 axis through the NLRC4-inflammasome that can induce a persistent, systemic

neutrophil activation.

FIR is associated with the impairment of adaptive immunity and thymic function through

the expansion of granulocytic myeloid-derived suppressor cells

The mechanism by which antenatal FIR leads to a postnatal systemic inflammation is unknown.

Among several plausible alternatives, Dammann and Leviton (2014)[8] suggested an alteration in

the number of T cells and their regulation. A closer look at the hierarchical clustering of genes in

Figure 3 reveals an enrichment of downregulated transcription-factor coding genes (one-tailed

Fisher's exact test *P* < 0.001, OR=2.9, Supp. Table S8) in FIR-affected ELGANs. Many of these

genes are regulators of T cell development and activation, such as *TBX21*, *GATA3* and *RORA*,

respectively involved in T-helper(h) fate of CD4+ T cells in Th1, Th2 and Th17 lineages (Fig.

2a). More generally, we found the downregulation of key genes associated with the IL-7 and IL-

2 signaling pathways (e.g., *IL7R, BCL2, PIK3AP1, IL2RB, CD3E, CD3G*) involved in T cell

development (Supp. Tables S2 and S9). The TCR signaling pathway (e.g., *CD3E, CD3G, LCK,*

FYN, ZAP70, CD247) and T cell differentiation co-stimulatory receptor genes (e.g., *CD28,*

ICOS, SLAMF1) were also downregulated[21]. Likewise, the inhibitory and stimulatory genes

KLRC1, KLRC3, KLRC4 that belong to the natural killer (NK) cell lectin-like receptors of the NK

gene complex, and genes involved in NK cell cytotoxicity, such as *PRF1, SH2D1B* genes, were

also downregulated in FIR-affected ELGANs[21]. Interestingly, we observed a significant overlap

of downregulated genes with an NLRC4-infantile-onset MAS signature (one-tailed Fisher's exact

P < 2.2x10[-16], OR=21.5)[16], which is associated with cytotoxic and regulatory T cell dysfunction.

Recent thymic emigrants consist of phenotypically and functionally immature T cells that have

recently egressed from the thymus, which predominate in the fetus and newborns. These cells

can be identified by their content of T cell receptor excision circles (TRECs)[22], which are DNA

fragments excised during the rearrangement of segments of T cell receptor (TCR) genes in the

thymus. TRECs are used as a measure of the number of T lymphocytes in newborns and,

indirectly, as an indicator of a possible thymus dysfunction. We found downregulation of the

genes coding for the PTK7 and S1PR1 proteins in FIR-affected ELGANs. PTK7 is a cell marker

of CD4+ T cell recent thymic emigrants used for assessing thymic activity[23], while the S1PR1

receptor expression in CD4+ T cells is required for their survival and thymic emigration[24].

Consistent with these results, FIR-affected ELGANs had significantly lower levels of TRECs in

the analyzed DBSs (P=0.0432; Fig. 4a, see Methods), a significantly higher postnatal neutrophil

to lymphocyte ratio (P=0.002; Fig. 4b), and a significantly lower percent of all leukocytes that

are lymphocytes at birth (P=0.012; Table 1) and during the first postnatal week[25] (P=0.003; Fig.

4c).

Myeloid-derived suppressor cells (MDSCs) develop under pathological conditions from myeloid

precursors. According to their phenotype, MDSCs can be classified into monocytic(M)-MDSCs

or granulocytic(G)-MDSCs. Postnatally, there is a high level of circulating MDSCs,

predominantly G-MDSCs, which correlates with inflammatory markers of perinatal infection[26,27].

Neonatal MDSCs display reduced apoptosis and immunosuppressive activity after bacterial

infection[28]. Indeed, these cells are potent inhibitors of adaptive immune responses, particularly

of T and NK cells, by different mechanisms, including the production of reactive oxygen species

(ROS)[29–31], prostaglandin E2 (PGE2), and the depletion of L-arginine by the arginase 1 enzyme

(ARG1)[32,33]. In our data, we found that the *ARG1* gene was 2.5-fold upregulated in FIR-affected

ELGANs, as well as the *OLR1* gene, encoding the lectin-type oxidized LDL receptor 1 (LOX1), a

marker of G-MDSCs[17], which was over 10-fold upregulated. We also found a significant overlap

(one-tailed Fisher's exact P < 2.2e-16, OR=26.4) between FIR upregulated genes and a G-

MDSC expression signature[17] (Fig. 2a).

Figure 4: Association of FIR with clinical and molecular signatures. (a) TREC corrected fluorescent counts on logarithmic scale by FIR status, after adjusting for sex, maximum lymphocyte count during the first postnatal week and batch. Diamonds indicate mean values. There are multiple measurements per newborn. (b) Postnatal ratio of neutrophil absolute count to absolute lymphocyte count, during first postnatal week, by FIR status. Diamonds indicate mean values. (c) Percent of all leukocytes that are lymphocytes during first postnatal week on the *y*-axis as function of the postnatal white blood cell count on the *x*-axis. (d) Fold changes in this study on the *y*-axis as function of FIR fold changes in UC tissue on the *x*-axis.

One of the mechanisms by which G-MDSCs suppress T cell responses is the release of ROS molecules. We found an upregulation of genes coding for proteins essential to ROS production such as *CYBB, NCF1, NCF4, MPO, RAB27A*, as well as genes coding for antioxidant enzymes (e.g., *SOD2, TXN* and the Msr family -*MsrA,MsrB1, MsrB2, MsrB3*). Genes encoding Msr proteins reduce protein methionine oxidation, a post-translational alteration that correlates with oxidative stress[34]. In summary, our data suggest the expansion of G-MDSCs as a mechanism

2   contributing to the impairment of adaptive immunity in ELGANs affected by FIR, and

3   consequently, to a postnatal systemic inflammation sustained over time after birth.

Discussion

We have described the largest catalog to date of postnatal transcriptomic and proteomic

changes associated with FIR in archived DBSs from ELGANs. This catalog demonstrates that

archived neonatal DBSs constitute a valuable source of genome-wide molecular information

about the perinatal period. More importantly, we have shown that molecular changes in this

catalog provide clues to the underlying relationships between FIR and ISSI.

Our data provide molecular evidence of the postnatal activation of NLRC4-inflammasome

dependent mechanisms, which may contribute to the development of a postnatal innate immune

response in FIR-affected ELGANs (Fig. 5a). The NLRC4-inflammasome hyperactivity can cause

diverse auto-inflammatory diseases, such as the infantile-onset MAS syndrome and

enterocolitis. These disorders are characterized by a huge elevation of IL-18 in the blood[16].

Recently, this protein has been proposed as a therapeutic target in neonatal sepsis[35], and in

fact, we observed a 2-fold overexpression of the *IL18* gene in FIR-affected infants. In preterm

newborns, Liang et. al (2017)[36] reported the case of a 28-week preterm neonate with a *de novo*

gain-of-function mutation in the *NLRC4* gene, who died of a fatal syndrome of excessive

immune activation likely to have begun in utero. NLRC4 dependent mechanisms can also

contribute to brain injury induced by cerebral ischemia, as has recently been described in a

rodent-based model of stroke[37]. Interestingly, caffeine therapy for the apnea of prematurity,

which reduces the incidence of moderate to severe neurodevelopmental disabilities in very low

birth weight infants[38], can also reduce NLRC4-inflammasome activation[39].

Figure 5: Model for the association between FIR, ISSI and perinatal brain injury. (a) FIR and postnatal

diseases can induce the postnatal activation of NLRC4-inflammasome dependent mechanisms in innate

immune cells that trigger the release of proinflammatory cytokines IL-1β and IL-18. (b) These cytokines

have a major role in neutrophil expansion and activation. (c) The neutrophil proliferation under

pathological conditions can induce the expansion of myeloid-derived suppressor cells (MDSCs). These

cells are potent suppressors of T and NK cells and strong producers of radical oxygen species (ROS). (d)

Robust production of calgranulins and IL-1β/ IL-18 cytokines can lead to an innate autoinflammatory loop

by autocrine feedback-loops. (e) The innate autoinflammatory loop, the immune adaptive suppression

and oxidative stress, can lead to ISSI. (f) ISSI can cause the infiltration of innate immune cells into the

brain, such as neutrophils, induced by diverse inflammatory mediators, such as calgranulins, leading to

perinatal brain injury in preterm newborns.

The release of the proinflammatory cytokines IL-18 and IL-1β after NLRC4 inflammasome

activation induces the expansion, migration and activation of neutrophils (Fig. 5b). Molecular

alterations in neonatal neutrophils, such as a reduced expression of Fas receptors, can lead to

delayed neutrophil apoptosis[40,41] and the persistence of proinflammatory responses[42]. We found

a significantly higher neutrophil to lymphocyte ratio in FIR-affected ELGANs (Fig. 4b), the

overexpression of *S100A8*, *S100A9*, *IL18*, *MMP8* and *OLFM4*, and the underexpression of

gasdermin B genes (Supp. Table S2). Calgranulins S100A9 and S100A8, and IL-18 can inhibit

neutrophil apoptosis[43,44] and neutrophils can evade pyroptosis, which is mediated by gasdermin

family proteins[45]. Furthermore, the robust production of calgranulins and the activation of the IL-

1/IL-18 axis can trigger a neutrophil-associated inflammation-boosting loop binding to PRRs and

triggering downstream inflammatory signaling pathways, just as in auto-inflammatory

disorders[46]. Among children with septic shock, elevated *MMP8* and *OLFM4* blood gene

expression levels, are associated with higher rates of mortality and organ failure[47,48]. Similarly,

other studies have also shown that preterm neonates with abnormal neuroimaging studies had

increased blood neutrophil activation markers[49,50].

The analysis of downregulated genes supports the inhibition of adaptive immune responses in

FIR (Fig. 1a and 3), which is consistent with the relationships we see between TRECs and FIR

(Fig. 4a), as well as the lymphocyte percent among ELGANs with and without FIR and MIR (Fig.

4c). The reduction of TRECs and the downregulation of the *PTK7* and *S1PR1* genes suggest a

thymic involution associated with FIR[51], which may lead to a relative increase in

proinflammatory CD31-CD4+ T cells producing TNF[52]. Perhaps that is why thymic involution has

also been associated with increased risk of brain damage in ELGANs[53]. On the other hand, in

our earlier investigation[4] of UC tissue from ELGANs, we found multiple upregulated genes in

FIR that code for proteins involved in the expansion of MDSCs (Fig. 4d). In the present study,

we found a significant overlap between FIR DE genes and a previously published G-MDSC

signature[17] (Fig. 5c).

In newborns, the LTF protein appears to promote the secretion of S100A9[54], which can induce

the production of PGE2, thereby repressing the expression of LCK and ZAP70 proteins in CD4+

T cells, and impairing their activation[30,31,55]. Overall, our results suggest that FIR may induce an

initial prenatal pro-inflammatory-driven response (IL-6, IL1-b, CXCL8) evident in UC tissue and

UC blood, followed by neonatal innate immune activation mediated by the NLRC4-

inflammasome. As a result of these events, FIR-affected ELGANs display a robust neutrophil

and G-MDSC expansion during the first postnatal days, which can lead to adaptive immune

suppression and the onset of an auto-inflammatory loop, contributing to ISSI and consequently

increasing the risk of perinatal brain injury (Fig. 5d).

An important feature of G-MDSCs is the production of ROS molecules, which can lead to

oxidative stress and damage. Moreover, preterm newborns have a reduced antioxidant capacity

that increases the risk of ROS-induced damage, which is an important factor in numerous

diseases in premature infants, such as perinatal brain damage[56]. In fact, ELGANs are born with

deficient levels of selenium[57], a cofactor for the MsrB and thioredoxin enzyme activity required

for proper antioxidant function. We found a 10-fold overexpression in FIR of both the *OLR1*

gene, which encodes LOX1, a marker of G-MDSCs[17], and the gene coding for the heat shock

protein HSPA1A. HSPA1A can activate LOX1 in UC neutrophils in response to in vitro exposure

to microbial peptidoglycan[58], leading to the downstream production of ROS. However, the LTF

protein, which we found to be significantly overexpressed in both transcriptomics and

proteomics data, can protect newborns from oxidative stress and damage by attenuating

inflammatory responses and controlling oxidative cell injury induced by innate immune

activation[59].

Using well-established biomarkers, it has been shown that antenatal and postnatal

inflammation, the so-called two-inflammatory hit, increases the risk of perinatal brain damage in

preterm newborns[10]. Here, we found the overexpression of multiple genes encoding proteins

that can be associated with perinatal brain damage in FIR-exposed newborns (Fig. 5e), such as

LCN2, CXCL1, CXCR2, S100A8 and S100A9, as well as histone-related genes (e.g.,

HIST1H2BL). LCN2 is an iron-binding protein contained in neutrophil granules that is associated

with neuroinflammation[60]. Using an animal model of chorioamnionitis, Yellowhair et al. (2018)[61]

found that chorioamnionitis-induced fetal brain damage was associated with increased

expression of CXCL1 and CXCR2 in placental and fetal brain, and with an elevated number of

cerebral CXCR2+ neutrophils. With respect to calgranulins, it has been observed that the brain

expression of S100A8 and S100A9 proteins was increased in the brain of patients dying of

sepsis; specifically, S100A9 expression was required for the cerebral recruitment of neutrophils

and microglia activation in an animal model of sepsis[62]. Finally, extracellular histones can act as

damage-associated molecular patterns, such as calgranulins, activating inflammatory signaling

pathways and inducing multiple organ-specific damage[63]. In an animal study, histone H1 was

found to induce proinflammatory responses in microglia and astrocytes[64,65]. Remarkably, we

found that both the gene and protein expression of S100A9, S100A8 and HIST1H2BL, was

significantly higher in FIR-affected ELGANs, while our proteomics data also revealed the

significant overexpression of HIST1H4A and HIST1H1C (Fig. 2c,d). In summary, our findings

provide new insights into the molecular mechanisms that trigger ISSI after FIR, which may help

to find new diagnostic biomarkers and therapeutic targets for pathological conditions derived

from extreme prematurity, such as perinatal brain injury.

Methods

Study design

We performed a retrospective chart review of extreme preterm births (< 28 weeks of

gestation) at the *Hospital Clinic de Barcelona* admitted to the neonatal intensive care unit

(NICU) between January 2010 and October 2016. Two hundred thirty-four cases born alive

were found. From this initial cohort, we excluded newborns from multiple pregnancies, outborn

deliveries and those who either died in the delivery room or within the first week after birth,

presented major congenital anomalies, received a blood transfusion before collecting the DBS,

had no histopathological examination of the placenta and umbilical cord, or whose DBS was

collected later than 8 days after birth. We attempted to maximize the number of individuals with

available biomarkers of intra-amniotic infection and inflammation, while recruiting a similar

number of FIR and non-FIR cases within a total maximum of 25. A final number of n=21 infants

met the eligibility criteria and their families provided written consent to use their DBS samples

for this study. At the time of processing these DBS samples, they had been stored at room

temperature by the Catalan newborn screening program for between 1 and 7 years. The study

protocol was approved by the Institutional Review Board from the *Hospital Clínic de Barcelona*

(September 16[th], 2016; ref. HCB/2016/0713).

Clinical outcomes

GA at birth was defined according to the first-trimester ultrasound examination. Preterm birth

was classified by the clinical presentation: preterm labor, preterm pre-labor rupture of

membranes, cervical insufficiency, placenta abruption, and preeclampsia. The laboratory results

of AF cultures (genital mycoplasma, aerobic and anaerobic) and IL-6 levels, as well as the

mother's blood CRP, WBC and ANC at admission were also recorded; see Table 1. Clinical

chorioamnionitis was defined following the criteria of Gibbs[66]. Slides from the placenta and

umbilical cord were independently reviewed by two experienced pathologists blinded to

experimental results and previous diagnosis. Maternal (MIR) and fetal (FIR) reactions were

staged and graded according to published criteria[1,67]. Cases read discrepantly by the two

pathologists were simultaneously re-reviewed on a two-headed microscope in order to reach a

consensus (Supp. Table S1 and Supp. Fig. S1). The annotated neonatal morbidity included:

respiratory distress syndrome (RDS); intraventricular hemorrhage (IVH); white matter disease

(WMD); retinopathy of prematurity (ROP); sepsis; necrotizing enterocolitis (NEC); patent ductus

arteriosus (PDA); and laboratory investigations at birth, and during the first 8 days after birth

(CRP concentration, WBC and ANC). Hypothesis tests on clinical variables used in Table 1

include two-tailed Fisher's exact test for categorical variables and *t*-tests for those that were

continuous.

Analysis of T cell receptor excision circles

TREC data were obtained using the EnLite™ Neonatal TREC kit (PerkinElmer, Turku, Finland),

which is a combination of PCR-based nucleic acid amplification and time-resolved fluorescence

resonance energy transfer (TR-FRET) based detection. EnLite™ simultaneously detects two

targets, TREC and beta-actin, the latter used as an internal control for monitoring specimen

amplification in each test. The assay involves DNA elution from 1.5mm DBS punches,

amplification and hybridization with target-sequence specific probes, and quantifies TREC levels

by measuring probe fluorescence with the Victor Enlite™ fluorometer (PerkinElmer). A full

calibration curve with blanks and three DBS calibrators was run in triplicate on each plate. A

low-TREC control, a no-TREC control, a high-TREC control, and a blank paper disk (containing

no blood), were used as quality controls for each plate. Raw fluorescence counts, measured at

615 nm, 665 nm, and 780 nm, were processed by the EnLite™ workstation software to produce

corrected fluorescence counts. TREC and beta-actin copies/µL were predicted from corrected

fluorescent counts by the EnLite™ workstation software using an unweighted linear regression

model on the ArcSinh transformation of the DBS calibrator copies/µL response.

The values for the number of TREC copies/µL are sensitive to technical effects associated with

the run in which they are obtained; for this reason, we generated technical replicates for the

same individuals in different runs. Because of limited DBS availability, we could not produce

more than one TREC read-out for all individuals and not all individuals with multiple readouts

had the same number of replicated measurements. For this reason, we used a linear mixed-

effects model to test the effect of FIR on the TREC read-out, using the newborn identifier as a

random effect variable to account for replicated measurements across individuals. We used the

logarithm of the corrected fluorescent counts as TREC read-out instead of the predicted number

of copies/µL, as these more accurately reflect the measurements made by the instrument and

its technical biases. In this linear mixed-effects model, the FIR stage was the main explanatory

variable, and aside from the random effect of the newborn, other covariates that entered as

main factors in the model were sex, maximum lymphocyte count during the first postnatal week,

and a batch indicator variable identifying the run of the TREC measurement. This linear mixed-

effects model, as well as a null one without the FIR stage, were fitted to the data using the R

package lme4[68].

A chi-squared test with one degree of freedom between the two models gave a *P* value=0.0432,

thus rejecting the hypothesis of no FIR effect at a significance level α < 0.05. Adjusted and

corrected fluorescent counts on a logarithmic scale (shown in Figure 4a) were obtained after

removing the effect of the previous covariates using a linear model.

RNA extraction, library preparation and sequencing

We used a scalpel to cut a surface of approximately 25 square millimeters from DBS samples in

Guthrie cards, using a different scalpel blade for each sample to avoid contamination. RNA was

then extracted using the illustra™ RNAspin Mini Isolation Kit (GE Healthcare) following the

supplier's protocol and eluting the sample in a final volume of 30 µl. The quality and quantity of

the RNA was assessed with an RNA Pico chip in an Agilent 2100 Bioanalyzer equipment. Next,

globin mRNA and ribosomal RNA were removed using the Globin Zero Gold rRNA Removal Kit

(Illumina), and its performance was checked using with an RNA Pico chip in an Agilent 2100

Bioanalyzer equipment. Libraries were prepared using the NebNext UltraTM II Directional RNA

Library Prep Kit (New England Biolabs), following the specific protocol for rRNA depleted FFPE

RNA and using 12 PCR cycles for library amplification. Libraries were validated and their

concentration was measured with a High Sensitivity DNA chip in an Agilent 2100 Bioanalyzer.

Finally, we prepared a pooled library with a normalized concentration for each sample. The final

concentration of the pool was measured by Real-Time PCR using the NGS Library

Quantification Kit (Takara). We loaded a final concentration of 1.9 pM into a NextSeq-500 High

Output run with 2×75 cycles using 4 sequencing lanes per library. The resulting 21×4=84

paired-end FASTQ files have been deposited to the European Genome-Phenome Archive with

the study identifier EGAS00001003635.

Pre-processing and differential expression analysis of transcriptomics RNA-seq data

We performed quality control on the 84 raw paired-end reads using FastQC and did not detect

artifacts in the sequence data. We did observe, however, that one of the samples, BS13, was

sequenced at about 4 times more depth than the rest of the samples. The reason for this higher

depth was a wrong normalization step during library preparation that led to a higher

concentration of RNA in that sequenced library. To adjust for this bias in sequencing depth we

downsampled uniformly at random the paired-end reads from the library to ¼ of its original

depth.

Raw paired-end reads in FASTQ files, including the downsampled version of BS13, were

aligned to the GRCh38 version of the reference human genome, without alternate locus

scaffolds (GCA_000001405.15) and including human decoy sequences from hs38d1

(GCA_000786075.2), using STAR[69] version 2.6.0c with default parameters except for `--`

`peOverlapNbasesMin 10` and `--sjdbOverhang 74`, producing an initial set of 84 BAM

files. Aligned reads in BAM files were reduced to a table of counts of 25,221 genes by 84

samples using gene annotations from GENCODE v24 and the R/Bioconductor package

GenomeAlignments[70] version 1.16.0, and its function summarizeOverlaps. We used specific

arguments in the call to this function to restrict the count of genic reads to only those that fell

entirely within the exonic regions and aligned to a unique site on the genome, to reflect library

preparation protocols and avoid counting reads without a matching pair or overlapping multiple

features.

The rest of the analysis of transcriptomics data was based on the *edgeR*[71] and *limma*[72]

pipelines, and a few other R packages from the Bioconductor project[73] release version 3.7. A

multidimensional scaling (MDS) plot of the samples showed a nearly perfect agreement

between libraries from the same sample sequenced in each of the 4 different lanes. For this

reason, we merged BAM files with aligned reads from the same sample and built a new table of

counts of 25,221 genes by 21 samples, using the previously employed reduction procedure.

Considering that 6 out of the 21 infants were not diagnosed with MIR or FIR (Supp. Fig. S1),

and following previously established recommendations[74], we filtered out lowly-expressed genes

by discarding those that did not show a minimum reliable level of expression of 10 counts per

million reads of the smallest library size, in less than 6 samples. This filtering step led to a final

table of counts of 11,279 genes by 21 samples.

We used TMM normalization[75] to produce the MDS plot shown in Figure 1c, as well as to obtain

gene-level robust dispersion estimates[76] that helped to select the sex of the infant and surrogate

variables calculated with SVA[77], as covariates in the linear models employed for differential

expression analysis. Finally, we used the limma-voom[78] pipeline with quality weights, combined

with quantile normalization, to conduct the differential expression analysis using linear models.

The design matrix included the FIR status of the infants as a main explanatory variable and sex

and surrogate variables, estimated with SVA, as covariates.

The differential expression analysis across different combined stages of inflammation in MIR

and FIR (Supp. Fig. S6, Supp. Table S4) was based on our replacing the FIR status variable in

the linear model, with numerical values for MIR and FIR. Specifically, we assigned a 0 when

MIR and FIR were not present, 1 when MIR was present but FIR was not, 2 when MIR was

stage III and FIR was stage I and, finally, 3 when MIR was stage III and FIR was stage II or III.

Pre-processing and differential expression analysis of transcriptomics microarray data

We obtained from Nigel Paneth and collaborators at the Michigan State University raw

microarray gene expression data files from paired frozen and unfrozen DBS samples from nine

neonates of the US ELGAN cohort, where two were affected by FIR and seven were not. The

n=18 corresponding samples had been hybridized on a Agilent SurePrint G3 Human Gene

Expression 8x60K Microarray Kit, and their analysis comparing the number of expressed genes

between frozen and unfrozen samples was published by Wei *et al.* (2014)[15]. Batch processing

information was derived from the scanning timestamp stored in raw data files as previously

described[79], creating a batch indicator variable that divided the samples into two different

batches. The cross-classification of infants by FIR status and batch indicator showed no

correlation between the primary outcome of our analysis, FIR status, and sample batch

processing (Supp. Fig. S4). The MDS plot of the data shows the effect of storage temperature

and batch processing, but not sex (Supp. Fig. S4).

We corrected background expression intensities using the "normexp" method in the function

backgroundCorrect from the limma package[80], and used a quantile normalization approach to

obtain comparable gene expression distributions across the samples. We removed probesets

without annotation to Entrez gene identifiers, and probesets annotated to a common Entrez

gene, keeping the one with highest variability measured by interquartile range (IQR). We then

discarded the Agilent positive control probes and retained initially the negative control probes to

later estimate the fraction of expressed genes per sample. This resulted in an expression data

set of 21,952 probesets in one-to-one correspondence with Entrez genes after discarding

negative control probes, by n = 18 samples. We calculated the 95% quantile of the expression

level distribution observed in negative control probes for each sample. As a function of the

percentage of increase over the 95% quantile of negative probe expression (from 100% to

200% increasing by 10%), we found the number of expressed genes per sample separately by

storage condition, which we later used for non-specific filtering based on minimum expression.

Using the software package SVA[77] and the top 1% most variable (IQR) genes, we estimated

one surrogate variable covariate that captures variability unrelated to FIR, storage temperature

and batch processing. For each gene, we defined a linear mixed-effects model where the

expression profile of every gene is a linear function of FIR status (yes/no, main explanatory

variable), storage temperature (frozen/unfrozen), batch processing, the estimated surrogate

variable covariate and the infant as a random effect. We estimated array quality weights to

down-weight samples of worse quality in a graduated way[81] and to avoid discarding them. Using

limma, we fitted this linear mixed-effects model, including the array quality weights, to each

gene expression profile. We then we calculated moderated *t*-statistics, squeezing the genewise

residual variances towards a global trend, for the coefficient estimating the effect between FIR

and non–FIR-affected infants, and their corresponding *P* values for the null hypothesis of no-

differential expression. We selected initially different subsets of DE genes at 5% FDR according

to different thresholds of the non-specific filtering on minimum expression and, within each non-

specific filter, FDR-adjusted *P* values were re-calculated. We found that a 110% cutoff yielded

the highest number of DE genes, and using this non-specific filter we readjusted the raw *P*

values and obtained 33 genes that changed significantly at 5% FDR by 50% or more, between

newborns affected and unaffected by FIR.

Mass spectrometry proteomics

Sample preparation for mass spectrometry: DBS samples were excised and collected in a 2 mL

eppendorf, and extracted as described previously[82]. Briefly, each sample was soaked with 970

µl of a solution consisting of 758 µl of 25 mM ammonium bicarbonate, 113 µl of 10% (w/v)

sodium deoxycholate and 99 µl of 5 mM tris(2-carboxytethyl)phosphine, which was incubated at

60 ºC during 1h in constant agitation. Samples were alkylated with 10 µM iodacetamide (37ºC,

30 min) and quenched with 10 µM dithiothreitol (37ºC, 30 min). Samples were digested with

trypsin (40 ng/µl) overnight at 37 ºC. Digestion was stopped with formic acid, and samples were

vortexed and centrifuged (800 xg, 20 min) prior to supernatant desalting using spin C18

columns.

Mass spectrometry data acquisition: Samples were reconstituted in 0.1% formic acid and were

analyzed on a Orbitrap Fusion Lumos with an EASY-Spray nanosource coupled to a nano-

UPLC system (EASY-nanoLC 1000 liquid chromatograph) equipped with a 50-cm C18 column

(EASY-Spray; 75 μm id, PepMap RSLC C18, 2 mm particles, 45 ºC). Chromatographic

gradients were started at 5% buffer B with a flow rate of 300 nL/min and gradually increased to

22% buffer B in 79 min and to 32% in 11 minutes. After each analysis, the column was washed

for 10 min with 95% buffer B (Buffer A: 0.1% formic acid in water. Buffer B: 0.1% formic acid in

acetonitrile). The mass spectrometer was operated in data-dependent acquisition mode, with full

MS scans over a mass range of m/z 350–1500 with detection in the Orbitrap (120K resolution)

and with auto gain control (AGC) set to 100,000. In each cycle of data-dependent acquisition

analysis, following each survey scan, the most intense ions above a threshold ion count of

10,000 were selected for fragmentation with HCD at normalized collision energy of 28%. The

number of selected precursor ions for fragmentation was determined by the "Top Speed"

acquisition algorithm (max cycle time of 3 seconds), and a dynamic exclusion of 60 s was set.

Fragment ion spectra were acquired in the ion trap with an AGC of 10,000 and a maximum

injection time of 200 ms. One of the samples, BS04, could not be profiled for technical reasons.

All data were acquired with Xcalibur software v4.1.

Mass spectrometry protein expression quantification: The MaxQuant software suite (v1.6.2.6)

was used for peptide identification and label-free protein quantification[83]. The data were

searched against the Uniprot human database with decoy entries (as of April 2018, 42,518

entries). A precursor ion mass tolerance of 4.5 ppm at the MS1 level was used, and up to two

missed cleavages for trypsin were allowed. The fragment ion mass tolerance was set to 0.5 Da.

Oxidation of methionine, protein acetylation at the N-terminal were defined as variable

modification; whereas carbamidomethylation on cysteines was set as a fixed modification. The

minimum number of razor and unique peptides ("min. razor + unique") for a protein group to be

considered as identified was set to 1. Identified peptides and proteins were filtered using a 5%

FDR. The match between runs algorithm was activated with a tolerance of 0.7 min for the

matching time window and 20 min for the alignment time window. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE[84]

partner repository with accession number PXD011626.

Pre-processing and differential expression analysis of proteomics data

From the initial set of 649 quantified protein expression profiles on 20 samples, we discarded 26

whose protein identifier could not be mapped to one of the 25,221 genes profiled by RNA-seq.

From the remaining 624 proteins, we selected 303 with protein quantification values in at least 6

out of the 20 samples. We further discarded 48 proteins whose protein quantification values

were based on one single peptide, leading to a final protein quantification data matrix of 245

proteins by 20 samples. Raw protein quantifications were normalized using the vsn[85,86]

R/Bioconductor package. A complete normalized matrix of protein expression values was

obtained by imputing missing quantifications with a multivariate method[87] based on the

expectation-maximization (EM) algorithm implemented in the function impute.wrapper.MLE from

the R package *imputeLCMD*. Using this complete and normalized matrix of protein expression

values as an input to the edgeR[71] and limma-trend[76] pipelines, we produced the MDS plot in

Figure 1d and conducted a protein differential expression analysis with linear models shown in

Figure 2c. The design matrix included the FIR status of infants as the main explanatory variable

and sex as a covariate.

Functional enrichment analysis with the Gene Ontology database

We performed functional enrichment analysis with the GO database using the R/Bioconductor

package GOstats[88]. More concretely, we used the biological process ontology and the

conditional hypergeometric test to account for dependencies derived from the GO hierarchical

structure. Under this procedure, the GO hierarchy is traversed towards the root, applying a one-

tailed Fisher's exact test between the genes in the GO term and the DE gene set at hand. When

a GO term is considered to be significantly enriched by the DE gene set, its genes are removed

from parent GO terms before testing them. To consider a GO term as significantly enriched we

used a *P* value cutoff of 0.01, a minimum OR of 1.5, a minimum of five and a maximum of 300

genes in the GO term. The gene universe was defined by the entire set of 25,221 profiled genes

by RNA-seq. To report only the most robustly enriched biological processes, the final set of

significantly enriched GO terms was selected by first using a multiple testing correction of FDR

< 10% on the whole set of tested GO terms, then by further selecting GO terms with OR > 1.5

and a minimum number of five genes enriching the GO term.

Acknowledgments

This work was supported by the Spanish MINECO/FEDER (TIN2015-71079-P). The CRG/UPF

Proteomics Unit is part of the Spanish Infrastructure for Omics Technologies (ICTS Omics Tech)

and is a member of Proteored, PRB3, which is supported by grant PT17/0019 of the PE I+D+i 2013-2016, funded by ISCIII and ERDF. We acknowledge support from the Spanish Ministry of Economy and Competitiveness, "Centro de Excelencia Severo Ochoa 2013-2017" (SEV-2012-0208), "Unidad de Excelencia Maria de Maeztu" (MDM-2014-0370), and "Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat de Catalunya" (SGR17-595; SGR17-1020). The authors thank the ELGAN PAD committee, and Nigel Paneth, Sok Kean Khoo, Pete Haak, Madeleine Lenski and Elizabeth Allred, for granting access to the microarray expression and clinical data from the US ELGAN cohort. We would like to thank Aida Andrés, José Aramburu, Sergi Castellano, Alan Leviton and Cristina López-Rodríguez for their critical remarks on various parts of the manuscript.

Contributors

DC, TC and RC conceived the study. DC, NB, AS, JMG, ES, FC, CR, AN and JLM collected the data. DC and RC analyzed the data. DC, TC and RC drafted the manuscript. All authors revised and approved the manuscript.

Competing interests

The authors declare no competing interests.

Ethics approval

The study was approved by the Institutional Review Board from the Hospital Clínic de Barcelona (September 16th, 2016; ref. HCB/2016/0713). The parents of the analyzed infants provided written, informed consent.

Data sharing statement

The transcriptomics and clinical data reported in this paper are available through the European Genome-phenome Archive (EGA) under accession number EGAS00001003635. The mass spectrometry proteomics data are available through the PRIDE repository under accession number PXD011626.

References

1. Khong, T. Y. *et al.* Sampling and Definitions of Placental Lesions: Amsterdam Placental

2  Workshop Group Consensus Statement. *Arch. Pathol. Lab. Med.* **140**, 698–713 (2016).

3 2. Gantert, M. *et al.* Chorioamnionitis: a multiorgan disease of the fetus? *J. Perinatol.* **30**

4  **Suppl**, S21–30 (2010).

5 3. Dammann, O. & Leviton, A. Role of the fetus in perinatal infection and neonatal brain

6  damage. *Curr. Opin. Pediatr.* **12**, 99–104 (2000).

7 4. Costa, D. & Castelo, R. Umbilical cord gene expression reveals the molecular architecture

8  of the fetal inflammatory response in extremely preterm newborns. *Pediatr. Res.* **79**, 473–

9  481 (2016).

10 5. Tilley, S. K. *et al.* Genomic biomarkers of prenatal intrauterine inflammation in umbilical

11  cord tissue predict later life neurological outcomes. *PLoS One* **12**, e0176953 (2017).

12 6. Leviton, A. *et al.* Persistence after birth of systemic inflammation associated with umbilical

13  cord inflammation. *J. Reprod. Immunol.* **90**, 235–243 (2011).

14 7. O'Shea, T. M. *et al.* Inflammation-initiating illnesses, inflammation-related proteins, and

15  cognitive impairment in extremely preterm infants. *Brain Behav. Immun.* **29**, 104–112

16  (2013).

17 8. Dammann, O. & Leviton, A. Intermittent or sustained systemic inflammation and the

18  preterm brain. *Pediatr. Res.* **75**, 376–380 (2014).

19 9. Dammann, O. *et al.* Duration of Systemic Inflammation in the First Postnatal Month Among

20  Infants Born Before the 28th Week of Gestation. *Inflammation* **39**, 672–677 (2016).

21 10. Yanni, D. *et al.* Both antenatal and postnatal inflammation contribute information about the

22  risk of brain damage in extremely preterm newborns. *Pediatr. Res.* **82**, 691–696 (2017).

23 11. Korzeniewski, S. J. *et al.* A 'multi-hit' model of neonatal white matter injury: cumulative

24  contributions of chronic placental inflammation, acute fetal inflammation and postnatal

25  inflammatory events. *J. Perinat. Med.* **42**, 731–743 (2014).

26 12. Olin, A. *et al.* Stereotypic Immune System Development in Newborn Children. *Cell* **174**,

27  1277–1292.e14 (2018).

13. Papile, L. A., Burstein, J., Burstein, R. & Koffler, H. Incidence and evolution of

subependymal and intraventricular hemorrhage: a study of infants with birth weights less

than 1,500 gm. *J. Pediatr.* **92**, 529–534 (1978).

14. Sawyer, S., Krause, J., Guschanski, K., Savolainen, V. & Pääbo, S. Temporal patterns of

nucleotide misincorporations and DNA fragmentation in ancient DNA. *PLoS One* **7**, e34131

(2012).

15. Wei, C. *et al.* Comparison of frozen and unfrozen blood spots for gene expression studies.

J. Pediatr. **164**, 189–191.e1 (2014).

16. Canna, S. W. *et al.* Life-threatening NLRC4-associated hyperinflammation successfully

treated with IL-18 inhibition. *J. Allergy Clin. Immunol.* **139**, 1698–1701 (2017).

17. Condamine, T. *et al.* Lectin-type oxidized LDL receptor-1 distinguishes population of human

polymorphonuclear myeloid-derived suppressor cells in cancer patients. *Sci Immunol* **1**,

(2016).

18. Breuer, K. *et al.* InnateDB: systems biology of innate immunity and beyond--recent updates

and continuing curation. *Nucleic Acids Res.* **41**, D1228–33 (2013).

19. Duncan, J. A. & Canna, S. W. The NLRC4 Inflammasome. *Immunol. Rev.* **281**, 115–123

(2017).

20. Naranbhai, V. *et al.* Genomic modulators of gene expression in human neutrophils. *Nat.*

Commun. **6**, 7545 (2015).

21. Boyman, O., Krieg, C., Homann, D. & Sprent, J. Homeostatic maintenance of T cells and

natural killer cells. *Cell. Mol. Life Sci.* **69**, 1597–1608 (2012).

22. Haines, C. J. *et al.* Human CD4 T cell recent thymic emigrants are identified by protein

tyrosine kinase 7 and have reduced immune function. *J. Exp. Med.* **206**, 275–285 (2009).

23. Lewis, D. B., Haines, C. & Ross, D. Protein tyrosine kinase 7: a novel surface marker for

human recent thymic emigrants with potential clinical utility. *J. Perinatol.* **31 Suppl 1**, S72–

81 (2011).

24. Pérès, M., Montfort, A., Andrieu-Abadie, N., Colacios, C. & Ségui, B. S1P: the elixir of life

for naive T cells. *Cell. Mol. Immunol.* (2017). doi:10.1038/cmi.2017.110

25. Glavina-Durdov, M. *et al.* The grade of acute thymus involution in neonates correlates with

the duration of acute illness and with the percentage of lymphocytes in peripheral blood

smear. Pathological study. *Biol. Neonate* **83**, 229–234 (2003).

26. Schwarz, J. *et al.* Granulocytic myeloid-derived suppressor cells (GR-MDSC) accumulate in

cord blood of preterm infants and remain elevated during the neonatal period. *Clin. Exp.*

Immunol. **191**, 328–337 (2018).

27. Gervassi, A. *et al.* Myeloid derived suppressor cells are present at high frequency in

neonates and suppress in vitro T cell responses. *PLoS One* **9**, e107816 (2014).

28. Leiber, A. *et al.* Neonatal myeloid derived suppressor cells show reduced apoptosis and

immunosuppressive activity upon infection with Escherichia coli. *Eur. J. Immunol.* **47**,

1009–1021 (2017).

29. Schieber, M. & Chandel, N. S. ROS function in redox signaling and oxidative stress. *Curr.*

Biol. **24**, R453–62 (2014).

30. Veglia, F., Perego, M. & Gabrilovich, D. Myeloid-derived suppressor cells coming of age.

Nat. Immunol. **19**, 108–119 (2018).

31. Zhou, J., Nefedova, Y., Lei, A. & Gabrilovich, D. Neutrophils and PMN-MDSC: Their

biological role and interaction with stromal cells. *Semin. Immunol.* **35**, 19–28 (2018).

32. Ost, M. *et al.* Myeloid-Derived Suppressor Cells in Bacterial Infections. *Front. Cell. Infect.*

Microbiol. **6**, (2016).

33. Tamadaho, R. S. E., Hoerauf, A. & Layland, L. E. Immunomodulatory effects of myeloid-

derived suppressor cells in diseases: Role in cancer and infections. *Immunobiology* **223**,

432–442 (2018).

34. Moskovitz, J. Roles of methionine suldfoxide reductases in antioxidant defense, protein

regulation and survival. *Curr. Pharm. Des.* **11**, 1451–1457 (2005).

35. Wynn, J. L. *et al.* Targeting IL-17A attenuates neonatal sepsis mortality induced by IL-18. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E2627–35 (2016).

36. Liang, J. *et al.* Novel NLRC4 Mutation Causes a Syndrome of Perinatal Autoinflammation With Hemophagocytic Lymphohistiocytosis, Hepatosplenomegaly, Fetal Thrombotic Vasculopathy, and Congenital Anemia and Ascites. *Pediatr. Dev. Pathol.* **20**, 498–505 (2017).

37. Denes, A. *et al.* AIM2 and NLRC4 inflammasomes contribute with ASC to acute brain injury independently of NLRP3. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 4050–4055 (2015).

38. Ravichandran, S. *et al.* Higher daily doses of caffeine lowered the incidence of moderate to severe neurodevelopmental disabilities in very low birth weight infants. *Acta Paediatr.* **108**, 430–435 (2019).

39. Furman, D. *et al.* Expression of specific inflammasome gene modules stratifies older individuals into two extreme clinical and immunological states. *Nat. Med.* **23**, 174–184 (2017).

40. Hanna, N. *et al.* Mechanisms underlying reduced apoptosis in neonatal neutrophils. *Pediatr. Res.* **57**, 56–62 (2005).

41. Allgaier, B., Shi, M., Luo, D. & Koenig, J. M. Spontaneous and Fas-mediated apoptosis are diminished in umbilical cord blood neutrophils compared with adult neutrophils. *J. Leukoc. Biol.* **64**, 331–336 (1998).

42. Nguyen, C. N., Schnulle, P. M., Chegini, N., Luo, X. & Koenig, J. M. Neonatal neutrophils with prolonged survival secrete mediators associated with chronic inflammation. *Neonatology* **98**, 341–347 (2010).

43. Atallah, M. *et al.* Constitutive Neutrophil Apoptosis: Regulation by Cell Concentration via S100 A8/9 and the MEK – ERK Pathway. *PLoS One* **7**, e29333 (2012).

44. Hirata, J.-I. *et al.* A role for IL-18 in human neutrophil apoptosis. *Shock* **30**, 628–633 (2008).

45. Chen, K. W. *et al.* The neutrophil NLRC4 inflammasome selectively promotes IL-1β

maturation without pyroptosis during acute Salmonella challenge. *Cell Rep.* **8**, 570–582

(2014).

46. Kessel, C., Holzinger, D. & Foell, D. Phagocyte-derived S100 proteins in autoinflammation:

putative role in pathogenesis and usefulness as biomarkers. *Clin. Immunol.* **147**, 229–241

(2013).

47. Alder, M. N., Opoka, A. M., Lahni, P., Hildeman, D. A. & Wong, H. R. Olfactomedin-4 Is a

Candidate Marker for a Pathogenic Neutrophil Subset in Septic Shock. *Crit. Care Med.* **45**,

e426–e432 (2017).

48. Solan, P. D. *et al.* A novel role for matrix metalloproteinase-8 in sepsis. *Crit. Care Med.* **40**,

379–387 (2012).

49. O'Hare, F. M. *et al.* Persistent systemic monocyte and neutrophil activation in neonatal

encephalopathy. *J. Matern. Fetal. Neonatal Med.* **29**, 582–589 (2016).

50. Molloy, E. J. *et al.* Neonatal encephalopathy is associated with altered perinatal systemic

neutrophil apoptosis. *Am. J. Perinatol.* **24**, 525–530 (2007).

51. Di Naro, E. *et al.* Fetal thymic involution: a sonographic marker of the fetal inflammatory

response syndrome. *Am. J. Obstet. Gynecol.* **194**, 153–159 (2006).

52. Scheible, K. M. *et al.* T cell developmental arrest in former premature infants increases risk

of respiratory morbidity later in infancy. *JCI Insight* **3**, (2018).

53. Kuban, J. D., Allred, E. N. & Leviton, A. Thymus involution and cerebral white matter

damage in extremely low gestational age neonates. *Biol. Neonate* **90**, 252–257 (2006).

54. He, Y.-M. *et al.* Transitory presence of myeloid-derived suppressor cells in neonates is

critical for control of inflammation. *Nat. Med.* **24**, 224–231 (2018).

55. Chemnitz, J. M. *et al.* Prostaglandin E2 impairs CD4+ T cell activation by inhibition of lck:

implications in Hodgkin's lymphoma. *Cancer Res.* **66**, 1114–1122 (2006).

56. Ozsurekci, Y. & Aykac, K. Oxidative Stress Related Diseases in Newborns. *Oxid. Med.*

Cell. Longev. **2016**, 1–9 (2016).

57. Tindell, R. & Tipple, T. Selenium: implications for outcomes in extremely preterm infants. *J. Perinatol.* **38**, 197–202 (2018).

58. Fong, O. N. *et al.* Expression profile of cord blood neutrophils and dysregulation of HSPA1A and OLR1 upon challenge by bacterial peptidoglycan. *J. Leukoc. Biol.* **95**, 169–178 (2014).

59. Kruzel, M. L., Zimecki, M. & Actor, J. K. Lactoferrin in a Context of Inflammation-Induced Pathology. *Front. Immunol.* **8**, (2017).

60. Song, J. & Kim, O. Y. Perspectives in Lipocalin-2: Emerging Biomarker for Medical Diagnosis and Prognosis for Alzheimer's Disease. *Clin. Nutr. Res.* **7**, 1–10 (2018).

61. Yellowhair, T. R. *et al.* Preclinical chorioamnionitis dysregulates CXCL1/CXCR2 signaling throughout the placental-fetal-brain axis. *Exp. Neurol.* **301**, 110–119 (2018).

62. Denstaedt, S. J. *et al.* S100A8/A9 Drives Neuroinflammatory Priming and Protects against Anxiety-like Behavior after Sepsis. *J. Immunol.* **200**, 3188–3200 (2018).

63. Silk, E., Zhao, H., Weng, H. & Ma, D. The role of extracellular histone in organ injury. *Cell Death Dis.* **8**, e2812 (2017).

64. Gilthorpe, J. D. *et al.* Extracellular histone H1 is neurotoxic and drives a pro-inflammatory response in microglia. *F1000Res.* **2**, 148 (2013).

65. Hagberg, H. *et al.* The role of inflammation in perinatal brain injury. *Nat. Rev. Neurol.* **11**, 192–208 (2015).

66. Gibbs, R. S. & Duff, P. Progress in pathogenesis and management of clinical intraamniotic infection. *Am. J. Obstet. Gynecol.* **164**, 1317–1326 (1991).

67. Redline, R. W. *et al.* Amniotic infection syndrome: nosology and reproducibility of placental reaction patterns. *Pediatr. Dev. Pathol.* **6**, 435–448 (2003).

68. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software, Articles* **67**, 1–48 (2015).

69. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

70. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS Comput. Biol.* **9**, e1003118 (2013).

71. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

72. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

73. Huber, W. *et al.* Orchestrating high-throughput genomic analysis with Bioconductor. *Nat. Methods* **12**, 115–121 (2015).

74. Chen, Y., Lun, A. T. L. & Smyth, G. K. From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Res.* **5**, 1438 (2016).

75. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

76. Phipson, B., Lee, S., Majewski, I. J., Alexander, W. S. & Smyth, G. K. Robust hyperparameter estimation protects against hypervariable genes and improves power to detect differential expression. *Ann. Appl. Stat.* **10**, 946–963 (2016).

77. Leek, J. T. & Storey, J. D. Capturing heterogeneity in gene expression studies by surrogate variable analysis. *PLoS Genet.* **3**, 1724–1735 (2007).

78. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).

79. Leek, J. T. *et al.* Tackling the widespread and critical impact of batch effects in high-throughput data. *Nat. Rev. Genet.* **11**, 733–739 (2010).

80. Silver, J. D., Ritchie, M. E. & Smyth, G. K. Microarray background correction: maximum likelihood estimation for the normal-exponential convolution. *Biostatistics* **10**, 352–363 (2009).

81. Ritchie, M. E. *et al.* Empirical array quality weights in the analysis of microarray data. *BMC Bioinformatics* **7**, 261 (2006).

82. Chambers, A. G., Percy, A. J., Yang, J. & Borchers, C. H. Multiple Reaction Monitoring Enables Precise Quantification of 97 Proteins in Dried Blood Spots. *Mol. Cell. Proteomics* **14**, 3094–3104 (2015).

83. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* **26**, 1367–1372 (2008).

84. Vizcaíno, J. A. *et al.* The PRoteomics IDEntifications (PRIDE) database and associated tools: status in 2013. *Nucleic Acids Res.* **41**, D1063–9 (2013).

85. Huber, W., von Heydebreck, A., Sültmann, H., Poustka, A. & Vingron, M. Variance stabilization applied to microarray data calibration and to the quantification of differential expression. *Bioinformatics* **18 Suppl 1**, S96–104 (2002).

86. Välikangas, T., Suomi, T. & Elo, L. L. A systematic evaluation of normalization methods in quantitative label-free proteomics. *Brief. Bioinform.* **19**, 1–11 (2018).

87. Schafer, J. L. *Analysis of Incomplete Multivariate Data*. (CRC Press, 1997).

88. Falcon, S. & Gentleman, R. Using GOstats to test gene lists for GO term association. *Bioinformatics* **23**, 257–258 (2007).