Mediterranean Diet improves thrombosis biomarkers in high cardiovascular risk individuals: a randomized controlled trial

Short title: Mediterranean Diet and thrombosis mechanisms

Álvaro Hernáez,[1,2,*] Olga Castañer,[2,3] Anna Tresserra-Rimbau,[2,4] Xavier Pintó,[2,5] Montserrat Fitó,[2,3] Rosa Casas,[1,2] Miguel Ángel Martínez-González,[2,6,7] Dolores Corella,[2,8] Jordi Salas-Salvadó,[2,4] José Lapetra,[2,9] Enrique Gómez-Gracia,[2,10] Fernando Arós,[2,11] Miquel Fiol,[2,12] Lluis Serra-Majem,[2,13] Emilio Ros,[2,14], and Ramón Estruch[1,2,15]

[1]Cardiovascular Risk, Nutrition and Aging Research Unit, August Pi i Sunyer Biomedical Research Institute (IDIBAPS), Barcelona, Spain, [2]CIBER of Pathophysiology of Obesity and Nutrition (CIBEROBN), Instituto de Salud Carlos III, Madrid, Spain; [3]Cardiovascular Risk and Nutrition Research Group, Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain; [4]Universitat Rovira i Virgili, Departament de Bioquímica i Biotecnologia, Unitat de Nutrició Humana. Hospital Universitari San Joan de Reus, Institut d'Investigació Pere Virgili (IISPV). Reus, Spain; [5]Lipids and Vascular Risk Unit, Internal Medicine, Hospital Universitario de Bellvitge, Hospitalet de Llobregat, Barcelona, Spain; [6]Universidad de Navarra, Department of Preventive Medicine and Public Health, Pamplona, Spain; [7]Department of Nutrition, Harvard TH Chan School of Public Health, Boston, USA; [8]Department of Preventive Medicine, Universidad de Valencia, Valencia, Spain; [9]Department of Family Medicine, Research Unit, Distrito Sanitario Atención Primaria Sevilla, Sevilla, Spain; [10]Department of Preventive Medicine and Public Health, Universidad de Málaga, Málaga, Spain; [11]Department of Cardiology, Hospital Universitario de Álava, Vitoria, Spain; [12]Balearic Islands Health Research Institute, Hospital Son Espases, Palma de Mallorca, Spain; [13]Department of Clinical Sciences, Universidad de Las Palmas de Gran Canaria, Las Palmas, Spain; [14]Lipid Clinic, Endocrinology and Nutrition Service, IDIBAPS, Hospital Clínic, Barcelona, Spain; and [15]Internal Medicine Service, Hospital Clínic, Barcelona, Spain.

Word count (text): 3,430
Word count (abstract): 250
Figure/table count: 1 figure, 2 tables
Reference count: 36

KEY POINTS

- Following a Mediterranean diet improves thrombosis-related biomarkers in high cardiovascular risk individuals

ABSTRACT

Following a Traditional Mediterranean diet (TMD) decreases cardiovascular risk. However, its anti-thrombotic effects are unexplored. We aimed to assess whether this dietary pattern improved thrombosis-related biomarkers in high cardiovascular risk subjects. In 358 random PREDIMED Study volunteers (*Prevención con Dieta Mediterránea*), we assessed: 1) 1-year effects of a TMD, enriched in virgin olive oil (TMD-VOO; N=120) or nuts (TMD-Nuts; N=119) versus a low-fat diet (N=119), on thrombosis biomarkers; 2) the differences between individuals with greater TMD compliance increments (≥3 adherence score points) and those with compliance decrements; and 3) associations between 1-year changes in key TMD food groups and thrombosis signals. TMD-VOO and TMD-Nuts attenuated pro-thrombotic status (-5.1% –P=0.045– and -5.8% –P=0.025–, respectively). TMD-VOO decreased HDL-bound α_1-antitrypsin levels (HDL pro-inflammatory/pro-thrombotic properties, -6.3%, P=0.034) and increased HDL platelet activating factor-acetylhydrolase activity (+7.5%, P=0.044). TMD-Nuts diminished non-esterified fatty acid levels (-10.3%, P=0.022). Individuals with greater TMD adherence increments presented decreases in their pro-thrombotic status (-9.5%, P=0.032), fibrinogen levels (-9.1%, P=0.030), and non-esterified fatty acid concentrations (-18.1%, P=0.011). Only the low-fat diet was associated with increased platelet factor-4 concentrations (P=0.012) and thrombin activation (higher prothrombin factor$_{1+2}$ levels, P=0.003) versus baseline. Finally, in the secondary analyses, nut intake increases were associated with antithrombin increments, fruit and vegetable consumption increases were linked to decreases of thrombin activation, and processed meat intake decrements were related to increases in antithrombin and nitrite + nitrate levels (P<0.05). Summarizing, TMD and some of its typical food items were associated with improved thrombosis biomarkers in high cardiovascular risk individuals.

INTRODUCTION

Following a traditional Mediterranean Diet (TMD) prevents the development of cardiovascular outcomes[1,2]. As observed in the PREDIMED Study (*Prevención con Dieta Mediterránea)*, TMD effects on cardiovascular protection may be mediated by improvements in risk factors related to blood pressure, glucose metabolism, oxidative stress, and low-grade inflammation[3]. However, very scarce evidence related to TMD effects on other mechanisms such as thrombosis is available.

Thrombosis consists of the formation of a clot to avoid blood loss in the case of a vessel disruption. However, its excessive activation is also linked to an increased risk of atherosclerotic outcomes[4]. Thrombus generation is based on the occurrence of four phenomena (apparition of an endothelial lesion, platelet aggregation, activation of the coagulation cascade, and fibrinolysis) which, in turn, are constituted by a chain of intertwined biochemical responses[4]. Excessive levels of biomarkers related to these responses (or defective, in the case of fibrinolysis) are linked to an increased risk of suffering atherosclerotic events. This association has been particularly reported for plasma biomarkers of endothelial damage (von Willebrand factor[5]), platelet aggregation (P-selectin[6], platelet factor-4[7]), activation of coagulation (fibrinogen[8], thrombin formation – directly proportional to the levels of prothrombin fragment 1+2–[9], antithrombin[10]), and fibrinolysis (plasminogen activator inhibitor-1 –PAI1–[11], D-dimer[12]). Other substances (such as non-esterified fatty acids and nitrite + nitrate levels) have been shown to be involved in the development of cardiovascular diseases through their capacity to modulate endothelial stress and platelet activation[13,14]. Finally, high-density lipoproteins (HDLs) have also been attributed antithrombotic properties[15]. On the one hand, HDLs are deeply involved in the regulation of inflammatory responses, which have been shown to promote thrombosis[8]. On the other hand, HDLs carry enzymes able to metabolize pro-thrombotic

signals such as the platelet activating factor[16] (the platelet activating factor

acetylhydrolase, PAF-AH).

Very few interventions with healthy dietary patterns have been associated with changes in

thrombosis biomarkers. Following a TMD has only been associated with a decrease in P-

selectin levels in high cardiovascular risk subjects in the context of the PREDIMED Trial[17].

Beyond this study, a cross-sectional study reported an inverse association between

adherence to a Mediterranean-like diet and platelet levels[18]. Finally, other small-scale

prospective analysis with 21 young, healthy male volunteers assessed the impact of a

TMD-like diet for 90 days on thrombosis biomarkers, and reported that this dietary pattern

was associated with lower fibrinogen levels and a moderation of the coagulation

response[19]. However, no extensive analysis of the long-term TMD effects on thrombosis

mechanisms has been performed to date in a large-scale intervention trial.

Our main aim was to assess whether a 1-year intervention with TMD improved a

comprehensive set of endothelial stress, platelet, coagulation, and fibrinolysis-related

biomarkers in high cardiovascular risk subjects. Our secondary aim was to determine

whether there were associations between 1-year changes in the consumption of key food

group whose intake was modified in the TMD intervention (virgin olive oil, nuts, fruits and

vegetables, legumes, fish, and processed meat) and thrombosis biomarkers.

MATERIALS AND METHODS

Study protocol and population

Study subjects belonged to the PREDIMED Study. It was a large-scale, parallel,

multicenter, randomized controlled trial assessing the long-term effects of following a TMD

on the prevention of primary cardiovascular disease in a population at high cardiovascular

risk[1]. Volunteers were free of cardiovascular disease at baseline but presented type 2 diabetes mellitus or at least three of the following risk factors: high low-density lipoprotein cholesterol levels, low HDL cholesterol concentrations, hypertension, overweight/obesity, current smoking, or family history of premature heart disease. Individuals were randomly assigned to follow one of the three following intervention groups (in a 1:1:1 ratio): 1) a TMD supplemented with virgin olive oil (TMD-VOO); 2) a TMD supplemented with mixed nuts (TMD-Nuts); and 3) a low-fat control diet. Further details of inclusion and exclusion criteria and the dietary intervention are available in **Supplemental Methods**.

We selected a random subsample of 358 subjects (4.8% of the total PREDIMED population, belonging to Hospital Clinic and IMIM study sites, with biological samples at baseline and after 1 year of dietary intervention) for the present analyses. 120 volunteers were allocated to the TMD-VOO intervention group, 119 to the TMD-Nuts one, and 119 to the low-fat diet. In these individuals, age, sex and educational level data were gathered at baseline. At baseline and after 1 year of intervention, trained clinical personnel collected the following information: 1) body mass index, blood pressure, smoking, and medication use; 2) physical activity levels (using a validated Minnesota Leisure-Time Physical Activity Questionnaire; 3) adherence to a Mediterranean diet (the 14-item Mediterranean Diet adherence score[20], used as a continuous variable); and 4) intake of 137 food items in the previous year (using a validated food frequency questionnaire). From the information in the food frequency questionnaires, we calculated the intakes of key TMD food groups (**Supplemental Methods**). In addition, trained nurses collected plasma and serum samples, and stored at -80ºC upon use and to quantify the levels of fasting glucose, total and HDL cholesterol, and triglycerides by automatized methods[1,21].

The trial protocol was approved by local institutional ethic committees, registered under the International Standard Randomized Controlled Trial Number ISRCTN35739639 (http://www.isrctn.com/ISRCTN35739639), and thoroughly described in previous

publications[1,21]. All participants provided written informed consent before joining the study. A.H. analyzed the data of the present work and all authors had access to primary data results. Finally, the CONSORT checklist regarding our study is available in **Supplemental Table 1**.

Thrombosis biomarkers

We determined thrombosis biomarkers in biological samples collected at baseline and after 1 year of intervention. We used ELISA kits to quantify the levels of: fibrinogen (*Human Fibrinogen SimpleStep ELISA kit*, ref.: ab208036, abcam, UK), plasminogen activator inhibitor-1 (PAI1; *Human PAI1 SimpleStep ELISA kit (SERPINE1)*, ref.: ab184863, abcam, UK), platelet factor-4 (*PF4 (CXCL4) Human SimpleStep ELISA kit*, ref.:ab189573, abcam, UK), prothrombin fragment 1+2 (*Human F1+2 (prothrombin fragment 1+2) ELISA kit*, ref.: E-EL-H1793, Elabscience, USA), and von Willebrand factor (*Human von Willebrand factor SimpleStep ELISA kit*, ref.: ab223864, abcam, UK). We assessed the concentrations of antithrombin (*Antithrombin III*, ref.: SR-1102016, Spinreact, Spain) and D-dimer (*D-Dimer*, ref.: SR-1709231, Spinreact, Spain) by immunoturbidimetry, and of non-esterified fatty acids (*Non-Esterified Fatty Acids*, ref.: FA115, Randox, Spain) using a colorimetric technique in an ABX-Pentra 400 autoanalyzer (Horiba-ABX, France). We quantified nitrates + nitrites by means of a colorimetric kit (*Nitrate/Nitrite Colorimetric Assay Kit*, ref.: CAY-780001, Cayman Chemical, USA). All the previous analyses were performed in citrate plasma samples. In parallel, in apolipoprotein B-depleted plasma samples (specimen in which all lipoproteins but high-density lipoproteins –HDLs– were eliminated, prepared from citrate plasma samples[22]), we determined the levels of total cholesterol by colorimetry (*Cholesterol Enzymatic-Colorimetry*, ref.: SR-41022, Spinreact, Spain) and of α1-antitrypsin (*a-1-antitrypsin*, ref.: SR-1102054, Spinreact, Spain) by immunoturbidimetry in an ABX-Pentra 400 autoanalyzer (Horiba-ABX, France). With this

data, we calculated the α1-antitrypsin/cholesterol ratio. We also determined in these samples the activity of the PAF-AH enzyme by a colorimetric kit (*PAF Acetylhydrolase Activity Assay Kit*, ref.: K765-1000, Cayman Chemical, USA). An extended explanation of the quality control, inter-assay coefficients of variation, and the number of missing values of each determination is available in **Supplemental Methods** and **Supplemental Table 2**.

Sample size

Since the only thrombosis-related biomarker assessed in the PREDIMED Study was P-selectin, we assumed that its variability would be similar to that in our variables. Thus, a sample size of 119 individuals per group (total sample size: *N*=357) allowed ≥80% power to detect a 10% change relative to the mean baseline value of P-selectin levels (7.65 ng/mL) between post-and pre-intervention values, and a 14% change relative to the mean baseline value (10.8 ng/mL) in the differences among the three diets, considering a 2-sided type I error of 0.05, a loss rate of 5%, and the standard deviation of the differences in P-selectin values in this intervention (SD=28.9 ng/mL)[17].

Statistical analyses

We first checked the distribution of continuous variables in normality plots and by the Shapiro-Wilk test. We assessed whether there were differences in baseline values among the three intervention groups with one-way ANOVA tests for normally distributed continuous variables, Kruskal-Wallis tests for non-normally distributed continuous ones, and chi-squared tests (or exact Fisher tests when expected frequencies were <5) for categorical parameters.

We first calculated a continuous score reflecting the general activation of the thrombosis mechanisms. In biomarkers where high concentrations were associated with a more pro-thrombotic state (von Willebrand factor, platelet factor-4, fibrinogen, prothrombin fragment

1+2, D-dimer, PAI-1, non-esterified fatty acids, and α_1-antitrypsin in HDL), we assigned values from 1 to 5 to the volunteers located in increasing quintiles. On the contrary, in biomarkers where high levels were linked to a less pro-thrombotic state (antithrombin, nitrites+nitrates, and PAF-AH activity in HDL), we assigned values from 5 to 1 to the volunteers located in increasing quintiles. For each biomarker, the same cut points were used at baseline and after 1 year of intervention. By summing these scores, we obtained a general pro-thrombotic score ranging from 11 (all biomarkers located in the least pro-thrombotic quintile) to 55 (all biomarkers located in the most pro-thrombotic quintile), in which higher scores indicated a higher pro-thrombotic state. These scores were only calculated in those volunteers with no missing values in any of the thrombosis biomarker variables at both time points (*N*=299).

We analyzed differences between pre- and post-intervention values in every intervention by paired t tests in normally distributed variables and Wilcoxon signed rank tests in non-normally distributed parameters. We checked whether there were differences in the 1-year changes in thrombosis biomarkers (calculated by subtracting baseline to post-intervention values) in the TMD interventions relative to the low-fat control diet by multivariate linear regressions (intention-to-treat analyses). We also compared with these regression models the differences in the 1-year changes in the volunteers with greater increases in TMD adherence (individuals incrementing their compliance in ≥3 points of the 14-item Mediterranean Diet score, *N*=66) relative to those who decreased their adherence in relation to baseline values (high versus low compliance analyses). Finally, in a secondary exploratory approach, we studied whether there were associations between 1-year changes in the intake of food groups whose consumption was significantly modified in the TMD interventions relative to the low-fat diet (virgin olive oil, nuts, fruits and vegetables, legumes, fish, and processed meat; see the "*Study population*" paragraph in the Results section) and thrombosis biomarkers by additional multivariate linear regression analyses.

To promote the interpretability of data, results were presented as percentage changes relative to baseline values (to calculate them, we divided the adjusted difference coefficients obtained in the regression models by the baseline value of the variable in each of the groups). Models were adjusted for: age, sex, study site, educational level (primary education or lower/secondary or equivalent/higher education), baseline value of the variable, fasting glucose levels, use of antidiabetic drugs (yes/no), cholesterol concentrations, triglyceride values (log-transformed), use of lipid-lowering drugs (yes/no), systolic blood pressure, use of antihypertensive drugs (yes/no), use of antithrombotic drugs (yes/no), tobacco use (never/former/actual smoker), body mass index, leisure-time physical activity (log-transformed), and two propensity scores to correct for the theoretical deviations in the randomization process (calculated from 30 baseline variables)[1]. High versus low compliance and food group intake analyses were additionally adjusted for the TMD adherence score or the intake of each particular food group at baseline, respectively, and for the PREDIMED intervention group. Models were plotted using the "lme" package in R Software[23].

We considered any two-sided *P*-value<0.05 as significant and performed statistical analyses with R software version 3.5.0[24].

Data sharing statement

Considering that we do not have the explicit written consent of the study volunteers to yield their deidentified data at the conclusion of the study, individual participant data cannot be shared. The study protocol is available in the main publication of the study[1] and in the study website (http://www.predimed.es/uploads/8/0/5/1/8051451/_1estr_protocol_olf.pdf), and a summary of the dietary intervention is also available in **Supplemental Methods**.

RESULTS

Study population and dietary intervention

We found no significant differences in clinical variables at baseline among the individuals

assigned to the three intervention groups (**Table 1**). Regarding the non-included

PREDIMED Study subjects, there were 6.2% more women and 7.4% less antithrombotic

drug users among our volunteers and they presented higher glucose (6 mg/dL), total

cholesterol (5 mg/dL), and leisure-time physical activity levels (22 METs·min/d) ($P<0.05$)

(**Supplemental Table 3**). Physical activity did not change throughout the study ($P>0.05$).

Volunteers' compliance to the dietary intervention was correct according to the

Mediterranean diet adherence score: we observed a 1.10 and 1.35-point increment in

score values after 1 year of intervention in the TMD-VOO and TMD-Nuts interventions,

respectively ($P<0.001$, relative to the low-fat diet) (**Supplemental Table 4**). Volunteers

allocated in the TMD-VOO intervention group increased their compliance due to

increments in the consumption of virgin olive oil (+3.5 servings/day), fruits and vegetables

(+0.8 servings/d), legumes (+0.2 servings/day), and fish (+0.2 servings/d), whilst those in

the TMD-Nuts group incremented their adherence due to increases in the intake of nuts

(+1.2 servings/d), legumes (+0.1 servings/d –only marginally–), and fatty fish (+0.2

servings/d), and decreases in the consumption of processed meat (-0.2 servings/d)

($P<0.05$, relative to the low-fat diet). Volunteers with greater compliance increments

presented a 4.63 point increase in the TMD adherence score due to increases in the

consumption of virgin olive oil (+1.8 servings/d), nuts (+0.6 servings/d), fruits and

vegetables (+2.4 servings/d), legumes (+0.2 servings/d), and fish (+0.6 servings/d), and

decreases in red meat intake (-0.3 servings/d) ($P<0.05$, in relation to the volunteers with

decrements in TMD adherence). Finally, low-fat diet compliance was observed as a

reduction in the consumption of fats, based on a decrease of the intake of mixed nuts and

the substitution of red for white meat (red meat consumption decreased while meat did

not) and the substitution of fatty for non-fatty fish (total fish intake increased while fatty fish

did not) (P<0.05, relative to baseline).

TMD and thrombosis biomarkers

The thrombosis score (a measurement of the pro-thrombotic status) was attenuated by

both TMD-VOO and TMD-Nuts interventions (-5.1% –P=0.045– and -5.8% –P=0.025–,

respectively, relative to the low-fat diet) (**Figure 1A**). TMD-VOO decreased the levels of

α_1-antitrypsin (-6.3%, P=0.034, **Figure 1K**) and increased the activity of the platelet

activating factor-acetylhydrolase in HDLs (+7.5%, P=0.044, **Figure 1L**) in relation to the

control intervention. TMD-Nuts particularly diminished the concentrations of non-esterified

fatty acids versus the low-fat diet (-10.3%, P=0.022, **Figure 1J**). Volunteers with greater

TMD adherence increments decreased their overall pro-thrombotic status (-9.5%,

P=0.032, **Figure 1A**), fibrinogen levels (-9.1%, P=0.030, **Figure 1D**), and non-esterified

fatty acid concentrations (-18.1%, P=0.011, **Figure 1J**) relative to those who decreased

their TMD adherence. Finally, when assessing post- vs. pre-intervention changes, only the

low-fat diet was associated with increases in platelet factor-4 concentrations (P=0.012)

and thrombin activation (higher prothrombin factor$_{1+2}$ levels, P=0.003) (**Table 2**). Exact

coefficients are available in **Supplemental Table 5** (expressed as percentage changes

relative to baseline values) and **Supplemental Table 6** (expressed as linear differences).

In the secondary exploratory approach regarding food group intake: 1) increases in nut

intake were associated with increments in antithrombin levels; 2) increases in the

consumption of fruit and vegetables were linked to decreases in thrombin activation (lower

prothrombin factor$_{1+2}$ levels); and 3) decrements in the intake of processed meats were

related to increases in antithrombin and nitrite + nitrate levels (P<0.05). Exact coefficients

for these analyses are available in **Supplemental Table 7** (expressed as percentage

changes relative to baseline values) and **Supplemental Table 8** (expressed as linear

differences).

DISCUSSION

Our results show that 1 year of intervention with TMD improves pro-thrombotic state in

high cardiovascular risk individuals. To the best of our knowledge, this trial is the largest

(N=358) and more comprehensive study on the effects of a healthy dietary pattern on the

molecular mechanisms related to thrombosis.

Thrombosis is deeply affected by inflammation and oxidative stress[25,26]. TMD is a lifestyle

intervention known to improve these last two risk factors[3]. Our data indicate that following

this dietary pattern decreased the overall pro-thrombotic response of high cardiovascular

risk subjects (around 5-6% in the subjects in the TMD-VOO and TMD-Nuts groups, and up

to 10% in those with greater increases in TMD adherence) and we hypothesize that this

effect could be partially explained by a decrease in oxidative stress and low-grade

inflammation. On the one hand, following a TMD has been associated in our study with

beneficial changes in thrombosis biomarkers linked as well to low-grade inflammation such

as fibrinogen and α_1-antitrypsin levels in HDL[8,27]. On the other hand, some particular

dietary modifications of the TMD that have been associated with beneficial changes in

thrombosis biomarkers in our data (increases in the consumption of nuts, fruits and

vegetables; decreases in the intake of processed meat) have been broadly related to

antioxidant and anti-inflammatory effects in previous human studies[28–31].

HDLs have been shown to be able to induce beneficial effects on some hemostatic

mechanisms[15]. In this regard, the TMD-VOO intervention led to a particular decrease in α_1-

antitrypsin levels and an increase in PAF-AH activity in HDL particles, two potential HDL

antithrombotic properties. It is not the first time that we report that the TMD-VOO

intervention promotes HDL functionality in the PREDIMED Study[22]. However, our data

extend this protection to anti-inflammatory and anti-thrombotic HDL functions. Antioxidant

protection of typical TMD food items on HDL components could contribute to explaining an

improved activity of the lipoprotein protective enzymes (due to a greater preservation of

their non-oxidized structure)[32]. In addition, the capacity of this dietary pattern to decrease

the levels of inflammation biomarkers in circulation may be related to a decrement in the

concentrations of pro-inflammatory molecules bound to HDL particles[33].

We observed a decrease in non-esterified fatty acid levels after 1 year of TMD-Nuts

intervention (also present in subjects with greater TMD adherence increments). Non-

esterified fatty acids promote endothelial stress and platelet aggregation and, therefore,

are considered an emergent pro-thrombotic risk factor in some cardiovascular risk states

(e.g., obesity, hypertriglyceridemia)[13]. The capacity of polyunsaturated fatty acids (whose

intake was particularly increased in the TMD-Nuts group an among individuals with greater

TMD compliance) to bind to GPR120 receptors in adipose tissue in order to decrease

lipolysis could be an explanation for this effect[34,35]. In addition, TMD is a fiber-rich dietary

pattern, whose intestinal metabolism leads to blood peaks of short-chain fatty acids

(known to be able to bind other GPR family receptors with a similar anti-lipolytic effect[36]).

Thus, a synergistic effect between these two nutrients could be expected.

Our study has several strengths. First, it presents a large sample size (*N*=358). Second, it

is based on a randomized design with an active comparator (low-fat control diet) and a

long-term duration (1 year). Last, it is the first analysis of the effects of a dietary pattern on

a comprehensive, hypothesis-driven set of thrombosis-related biomarkers, performed after

a strict laboratory quality control. However, it also presents limitations. First, our volunteers

were elderly people at high cardiovascular risk and, therefore, the extrapolation of our

conclusions to other populations is hindered. Second, we report moderate changes after

the interventions. However, the PREDIMED trial is based on modest real-life dietary modifications and the control diet is already a well-known healthy dietary pattern. Finally, it was not possible to calculate the pro-thrombotic score in a random 16.5% of the volunteers due to the missing values in the individual determinations of thrombosis biomarkers (each marker presented missing values that were not overlapping with the others). Nevertheless, the amount of missing values was low for individual parameters (0% for fibrinogen and antithrombin; ≤1% for the von Willebrand factor, PAF-AH activity in HDLs, and non-esterified fatty acids; and ≤5% for the rest of the determinations excepting nitrites + nitrates –6.1%–) and the order in which samples were analyzed was randomly assigned before starting the laboratory procedures. Thus, we may assume that missing data occurred at random and did not consistently affect the rest of determinations.

In conclusion, 1 year of intervention with TMD improved the pro-thrombotic status in high cardiovascular risk individuals. The TMD-VOO intervention enhanced HDL antithrombotic properties, whilst the TMD-Nuts one decreased the levels of non-esterified fatty acids. High TMD adherences particularly improved the pro-thrombotic status and decreased fibrinogen and non-esterified fatty acid concentrations. The low-fat diet was the only dietary group linked to increments in the levels of platelet factor-4 and prothrombin activation. TMD effects could be particularly based on increments in the intake of nuts, fruits, and vegetables, and decreases in processed meat consumption. As far as we know, this is the largest, most comprehensive, hypothesis-based analysis of the effects of a healthy dietary pattern on thrombosis mechanisms in high cardiovascular risk subjects. Our results support the improvement of thrombosis status after following a TMD. Further studies are needed to confirm whether these improvements mediate the reported cardioprotective benefits of this dietary pattern.

ACKNOWLEDGEMENTS

This work was supported by grants of Instituto de Salud Carlos III [OBN17PI02, PIE14/00045_INFLAMES, CB06/03/0019, CB06/03/0028, CD17/00122 (A.H.), and JR17/00022 (O.C.)], and Agència de Gestió d'Ajuts Universitaris i de Recerca (2017 SGR 222). The sponsors of this study are public/nonprofit organizations that support science in general and had no role in gathering, analyzing, or interpreting the data. CIBER de Fisiopatología de la Obesidad y Nutrición (CIBEROBN) is an initiative of the Instituto de Salud Carlos III, Madrid, Spain.

Authors wish to thank Daniel Muñoz-Aguayo, Gemma Blanchart, and Sònia Gaixas for their assistance in the development of the present research project.

AUTHORSHIP

Contribution: A.H. and M. Fitó designed the research; A.H., O.C. A.T.-R., X.P., and R.E were involved in funding search; X.P., M. Fitó, M.-Á. M.-G., D.C., J.S.-S., J.L., E.G.-G., F.A., M. Fiol, L.S.-M., E.R., and R.E. conducted the clinical trial and provided biological samples; A.H., O.C., M. Fitó, and R.C. developed the experimental part of the project; A.H. analyzed results and wrote the paper; O.C. A.T.-R., X.P., M. Fitó, R.C., M.-Á. M.-G., D.C., J.S.-S., J.L., E.G.-G., F.A., M. Fiol, L.S.-M., E.R., and R.E. revised the manuscript and approved its final version.

Conflict-of-interest disclosure: X.P. reports personal fees from Abbott, Esteve, Lacer, Rubio, and Sanofi outside the submitted work. J.S.-S. reports to be a board member and grants from Nut and Dried Fruit; personal fees from Aguas Font Vella Lanjarón, Danone S.A., and Instituto Danone; grants from Eroski Distributors; and non-financial support from Nuts for Life outside the submitted work. F.A. reports personal fees from Menarini and AstraZeneca outside the submitted work. E.R. reports personal fees and non-financial support from Merck, Sharp & Dohme; grants, personal fees, and non-financial support

Correspondence: Álvaro Hernáez, August Pi i Sunyer Biomedical Research Institute (IDIBAPS), IDIBAPS Offices-Mallorca (Floor -1, PREDIMED Office), Carrer Mallorca 185, 08036, Barcelona, Spain; e-mail: alvaro.hernaez1@gmail.com.

REFERENCES

1.   Estruch R, Ros E, Salas-Salvadó J, et al. Primary Prevention of Cardiovascular Disease with a Mediterranean Diet Supplemented with Extra-Virgin Olive Oil or Nuts. *N. Engl. J. Med.* 2018;378(25):e34.

2.   Martínez-González MA, Gea A, Ruiz-Canela M. The Mediterranean Diet and Cardiovascular Health. *Circ. Res.* 2019;124(5):779–798.

3.   Martínez-González MA, Salas-Salvadó J, Estruch R, et al. Benefits of the Mediterranean Diet: Insights From the PREDIMED Study. *Prog. Cardiovasc. Dis.* 2015;58(1):50–60.

4.   Gresele P, Momi S, Migliacci R. Endothelium, venous thromboembolism and ischaemic cardiovascular events. *Thromb. Haemost.* 2010;103(1):56–61.

5.   Montalescot G, Philippe F, Ankri A, et al. Early increase of von Willebrand factor predicts adverse outcome in unstable coronary artery disease: beneficial effects of enoxaparin. French Investigators of the ESSENCE Trial. *Circulation.* 1998;98(4):294–9.

6.   Wassel CL, Berardi C, Pankow JS, et al. Soluble P-selectin predicts lower extremity peripheral artery disease incidence and change in the ankle brachial index: The Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis.* 2015;239(2):405–411.

7.   Altara R, Manca M, Brandao RD, et al. Emerging importance of chemokine receptor CXCR3 and its ligands in cardiovascular diseases. *Clin. Sci.* 2016;130(7):463–478.

8.   Feinbloom D, Bauer KA. Assessment of hemostatic risk factors in predicting arterial thrombotic events. *Arterioscler. Thromb. Vasc. Biol.* 2005;25(10):2043–53.

9.   Loeffen R, van Oerle R, Leers MPG, et al. Factor XIa and Thrombin Generation Are Elevated in Patients with Acute Coronary Syndrome and Predict Recurrent

Cardiovascular Events. *PLoS One.* 2016;11(7):e0158355.

10.   Thompson SG, Fechtrup C, Squire E, et al. Antithrombin III and fibrinogen as

      predictors of cardiac events in patients with angina pectoris. *Arterioscler. Thromb.*

      Vasc. Biol. 1996;16(3):357–62.

11.   Cortellaro M, Cofrancesco E, Boschetti C, et al. Increased fibrin turnover and high

      PAI-1 activity as predictors of ischemic events in atherosclerotic patients. A case-

      control study. The PLAT Group. *Arterioscler. Thromb. Vasc. Biol.*

      1993;13(10):1412–7.

12.   Lowe GD, Rumley A. Use of fibrinogen and fibrin D-dimer in prediction of arterial

      thrombotic events. *Thromb. Haemost.* 1999;82(2):667–72.

13.   Darvall KAL, Sam RC, Silverman SH, Bradbury AW, Adam DJ. Obesity and

      Thrombosis. *Eur. J. Vasc. Endovasc. Surg.* 2007;33(2):223–233.

14.   Moshage H, Kok B, Huizenga JR, Jansen PL. Nitrite and nitrate determinations in

      plasma: a critical evaluation. *Clin. Chem.* 1995;41(6):.

15.   van der Stoep M, Korporaal SJA, Van Eck M. High-density lipoprotein as a

      modulator of platelet and coagulation responses. *Cardiovasc. Res.*

      2014;103(3):362–71.

16.   Tellis CC, Tselepis AD. The role of lipoprotein-associated phospholipase A2 in

      atherosclerosis may depend on its lipoprotein carrier in plasma. *Biochim. Biophys.*

      Acta. 2009;1791(5):327–38.

17.   Casas R, Sacanella E, Urpí-Sardà M, et al. The effects of the mediterranean diet on

      biomarkers of vascular wall inflammation and plaque vulnerability in subjects with

      high risk for cardiovascular disease. A randomized trial. *PLoS One.*

      2014;9(6):e100084.

18.   Bonaccio M, Di Castelnuovo A, De Curtis A, et al. Adherence to the Mediterranean

      diet is associated with lower platelet and leukocyte counts: results from the Moli-

sani study. *Blood*. 2014;123(19):3037–3044.

19.   Mezzano D, Leighton F, Martínez C, et al. Complementary effects of Mediterranean
      diet and moderate red wine intake on haemostatic cardiovascular risk factors. *Eur.
      J. Clin. Nutr.* 2001;55(6):444–51.

20.   Schröder H, Fitó M, Estruch RR, et al. A short screener is valid for assessing
      Mediterranean diet adherence among older Spanish men and women. *J. Nutr.*
      2011;141(6):1140–5.

21.   Martínez-González MÁ, Corella D, Salas-Salvadó J, et al. Cohort profile: design and
      methods of the PREDIMED study. *Int. J. Epidemiol.* 2012;41(2):377–85.

22.   Hernaez A, Castañer O, Elosua R, et al. Mediterranean Diet Improves High-Density
      Lipoprotein Function in High-Cardiovascular-Risk Individuals: A Randomized
      Controlled Trial. *Circulation*. 2017;135(7):633–643.

23.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
      lme4. *J. Stat. Softw.* 2015;67(1):1–48.

24.   R Core Team. R: A language and environment for statistical computing. R
      Foundation for Statistical Computing, Vienna, Austria. 2014;

25.   Franco AT, Corken A, Ware J. Platelets at the interface of thrombosis, inflammation,
      and cancer. *Blood*. 2015;126(5):582–588.

26.   Samad F, Ruf W. Inflammation, obesity, and thrombosis. *Blood*.
      2013;122(20):3415–3422.

27.   Luyendyk JP, Schoenecker JG, Flick MJ. The multifaceted role of fibrinogen in
      tissue injury and inflammation. *Blood*. 2019;133(6):511–520.

28.   Hussain T, Tan B, Yin Y, et al. Oxidative Stress and Inflammation: What
      Polyphenols Can Do for Us? *Oxid. Med. Cell. Longev.* 2016;2016:7432797.

29.   Marion-Letellier R, Savoye G, Ghosh S. Polyunsaturated fatty acids and
      inflammation. *IUBMB Life*. 2015;67(9):659–667.

30. Ohira H, Tsutsui W, Fujioka Y. Are Short Chain Fatty Acids in Gut Microbiota Defensive Players for Inflammation and Atherosclerosis? *J. Atheroscler. Thromb.* 2017;24(7):660–672.

31. Chai W, Morimoto Y, Cooney R V, et al. Dietary Red and Processed Meat Intake and Markers of Adiposity and Inflammation: The Multiethnic Cohort Study. *J. Am. Coll. Nutr.* 2017;36(5):378–385.

32. Hernáez A, Farràs M, Fitó M. Olive oil phenolic compounds and high-density lipoprotein function. *Curr. Opin. Lipidol.* 2016;27(1):47–53.

33. Casas R, Sacanella E, Urpí-Sardà M, et al. Long-Term Immunomodulatory Effects of a Mediterranean Diet in Adults at High Risk of Cardiovascular Disease in the PREvención con DIeta MEDiterránea (PREDIMED) Randomized Controlled Trial. *J. Nutr.* 2016;146(9):1684–93.

34. Oh DY, Talukdar S, Bae EJ, et al. GPR120 is an omega-3 fatty acid receptor mediating potent anti-inflammatory and insulin-sensitizing effects. *Cell.* 2010;142(5):687–98.

35. Satapati S, Qian Y, Wu MS, et al. GPR120 suppresses adipose tissue lipolysis and synergizes with GPR40 in antidiabetic efficacy. *J. Lipid Res.* 2017;58(8):1561–1578.

36. den Besten G, van Eunen K, Groen AK, et al. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J. Lipid Res.* 2013;54(9):2325–40.

TABLES

Table 1. Description of the study population

	TMD-VOO (N=120)	TMD-Nuts (N=119)	Low-fat diet (N=119)	P-value
Sex (female) (n, %)	78 (65.0%)	75 (63.0%)	74 (62.2%)	0.898
Age (years)	67.3 (5.34)*	66.6 (5.79)	66.5 (6.35)	0.530
Educational level:				0.228
Primary or less (n, %)	95 (79.2%)	79 (66.4%)	81 (68.1%)	
Secondary (n, %)	15 (12.5%)	25 (21.0%)	21 (17.6%)	
Higher (n, %)	9 (7.50%)	14 (11.8%)	13 (10.9%)	
Non-available (n, %)	1 (0.83%)	1 (0.84%)	4 (3.36%)	
Glucose (mg/dL)	125 (39.9)	123 (46.7)	131 (46.4)	0.321
Glucose-lowering drug users (n, %)	41 (34.2%)	35 (29.4%)	44 (37.0%)	0.458
Total cholesterol (mg/dL)	213 (34.6)	214 (36.3)	207 (33.5)	0.298
HDL cholesterol (mg/dL)	53.4 (9.03)	51.2 (9.88)	51.0 (11.1)	0.134
LDL cholesterol (mg/dL)	134 (32.5)	137 (33.1)	131 (29.5)	0.414
Triglycerides (mg/dL)	114 [92.9; 156]†	109 [80.7; 156]	111 [84.5; 146]	0.672
Lipid-lowering drug users (n, %)	62 (51.7%)	51 (42.9%)	51 (42.9%)	0.287
Systolic blood pressure (mmHg)	154 (18.3)	155 (21.6)	153 (20.9)	0.668
Diastolic blood pressure (mmHg)	83.0 (10.5)	85.6 (11.3)	83.9 (8.90)	0.141
Antihypertensive drug users (n, %)	86 (71.7%)	79 (66.4%)	76 (63.9%)	0.423
Antithrombotic drug users (n, %)	19 (15.8%)	13 (10.9%)	14 (11.8%)	0.479
Tobacco use:				0.445
Never smoker (n, %)	83 (69.2%)	77 (64.7%)	69 (58.0%)	

Actual smoker (*n*, %)	16 (13.3%)	15 (12.6%)	20 (16.8%)	
Former smoker (*n*, %)	21 (17.5%)	27 (22.7%)	30 (25.2%)	
Body mass index (kg/m^2)	29.7 (3.37)	29.9 (3.50)	29.5 (3.34)	0.690
Leisure-time physical activity (METs·min/d)	215 [103; 367]	179 [76.9; 335]	202 [59.4; 377]	0.447
Mediterranean diet adherence score	8.61 (1.68)	8.69 (1.85)	8.99 (1.64)	0.197

*: Mean (standard deviation)

: Median [first quartile; third quartile]

HDL: high-density lipoprotein; *LDL*: low-density lipoprotein; *METs*: metabolic equivalents of task.

Table 2. Changes in thrombosis biomarkers throughout the 1-year intervention relative to baseline.

	Low-fat control diet (*N*=119)			TMD-VOO (*N*=120)			TMD-Nuts (*N*=119)		
Normally distributed variables (*paired t-tests*)									
	Post-int. value	Difference	*P*-value	Post-int. value	Difference	*P*-value	Post-int. value	Difference	*P*-value
	30.3	0.50		30.2	0.057		29.3	-0.52	
Thrombosis score	(6.54)	(6.52)	0.434	(5.69)	(5.32)	0.913	(5.82)	(5.87)	0.378
Von Willebrand factor	31.2	-2.03		32.2	0.22		30.9	1.00	
(μg/mL)	(14.3)	(14.8)	0.139	(14.5)	(13.6)	0.859	(12.9)	(11.9)	0.364
Fibrinogen	275	7.40		277	-0.094		272	1.92	
(mg/dL)	(65.1)	(64.2)	0.211	(63.5)	(68.0)	0.988	(62.1)	(57.8)	0.718
Antithrombin	36.1	0.38		37.0	0.33		36.1	-0.15	
(mg/dL)	(4.44)	(4.14)	0.321	(4.37)	(3.19)	0.254	(5.29)	(4.40)	0.718
Plasminogen activator	3.68	0.080		3.60	0.009		3.68	0.098	
inhibitor-1 (mg/dL)	(1.56)	(2.12)	0.683	(1.66)	(1.97)	0.962	(1.73)	(2.10)	0.619
D-dimer	1.12	0.057		1.13	0.063		1.25	0.089	
(μg/mL)	(0.54)	(0.67)	0.360	(0.59)	(0.77)	0.378	(0.70)	(0.83)	0.251

	Pre-int. value	Post-int. value	P-value	Pre-int. value	Post-int. value	P-value	Pre-int. value	Post-int. value	P-value
α1-antitrypsin in HDLs	2.09	0.073		1.98	0.028		2.00	-0.009	
(per mg of HDLc, no units)	(0.65)	(0.51)	0.130	(0.68)	(0.60)	0.615	(0.61)	(0.43)	0.825
PAF-AH activity in HDLs	1.00	-0.076		1.06	-0.036		1.05	-0.10	
(normalized units)	(0.35)	(0.33)	0.016	(0.38)	(0.38)	0.303	(0.36)	(0.42)	0.010
Non-esterified	0.53	-0.007		0.51	0.012		0.47	-0.036	
fatty acids (mmol/L)	(0.20)	(0.24)	0.736	(0.19)	(0.19)	0.495	(0.18)	(0.19)	0.038
Nitrites + nitrates	37.8	0.93		36.6	1.54		34.7	0.51	
(µM)	(33.8)	(40.1)	0.808	(24.8)	(24.2)	0.494	(24.3)	(27.9)	0.849

Non-normally distributed variables (*Wilcoxon signed rank tests*)									
	Pre-int. value	Post-int. value	*P*-value	Pre-int. value	Post-int. value	*P*-value	Pre-int. value	Post-int. value	*P*-value
Platelet factor-4	268	351		287	372		288	323	
(ng/mL)	[183; 431]	[211; 565]	0.012	[164; 498]	[202; 563]	0.168	[186; 486]	[205; 514]	0.901
Prothrombin fragment 1+2	261	319		246	312		275	289	
(pg/mL)	[149; 433]	[209; 538]	0.005	[162; 415]	[223; 487]	0.114	[165; 439]	[194; 443]	0.573

HDL: high-density lipoprotein; *PAF-AH*: platelet activating factor acetylhydrolase; *Pre-int.*: pre-intervention; *Post-int.*: post-intervention; *TMD-Nuts*: traditional Mediterranean diet enriched with mixed nuts; *TMD-VOO*: traditional Mediterranean diet enriched with virgin olive oil.

FIGURES

Figure 1. TMD effects on thrombosis biomarkers.

Effects of TMD-VOO (first forest plot of each panel) and TMD-Nuts interventions (second

forest plot of each panel) (versus the low-fat control diet), and of subjects with greater

increases in TMD adherence (versus those showing a decrease in their compliance to the

TMD intervention): thrombosis score (**A**), and levels of von Willebrand factor (**B**), platelet

factor-4 (**C**), fibrinogen (**D**), prothrombin fragment 1+2 (**E**), antithrombin (**F**), plasminogen

activator inhibitor-1 (**G**), D-dimer (**H**), nitrites + nitrates (**I**), non-esterified fatty acids (**J**), α_1-

antitrypsin in HDLs (**K**), and platelet activating factor acetylhydrolase activity in HDLs (**L**).

Results are shown as adjusted coefficients (percentage changes relative to baseline

values) with 95% confidence intervals.

*: *P*<0.05.

HDL: high-density lipoprotein; *PAF-AH*: platelet activating factor acetylhydrolase; *TMD*:

traditional Mediterranean diet; *TMD-Nuts*: traditional Mediterranean diet enriched with

mixed nuts; *TMD-VOO*: traditional Mediterranean diet enriched with virgin olive oil.