Population Structure Drives Differential Methicillin-resistant

***Staphylococcus aureus* Colonization Dynamics**

Matthew S. Mietchen MPH[1], Christopher T. Short PhD[1], Matthew Samore MD[2], Eric

T. Lofgren MSPH, PhD[1] for the CDC Modeling Infectious Diseases in Healthcare

Program (MInD-Healthcare)

[1]Paul G. Allen School for Global Animal Health, College of Veterinary Medicine,

Washington State University, Pullman, WA

[2]Department of Internal Medicine, University of Utah School of Medicine, University of

Utah, Salt Lake City, UT

Corresponding Author Email: Eric.Lofgren@wsu.edu

Keywords: MRSA, colonization, ICU, mathematical model, metapopulation

Abstract

Background: Using a model of methicillin-resistant *Staphylococcus aureus* (MRSA)

within an intensive care unit (ICU), we explore how differing hospital population

structures impact these infection dynamics.

Methods: Using a stochastic compartmental model of an 18-bed ICU, we compared the

rates of MRSA acquisition across three potential population structures: a Single Staff

Type (SST) model with nurses and physicians as a single staff type, a model with

separate staff types for nurses and physicians (Nurse-MD model), and a

Metapopulation model where each nurse was assigned a group of patients. By varying

the proportion of time spent with the assigned patient group (γ) within the

Metapopulation model, we explored whether simpler models may be acceptable

approximations to more realistic patient-healthcare staff contact patterns.

Results: The SST, Nurse-MD, and Metapopulation models had a mean annual number

of cumulative MRSA acquisitions of 40.6, 32.2 and 19.6 respectively. All models were

sensitive to the same parameters in the same direction, although the Metapopulation

model was less sensitive. The number of acquisitions varied non-linearly by values of γ,

with values below 0.40 resembling the Nurse-MD model, while values above that

converged toward the metapopulation structure.

Discussion: The population structure of a modeled hospital has considerable impact on

model results, with the SST model having more than double the acquisition rate of the

more structured Metapopulation model. While the direction of parameter sensitivity

remained the same, the magnitude of these differences varied, producing different

infection rates across relatively similar populations. The non-linearity of the model's

response to differing values of γ suggests only a narrow space of relatively dispersed

nursing assignments where simple model approximations are appropriate.

Conclusion: Simplifying assumptions around how a hospital population is modeled,

especially assuming random mixing, may overestimate infection rates and the impact of

interventions.

Introduction

The Centers for Disease Control and Prevention (CDC) considers methicillin-resistant *Staphylococcus aureus* (MRSA) to be a serious threat to patient safety, and recent trends show that the incidence of infection is no longer declining as aggressively as it has for much of the last decade [1]. The evidence base for interventions to successfully address MRSA is mixed. For example, it has been difficult to quantify the effectiveness of MRSA screening and contact precautions, which has led to disagreement over their benefit [2]–[5]. Other efforts, such as improved hand hygiene are often successful [6], [7], but reducing MRSA acquisitions by means of improved transmission prevention continues to be a focus for hospital infection control efforts.

Preventing the spread of MRSA within intensive care units (ICUs) is especially critical, due to the vulnerable nature of the patients and the difficulty in treating severe infections. Patients admitted to the ICU have been found to have persistent colonization with MRSA 12-14 days after discharge from the hospital [8]. There is strong evidence that frequency and patterns of interaction between staff and patients play a critical role in transmission [9]. The ratio of nurses to patients has been found to contribute to the overall level of colonization and transmission within healthcare settings and ICUs. Lower nurse-to-patient ratios have a positive correlation with increased transmission and poor health outcomes [10]. Beyond this, clinical practices such as the use of consistent care teams, grouping particular types of patients together, or even the hospital's built environment may create distinctive sub-populations even within a single unit. These sub-populations then form a larger, interconnected "metapopulation", the infection dynamics of which, in a hospital setting, remain largely unexplored. In other

areas of infectious disease epidemiology, metapopulations and other complex

population structures have been shown to be important [11], [12].

Studying the transmission of MRSA within an ICU is often a difficult task due to

the highly constrained environment. The patient population within any given ICU is

small, with patients sharing care team, being impacted by the same policies, and

sharing the same environment, which violates the independence assumptions

underlying many statistical techniques. There are limited or no control groups due to co-

occurring interventions and the clinical care mission, and many potential interventions,

such as policy changes, are difficult if not impossible to effectively blind as part of a

clinical trial.

Using mathematical models can help alleviate many of these issues. Widely used

models generally which assume random mixing between healthcare workers and

patients, where all patients are cared for by all healthcare workers, but clinical practices

such as the use of consistent care teams, grouping particular types of patients together,

etc. naturally creates a metapopulation like structure within hospital units. Patient

isolation and limited interaction with other patients creates sub-populations within the

larger patient population forming a weakly connected metapopulation.

Evidence suggests that modeling disease transmission dynamics within

metapopulations has an important role in understanding the underlying patterns that

govern disease spread. One previous study [13] suggested that nurses caring for

specific patients can reduce the basic reproductive number (R_0) of a healthcare-

associated pathogen below 1.0, though this model examined a particularly strict form of

non-random mixing, examining the circumstance of a nurse visiting the *same* patient as

they previously cared for rather than visiting another patient. More specifically, the

spatial patterns and aggregation of host populations are important in identifying how a

pathogen is introduced and dispersed within a population [14]. Rarely, however, are

healthcare settings modeled as metapopulations.

Using a stochastic compartmental model of an 18-bed ICU, we compared three

potential population structures: a single staff type model with nurses and physicians as

the same type of healthcare worker, a model with separate staff types for nurses and

physicians where they are different types of healthcare workers, and a highly structured

model where each nurse is assigned a specific group of patients, in order to examine

the impact of treating an ICU with a metapopulation type structure. These models are

hereafter referred to as "SST", "Nurse-MD" and "Metapopulation" respectively. We

explored the impact on the estimated number of MRSA acquisitions and the sensitivity

of the different models to changes in their underlying parameters.

We then developed a hybrid model that allows a nurse to randomly interact with

patients not originally under their direct care over some proportion of their workday. This

hybrid model allows for a population that is primarily, but not exclusively, organized into

distinct sub-populations, but still allows for some interaction between nurses and all

patients. The limited random interaction represents variance from patient assignments

often seen in the ICU environment, such as cross-coverage during breaks or complex

procedures that require higher numbers of healthcare workers.

Methods

Model Structure

MRSA transmission was simulated in an 18-bed ICU that included six nurses and

a dedicated critical care physician based on a previously published model [15]. Hospital

staff are either uncontaminated (S_U) or contaminated (S_C), representing the presence of

infectious material on their hands or person. Patients, similarly, are either uncolonized

(P_U) or colonized (P_C).In the baseline model, patients are assumed to mix randomly with

healthcare workers (HCWs) in the ICU, with no distinction between the nurses or the

intensivist (Fig. 1).

Three alternative models were created to compare how different population

structures within the ICU impacted MRSA acquisitions when relaxing the random mixing

assumption. The "Nurse-MD" model retained random mixing but separated the

intensivist from the nursing staff (Fig. 2). Separation of the physician also allowed the

interactions between healthcare workers and the patients to be more realistic, using

role-specific contact rates with patients, rather than a generic behavior that was the

weighted average of nurse and physician contact rates. This resulted in physicians

having less direct care tasks (touching the patient or their immediate surrounding

environment) involving patients when compared to either nurses or the generic

healthcare workers in the SST model. This model thus had six compartments within it:

the number of patients either colonized (P_C) or uncolonized (P_U), the number of nurses

either contaminated (N_C) or uncontaminated (N_U), and the two additional compartments

representing the physician as either contaminated (D_C) or uncontaminated (D_U).

The second alternative model, the Metapopulation model, further segregated the

healthcare workers by assigning each nurse a specific group of patients (one nurse for

every three patients) and assuming the nurse cared exclusively for those patients. This

practice is common in many ICUs for continuity of care, familiarity, and scheduling

purposes, and may be assumed to predominantly care for those patients. The model's

compartments thus become further divided into six subpopulations, with the physician

acting as a bridge between them by visiting all patients. (Fig. 3). This model structure

creates a metapopulation that is closer to the actual organizational structure. This

model does not assume full random mixing, but rather only assumes that the nurse

visits each patient in their assigned group randomly.

Even where nurses are assigned a specific group of patients, a nurse will have

some interaction with patients outside their assigned group due to cross coverage, staff

breaks, or patient care tasks that require more than one nurse to perform. The final

extension of this model adds an additional parameter, γ, to represent the amount of time

a nurse spends in their assigned group, with the remainder of the time spent moving

randomly among patients outside their assignment. In doing so, the expanded model

effectively hybridizes the previous models, allowing the exploration of intermediate

population structures between a purely random mixing model and one with strict

assignment. In essence, when $\gamma = 1/6$, this model replicates the Nurse-MD model, as a

nurse is no more likely to spend time with their assigned patients as they are any other

five patient groups not assigned to them. Similarly, when $\gamma = 1$, the model replicates the

Metapopulation model, where nurses are strictly confined to treating their assigned

patients. The state transitions for each model, and the equations that govern those

transitions, may be found in the Supplemental Material as Tables S1, S2 and S3

respectively for the SST, Nurse-MD and the expanded Metapopulation model with γ

respectively.

160   Several assumptions underlie all four models. First, patients are assumed to

161 have a single-occupancy room and not to interact with other patients. The nurses and

162 the physician only interact with the patients and do not interact with each other – or

163 more specifically, do not interact in ways relevant to pathogen transmission. The ICU is

164 considered a "closed ICU" – physicians from outside the ICU do not see patients. The

165 ICU is also considered to be at 100% capacity – if a patient is discharged it is assumed

166 another patient is admitted to the bed immediately [16]. A hand hygiene opportunity

167 occurs after every direct care task and personal protective equipment such as gowns

168 and gloves are changed on entry and exit from the rooms of colonized patients. Note

169 both of these are performed with imperfect compliance (Table 1). Lastly, we assumed

170 that MRSA colonization was detected instantly and with perfect sensitivity and

171 specificity to simplify the model.

173 *Parameterization*

174   Parameter values were obtained predominantly from a previously published

175 model of MRSA transmission in an ICU [17] and are described in Table 1. The

176 alternative models introduce new interactions between the patient and their healthcare

177 team, which required rederivation of some parameters from their original sources [18]–

178 [20]. Specifically, the hand hygiene and gown/glove change rates incorporate nurse and

179 physician specific contact rates, which were recalculated using the same methods as in

180 the previous work.

181   Contact rates between patients and healthcare workers were represented by

182 direct care tasks per hour for each healthcare worker type. Direct care tasks are defined

as the physical interaction of the healthcare worker with the patient or their surrounding

environment [20]. Effective hand-decontaminations per hour (ι) were calculated by the

number of direct care tasks and taking into consideration the compliance rate and

handwashing efficacy. Effective gown and glove changes per hour (τ) were calculated

based on the number of visits to a patient per hour and a compliance rate – changing

gowns and gloves was assumed to be 100% effective at removing contamination from a

healthcare worker.

190   One additional parameter was added to the model differing from previously

191   published work. A natural decolonization rate based on results from the STAR*ICU Trial

192   was added based on evidence that colonization of MRSA is limited, and natural

193   decolonization can occur without targeted treatment or decontamination efforts, moving

194   patients from P_C to P_U at a low rate absent any direct intervention [8], [21].

196   *Model Simulation*

197   The SST, Nurse-MD and Metapopulation models were simulated to count the

number of patients who transitioned to the colonized state (P_C) in order to compare the

average number of MRSA acquisitions. The models were stochastically simulated using

Gillespie's Direct Method [25] in Python 3.6 using the StochPy package [26] for 1000

iterations per model. The initial conditions for each model was set to have no

contaminated healthcare workers, either nurses or the physician, and no colonized

patients, with initial MRSA infections being seeded from colonized members of the

community being admitted to the ICU. Each iteration was run for a single year. The

distribution of the acquisitions for each model's 1000 iterations was visualized in R

v3.5.1 using the vioplot package [27], and the difference between them assessed using

a Kruskal-Wallis test. The code for the model simulation and subsequent analysis may

be found at github.com/epimodels/Metapopulation_MRSA.

Model Recalibration

In addition to considering model outcomes using a single set of parameters

(originally calibrated to the SST model), we also examined the difference in the

estimated value of a single free parameter which could be fit within each model. The

purpose of this recalibration is two-fold. First, it allows for a comparison of the models in

a setting where their outcomes are equal. Second, it allows us to examine how each

model form might influence the value of an estimated parameter – important information

in a setting where models may be used to perform statistical inference and estimate

intervention efficacy. The parameter chosen for this recalibration, ψ, is the probability of

an effective colonization of an uncolonized patient from contact between a

contaminated healthcare worker.

Approximate Bayesian Computation (ABC) [28] was used for the parameter

fitting and to obtain an approximate Bayesian posterior of ψ for the SST, Nurse-MD and

Metapopulation models. This method samples a candidate value from a prior

distribution, performs the model simulation using that candidate, and compares a

summary statistic from that simulation to a target statistic. The candidate value is

accepted if the simulation's summary statistic equals the target statistic ± an error term

ε. This is performed repeatedly, and the resulting distribution of accepted candidates is

an approximation of a Bayesian posterior distribution.

For this analysis, the target number of acquisitions was set to 5.94 acquisitions

per 1,000 person-days with an ε of 15%, matching the rate seen in the control arm of a

large randomized clinical trial on MRSA prevention during the study period [17]. A

uniform prior bounded by 0.0 and 1.0 was used, and 1,000,000 candidate parameters

were drawn from this distribution to obtain the approximated Bayesian posterior of ψ for

each model, using a simulation procedure similar to the one described above. For

comparison between models, the median of this distribution was used as the value for

ψ.

Parameter Sensitivity Analysis

In addition to assessing the difference in raw acquisitions in each model, we

assessed the sensitivity of this outcome to changes in the model's parameters. All

parameters in the model were allowed to vary uniformly $\pm50\%$ of their original values,

and 100,000 parameter combinations were simulated for each model. For each model,

the recalibrated value for ψ was used in order to ensure the models were compared

against a consistent acquisition rate. The number of acquisitions in each simulation was

then normalized as a percentage-change from the mean number of acquisitions. Linear

regression was used on the normalized acquisition rate to determine the percentage

change in acquisitions due to a single-percentage change in each parameter value.

A more structural sensitivity question within the Metapopulation model was

explored by varying the amount of time a nurse spends exclusively with their assigned

group vs. other patients on the ward, γ. The Metapopulation model including γ was

simulated 10,000 times, drawing a value of γ for each iteration from a uniform

distribution bounded by 1/6 and 1. A segmented Poisson regression model was then fit

to detect any thresholds in the value of γ where it's relationship to the rate of MRSA

acquisitions notably changed, or if the transition between the Nurse-MD model ($\gamma=1/6$)

and the Metapopulation model ($\gamma=1$) was linear. This model incorporated linear and

quadratic terms for γ and allowed the model to choose any number of break points.

Results

Model Comparison

When using the same parameter set (calibrated to the SST model), the

probability density and average number of MRSA acquisition were significantly different

between the SST, Nurse-MD and Metapopulation models (χ^2 = 1796.8, df = 2, p >

0.001) (Fig. 4). Using the SST model as the baseline for comparison, a decrease in the

average number of MRSA acquisitions were observed in both the separate Nurse-MD

model and the Metapopulation model. By separating the physician from the nurses, the

mean acquisitions decreased 20.7% from 40.6 acquisitions to 32.2 acquisitions,

respectively. The Metapopulation model experienced a 51.7% decrease from the

original SST model at 19.6 acquisitions.

Model Recalibration

The model parameter ψ (psi) (the probability of effective colonization of an

uncolonized patient from contact with a contaminated healthcare worker) was used to

calibrate the models. Calibration of the SST model resulted in the median value of the

parameter of 0.024 (95% Credible Interval: 0.016, 0.034). The Nurse-MD model results

were very similar to the SST model, with a median value of 0.029 (95% Credible

Interval: 0.019,0.042). In contrast, the Metapopulation model had a median ψ value of

0.046 (95% Credible Interval: 0.032, 0.07), both a substantially higher estimate than the

other models and one in which the bounds of the credible interval did not contain the

other estimates. The altered contact patterns in the Metapopulation model thus needs

substantially higher per-contact colonization probabilities to sustain the same level of

contact.

Sensitivity Analysis

While the Metapopulation model resulted in fewer acquisitions, certain

parameters were found to affect the model outcomes to a larger magnitude when

compared to the other models. The three parameters in SST model showing the largest

proportional change (> 0.20) in cumulative acquisitions (Fig. 5a): contact rate (ρ),

probability of patient colonization (ψ), and hand-decontamination (ι). Similar findings

were also found in the Nurse-MD model, though generally only for the nurse-specific

parameters (Fig. 5b). The doctor-specific parameters had little effect on the model

outcomes. Only one parameter of the Metapopulation model had a large change in

cumulative acquisitions > 0.20 – the nurse-specific contact rate. However, the

parameters with the largest effects were consistent with the previous two models (Fig.

5c).

The directionality of the overall change in cumulative acquisitions by parameter is

an important measure of model stability and correct parameter estimates, as this

reflects whether the models *qualitatively* give the same results as to whether or not a

particular parameter value changing results in an increase or decrease in MRSA

acquisitions, even if the models disagree as to the specific value of that change. All of

the parameters between the models are consistent in terms of directionality, with the

Metapopulation model having a smaller change in magnitude of the cumulative

acquisitions (Fig. 5d).

Metapopulation Interactions

The relationship between γ and MRSA acquisitions was non-linear (Fig. 6), with

progressively higher values of γ resulting in drastically reduced rates of MRSA

acquisition. The segmented Poisson regression model identified a single change point,

γ^*, at 0.40 (95% Confidence Interval: 0.37, 0.42). Values below γ^* were well

approximated by the Nurse-MD model, and values above it rapidly approached the

stricter assignment of the Metapopulation model.

Discussion

Developing interventions to reduce the spread or transmission of antibiotic-

resistant pathogens, such as MRSA, is an important public health and healthcare goal

due to costly on-going hospitalizations and difficulty of treatment. Mathematical models

are an important tool in the development of these interventions, but only if they can

represent the population and transmission dynamics of a hospital.

One important aspect of the highly structured hospital population is the

interaction between patients and healthcare workers. Nurses in particular are typically

assigned to a group of patients within a particular shift, suggesting the possibility of

modeling a hospital unit as a metapopulation. We developed a model using a

metapopulation structure that explores how nurses-to-patient interactions may affect the

transmission of pathogens in an ICU setting.

As compared to a model that assumes random mixing, which dominate the

epidemiological compartmental model literature, a more structured metapopulation

appears to be warranted when modeling MRSA within an ICU. The random mixing

assumption allows a higher degree of interaction with the patients and healthcare

workers, resulting in an over-estimation of both the overall rate of MRSA acquisition and

an over-estimation of the impact of interventions.

While the models considered in this study had similar parameter sensitivity in

terms of the *direction* of changes, the more highly structured models were relatively less

sensitive. In all cases, the contact rate (ρ), probability of patient colonization (ψ), and

hand-decontamination (ι) parameters had the largest impact, consistent with many of

the known drivers of infection rates within hospitals.

Conclusion

When combined, these results suggest that while compartmental models that

assume random mixing and those that have more structured populations may give

qualitatively the same answer as to the benefit of an intervention, the magnitude of

these estimates may vary considerably, which has implications for cost-effectiveness

models and other studies that rely on these estimates. Additionally, if the interventions

suggested by the model are implemented in practice, the performance of the

intervention may differ from the model's predictions due to the choice of population

structure. Finally, these results show that fitted parameters can vary considerably even

for very similar models, suggesting even mild changes in model form necessitate

refitting, and parameter estimates are not transportable from one model to another

model with a differing population structure.

Even within a highly structured population, there are events that will lead to the

interactions outside the normal structure. In an ICU setting, it is foreseeable that events

like emergencies, breaks, or cross-coverage of nurses will occur with a reasonable

degree of frequency. Our model suggests that even relatively small increases in the rate

at which these interactions occur can have outsized impacts on MRSA rates. This

finding has broad implications for staffing levels and hospital policy and may provide an

avenue for reducing MRSA rates that does not rely on individual-level actions.

This study has several limitations. While the Metapopulation model is a more

granular representation of a hospital population than the more-common SST model, it

too is a simplification. Similarly, the parameter estimates used in the model are

imperfect. In particular, it is likely that the hand hygiene rate is likely higher than the

rates occurring in many hospitals, as reported rates are often substantially inflated.

However, these estimates are drawn primarily from the established literature, and

represent the field's best understanding of the underlying processes.

Other limitations include the structure of the model – it focuses specifically on

healthcare worker and patient interactions and does not account for interactions with

individuals other than nurses and the physician. For example, interactions among

patients, visitation by family and friends, medical or radiological technicians performing

a specific procedure, etc. are not represented. Similarly, transmission purely through

environmental contamination is not represented.

Hospital settings are characterized by highly structured populations and very

limited types of interactions. Despite this, these environments are rarely modeled as

metapopulations, especially at the ward level. This study shows that the random mixing

assumption results in an over-estimation of both the overall rate of MRSA acquisition

and an over-estimation of the impact of interventions as expressed as changes in model

parameter values. In contrast, a metapopulation model tends to result in proportionately

lower acquisition rates and somewhat more attenuated responses to changes in

parameters. These findings are consistent with observations of the effect of

metapopulations in the disease ecology literature. These results suggest that, even at

small scales, models that assume random mixing may be inappropriate and result in an

overestimation in both acquisition rates and the impact of interventions.

Acknowledgements

Funding: This work was supported by the CDC Cooperative Agreement RFA-CK-17-

001-Modeling Infectious Diseases in Healthcare Program (MInD-Healthcare). The

authors would also like to acknowledge Justin O'Hagan for his thoughtful input.

Conflict of Interest Statement

The authors declare they have no conflicts of interest.

References

[1]   A. Kourtis, K. Hatfield, and J. Baggs, "Vital Signs: Epidemiology and Recent Trends in Methicillin-Resistant and in Methicillin-Susceptible Staphylococcus aureus Bloodstream Infections — United States.," *MMWR Morb Mortal Wkly Rep*, no. 68, pp. 214–219, 2019.

[2]   G. Fätkenheuer, B. Hirschel, and S. Harbarth, "Screening and isolation to control meticillin-resistant Staphylococcus aureus: sense, nonsense, and evidence," *The Lancet*, vol. 385, no. 9973, pp. 1146–1149, Mar. 2015.

[3]   D. J. Morgan, P. Malani, and D. J. Diekema, "Diagnostic Stewardship—Leveraging the Laboratory to Improve Antimicrobial Use," *JAMA*, vol. 318, no. 7, p. 607, Aug. 2017.

[4]   D. J. Morgan, R. P. Wenzel, and G. Bearman, "Contact Precautions for Endemic MRSA and VRE: Time to Retire Legal Mandates," *JAMA*, vol. 318, no. 4, p. 329, Jul. 2017.

[5]   M. A. Rubin, M. H. Samore, and A. D. Harris, "The Importance of Contact Precautions for Endemic Methicillin-Resistant Staphylococcus aureus and Vancomycin-Resistant Enterococci," *JAMA*, vol. 319, no. 9, pp. 863–864, Mar. 2018.

[6]   L. P. G. Derde *et al.*, "Interventions to reduce colonisation and transmission of antimicrobial-resistant bacteria in intensive care units: an interrupted time series study and cluster randomised trial," *Lancet Infect. Dis.*, vol. 14, no. 1, pp. 31–39, Jan. 2014.

[7]   J. M. Boyce and D. Pittet, "Guideline for Hand Hygiene in Health-Care Settings: Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force," *Infect. Control Hosp. Epidemiol.*, vol. 23, no. S12, pp. S3–S40, Dec. 2002.

[8]   M. R. Haverkate, L. P. G. Derde, C. Brun-Buisson, M. J. M. Bonten, and M. C. J. Bootsma, "Duration of colonization with antimicrobial-resistant bacteria after ICU discharge," *Intensive Care Med.*, vol. 40, no. 4, pp. 564–571, Apr. 2014.

[9]   B. G. Mitchell, A. Gardner, P. W. Stone, L. Hall, and M. Pogorzelska-Maziarz, "Hospital Staffing and Health Care–Associated Infections: A Systematic Review of the Literature," *Jt. Comm. J. Qual. Patient Saf.*, Jun. 2018.

[10]   A. Lee, Y. S. L. Cheung, G. M. Joynt, C. C. H. Leung, W.-T. Wong, and C. D. Gomersall, "Are high nurse workload/staffing ratios associated with decreased survival in critically ill patients? A cohort study," *Ann. Intensive Care*, vol. 7, no. 1, Dec. 2017.

[11]   G. Hess, "Disease in Metapopulation Models: Implications for Conservation," *Ecology*, vol. 77, no. 5, pp. 1617–1632, Jul. 1996.

[12]   M. Jesse, P. Ezanno, S. Davis, and J. A. P. Heesterbeek, "A fully coupled, mechanistic model for infectious disease dynamics in a metapopulation: Movement and epidemic duration," *J. Theor. Biol.*, vol. 254, no. 2, pp. 331–338, Sep. 2008.

[13]   D. J. Austin and R. M. Anderson, "Studies of antibiotic resistance within the patient, hospitals and the community using simple mathematical models," *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, vol. 354, no. 1384, pp. 721–738, Apr. 1999.

[14]   A. R. North and H. C. J. Godfray, "The dynamics of disease in a metapopulation: The role of dispersal range," *J. Theor. Biol.*, vol. 418, pp. 57–65, Apr. 2017.

[15]   E. T. Lofgren, "Estimating the impact of post randomization changes in staff behavior in infection prevention trials: a mathematical modeling approach," *BMC Infect. Dis.*, vol. 17, no. 1, Dec. 2017.

[16]   E. T. Lofgren, "Pools versus Queues: The Variable Dynamics of Stochastic 'Steady
States,'" *PLOS ONE*, vol. 10, no. 6, p. e0130574, Jun. 2015.
[17]   A. D. Harris, "Universal Glove and Gown Use and Acquisition of Antibiotic-Resistant
Bacteria in the ICU: A Randomized Trial," *JAMA*, Oct. 2013.
[18]   J. Westbrook, A. Ampt, and L. Kearney, "Work observation method by activity timing
(WOMBAT) manual." Health Informatics Research & Evaluation Unit, The University of Sydney,
Sydney, 2007.
[19]   J. I. Westbrook and A. Ampt, "Design, application and testing of the Work Observation
Method by Activity Timing (WOMBAT) to measure clinicians' patterns of work and
communication☆," *Int. J. Med. Inf.*, vol. 78, pp. S25–S33, Apr. 2009.
[20]   M. A. Ballermann, N. T. Shaw, D. C. Mayes, R. N. Gibney, and J. I. Westbrook, "Validation
of the Work Observation Method By Activity Timing (WOMBAT) method of conducting time-
motion observations in critical care settings: an observational study," *BMC Med. Inform. Decis.
Mak.*, vol. 11, no. 1, Dec. 2011.
[21]   K. Khader *et al.*, "A Dynamic Transmission Model to Evaluate the Effectiveness of
Infection Control Strategies," *Open Forum Infect. Dis.*, p. ofw247, Dec. 2016.
[22]   J. I. Westbrook, C. Duffield, L. Li, and N. J. Creswick, "How much time do nurses have for
patients? a longitudinal study quantifying hospital nurses' patterns of task time distribution and
interactions with health professionals," *BMC Health Serv. Res.*, vol. 11, no. 1, Dec. 2011.
[23]   E. T. Lofgren, R. W. Moehring, D. J. Anderson, D. J. Weber, and N. H. Fefferman, "A
Mathematical Model to Evaluate the Routine Use of Fecal Microbiota Transplantation to
Prevent Incident and Recurrent *Clostridium difficile* Infection," *Infect. Control Hosp. Epidemiol.*,
vol. 35, no. 1, pp. 18–27, Jan. 2014.
[24]   E. Sickbertbennett, D. Weber, M. Gergenteague, M. Sobsey, G. Samsa, and W. Rutala,
"Comparative efficacy of hand hygiene agents in the reduction of bacteria and viruses," *Am. J.
Infect. Control*, vol. 33, no. 2, pp. 67–77, Mar. 2005.
[25]   D. T. Gillespie, "Exact stochastic simulation of coupled chemical reactions," *J. Phys.
Chem.*, vol. 81, no. 25, pp. 2340–2361, Dec. 1977.
[26]   T. R. Maarleveld, B. G. Olivier, and F. J. Bruggeman, "StochPy: A Comprehensive, User-
Friendly Tool for Simulating Stochastic Biological Processes," *PLoS ONE*, vol. 8, no. 11, p.
e79345, Nov. 2013.
[27]   D. Adler, *vioplot: Violin plot. R package.* 2005.
[28]   T. Toni, D. Welch, N. Strelkowa, A. Ipsen, and M. P. H. Stumpf, "Approximate Bayesian
computation scheme for parameter inference and model selection in dynamical systems," *J. R.
Soc. Interface*, vol. 6, no. 31, pp. 187–202, Feb. 2009.
467

468
Figure 1. Schematic representation of the compartmental flow of a mathematical model
of methicillin-resistant *Staphylococcus aureus* (MRSA) acquisition with a single type of
staff. Solid arrows indicate possible transition states, while dashed arrows indicate
potential routes of MRSA contamination or colonization. Healthcare staff are classified
as uncontaminated (S_U) or contaminated (S_c), while patients are classified as
uncolonized (P_U) or colonized (P_C).

475
476

477
Figure 2. Schematic representation of the compartmental flow of a mathematical model
of methicillin-resistant *Staphylococcus aureus* (MRSA) acquisition with nurses and
intensivists separated into different staff types. Solid arrows indicate possible transition
states, while dashed arrows indicate potential routes of MRSA contamination or
colonization. Nurses and doctors are classified as uncontaminated (N_U or D_U) and
contaminated (N_C and D_C), while patients are classified as uncolonized (P_U) or
colonized (P_C).

485
486

487
Figure 3. Schematic representation of a Metapopulation model of methicillin-resistant
Staphylococcus aureus (MRSA) acquisition. Patients (blue) are treated by a single
assigned nurse (orange). A single intensivist (red) randomly treats all patients.
491

492
493 **Figure 4.** Distribution of cumulative MRSA acquisitions in 3,000 simulated 18-bed
494 intensive care units under three theoretical population structures.
495

Figure 5. Global parameter sensitivity of three modeled ICU population structures. Panel A depicts the change in proportional change in cumulative MRSA acquisitions per one-percent change in the value of a specific parameter, with light bars indicating increased acquisitions, and dark bars indicating decreased acquisitions for a model assuming random mixing and with a single staff type for both nurses and physicians. Panel B depicts the same for a model that separates nurses and physicians into different staff types, while Panel C depicts the same for a metapopulation model where nurses were assigned to a strict subpopulation of patients. Panel D depicts the difference in proportional changes between the Metapopulation and Nurse-MD models.

507
Figure 6. Relationship between the proportion of time nurses spend treating patients
outside their assigned group (γ) and cumulative MRSA acquisitions over 10,000
simulations, randomly sampling γ from a uniform distribution between 1/6 and 1. Grey
dots show an individual simulation, while the black line shows a segmented Poisson
regression fit with linear and quadratic terms for γ. The vertical dashed line depicts the
single segmentation point, γ^*, to the left of which these more complicated models are
adequately approximated by the Nurse-MD model where random mixing occurs. The
shaded area shows the corresponding confidence interval.
516

Table 1. Parameters for modeling the acquisition of methicillin-resistant *Staplylococcus aureus* in an Intensive Care Unit

Parameter	Parameter Description	Parameter Value	Source(s)
ρ	Contact rate between patients and HCWs	4.154 (# of direct care tasks/hour)	[22], [23]
ρ_N	Contact rate between patients and nurses	3.973 (# of nurse direct care tasks/hour)	[22], [23]
ρ_D	Contact rate between patients and physician	0.181 (# of physician direct care tasks/hour)	[22], [23]
σ	Probability that a HCW's hands are contaminated from a single contact with a colonized patient	0.054	[13]
ψ_{SST}	Probability of successful colonization of an uncolonized patient due to contact with a contaminated HCW when randomly mixed	0.1494	Fitted to [17]
$\psi_{Nurse\text{-}MD}$	Probability of successful colonization of an uncolonized patient due to contact with a contaminated HCW with physician separated	0.1660	Fitted to [17]
$\psi_{Metapopulation}$	Probability of successful colonization of an uncolonized patient due to contact with a contaminated HCW in metapopulation structure	0.4481	Fitted to [17]
θ	Probability of discharge	4.39 days^{-1}	[17]
ν_u	Proportion of admissions uncolonized with MRSA	0.9221	[17]
ν_c	Proportion of admissions colonized with MRSA	0.0779	[17]
ι	Effective hand-decontaminations/hour (direct care tasks × hand hygiene compliance × efficacy)	5.740 (10.682 direct care tasks/hour × 56.55% compliance × ~ 95% efficacy)	[17], [22]–[24]
ι_N	Effective nurse hand-decontaminations/hour	6.404 (11.92 direct care tasks/hour × 56.55% compliance × ~ 95% efficacy)	[17], [22]–[24]
ι_D	Effective physician hand-decontaminations/hour	1.748 (3.253 direct care tasks/hour × 56.55% compliance × ~ 95% efficacy)	[17], [22]–[24]
τ	Effective gown or glove changes/hour (2 × # of visits × compliance)	2.445 (2.957 changes/hour × 82.66% compliance)	[13], [17], [20]
τ_N	Effective nurse gown or glove changes/hour	2.728 (3.30 changes/hour × 82.66% compliance)	[13], [17], [20]
τ_D	Effective physician gown or glove changes/hour	0.744 (0.90 changes/hour × 82.66% compliance)	[13], [17], [20]
μ	Natural decolonization rate	20.0 days^{-1}	[21]