Disruptive Mood Dysregulation Disorder: symptomatic and syndromic thresholds and diagnostic operationalization

Running title: DMDD: diagnostic thresholds and operationalization

Paola Paganella Laporte[1,2,3], Alicia Matijasevich[4,5], Tiago N. Munhoz[5,6], Iná S. Santos[5], Aluísio J. D. Barros[5], Daniel Samuel Pine[7], Luis Augusto Rohde[1,2,3], Ellen Leibenluft[7], Giovanni Abrahão Salum[1,2,3]

[1] Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Graduate Program in Psychiatry and Behavioral Sciences, Brazil

[2] Section on Negative Affect and Social Processes, Hospital de Clínicas de Porto Alegre, Brazil

[3] National Institute of Developmental Psychiatry for Children and Adolescents, CNPq, Brazil

[4] Department of Preventive Medicine, Faculty of Medicine FMUSP, University of São Paulo, São Paulo, Brazil

[5] Postgraduate Program in Epidemiology, Federal University of Pelotas, Brazil

[6] Faculty of Psychology, Federal University of Pelotas, Pelotas, Brazil

[7] National Institute of Mental Health, Intramural Research Program, Bethesda, MD, USA

Address correspondence and reprint requests

Paola Paganella Laporte

Hospital de Clínicas de Porto Alegre

Ramiro Barcelos, 2350 – room 2202; Porto Alegre, Brazil – 90035-003;

E-mail: paola_laporte@yahoo.com.br - Phone/Fax: +55 51 3359 8094

Acknowledgements

Acknowledgements

The authors thank the children and families for their participation, which made this research possible. We also would like to thank the research team from the Pelotas 2004 Birth Cohort. The Pelotas 2004 Birth Cohort is conducted by the Graduate Program in Epidemiology of Universidade Federal de Pelotas, supported by *Associação Brasileira de Saúde Coletiva* (ABRASCO – Brazilian Association of Public Health). From 2009 to 2013, the 2004 birth cohort was funded by the Wellcome Trust (United Kingdom). The 11-year follow-up was funded by the Sao Paulo Research Foundation-FAPESP (grant number 2014/13864-6). Previous phases of the study were funded by the World Health Organization, *Programa de Apoio a Núcleos de Excelência* (PRONEX – Support Program for Excellence Centers), *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq – National Council for Scientific and Technological Development), the Brazilian Ministry of Health, and *Pastoral da Criança* (Child's Pastoral).

Financial Disclosures

Dr Laporte, Matijasevich, Munhoz, Santos, Barros, Pine, Leibenluft and Salum report no biomedical financial interests or potential conflicts of interest. Dr. Laporte, Matijasevich, Santos, Barros, Rohde and Salum are supported by the CNPq. Dr. Luis Augusto Rohde was on the speakers' bureau/advisory board and/or acted as consultant for Eli-Lilly, Janssen-Cilag, Novartis, Medice and Shire in the last three years. He receives authorship royalties

Abstract

Objective: The aim of this study is to identify the most appropriate threshold for Disruptive Mood Dysregulation Disorder (DMDD) diagnosis and the impact of potential changes in diagnostic rules on prevalence levels in the community.

Methods: Trained psychologists evaluated 3,562 adolescents from the 2004 Pelotas Birth Cohort with the Development and Well-Being Behavior Assessment (DAWBA). The clinical threshold was assessed in three stages: symptomatic, syndromic and clinical operationalization. The symptomatic threshold identified the response category in each DAWBA item which separates normative misbehavior from a clinical indicator. The syndromic threshold identified the number of irritable mood and outbursts needed to capture children with high symptom levels. Clinical operationalization compared the impact of AND/OR rules for combining irritable mood and outbursts on impairment and levels of psychopathology.

Results: At the symptomatic threshold, most irritable mood items were normative in their lowest response categories and clinically significant in their highest response categories. For outbursts some indicated a symptom even when present at only a mild level, while others did not indicate symptoms at any level. At the syndromic level, a combination of 2 out of 7 irritable mood and 3 out of 8 outburst indicators accurately captured a cluster of individuals with high level of symptoms. Analysis combining irritable mood and outbursts delineated non-overlapping aspects of DMDD, providing support for the OR rule in clinical operationalization. The best DMDD criteria resulted in a prevalence of 4%.

Conclusions: Results provide information for initiatives aiming to provide data-driven and clinically-oriented operationalized criteria for DMDD.

Introduction

Temper outbursts and irritable mood are common manifestations of typical development. When outbursts and irritable mood are intense, frequent, last for significant periods, occur in several contexts, and are associated with behaviors not seen in typically developing children, they often require clinical attention [1–3]. Disruptive Mood Dysregulation Disorder (DMDD) is a new diagnosis designed to capture pathological manifestations of irritable mood and temper outbursts [4]. Given the newness of DMDD, data-driven approaches based on epidemiological evidence are needed to evaluate appropriate thresholds for DMDD and consider the need to refine criteria. The current report provides such data.

DMDD has its origins in the mid-2000s when Leibenluft and colleagues [5,6] defined a syndrome called severe mood dysregulation (SMD). SMD involved severe, chronic grouchy mood and heightened reactivity, along with symptoms of hyperarousal [6]. The syndrome was defined to distinguish children with severe irritability from those with classic bipolar disorder (BD), in light of increasing numbers of children diagnosed with BD [7,8]. The results of those studies converged to differentiate SMD from classic bipolar disorder based on course and familial aggregation [9–11]. For DSM-5, SMD was modified to create DMDD.

Alternative thresholds for defining DMDD have been only partially considered in the current literature. Some previous studies have focused on irritability as a dimensional trait, which is broader than DMDD as a diagnostic entity. These studies provide an important framework for investigating clinically-relevant thresholds for specific behaviors. Wakschlag and collaborators [12] used item response theory analysis to disentangle normative misbehavior from clinically significant problems by studying the 'symptomatic threshold', i.e., investigating which response category in each item from a questionnaire separates normative misbehavior from a clinical indicator. They found that some behaviors are normative and

only represent problems when their frequency is high or very high, whereas others always indicate a significant problem that requires clinical attention. This and similar research efforts in preschoolers [13] inform attempts to evaluate varying boundaries for the definition of DMDD. Other studies focused more specifically on varying DSM-5 criteria for DMDD in pre-adolescents [14,15] and adolescents [14–16]. They found the prevalence of temper outbursts and negative mood are much lower than what is found in preschoolers and that applying exclusion criteria such as frequency and hierarchical diagnostic rules affects DMDD prevalence rates considerably [14–16]. There was no evidence that clinical markers changed between pre-adolescents/early adolescents (9-12) to middle adolescents (13-16) [15]. Nonetheless, it is important to continue to identify appropriate diagnostic thresholds for distinct developmental periods, given that normative levels of irritability clearly vary across the lifespan [14–16].

Another important step towards evaluating such varying boundaries involves quantifying the number of abnormal behaviors required to characterize a valid diagnosis i.e., identifying the 'syndromic threshold' for a given diagnosis. Data-driven clustering approaches such as latent class analysis derive groups that differ in the number of clinical indicators endorsed [17] and thus inform attempts to set syndromic thresholds. Such efforts need to be balanced with clinical applicability in real world settings, which require practical decisions such as how to combine clinical indicators from distinct domains (i.e., irritable mood and outbursts). The latter can be achieved by investigating whether domains explain overlapping or distinct aspects of DMDD latent structure and related impairment, thus determining whether "and" or "or" rules should be used to provide a 'clinical operationalization' of the diagnosis. Previous research in pre-adolescents and adolescents suggests irritable mood and temper outbursts predict each other over time. However, while

each of them are associated with increased risk for disrupted functioning in adolescents[15], current criteria require both to be present for a diagnosis to be assigned.

The aim of this study is to evaluate alternative clinical thresholds for the DMDD diagnosis (see Figure 1 for an overview of the analytic strategy and Methods for details). We investigate 3,562 pre-adolescents/early adolescents aged 10-12. First, we used Confirmatory Factor Analysis (CFA) to identify item-level thresholds differentiating normative from clinical problems (the symptomatic threshold). This was used to dichotomize response levels as clinically significant or not. We next used these binary clinical indicators as input to a latent class analysis (LCA) that assigned individuals into clusters with high and low levels of clinical indicators for each domain. This was followed by receiver operating curves (ROC) to detect the number of clinical indicators needed to predict class membership from the LCA and to translate the data-driven results to DSM-5 symptom counts (the syndromic threshold). We then compare the impact of AND/OR rules on impairment and dimensional levels of psychopathology (clinical operationalization). Finally, we investigate the impact of varying definitions on DMDD prevalence and comorbidity profiles in a population-based sample.

Figure 1 around here

Methods

Participants

Participants of this study were pre-adolescents/early adolescents aged 10-12 from the 2004 Pelotas Birth Cohort Study. All births occurring in the city of Pelotas, from January 1st to December 31st, 2004 were enrolled and followed over time. Pelotas is in southern Brazil and has a population of 328,000. For a full description of the methods, see [18]. Briefly, all

4231 live births in the city in 2004 whose mothers lived in the urban area and agreed to participate in the longitudinal study were considered eligible. Follow-up home visits were performed when the subjects had reached the ages of 3.0 months (SD=0.1), 11.9 months (SD=0.2), 23.9 months (SD=0.4), and 49.5 months (SD=1.7). When the subjects were, on average, 6.8 years old (SD=0.3) and 11.0 years old (SD=0.4), additional follow-up visits were conducted at a research clinic run by the Postgraduate Program of Epidemiology (Faculty of Medicine, Federal University of Pelotas, Brazil). Of the 4,231 subjects in the original birth cohort, 3,562 (84.1%) were included in our analysis, which used all available data from the 10-12 years of age assessment. The sample comprises 2,353 participants aged 10, 1,206 aged 11 and 4 aged 12. The prevalence of DMDD in this sample using current criteria associated with clinical ratings was 2.5% (95% CI=2.0–3.0) [19]. The study was approved by the Research Ethics Committee of the Federal University of Pelotas and by the Research Committee of the University of São Paulo School of Medicine. Written informed consent was obtained from all subjects.

Instruments and Diagnostic Assessment

The parent-version of the DMDD section from the Development and Well-Being Assessment (DAWBA) questionnaire [20] was administered by certified psychologists. This questionnaire uses open and closed ended questions to identify the occurrence of clinical indicators in children and adolescents aged 5-17, based on the DSM criteria. The closed ended questions start with two skip questions about the frequency of temper outbursts and irritable mood. Parents who answered that temper outbursts and/or irritable mood occurred at least once a week were probed to answer specific questions that characterize all DSM-5 criteria for DMDD.

A total of 593 parents of participants answered the DMDD section on irritable mood, representing the top 17% of irritable mood frequency. This section includes 9 items characterizing the threshold for experiencing anger, intensity of anger if compared to peers of the same age, duration of anger during the day, whether irritable mood is perceived by others, setting in which anger occurs (at home, at school, with peers) and number of anger weeks throughout the year. A total of 425 parents answered the DMDD section on outbursts, representing the top 12% of frequency of outbursts. This section includes 15 items describing behavior during outbursts (slamming doors, shouting, swearing, saying mean things to others, saying negative things about self, physical aggression to others, deliberate self-harm, breaking things), setting in which outbursts occurred (at home, at school, with peers) and triggers (recognizable and easily triggered). We do not use the item "outbursts free-gap in the last year" in our analysis (DSM requires that there is not a period higher than 3 or more consecutive months without irritable mood and temper outbursts). The rationale for excluding this item is that it is unclear whether we would expect this item to be monotonically related to the overall latent construct given short periods of irritability with large gaps could also inform episodes of irritability (a marker of severity and bipolar disorder).

Lastly, 686 mothers or caregivers that completed either the outburst or irritable mood sections were asked to also complete 4 items about impairment (functioning in daily activities, friendship, learning, and leisure activities). After the impairment questions, mothers or caregivers answered the open-ended questions that allow qualitative description of the symptoms, frequency, and other characteristics of the disorder. All questions and response categories from the DMDD section are depicted in Supplemental Table S7.

The DAWBA was administered to mothers or caregivers by trained psychologists. The forty-hour training included lectures, role playing, and supervised clinical interviews

with pediatric and mental health outpatients at the Federal University of Pelotas. The clinical evaluation of the total sample was performed by a psychologist, and a second independent psychologist evaluated 10% of the study sample. Both were trained in how to apply the DAWBA, in a standardized manner, by the child psychiatrist who had translated and validated the questionnaire for use in Brazil [21]. The inter-rater agreement was 91.2% for the presence of any psychiatric disorder, 75.9% for any anxiety disorder, 73.5% for any depressive disorder, 72.7% for ADHD, 72.9% for conduct disorder, 85.6% for any autism spectrum disorder, 59.5% for any eating disorder, and 52.4% for any tic disorder. Details of the questionnaire can be found online and in other studies [22].

The Strengths and Difficulties Questionnaire (SDQ) was used to measure dimensional psychopathology. The SDQ is a 25-item behavioral screening questionnaire with five domains, each of which contain five items (emotional, conduct, hyperkinetic, peer relationships, prosocial behaviors and impact scores). The overall SDQ total scores had a Cronbach's alpha of 0.82, which is considered high. Internal consistency for the SDQ subscales were low to moderate ranging from 0.48 (peer relationships) to 0.78 (hyperkinetic). Despite low reliability, we maintained results from subscales in the supplemental material for their descriptive nature.

Statistical Analysis

Symptomatic threshold

The fourteen items on outbursts, eight items on irritable mood and four items on impairment were included in three CFAs testing unidimensional models for each construct (n= 593, 425 and 685 respectively). CFA models estimate item level factor loadings (λ) and response category thresholds. Factor loadings represent the strength of the relationship

between the latent trait and the item, i.e., they indicate how well each item discriminates different severity levels of a given construct. Category thresholds indicate the expected value of the latent factor at which there is a 50% probability of endorsing a given category or higher i.e., the category threshold indicates the severity level at which the transition from one response category to the next is likely to happen (e.g., from 'No' to 'A little' or higher, or from 'A little' to 'A lot').

To distinguish normative misbehavior from behavior that would meet a diagnostic criterion, we used category thresholds from the CFA. CFAs were performed only in subjects with a frequency of irritable mood and outbursts greater than once a week. In this sample, a value of 0.5 represents a half standard deviation above the mean of the distribution of subjects with a frequency of irritable mood and outbursts greater than once a week. Therefore, we interpreted values below 0.5 as typical development (normative) and values at or above 0.5 as 'clinical indicators' (a proxy for symptoms or problem indicators). The latter represents approximately the top 5% most symptomatic pre-adolescents/early adolescents in the population, which is a threshold used in other diagnostic investigations [12]. For details about the CFA, see Supplemental Material.

Syndromic threshold

Before data analysis, each questionnaire item was dichotomized at the value of the category threshold defined in the symptomatic threshold analysis described above in subjects with at least one clinical indicator. Dichotomized items were chosen to enter the LCA analysis because our intention was not to characterize varying levels of irritability in the community, but to identify groups that differ in their number of clinical indicators. Three Latent Class Analyses (LCA) were used to create empirically derived groups with different

levels of clinical indicators for irritable mood, outbursts and impairment. Next, we used three Receiving Operating Curves to predict the most accurate number of clinical indicators for detecting participants with high levels of symptoms (as defined by latent class analysis) with regard to irritable mood, outburst and impairment. ROC analysis was used as a way to translate results from the syndromic thresholds of the LCA to the reality of clinical practice, which uses symptom counts. Thus, the ROC identifies a simple rule to allow the identification of patients that are likely to be members of the cluster that exhibit a high level of clinical indicators. The optimal cut-off was estimated using the Younden's J Statistic, which maximizes both sensitivity and specificity [23].

Clinical operationalization

Four analytical strategies were used to determine the most appropriate rule for clinical operationalization: the 'OR' rule vs. the 'AND' rule. First, we compared the fit of CFA models (n=398), putting the selected dichotomized clinical indicators into a unidimensional model of irritability and a correlated model of irritability with two domains (irritable mood and outbursts). Second, we tested whether meeting criteria for the irritable mood group and/or for the outbursts group have distinct or overlapping associations with the impaired functioning group using a multiple logistic regression. Third, we used left censored regressions to compare skewed SDQ dimensional scores between subjects meeting criteria only for irritable mood, only outbursts, either, or both and compared with a group of participants with other DSM disorders except for DMDD and typically developing comparisons. Fourth, for both OR and AND groups, we used a matching procedure to compare levels of SDQ scores between a group that differed in DMDD status (yes vs. no DMDD) but were otherwise fully matched for comorbidities.

Epidemiological impact

Finally, using the relative frequency, we investigated the impact of these AND/OR rules and combinations for impairment requirements on the prevalence rates of DMDD in the community and on the comorbidity profile.

All analysis were performed in R version 3.6.1 [24], including applications implemented in the packages lavaan 0.6-5 [25], poLCA 1.4.1 [26], pROC 1.15.3 [27], CensReg 0.5-26 [28] and MatchIt 3.0.2 [29]. The R markdown codes for the symptomatic, syndromic and clinical operationalization thresholds of current analysis can be found in the Supplemental Material.

Results

Symptomatic threshold

All eight items of irritable mood were found to be normative in their lowest thresholds and clinical indicators (proxy for symptoms) in the highest thresholds, except "irritable mood that happens at home", which was found to be normative in all response categories. For the six items that describe intensity, the response option "A little" indicated normative behavior, while the response option "A lot" or "A great deal" indicated a symptom. For the duration item, irritable mood lasting less than an hour indicated normative behavior, whereas irritability lasting a few hours or most of the day indicated a symptom. For the frequency item, irritable mood occurring less than 3 times a week indicated normative behavior, whereas irritable mood occurring every day indicated a symptom (Supplemental Table S2).

For outbursts, threshold varied substantially across items. For some items, their occurrence even at mild levels indicated a symptom, while other items did not indicate a symptom at any level. Outbursts that include self-harm, breaking things or saying negative things about self, or those that occurred in the classroom are indicative of a symptom if they occur at any level (i.e., "A little" or "A lot"). Outbursts that occurred with peers, include physical aggression, or are easily triggered indicate a symptom when they occurred "A lot", but were normative when they occurred "A little". Outbursts that occurred at home and included the pre-adolescents/early adolescents saying mean things, slamming doors, shouting, or swearing did not indicate a symptom irrespective of the level endorsed. Also, whether the triggers were recognizable or not was not relevant to symptom designation. Regarding frequency, only outbursts that occurred daily indicated a symptom (Supplemental Table S3).

For impairment, "impact on family life" is normative when "A little" and indicate a symptom when "A medium amount" or "A lot". Impairment that occurs in the other settings (friendship, learning or leisure) indicate a symptom at any level (Supplemental Table S4). See Figure 2.

Figure 2 around here

Syndromic threshold

The prevalence of each clinical indicator is presented in Table 1. The three LCA analyses (irritable mood, outbursts and impairment) indicated that the two-class solution was the best for each of the three domains (Supplemental Table S5). This indicates that, in each of the three domains, the population is divided into two groups characterized by high vs. low

symptoms (Supplemental Figure S1). We next performed three Receiver Operating Curve analyses (irritable mood, outburst, impairment) to determine the best number of clinical indicators (i.e., those items identified by the CFA) to use to predict membership in the high vs. low symptom classes identified by the LCA. Younden's J demonstrates that subjects in the high symptom irritable mood and outburst classes are most accurately characterized by 2 out of 7 irritable mood symptoms and 3 out of 8 outburst symptoms. As for impairment, the subjects with high level of symptoms in LCA are most accurately characterized by significant impairment in at least two settings (Table 2 and Supplemental Figure S2).

Table 1 around here

Table 2 around here

Clinical operationalization

First, a model with two correlated domains (irritable mood and outbursts) provided a better fit than a unidimensional model encompassing both domains (χ^2_{diff}=7.3, df=1, p=0.007; Supplemental Table S6). Second, both irritable mood and outbursts were associated with clinical impairment, even after adjusting for each other in a multiple model (irritable mood crude OR=41.71, adjusted OR=18.2, p<0.001; outbursts crude OR=76.1, adjusted OR=23.63, p<0.001). Third, comparisons between irritable only, outbursts only and combined groups with typically developing comparisons and with a group of patients with other DSM disorders (except for DMDD) showed all three DMDD groups had higher scores on all SDQ scales than typically developing comparisons and higher total SDQ total scores than subjects with other DSM diagnosis (Supplemental Figure S3; Supplemental Table S8). Fourth, left-censored regressions comparing groups matched for comorbidity (any anxiety,

any mood, any hyperkinetic and any disruptive behavior disorder) showed that, using either the OR or the AND rule, the DMDD group showed higher total, emotional, conduct, hyperactivity, peer relationship and impact scores than did the non-DMDD group with matched comorbidities (Supplemental Figure S4; Supplemental Figure S5).

Epidemiological impact

When using an "OR" rule, the optimal criteria from the ROC analysis (2 of 7 irritable mood symptoms, 3 of 8 outburst symptoms, impairment in at least two settings) resulted in a prevalence of 3.0%: 1.12% have only irritable mood, 0.64% only outbursts and 1.23% have both irritable mood and outbursts (Table 3). Both the "OR rule" and the "AND rule" resulted in higher levels of psychiatric comorbidities compared to the current DMDD clinical criteria (Supplemental Table S9).

Table 3 around here

Discussion

This study provides important information to guide a revision of the diagnostic criteria for DMDD. Using CFA, we found that seven of the eight irritable mood items were normative when endorsed in the low response categories and clinical indicators in the high response categories. The one exception was "irritable mood that happens at home", which was always normative. For outbursts, the threshold for a clinical indicator varied substantially across items. For some items, such as outbursts with self-harm, their presence indicated a problem even at only mild levels. Others, such as shouting, were not clinical indicators even when present at the highest threshold. ROC analyses indicated that a combination of 2 of 7 irritable mood symptoms, 3 of 8 outburst symptoms and significant impairment in at least

two settings would best predict membership in the "high" vs. "low" LCA-based symptom classes. The four clinical operationalization analysis converge to demonstrate that the two domains differ from a latent perspective; they are independently associated with impairment; and OR-rule groups show comparable or even higher levels of impairment then other DSM disorders. Matched analysis showed that results cannot be attributed to comorbidity. The most accurate solution resulted in a prevalence of 3% in the fully automated operationalized criteria (1.12% only irritable mood, 0.64% only outbursts and 1.23% combined).

Our findings are consistent with the limited literature examining irritability dimensionally in the population. Each set of findings suggest that normative outbursts differ from clinical indicators in frequency, duration, quality, context, and triggering events [30–33]. Wakschlag et al. [12] found that outbursts characterized by high frequency, "long duration", or "aggressive components", or those that occurred "with nonparental adults" or "out of the blue" were clinical indicators. Wiggins and collaborators [13] also used an empirical approach to identify irritable behaviors indicative of problems in preschoolers. They examined 22 temper loss behaviors from the criteria for oppositional defiant disorder, DMDD and other depressive disorders in the DSM-5 and found two informative items. Similar to our work, the item "easily frustrated" indicated a symptom only when present nearly every day, but the item "break/destroy" indicated a symptom even when at lower frequencies. Nevertheless, those thresholds might vary substantially in distinct age ranges and cultures, which highlights the need for developmentally sensitive studies.

Clinical operationalization analysis suggests that an OR rule is most appropriate to capture cases in need of treatment. This algorithm identified pre-adolescents/early adolescents with either irritable mood or outbursts who manifested associated impairment, elevated symptoms, and functional impairment. This resulted in a prevalence rate of 3%,

which is higher than the prevalence rate of 2.5% by the current diagnostic criteria. Of course, it is not possible to identify the "true" prevalence of DMDD in the population with one study; rather the current analyses inform nosologists' attempts to weigh the strengths and weaknesses of various diagnostic thresholds.

Our study has important strengths. *First*, we relied on a large representative population sample and implemented assessment methods that could mimic clinical assessment in the real world, as far as possible in an epidemiological investigation. *Second*, we applied Confirmatory Factor Analysis, Latent Class Analysis and Receiver Operating Curves Analysis, applying a similar framework used in other disorders [34,35] to a new syndrome that lacks empirical investigations to guide operationalization. However, this work has also some important limitations. First, our analysis is focused on internal validators. Further studies investigating course, family history, treatment response, and other external validators are needed to demonstrate the validity of the operationalized syndrome. Since associations between symptoms and irritability-related impairment were investigated using the same DAWBA DMDD section, the size of the associations is likely to be overestimated. However, the value of these odds ratios may be helpful in understanding whether the two aspects of irritability capture distinct or overlapping aspects of irritability-related impairment. Second, our subjects were all 10-12 years old, and our data might not be generalized to other developmental stages. Third, because of the skip rule questions, the CFA parameters were estimated for subjects with irritable mood or outbursts that occurred at least once a week. Analysis were modeled to consider these characteristics, but this might have biased the parameter estimates for some items. Also, our approach assumes irritable mood, outbursts and impairment are distinct domains, which is still an empirical question to be further tested. Fourth, our analysis is restricted to parent reports, and no information was

acquired from pre-adolescents/early adolescents themselves. Lastly, our approach is restricted to a single sample and it is unclear whether those results can be replicated in other samples.

Advancing understanding about DMDD diagnostic criteria is a major concern in children and adolescent psychiatric practice. Our findings are a first step towards defining parameters to alert the clinician when to be (and when not to be) concerned with irritable mood and outbursts as a way to inform the presence of DMDD diagnosis. Future research that examines these patterns with measures specifically designed to differentiate normative versus non-normative patterns and that include developmentally sensitive items might be an important step forward to advance the field of DMDD, specifically. Prospective longitudinal investigation that applies this framework beginning at earlier ages can elucidate the origins of pathologic irritability and inform developmentally-based prevention.

References

1.   Wakschlag LS, Estabrook R, Petitclerc A, et al. Clinical Implications of a Dimensional Approach: The Normal:Abnormal Spectrum of Early Irritability. *J Am Acad Child Adolesc Psychiatry*. 2015;54(8):626-634. doi:10.1016/j.jaac.2015.05.016

2.   Petitclerc A, Briggs-Gowan MJ, Estabrook R, et al. Contextual variation in young children's observed disruptive behavior on the DB-DOS: implications for early identification. *J Child Psychol Psychiatry*. 2015;56(9):1008-1016. doi:10.1111/jcpp.12430

3.   Perlman SB, Jones BM, Wakschlag LS, Axelson D, Birmaher B, Phillips ML. Neural substrates of child irritability in typically developing and psychiatric populations. *Dev Cogn Neurosci*. 2015;14:71-80. doi:10.1016/j.dcn.2015.07.003

4.   DSM-5. https://www.psychiatry.org/psychiatrists/practice/dsm. Accessed July 1, 2018.

5.   Leibenluft E. Severe mood dysregulation, irritability, and the diagnostic boundaries of bipolar disorder in youths. *Am J Psychiatry*. 2011;168(2):129-142. doi:10.1176/appi.ajp.2010.10050766

6.   Leibenluft E, Blair RJR, Charney DS, Pine DS. Irritability in pediatric mania and other childhood psychopathology. *Ann N Y Acad Sci*. 2003;1008:201-218.

7.   Blader JC, Carlson GA. Increased rates of bipolar disorder diagnoses among U.S. child, adolescent, and adult inpatients, 1996-2004. *Biol Psychiatry*. 2007;62(2):107-114. doi:10.1016/j.biopsych.2006.11.006

8.   Olfson M, Blanco C, Liu L, Moreno C, Laje G. National trends in the outpatient treatment of children and adolescents with antipsychotic drugs. *Arch Gen Psychiatry*. 2006;63(6):679-685. doi:10.1001/archpsyc.63.6.679

9.   Stringaris A, Cohen P, Pine DS, Leibenluft E. Adult outcomes of youth irritability:

a 20-year prospective community-based study. *Am J Psychiatry*. 2009;166(9):1048-1054. doi:10.1176/appi.ajp.2009.08121849

10.   Leibenluft E, Cohen P, Gorrindo T, Brook JS, Pine DS. Chronic versus episodic irritability in youth: a community-based, longitudinal study of clinical and diagnostic associations. *J Child Adolesc Psychopharmacol*. 2006;16(4):456-466. doi:10.1089/cap.2006.16.456

11.   Stringaris A, Baroni A, Haimm C, et al. Pediatric bipolar disorder versus severe mood dysregulation: risk for manic episodes on follow-up. *J Am Acad Child Adolesc Psychiatry*. 2010;49(4):397-405.

12.   Wakschlag LS, Choi SW, Carter AS, et al. Defining the developmental parameters of temper loss in early childhood: implications for developmental psychopathology. *J Child Psychol Psychiatry*. 2012;53(11):1099-1108. doi:10.1111/j.1469-7610.2012.02595.x

13.   Wiggins JL, Briggs-Gowan MJ, Estabrook R, et al. Identifying Clinically Significant Irritability in Early Childhood. *J Am Acad Child Adolesc Psychiatry*. 2018;57(3):191-199.e2. doi:10.1016/j.jaac.2017.12.008

14.   Copeland WE, Angold A, Costello EJ, Egger H. Prevalence, comorbidity, and correlates of DSM-5 proposed disruptive mood dysregulation disorder. *Am J Psychiatry*. 2013;170(2):173-179. doi:10.1176/appi.ajp.2012.12010132

15.   Copeland WE, Brotman MA, Costello EJ. Normative Irritability in Youth: Developmental Findings From the Great Smoky Mountains Study. *J Am Acad Child Adolesc Psychiatry*. 2015;54(8):635-642. doi:10.1016/j.jaac.2015.05.008

16.   Althoff RR, Crehan ET, He J-P, Burstein M, Hudziak JJ, Merikangas KR. Disruptive Mood Dysregulation Disorder at Ages 13-18: Results from the National Comorbidity Survey-Adolescent Supplement. *J Child Adolesc Psychopharmacol*.

2016;26(2):107-113. doi:10.1089/cap.2015.0038

17.   Borsboom D, Rhemtulla M, Cramer AOJ, van der Maas HLJ, Scheffer M, Dolan CV. Kinds versus continua: a review of psychometric approaches to uncover the structure of psychiatric constructs. *Psychol Med*. 2016;46(8):1567-1579. doi:10.1017/S0033291715001944

18.   Santos IS, Barros AJD, Matijasevich A, et al. Cohort profile update: 2004 Pelotas (Brazil) Birth Cohort Study. Body composition, mental health and genetic assessment at the 6 years follow-up. *Int J Epidemiol*. 2014;43(5):1437-1437a-f. doi:10.1093/ije/dyu144

19.   Munhoz TN, Santos IS, Barros AJD, Anselmi L, Barros FC, Matijasevich A. Perinatal and postnatal risk factors for disruptive mood dysregulation disorder at age 11: 2004 Pelotas Birth Cohort Study. *J Affect Disord*. 2017;215:263-268. doi:10.1016/j.jad.2017.03.040

20.   Goodman R, Ford T, Richards H, Gatward R, Meltzer H. The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology. *J Child Psychol Psychiatry*. 2000;41(5):645-655.

21.   Fleitlich-Bilyk B, Goodman R. Prevalence of child and adolescent psychiatric disorders in southeast Brazil. *J Am Acad Child Adolesc Psychiatry*. 2004;43(6):727-734. doi:10.1097/01.chi.0000120021.14101.ca

22.   La Maison C, Munhoz TN, Santos IS, Anselmi L, Barros FC, Matijasevich A. Prevalence and risk factors of psychiatric disorders in early adolescence: 2004 Pelotas (Brazil) birth cohort. *Soc Psychiatry Psychiatr Epidemiol*. 2018;53(7):685-697. doi:10.1007/s00127-018-1516-z

23.   Youden WJ. Index for rating diagnostic tests. *Cancer*. 1950;3(1):32-35.

24.   R: The R Project for Statistical Computing. https://www.r-project.org/. Accessed

August 26, 2018.

25.   Rosseel Y, Oberski D, Byrnes J, et al. *Lavaan: Latent Variable Analysis*.; 2018.

https://CRAN.R-project.org/package=lavaan. Accessed August 26, 2018.

26.   poLCA.pdf. https://cran.r-project.org/web/packages/poLCA/poLCA.pdf. Accessed

September 19, 2019.

27.   Robin X, Turck N, Hainard A, et al. pROC: an open-source package for R and S+ to

analyze and compare ROC curves. *BMC Bioinformatics*. 2011;12:77. doi:10.1186/1471-

2105-12-77

28.   CRAN - Package censReg. https://cran.r-

project.org/web/packages/censReg/index.html. Accessed May 27, 2019.

29.   Ho D, Imai K, King G, Stuart E, Whitworth A. *MatchIt: Nonparametric*

Preprocessing for Parametric Causal Inference.; 2018. https://CRAN.R-

project.org/package=MatchIt. Accessed September 19, 2019.

30.   Belden AC, Thomson NR, Luby JL. Temper tantrums in healthy versus depressed

and disruptive preschoolers: defining tantrum behaviors associated with clinical problems.

J Pediatr. 2008;152(1):117-122. doi:10.1016/j.jpeds.2007.06.030

31.   Bhatia MS, Dhar NK, Singhal PK, Nigam VR, Malik SC, Mullick DN. Temper

tantrums. Prevalence and etiology in a non-referral outpatient setting. *Clin Pediatr (Phila)*.

1990;29(6):311-315. doi:10.1177/000992289002900603

32.   Osterman K, Björkqvist K. A cross-sectional study of onset, cessation, frequency,

and duration of children's temper tantrums in a nonclinical sample. *Psychol Rep*.

2010;106(2):448-454. doi:10.2466/PR0.106.2.448-454

33.   Wakschlag LS, Briggs-Gowan MJ, Carter AS, et al. A developmental framework

for distinguishing disruptive behavior from normative misbehavior in preschool children. *J*

Child Psychol Psychiatry. 2007;48(10):976-987. doi:10.1111/j.1469-7610.2007.01786.x

34.   Kessler RC, Green JG, Adler LA, et al. Structure and diagnosis of adult attention-deficit/hyperactivity disorder: analysis of expanded symptom criteria from the Adult ADHD Clinical Diagnostic Scale. *Arch Gen Psychiatry*. 2010;67(11):1168-1178. doi:10.1001/archgenpsychiatry.2010.146

35.   Matte B, Anselmi L, Salum GA, et al. ADHD in DSM-5: a field trial in a large, representative sample of 18- to 19-year-old adults. *Psychol Med*. 2015;45(2):361-373. doi:10.1017/S0033291714001470

Table 1- Prevalence* of each Disruptive Mood Dysregulation Disorder item written in combination with the response category that defines a clinical indicator

	Irritable Mood	Prevalence Estimation of Clinical Indicators (%)
	Frequency/duration	
1	Irritable mood **occurring every day**	2.1
2	Irritable mood that **lasts more than a few hours**	2.7
	Characteristics	
3	Easily irritated, annoyed or angry **a lot**	3.6
4	Intense irritable mood **a lot**	2.8
	Settings	
5	Irritable mood occurs in the classroom **a lot**	1.1
6	Irritable mood occurs with peers **a lot**	0.9
7	Irritable mood is evident to others **a great deal**	1.7
	Temper Outbursts	Prevalence Estimation of Clinical Indicators (%)
	Frequency/duration	
1	Outbursts **occurring every day**	1.5
	Characteristics of the outbursts	
2	Saying **any** negative thing about self	3.1
3	**Any** physical aggression to others	1.7
4	**Any** form of deliberate self-harm	1.3
5	Breaking things (**any**)	3.4

	Settings	
6	*Any* outburst in the classroom	3.3
7	Outbursts occurs with peers *a lot*	0.7
	Triggers	
8	Easily triggered *a lot*	2.6

* Prevalence estimates assume that pre-adolescents/early adolescents whose irritable mood and outbursts occurred less than once per week (and who therefore did not complete these items) do not have any of these problems to a significant degree.

Table 2- Receiver Operating Curves Parameters investigating the best number of clinical indicators to capture latent class groups

Threshold	Prediction of Latent Class Groups														
	Irritable Mood					Outbursts					Severity of Impairment				
	ACC	Sens	Spe	PPV	NPV	ACC	Sens	Spe	PPV	NPV	ACC	Sens	Spe	PPV	NPV
0	0.25	1.00	0.00	0.25	-	0.12	1.00	0.00	0.12	-	0.36	1.00	0.00	0.12	-
1	0.56	1.00	0.41	0.36	1.00	0.40	1.00	0.32	0.17	1.00	0.73	1.00	0.58	0.57	1.00
2	**0.86**	**0.95**	**0.83**	**0.65**	**0.98**	0.69	1.00	0.64	0.28	1.00	**0.93**	**1.00**	**0.89**	**0.83**	**1.00**
3	0.90	0.64	0.99	0.95	0.89	**0.90**	**1.00**	**0.88**	**0.54**	**1.00**	0.90	0.72	1.00	1.00	0.87
4	0.83	0.32	1.00	1.00	0.82	0.98	0.86	0.99	0.98	0.98	0.74	0.28	1.00	1.00	0.71
5	0.78	0.11	1.00	1.00	0.77	0.93	0.43	1.00	1.00	0.92					
6	0.76	0.04	1.00	1.00	0.76	0.89	0.14	1.00	1.00	0.89					
7	0.75	0.00	1.00	-	0.75	0.89	0.12	1.00	1.00	0.89					
8						0.88	0.02	1.00	1.00	0.88					

Note: ACC, Accuracy; Sens, Sensitivity; Spe, Specificity; PPV, Positive Predictive Value; NPV, Negative Predictive Value.

Table 3 – Impact of different rules for combining irritable mood and temper outburst clinical indicators and impairment requirements on prevalence rates of DMDD

	Irritable Mood	Outbursts	AND Rule	OR Rule
No impairment requirement	2.41	0.9	1.63	4.94
At least one setting	1.80	0.70	1.46	3.96
At least two settings (optimal)	**1.12**	**0.64**	**1.23**	**3.00**
At least three settings	0.61	0.48	0.87	1.96
All four settings	0.17	0.17	0.53	0.87

Figure 1 – Symptomatic and syndromic thresholds and clinical operationalizations

Figure 2 – Symptomatic threshold for each irritable mood and outbursts item in the

Confirmatory Factor Analysis

Panel A - Irritable Mood

Panel B - Outbursts

Figure 1 – Symptomatic and syndromic thresholds and clinical operationalizations

Figure 2 – Symptomatic threshold for each irritable mood and outbursts item in the

Confirmatory Factor Analysis

Irritable Mood

Outbursts