Title: Alcohol use and cognitive functioning in young adults: an observational and Mendelian

randomisation study

Liam Mahedy[1,2,3] (Ph.D.), Steph Suddell[1,2,3] (MSc), Caroline Skirrow[1,4], (Ph.D.), Gwen S.

Fernandes[5] (Ph.D.), Matt Field[6] (Ph.D.), Jon Heron[5] (Ph.D.), Matthew Hickman[5] (Ph.D.),

Robyn Wootton[1,2,3] (Ph.D.), Marcus R. Munafò[1,2,3] (Ph.D.)

Affiliations

[1]School of Psychological Science, University of Bristol, Bristol, UK.

[2]MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK.

[3]National Institute for Health Research Biomedical Research Centre at the University

Hospitals Bristol NHS Foundation Trust and the University of Bristol, Bristol, UK.

[4]Cambridge Cognition, Cambridge, UK.

[5]Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[6]Department of Psychology, University of Sheffield, Sheffield, UK.

Corresponding author: Liam Mahedy, School of Psychological Science, University of Bristol,

12a Priory Road, Bristol, UK, BS8 1TU.

Tel: +44 (0) 117 33 10616; Email: liam.mahedy@bristol.ac.uk

Abstract

Background and Aims: There have been few prospective studies of association between alcohol use and cognitive functioning in young people. We aim to examine whether alcohol use is a risk factor for deficient cognitive functioning in young adults.

Design: Longitudinal latent class analysis was used to derive patterns of binge drinking. Linear regression was used to examine the relationship between binge drinking and subsequent cognitive functioning. Two-sample Mendelian randomisation (MR) tested evidence for the causal relationship between alcohol use and cognitive functioning.

Setting: South West England.

Participants: The observational study comprised of 2086 adolescents and their parents (fully adjusted models) from the Avon Longitudinal Study of Parents and Children.

Measurements: Binge drinking was assessed on five occasions from age 16 to 23 years. Cognitive functioning comprised working memory, response inhibition, and emotion recognition assessed at 24 years of age. 99-independent genome-wide significant SNPs associated with 'number of drinks per week' was used as the genetic instrument for alcohol consumption. Potential confounders were included.

Findings: Four binge drinking classes were identified: 'low-risk' (41%), 'adolescent-limited' (19%), 'adult-regular' (23%), and 'early-onset regular' (17%). There was not a consistent association between binge drinking and subsequent cognitive functioning. For example, we found insufficient evidence to support an association between early-onset regular binge drinking and cognitive functioning in the fully adjusted models: working memory (b=0.09, 95% CI=-0.10 to 0.28), response inhibition (b=0.70, 95% CI=-10.55 to 11.95), and emotion recognition (b=0.01, 95% CI=-0.01 to 0.02) compared to low-risk drinkers. Two-sample MR analyses similarly provided little evidence that alcohol use is associated with deficits in

49   working memory (b=0.14, 95% CI=-0.45 to 0.72), response inhibition (b=-0.31, 95% CI=-0.92

50   to 0.29), and emotion recognition (0.03, 95% CI=-0.55 to 0.61).

51   **Conclusions:** Binge drinking in adolescence and early adulthood may not be a strong predictor

52   for deficient cognitive functioning in young adults.

INTRODUCTION

Alcohol use during adolescence, when the brain is still developing and undergoing considerable structural and functional changes (1) is a major public health concern. The association between binge drinking and cognitive functioning (i.e., working memory, response inhibition, and emotion recognition) has received particular attention because some cognitive functions do not peak until early adulthood (2–5) in parallel with maturation of the prefrontal cortex (6,7).

There is a wealth of evidence from animal (8,9), neuroimaging (10–12), twin (13,14), and cognitive neuroscience (15,16) studies suggesting that adolescent binge drinking is negatively associated with cognitive functioning. However, the direction of this association remains unclear as many of these results are based on evidence from small cross-sectional studies. Studies that have examined this association using prospective data have largely revealed mixed findings. For example, some studies have found that alcohol use preceded deficits in domains of cognitive functioning (17–20), while other studies have provided support for the opposite direction (21–23).

In an effort to strengthen the evidence we used a triangulation approach with observational and genetic epidemiological methods to better understand the causal relationships between drinking patterns and cognitive functioning in young adults using data from Avon Longitudinal Study of Parents and Children (ALSPAC). The aims were to investigate (1) whether patterns of binge drinking (assessed between 16 to 23 years) were associated with working memory, response inhibition, and emotion recognition assessed at age 24, and (2) whether alcohol use was associated with cognitive functioning using two-sample Mendelian randomisation (MR) (24). MR can reduce bias from residual confounding and reverse causation by using genetic variants that are known to be associated with the exposure

(25). We expected to find that more regular binge drinking would be associated with deficient

cognitive functioning, and that this association would be supported by the MR analyses.

METHODS

Design

Longitudinal latent class analysis was used to derive heterogenous patterns of binge

drinking from ages 16 to 23 years. Linear regression was used to examine the relationship

between patterns of binge drinking and subsequent cognitive functioning. The young person

provided self-reported information on binge drinking and cognitive functioning. The clear

temporal ordering of exposure, outcome and confounders helps to rule out the possibility of

reverse causality. Two-sample MR tested evidence for the causal relationship between

alcohol use and cognitive functioning.

Observational analyses

Participants and Procedure

We used data from the Avon Longitudinal Study of Parents and Children (ALSPAC), an

ongoing population-based study that contains a wide range of phenotypic and environmental

measures, genetic information and linkage to health and administrative records. A fully

searchable　　data　　dictionary　　is　　available　　on　　the　　study's　　website

http://www.bristol.ac.uk/alspac/researchers/our-data/. Approval for the study was obtained

from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

Informed consent for the use of data collected via questionnaires and clinics was obtained

from participants following recommendations of the ALSPAC Ethics and Law Committee at

the time. Consent for biological samples has been collected in accordance with the Human

Tissue Act (2004). All pregnant women residing in the former Avon Health Authority in the

south-west of England and had an estimated date of delivery between 1 April 1991 and

December 1992 were eligible for the study (Phase I consisted of *N*=14,541). Of the 13,988

offspring alive at one year, a small number of participants have withdrawn fully from the

study (*n*=41), leaving an eligible sample of 13,947. Detailed information about ALSPAC is

available online www.bris.ac.uk/alspac and in the cohort profiles (26–28). A detailed overview

of our study population, including attrition at the different measurement occasions is

presented in Supplementary Material Figure S1.

Measures

A timeline of data collection is presented in Supplementary Material Figure S2.

Exposure: Binge drinking

Information on binge drinking was collected on five occasions via a questionnaire (Q)

or during attendance at a study clinic (C). Median ages at response were: 16y 7m (Q), 17y 9m

(C), 18y 8m (Q) 21y (Q), and 23y (Q) using the following question reflecting drinking over the

past year "How often do you have six or more drinks on one occasion?". One drink was

specified as ½ pint (568 ml) average strength beer/lager, one glass of wine, or one single

measure (25ml) of spirits. Responses at each time point were used to derive a repeated 3

level ordinal variable with categories "Never/Occasional" (comprising of "Never" and "Less

than monthly"), "Monthly" and "Weekly". Daily or almost daily was collapsed into the

"Weekly" group.

Outcome variables

At 24 years of age (*M*=24.0 years; *SD*=9.8 months) participants attended a clinic-based

assessment which included computerised cognitive assessments as part of a broader

assessment battery of mental and physical health and behaviour. Data collection for the

online questionnaires was collected and manged by REDcap electronic data capture tools

(29,30).

Working memory

The *N*-back task (2-back condition) was used to assess working memory. Participants

were asked to monitor the series of numbers (0-9) and respond with a '1' keystroke when any

number was the same as the one presented 2 trials previously, and a '2' keystroke if it was

different. Forty-eight trials were presented, including eight 2-back matches. The *N*-back task

(31) is widely used to measure working memory (17,32,33). A measure of discriminability (*d'*)

was chosen as the primary outcome measure given it is an overall performance estimate. High

scores on *number of hits* indicated more accurate identification, while high scores on *false*

alarms indicated less accurate identification. High scores on *d'*, therefore, indicated a greater

ability to distinguish signal from noise. Of the participants assessed with cognitive tasks at age

24 (*n*=3,312), *n*=182 did not provide any data on the task; *n*=70 were omitted due to negative

d' scores and/or not responding to over 50% of the trials, leaving a sample of *n*=3,242.

Response inhibition

The Stop Signal Task (34) was used to assess response inhibition – the ability to

prevent an ongoing motor response. Participants were asked to respond to visual stimuli (X

and O response keys to X and O letters presented sequentially on a screen) unless a tone was

heard (the stop signal) after the stimulus presentation, at which point they were required to

withhold responding. The task consisted of 256 trials, which included a 4:1 ratio of trials

without stop signals to trials with stop signals. Mean response times were calculated. An

estimate of *stop signal reaction time* (SSRT) was calculated and used as the primary outcome

as it is a reliable measure of inhibitory control, with shorter SSRT's indicating faster inhibition.

SSRT data were available for *n*=3,201 participants.

Emotion recognition

Emotion recognition was assessed using a six alternative forced choice (6AFC) emotion recognition task (35) comprising of 96 trials (16 for each emotion) which measures the ability to identify emotions in facial expressions. In each trial, participants were presented with a face displaying one of six emotions: anger, disgust, fear, happiness, sadness, or surprise. Participants were required to select the descriptor that best described the emotion that was present in the face, using the computer mouse. Emotion intensity varied across 8 levels within each emotion from the prototypical emotion to an almost neutral face. Each individual stimulus was presented twice, giving a total of 96 trials. An overall measure of ER (the number of facial emotions accurately identified) was used as the primary outcome. ER data were available for n=3,368 participants. Further information on all three cognitive tasks is presented in Supplementary material.

Potential confounders

We identified confounders from established risk factors for cognitive functioning that could plausibly have a causal predictive relationship with earlier binge drinking including income, maternal education, socioeconomic position, housing tenure, sex, and maternal smoking during first trimester in pregnancy. Two measures were included to control for cognitive function prior to alcohol initiation. Working memory at approximately 11 years and experience of a head injury/unconsciousness up to 11 years. Finally, measures of cigarette and cannabis use were collected at 4 timepoints between ages ~14 and ~16.5 years (up to the first assessment of binge drinking). Further information is presented in Supplementary material.

Statistical methods

Observational analyses

The observational analyses were conducted in two stages. First, longitudinal latent

class analysis was used to derive trajectories of binge drinking for individuals having at least

one measure of binge drinking (*n*=6,888). Starting with a single latent class, additional classes

were added until model fit was optimised. See supplementary material for a description of

model fit. Analyses were carried out using Mplus 8.1 (36).

Class membership was then related to covariates using the three-step method using

the Bolck-Croon-Hagenaars (37) method. The first stage estimated the latent class

measurement model and saves the weights. While, the second stage involved using these

weights which reflect the measurement error of the latent class variable. Linear regression

was used to examine the association between the continuous distal outcomes and latent class

membership controlling for the confounding variables. Results are reported as

unstandardised beta coefficients and 95% confidence intervals.

Missing data

6,888 participants (49%) provided self-report information on binge drinking on at least

one occasion. Of these, *n*=3,755 (55%) had available information on all covariates, while

n=2,086 (30%) had data on at least one outcome at age 24 years and all confounders. See

Table S1 for a detailed description of attrition.

Mendelian randomisation (MR)

Two-sample MR was used to test the hypothesised causal effect of alcohol use on

cognitive functioning. The two-sample MR approach requires summary level data from

GWAS, enabling SNP-outcome and SNP-exposure effects to be derived from different data

sources. As the genetic instrument for alcohol consumption we used the 99-independent

genome-wide significant SNPs associated with 'number of drinks per week', identified by the

GWAS & Sequencing Consortium of Alcohol and Nicotine use (GSCAN

https://gscan.sph.umich.edu/) based on a sample of $N\sim1,000,000$. As outcomes, we used

GWAS conducted in ALSPAC for each of our three primary outcome measures: i) working

memory assessed using *d'*; ii) response inhibition assessed using SSRT; and iii) emotion

recognition assessed using total number of recognised emotions.

Analyses were performed using the TwoSampleMR R package, part of MR-Base

(38). The inverse-variance weighted (IVW) approach was used as a primary analysis, with

three complementary estimation methods as sensitivity analyses which each make different

assumptions about the nature of horizontal pleiotropy (where the genetic variant associates

with the outcome via an independent pathway to the exposure): MR Egger (39), weighted

median (40), and weighted mode (41). To test the suitability of the MR Egger method, the I^2_{GX}

statistic was calculated to quantify the degree of regression dilution bias due to measurement

error of SNP-exposure effects (42). The mean F statistic as an indicator of instrument strength

was also calculated. Steiger filtering was conducted to confirm the direction of effect (43). A

consistent effect across all of these methods would provide the most confidence that any

observed effects are not due to pleiotropy.

RESULTS

Observational analyses

Patterns of binge drinking

The prevalence of both monthly and weekly binge drinking increased across time

apart from a slight decrease at age 23 years (Table S2). There was good agreement that a

four-class solution was adequate in explaining the heterogeneity in binge drinking based on

increasing Bayesian Information Criterion (BIC) (44) and sample size adjusted Bayesian

Information Criterion (SSABIC) (45) values in the five-class model and an LRT value of *p*=0.05.

See Table S3 in the Supplementary Material for a comparison of model fit indices. The four-

class solution (Figure 1) comprised patterns of binge drinking that were labelled as 'low-risk'

(47.1%), 'adolescent-limited' (19.0%), 'adult-regular' (18.7%), and 'early-onset regular'

(19.0%). See Table S4 in the Supplementary Material for class validation. See Table S5 in the

Supplementary Material for a detailed description of confounding factors associated with

binge drinking class membership.

226   [Figure 1]

Working memory – 2-back task

Table 1 presents unadjusted and adjusted associations between patterns of binge

drinking from 16 to 23 years and working memory at age 24. There were unadjusted

associations for adult-regular (b=0.21, 95% CI=0.03 to 0.40) and early-onset regular (b=0.19,

95% CI=0.01 to 0.36) being associated with better working memory performance compared

to the low-risk binge drinking group. These associations were attenuated after adjustment for

sociodemographic confounders (adult-regular: b=0.14, 95% CI=-0.05 to 0.33; early-onset

regular: b=0.14, 95% CI=-0.07 to 0.30); working memory at age 11 years and participants who

experienced a head injury/unconsciousness before age 11 (adult-regular: b=0.12, 95% CI=-

0.06, 0.31; early-onset regular: b=0.06, 95% CI=-0.12 to 0.24), and cannabis and/or tobacco

use (adult-regular: b=0.12, 95% CI=-0.07 to 0.31; early-onset regular: b=0.09, 95% CI=-0.10 to

0.28). There was no evidence of an association between other patterns of binge drinking and

working memory performance. Furthermore, there was little evidence to suggest that

patterns of binge drinking were associated with the secondary outcomes (i.e., number of hits

and false alarms). Results of the secondary analyses are provided in Table S6 in the

Supplementary Material.

243   [Table 1]

Response inhibition - stop signal task

Table 2 presents unadjusted and adjusted associations between patterns of binge

drinking and an overall measure of response inhibition. Although, there was evidence for

adolescent-limited being associated with longer SSRT, indicating slower ability to inhibit

responses (b=19.65, 95% CI=3.20 to 36.09) compared to the low-risk binge drinking group in

the unadjusted models, omnibus Wald test values provided weak evidence of an overall

association between binge drinking patterns and response inhibition in the fully adjusted

model (Wald test=6.51 (3) p=0.09). There was no evidence of an association between other

patterns of binge drinking and ability to inhibit responses. Furthermore, there was little

evidence to suggest that patterns of binge drinking were associated with any of the secondary

outcomes (i.e., Go reaction time, Go accuracy, and Stop accuracy). Results of the secondary

analyses are provided in Table S7 in the Supplementary Material.

256   [Table 2]

Emotion recognition – 6AFC task

Table 3 presents unadjusted and adjusted associations between patterns of binge

drinking and number of correctly identified emotions. Although there was evidence for adult-

regular being associated with better performance on the task (b=0.02, 95% CI=0.00 to 0.03)

compared to the low-risk binge drinking group in the unadjusted models, these associations

were attenuated when controlling for confounders (fully adjusted model: b=0.01, 95% CI=-

0.00 to 0.03). There was no evidence of an association between other patterns of binge

drinking and number of correct hits. Furthermore, there was little evidence to suggest that

patterns of binge drinking were associated with any of the secondary outcomes (i.e., anger,

disgust, fear, happy, sad, and surprise). Results of the secondary analyses are provided in

Table S8 in the Supplementary Material.

268   [Table 3]

Mendelian randomisation

GWAS did not identify any genome-wide significant SNPs (at the $p<5\times10^{-8}$ level) for

any of the three cognitive outcomes, therefore we report the lead SNPs at a $p<5\times10^{-6}$

threshold (Tables S9-S11). The full distribution of results can be seen in Manhattan plots

(Figures S3-S5). QQ plots, and corresponding lambda (λ) statistics, are displayed in Figures S6-

S8. The lead SNP for working memory was rs7992404 (chromosome 13, b=-0.17, CI=-0.20 to

-0.14, effect allele=T, p=8.76x10^{-8}, intergenic variant), emotion recognition: rs199934102

(chromosome 6, b=0.34, CI=0.28 to 0.40, effect allele=A, p=8.19 x 10^{-8}, intronic variant within

gene CNKSR3), and response inhibition: rs9853329 (chromosome 3, b=-0.16, CI=-0.19 to-0.13,

effect allele=T, p=5.33x10^{-7}, intergenic variant).

The two-sample MR method provides little evidence to suggest that alcohol use (SNPs

associated with number of drinks per week) is a causal risk factor for deficits in cognitive

functioning (Table 4). Focusing on the IVW estimate as the primary measure, SNPs associated

with the number of alcoholic drinks per week were not associated with *d'* on the working

memory task (b=0.137 95% CI=-0.45 to 0.72; p=0.65); SSRT on the response inhibition task

(b=-0.313 95%CI=-0.92 to 0.29; p=0.31); or total hits in the emotion recognition task (b=0.028

95% CI=-0.55 to 0.61; p=0.93). There was no evidence of heterogeneity in the SNP-exposure

association (Tables S12 and S13 in the Supplementary material).

287   [Table 4]

DISCUSSION

We found insufficient evidence to suggest an association between binge drinking

between the ages of 16 and 23 and cognitive deficits at age 24 using a combination of

observational and genetic approaches. In the observational analyses, four patterns of binge

drinking across adolescence and early adulthood were identified. These patterns were not

associated with later measures of cognitive functioning in either the unadjusted or adjusted

models. In the genetic analyses, there was no evidence to suggest that number of drinks

consumed per week was a causal risk factor for deficits in cognitive functioning in the ALSPAC

sample.

Comparison with previous studies

To the best of our knowledge, this is the first study to assess the relationship between

alcohol use and cognitive functioning using a combination of observational and genetic

epidemiological approaches. Unlike the studies that have demonstrated a prospective

association between binge drinking and cognitive functioning in adolescence e.g. (17–20), we

found little evidence in support of this association. The contrast in findings could be due to a

number of possibilities. First, our study used a large population sample incorporating data

from over 2,000 participants spanning maternal pregnancy to 24 years of age. Previous

studies (19,20) used functional magnetic resonance imaging based on youths at high-risk for

substance use disorders (*n*=40 for both studies), while the study by Peeters and colleagues

(18) used a sample at high-risk for externalising problems (*n*=374 at baseline). Second, this

study assessed cognitive functioning in early adulthood, a time when cognitions are thought

to reach maturity (2–5), in comparison to all four studies who assessed cognitive functioning

up to 19 years of age. Examining peak levels of cognitive functioning helps to reduce the

possibility that cognitive functioning is influencing earlier alcohol use (i.e., reverse causation).

Third, as alcohol use behaviours typically change over time (46), repeated measures of binge

drinking were used in this study to capture heterogenous patterns across this sensitive period

in comparison to our previous study which assessed alcohol use on one occasion (17). Finally,

most of the previous studies (18–20) assessed cognitive functioning using a range of measures

compared to the more widely used measures in this study (i.e., *N*-back task, Stop signal task,

and 6AFC task).

Our findings support and extend those of Boelema and colleagues (47) who found

insufficient evidence to suggest that alcohol use prospectively affected maturation of

cognitive functions in a large prospective study of Dutch adolescents (48). Frist, Boelema and

colleagues examined cognitive functioning across adolescence, while we were able to

examine peak levels of cognitive functioning. Second, assessing binge drinking in young

adulthood allowed us to capture heterogenous patterns during the sensitive period (i.e.,

going to University or in full-time paid employment). Finally, WM performance was measured

in reaction times only, as opposed to the more comprehensive approach used in our study

(i.e., number of hits, false alarms, and *d'*).

Limitations

There are a number of limitations to this study. First, the ALSPAC cohort suffers from

attrition which reduces study power and also is higher among the socially disadvantaged (49)

(Table S1). We attempted to minimise the impact of attrition using sensitivity analyses. It is

possible that both the observational and two-sample MR analyses are underpowered. For

example, poor entropy (a measure of class separation) in the observational analyses could

indicate poor misclassification in the latent classes resulting in dilution of effect estimates.

Small sample size in the cognitive functioning GWAS is likely to be underpowered in the two-

sample MR analyses. Future studies should use larger observational samples and meta-

analyses of related cognitive measures in GWAS to help to increase power. Second, although

it is possible to either under- or over-estimate drinking behaviour using self-reported data,

participants completed questionnaires individually and were assured of their anonymity.

Third, although there is some debate in the literature surrounding the construct validity of

performance on the *N*-back task as an indicator of working memory ability, it has been argued

that using *N*-back performance indices from a signal-detection framework (i.e. d') may reveal

clearer insights about its validity as a measure of working memory performance (50–52).

Fourth, as we examined one potential causal pathway, it is possible that the

association could work in the opposite direction, that is, impairments in cognitive functioning

may precede (and increase the risk of developing) alcohol problems (18,23). We were

however able to include a number of measures to maximise the robustness of our findings:

(i) ascertaining the time order of exposures and outcomes; (ii) controlling for premorbid

working memory function and brain insults prior to the onset of alcohol use; and (iii) a number

of relevant confounders were included to help reduce the possibility of residual confounding.

It is possible that a common risk factor is influencing both binge drinking and deficits in

cognition, however the MR analyses helps to protect against this possibility by minimising

bias from reverse causation and residual confounding. Finally, genetic variants were based on

number of drinks per week, whereas the observational analyses used frequency of binge

drinking. Although not directly comparable, there was evidence of a dose-response

relationship between binge drinking patterns and the AUDIT-C measure, which taps into

quantity and frequency (see Supplementary material).

Implications and Conclusions

In order to rule out the possibility of deficient cognitive functioning preceding binge

drinking in adolescence, future research should use an equally robust approach to examine

the alternate hypothesis. We found insufficient evidence to suggest an association between

binge drinking between the ages of 16 and 23 and cognitive deficits at age 24 using a

combination of observational and genetic approaches, although both approaches are likely

to be underpowered.

Acknowledgements: The work was undertaken with the support of the MRC and Alcohol

Research UK (grant number MR/L022206/1). We acknowledge also support from The Centre

for the Development and Evaluation of Complex Interventions for Public Health Improvement

(DECIPHer); a UKCRC Public Health Research Centre of Excellence (joint funding (grant

number MR/KO232331/1) from the British Heart Foundation, Cancer Research UK, Economic

and Social Research Council, Medical Research Council, the Welsh Government and the

Wellcome Trust, under the auspices of the UK Clinical Research Collaboration); and the NIHR

School of Public Health Research. Support was also provided by the UK Medical Research

Council Integrative Epidemiology Unit at the University of Bristol (MM_UU_00011/7). LM,

RW, SS, and MRM are members of the UK Centre for Tobacco and Alcohol Studies, a UKCRC

Public Health Research Centre of Excellence. LM, REW, SS, and MRM are supported by the

NIHR Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust

and the University of Bristol. The views expressed in this publication are those of the authors

and not necessarily those of the NHS, the National Institute for Health Research or the

Department of Health and Social Care. GWAS data were generated by Sample Logistics and

Genotyping Facilities at Wellcome Sanger Institute and LabCorp (Laboratory Corporation of

America) using support from 23andMe. The UK Medical Research Council and Wellcome

(Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This

publication is the work of the authors and LM and MRM will serve as guarantors for the

contents of this paper. A comprehensive list of grants funding is available on the website

(http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf).   We

are extremely grateful to all the families who took part in this study, the midwives for their

help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer

and laboratory technicians, clerical workers, research scientists, volunteers, managers,

receptionists and nurses. This publication is the work of the authors who will serve as

guarantors for the contents of this paper.

REFERENCES

1.   De Bellis MD, Clark DB, Beers SR, Soloff PH, Boring AM, Hall J, et al. Hippocampal

volume in adolescent-onset alcohol use disorders. Am J Psychiatry. 2000;157(5):737–

44.

2.   Fry AF, Hale S. Relationships among processing speed, working memory, and fluid

intelligence in children. Biol Psychol. 2000;54(1–3):1–34.

3.   Kramer AF, de Sather JCMG, Cassavaugh ND. Development of attentional and

oculomotor control. Dev Psychol. 2005;41(5):760–72.

4.   Davidson MC, Amso D, Anderson LC, Diamond A. Development of cognitive control

and executive functions from 4 to 13 years: evidence from manipulations of memory,

inhibition, and task switching. Neuropsychologia. 2006;44(11):2037–78.

5.   Thomas LA, De Bellis MD, Graham R, LaBar KS. Development of emotional facial

recognition in late childhood and adolescence. Dev Sci. 2007;10(5):547–58.

6.   Sowell ER, Delis D, Stiles J, Jernigan TL. Improved memory functioning and frontal

lobe maturation between childhood and adolescence: a structural MRI study. J Int

Neuropsychol Soc. 2001;7(3):312–22.

7.   Sowell ER, Thompson PM, Holmes CJ, Jernigan TL, Toga AW. In vivo evidence for post-

adolescent brain maturation in frontal and striatal regions. Nat Neurosci. 1999;2:859.

8.   White AM, Ghia AJ, Levin ED, Swartzwelder HS. Binge pattern ethanol exposure in

adolescent and adult rats: differential impact on subsequent responsiveness to

ethanol. Alcohol Clin Exp Res. 2000;24(8):1251–6.

9.   Risher M-L, Fleming RL, Boutros N, Semenova S, Wilson WA, Levin ED, et al. Long-

term effects of chronic intermittent ethanol exposure in adolescent and adult rats:

radial-arm maze performance and operant food reinforced responding. PLoS One.

2013;8(5):e62940–e62940.

10.   Crews F, He J, Hodge C. Adolescent cortical development: a critical period of

vulnerability for addiction. Pharmacol Biochem Behav. 2007;86(2):189–99.

11.   Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in

adolescence: a critical period of addiction vulnerability. Am J Psychiatry.

2003;160(6):1041–52.

12.   Squeglia LM, Schweinsburg AD, Pulido C, Tapert SF. Adolescent binge drinking linked

to abnormal spatial working memory brain activation: differential gender effects.

Alcohol Clin Exp Res. 2011;35(10):1831–41.

13.   Wilson S, Malone SM, Thomas KM, Iacono WG. Adolescent drinking and brain

morphometry: A co-twin control analysis. Dev Cogn Neurosci. 2015;16:130–8.

14.   Malone SM, Luciana M, Wilson S, Sparks JC, Hunt RH, Thomas KM, et al. Adolescent

drinking and motivated decision-making: a cotwin-control investigation with

monozygotic twins. Behav Genet. 2014;44(4):407–18.

15.   Parada M, Corral M, Mota N, Crego A, Rodriguez Holguin S, Cadaveira F. Executive

functioning and alcohol binge drinking in university students. Addict Behav.

2012;37(2):167–72.

16.   Gullo MJ, Loxton NJ, Price T, Voisey J, Young RM, Connor JP. A laboratory model of

impulsivity and alcohol use in late adolescence. Behav Res Ther. 2017;97:52–63.

17.   Mahedy L, Field M, Gage S, Hammerton G, Heron J, Hickman M, et al. Alcohol use in

adolescence and later working memory: Findings from a large population-based birth

cohort. Alcohol Alcohol. 2018;53:251–8.

18.   Peeters M, Monshouwer K, Janssen T, Wiers RW, Vollebergh WAM. Working memory

and alcohol use in at-risk adolescents: a 2-year follow-up. Alcohol Clin Exp Res.

2014;38(4):1176–83.

19.   Squeglia LM, Pulido C, Wetherill RR, Jacobus J, Brown GG, Tapert SF. Brain response

to working memory over three years of adolescence: influence of initiating heavy

drinking. J Stud Alcohol Drugs. 2012;73(5):749–60.

20.   Wetherill RR, Squeglia LM, Yang TT, Tapert SF. A longitudinal examination of

adolescent response inhibition: neural differences before and after the initiation of

heavy drinking. Psychopharmacology (Berl). 2013;230(4):663–71.

21.   Peeters M, Janssen T, Monshouwer K, Boendermaker W, Pronk T, Wiers R, et al.

Weaknesses in executive functioning predict the initiating of adolescents' alcohol use.

Dev Cogn Neurosci. 2015;16:139–46.

22.   Squeglia LM, Jacobus J, Nguyen-Louie TT, Tapert SF. Inhibition during early

adolescence predicts alcohol and marijuana use by late adolescence.

Neuropsychology. 2014;28(5):782–90.

23.   Fernie G, Peeters M, Gullo MJ, Christiansen P, Cole JC, Sumnall H, et al. Multiple

behavioural impulsivity tasks predict prospective alcohol involvement in adolescents.

Addiction. 2013;108(11):1916–23.

24.   Liu M, Jiang Y, Wedow R, Li Y, Brazel DM, Chen F, et al. Association studies of up to

1.2 million individuals yield new insights into the genetic etiology of tobacco and

alcohol use. Nat Genet. 2019;51(2):237–44.

25.   Lawlor DA, Harbord RM, Sterne JAC, Timpson N, Davey Smith G. Mendelian

randomization: using genes as instruments for making causal inferences in

epidemiology. Stat Med. 2008;27(8):1133–63.

26.   Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile:

the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of

Parents and Children. Int J Epidemiol. 2013;42(1):111–27.

27.   Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al.

Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers

cohort. Int J Epidemiol. 2013;42(1):97–110.

28.   Northstone K, Lewcock M, Groom A, Boyd A, Macleod J, Timpson N, et al. The Avon

Longitudinal Study of Parents and Children (ALSPAC): an update on the enrolled

sample of index children in 2019. Wellcome Open Res. 2019;4:51.

29.   Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic

data capture (REDCap)--a metadata-driven methodology and workflow process for

providing translational research informatics support. J Biomed Inform.

2009;42(2):377–81.

30.   Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, Neal LO, et al. The REDCap

consortium: Building an international community of software platform partners. J

Biomed Inform. 2019;95(5):103208.

31.   Kirchner WK. Age differences in short-term retention of rapidly changing information.

J Exp Psychol. 1958;55(4):352–8.

32.   Wardle MC, de Wit H, Penton-Voak I, Lewis G, Munafo MR. Lack of association

between COMT and working memory in a population-based cohort of healthy young

adults. Neuropsychopharmacology. 2013;38(7):1253–63.

33.   Rossi R, Zammit S, Button KS, Munafò MR, Lewis G, David AS. Psychotic experiences

and working memory: a population-based study using signal-detection analysis. PLoS

One. 2016;11(4):e0153148–e0153148.

34.   Logan GD, Cowan WB, Davis KA. On the ability to inhibit simple and choice reaction

time responses: a model and a method. J Exp Psychol Hum Percept Perform.

1984;10(2):276–91.

35.   Penton-Voak IS, Bate H, Lewis G, Munafo MR. Effects of emotion perception training

on mood in undergraduate students: randomised controlled trial. Br J Psychiatry.

2012;201(1):71–2.

36.   Muthén LK, Muthén BO. User's Guide. 7th ed. Muthén LK, Muthén BO, editors. Los

Angeles, CA: Muthén & Muthén; 2016.

37.   Bolck A, Croon M, Hagenaars J. Estimating latent structure models with categorical

variables: one-step versus three-step estimators. Polit Anal. 2004;12(1):3–27.

38.   Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, et al. The MR-Base

platform supports systematic causal inference across the human phenome. Elife.

2018 May;7.

39.   Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid

instruments: effect estimation and bias detection through Egger regression. Int J

Epidemiol. 2015;44(2):512–25.

40.   Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in Mendelian

randomization with some invalid instruments using a weighted median estimator.

Genet Epidemiol. 2016;40(4):304–14.

41.   Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian

randomization via the zero modal pleiotropy assumption. Int J Epidemiol.

2017;46(6):1985–98.

42.   Bowden J, Del Greco M F, Minelli C, Davey Smith G, Sheehan NA, Thompson JR.

Assessing the suitability of summary data for two-sample Mendelian randomization

analyses using MR-Egger regression: the role of the I2 statistic. Int J Epidemiol.

2016;45(6):1961–74.

43.   Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between

512   imprecisely measured traits using GWAS summary data. PLOS Genet.

513   2017;13(11):e1007081.

514   44.   Schwarz G. Estimating the dimension of a model. Ann Stat. 1978;6:461–4.

515   45.   Sclove L. Application of model-selection criteria to some problems in multivariate

516   analysis. Psychometrika. 1987;52:333–43.

517   46.   Faden VB. Trends in initiation of alcohol use in the United States 1975 to 2003.

518   Alcohol Clin Exp Res. 2006;30(6):1011–22.

519   47.   Boelema SR, Harakeh Z, Van Zandvoort MJE, Reijneveld SA, Verhulst FC, Ormel J, et al.

520   Adolescent heavy drinking does not affect maturation of basic executive functioning:

521   Longitudinal findings from the TRAILS study. PLoS One. 2015;10(10):1–15.

522   48.   Ormel J, Oldehinkel AJ, Sijtsema J, van Oort F, Raven D, Veenstra R, et al. The

523   TRacking Adolescents' Individual Lives Survey (TRAILS): design, current status, and

524   selected findings. J Am Acad Child Adolesc Psychiatry. 2012;51(10):1020–36.

525   49.   Wolke D, Waylen A, Samara M, Steer C, Goodman R, Ford T, et al. Selective drop-out

526   in longitudinal studies and non-biased prediction of behaviour disorders. Br J

527   Psychiatry. 2009;195(3):249–56.

528   50.   Kane MJ, Conway ARA, Miura TK, Colflesh GJH. Working memory, attention control,

529   and the N-back task: a question of construct validity. J Exp Psychol Learn Mem Cogn.

530   2007;33(3):615–22.

531   51.   Meule A. Reporting and interpreting working memory performance in n-back tasks.

532   Front Psychol. 2017;8:352.

533   52.   Haatveit BC, Sundet K, Hugdahl K, Ueland T, Melle I, Andreassen OA. The validity of d

534   prime as a working memory index: results from the "Bergen n-back task". J Clin Exp

535   Neuropsychol. 2010;32(8):871–80.

Figure 1. Distribution of binge drinking response across latent classes at each timepoint (*n*=6,888). Class proportions based on estimated posterior probability[1]

---

[1] Overall, the low-risk group reported a low probability of binge drinking across all measurement occasions; adolescent-limited binge drinkers was mostly characterised by binge drinking in the earlier measurement occasions (but not later ones), adult-regular binge drinkers were mostly characterised by binge drinking in the later measurement occasions (but not earlier ones), while early-onset regular binge drinking was mostly characterised by binge drinking across all timepoints.

Table 1. Patterns of binge drinking from 16 to 23 years and working memory at age 24 (higher *d'* scores reflect better performance)

	Low risk	Adolescence limited	Adult limited	Early onset regular	
n=2,086 for all models	Reference group	*b* (95% CI)	*b* (95% CI)	*b* (95% CI)	Wald (df) *p* value
Unadjusted models					
Working memory *d'*	-	-0.15 (-.46, .17)	0.21 (.03, .40)	0.19 (.01, .36)	9.56 (3) *p*=0.02
Adjusted for SES					
Working memory *d'*	-	-0.15 (-.47, .17)	0.14 (-.05, .33)	0.14 (-.07, .30)	3.82 (3) *p*=0.28
Adjusted for SES/WM/HI					
Working memory *d'*	-	-0.17 (-.49, .15)	0.12 (-.06, .31)	0.06 (-.12, .24)	3.24 (3) *p*=0.36
Fully adjusted models					
Working memory *d'*	-	-0.15 (-.47, .17)	0.12 (-.07, .31)	0.09 (-.10, .28)	3.20 (3) *p*=0.36

Note. SES: socioeconomic status; WM: working memory at age ~11 years; HI: head injury/ unconsciousness up to age 11 years

Table 2. Patterns of binge drinking from 16 to 23 years and response inhibition at age 24 (shorter scores reflect faster reaction times)

n=2,086 for all models	Low risk Reference group	Adolescence limited b (95% CI)	Adult limited b (95% CI)	Early onset regular b (95% CI)	Wald (df) p value
Unadjusted models					
Stop signal reaction time	-	18.58 (2.62, 34.54)	-7.73 (-17.25, 2.09)	-5.42 (-15.62, 4.77)	10.14 (3) p=0.02
Adjusted for SES					
Stop signal reaction time	-	19.39 (3.33, 35.44)	-4.07 (-14.48, 6.34)	-2.66 (-13.30, 7.99)	6.92 (3) p=0.08
Adjusted for SES/WM/HI					
Stop signal reaction time	-	20.06 (4.07, 36.05)	-3.57 (-13.82, 6.69)	-0.05 (-10.72, 10.61)	7.40 (3) p=0.06
Fully adjusted model					
Stop signal reaction time	-	19.65 (3.20, 36.09)	-3.57 (-13.82, 6.69)	-0.70 (-11.95, 10.55)	6.51 (3) p=0.09

Note. SES: socioeconomic status; WM: working memory at age ~11 years; HI: head injury/ unconsciousness up to age 11 years

Table 3. Patterns of binge drinking from 16 to 22 and emotion recognition at age 24 (higher scores reflect better performance)

n=2,086 for all models	Low risk Reference group	Adolescence limited *b* (95% CI)	Adult limited *b* (95% CI)	Early onset regular *b* (95% CI)	Wald (df) *p* value
Unadjusted models					
Total hits	-	-0.02 (-.04, .01)	0.02 (.00, .03)	0.01 (-.01, .02)	8.14 (3) *p*=0.04
Adjusted for SES					
Total hits	-	-0.01 (-.04, .01)	0.01 (-.00, .02)	0.01 (-.01, .02)	4.17 (3) *p*=0.24
Adjusted for SES/WM/HI					
Total hits	-	-0.01 (-.04, .01)	0.01 (-.00, .02)	0.00 (-.01, .02)	3.89 (3) *p*=0.27
Fully adjusted model					
Total hits	-	-0.01 (-.04, .01)	0.01 (-.00, .03)	0.01 (-.01, .02)	3.68 (3) *p*=0.30

Note. SES: socioeconomic status; WM: working memory at age ~11 years; HI: head injury/ unconsciousness up to age 11 years

Table 4. Two-sample Mendelian randomization analyses of the effects of alcohol use on cognitive functioning

Exposure	Outcome	Method	N SNPs	Beta (95% CI)	P-value
Drinks per week	**Working memory**	Inverse-Variance Weighted	88	0.137 (-0.45, 0.72)	0.65
		MR Egger (SIMEX)	88	-0.381 (-2.13, 01.37)	0.67
		Weighted Median	88	0.421 (-0.52, 1.37)	0.38
		Weighted Mode	88	0.767 (-0.79, 2.32)	0.34
Drinks per week	**Response inhibition**	Inverse-Variance Weighted	88	-0.313 (-0.92, 0.29)	0.31
		MR Egger (SIMEX)	88	-0.729 (-2.63, 1.17)	0.46
		Weighted Median	88	-0.728 (-1.67, 0.21)	0.13
		Weighted Mode	88	-0.895 (-2.47, 0.68)	0.27
Drinks per week	**Emotion recognition**	Inverse-Variance Weighted	88	0.028 (-0.55, 0.61)	0.93
		MR Egger (SIMEX)	88	0.445 (-1.14, 2.03)	0.58
		Weighted Median	88	-0.157 (-1.01, 0.69)	0.72
		Weighted Mode	88	-0.180 (-1.78, 1.42)	0.82

Note: SIMEX = simulation extrapolation. SIMEX-corrected estimates were used based on regression dilution I^2_{GX} for number of drinks per week values between 0.3 and 0.9 (see Supplementary Table S9). SIMEX-corrected estimates are unweighted.