Eating behavior trajectories in the first ten years of life and their relationship with BMI

Dr Moritz Herle[1,2], Prof Bianca De Stavola[1], Christopher Hübel[3,4,5], Dr Diana L Santos

Ferreira[6,7], Mohamed Abdulkadir[8], Dr Zeynep Yilmaz[9], Prof Ruth Loos[10], Dr Rachel Bryant-

Waugh[1,11], Prof Cynthia M. Bulik[5,9,12] & Dr Nadia Micali[1,8,12]

[1] Great Ormond Street Institute of Child Health, University College London, London, UK

[2] Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology &
Neuroscience at King's College London, United Kingdom

[3] Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology &
Neuroscience, King's College London, UK

[4] UK National Institute for Health Research (NIHR) Biomedical Research Centre for Mental
Health, South London and Maudsley Hospital, London, UK

[5] Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,
Sweden

[6] Medical Research Council Integrative Epidemiology, University of Bristol, Bristol, UK
[7] Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
[8] Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva, Switzerland

[9] Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC,
USA

[10] Icahn Mount Sinai School of Medicine, New York, NY, USA

[11] Great Ormond Street Hospital, London, UK

[12] Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC,
USA

[12] Child and Adolescent Psychiatry Division, Department of Child and Adolescent Health,
Geneva University Hospital, Geneva, Switzerland

Names for PubMed Index: Herle, De Stavola, Hübel, Santos Ferreira, Abdulkadir, Yilmaz,
Loos, Bryant-Waugh, Bulik, Micali

Corresponding author:

Dr Nadia Micali MD, MRCPsych, PhD, FAED

Senior Lecturer and Honorary Consultant Psychiatrist
Child and Adolescent Mental Health, Palliative Care and Paediatrics Section
Population, Policy and Practice Research Theme
UCL Institute of Child Health
30 Guilford Street
London WC1N 1EH

Tel: 020 7905 2163
Fax: 020 7831 7050
e-mail: n.micali@ucl.ac.uk

Running title: Patterns of child eating behaviors and later BMI

Sources of support: This work was specifically funded by the UK Medical Research Council
and the Medical Research Foundation (ref: MR/R004803/1).

Financial Disclosure: Bulik reports: Shire (grant recipient, Scientific Advisory Board
member) and Pearson and Walker (author, royalty recipient).

Abbreviations: Eating behaviors: EB, Body Mass Index: BMI, Avon Longitudinal Study of
Parents and Children: ALSPAC, Latent Class Growth Trajectories: LCGA,
Avoidant/restrictive food intake disorder: ARFID

Abstract

Background: Child eating behaviors are highly heterogeneous and their longitudinal impact

on childhood weight is unclear. The objective of this study was to characterize eating

behaviors during the first ten years of life and evaluate associations with BMI at age 11 years.

Method: Data were parental reports of eating behaviors from 15 months to age 10 years

(n=12,048) and standardized body mass index (zBMI) at age 11 years (n=4884) from the

Avon Longitudinal Study of Parents and Children. Latent class growth analysis was used to

derive latent classes of over-, under-, and fussy eating. Linear regression models for zBMI at

11 years on each set of classes were fitted to assess associations with eating behavior

trajectories.

Results: We identified four classes of overeating; "low stable" (70%), "low transient" (15%),

"late increasing" (11%), and "early increasing" (6%). The "early increasing" class was

associated with higher zBMI (boys: β=0.83, 95%CI:0.65, 1.02; girls: β=1.1; 0.92, 1.28)

compared to "low stable". Six classes were found for undereating; "low stable" (25%), "low

transient" (37%), "low decreasing" (21%), "high transient" (11%), "high decreasing" (4%),

and "high stable" (2%). The latter was associated with lower zBMI (boys: β=-0.79; -1.15, -

0.42; girls: β=-0.76; -1.06, -0.45). Six classes were found for fussy eating; "low stable"

(23%), "low transient" (15%), "low increasing" (28%), "high decreasing" (14%), "low

increasing" (13%), "high stable" (8%). The "high stable" class was associated with lower

zBMI (boys: β =-0.49; -0.68 -0.30; girls: β =-0.35; -0.52, -0.18).

Conclusions: Early increasing overeating during childhood is associated with higher zBMI at

age 11. High persistent levels of undereating and fussy eating are associated with lower

zBMI. Longitudinal trajectories of eating behaviors may help identify children potentially at

risk of adverse weight outcomes.

Introduction

Child eating behaviors have received attention, especially due to their potential association

with weight. However, previous cross-sectional and a limited number of longitudinal studies

produced inconsistent findings. Previous research has suggested that some eating behaviors

are stable across childhood, as indicated by moderate correlations between eating behaviors

at age 4 and 10 in English (1) and Dutch samples (2), as well as in younger children, between

two and five years (3). However, these studies only had access to two data points, precluding

a comprehensive examination stability and change. Some eating behaviors, such as fussy

eating, which is the tendency to eat only certain foods and to refuse to try new foods, are

common and potentially more transient (1). A previous study reported that one third of

children exhibit some fussiness during the first four years of life, but many tend to remit by

age six with about 4% being persistently fussy.(5) More recently, a study of the same cohort

as discussed in this paper, the Avon Longitudinal Study of Parents and Children, found that

mothers indicated that more than half of the children at 15 months were fussy about what

foods to eat.(2)

Cross-sectional studies (3-5) have primarily suggested that eating behaviors, such as

responsiveness to external food cues or emotional overeating are associated with higher child

weight. Other eating behaviors, such as fussy eating and responsiveness to internal satiety

cues are associated with lower weight (6-8). However, other cross-sectional studies have not

replicated these findings (9, 10). Longitudinally, eating behaviors measured at 5-6 years are

weakly associated with body mass index (BMI) at about 6-8 years (11) . In earlier ages,

between 3 months and 9-15 months, a bidirectional association between child eating and

weight has been reported.(12) More recently, the bidirectional association between child

eating and later BMI was replicated in a sample of Norwegian children, aged 4 to 8 years(13).

Furthermore, children who display fussy eating appear to be at higher risk for developing

underweight in childhood, but may be at increased risk for later overweight(14, 15) .

However, some studies report no or only weak longitudinal relationships (16-18).

Overall, childhood eating behaviors and childhood weight outcomes and the longitudinal

development of child eating behaviors remains poorly understood. Longitudinal studies often

focus on overall mean scores, ignoring heterogeneity and transience of child eating behaviors.

We, therefore, aimed to investigate repeatedly measured eating behaviors in a large

population-based birth cohort using latent class modeling to identify longitudinal trajectories

during the first ten years of life. Furthermore, we examined their relationship with age- and

sex-standardized zBMI at age 11. This age was selected as the outcome measures, due to the

proximity to the derived trajectories and to ensure the largest and most representative sample

of prepubertal children. Our hypothesis was that persistent EB patterns in childhood would be

more strongly associated with child zBMI than transient ones.

Methods

Participants

Data from the Avon Longitudinal Study of Parents and Children (ALSPAC), a population

based, longitudinal cohort of mothers and their children born in the southwest of England (19,

20) were analyzed. All pregnant women expected to have children between the 1[st] April 1991

and 31[st] December 1992 were invited to enroll in the study, providing informed written

consent. From all pregnancies (n = 14,676), 14,451 mothers opted to take part; by one year

13,988 children were alive. When the oldest children were approximately 7 years of age, an

attempt was made to bolster the initial sample with eligible cases who had failed to join the

study originally (referred to as Phase 2), however these participants were not included in

these analyses. The phases of enrolment are described in more detail in the cohort profile

papers (19, 20). One sibling per set of multiple births, was randomly excluded from these

analyses to guarantee independence of participants.

Eating behaviors

Repeated measures of parent-reported child eating behaviors were available at a maximum of

eight time points around the age of 15, 24, 38, 54, 62, 81, 105, and 116 months. Parents were

asked if they were worried about their child overeating ("How worried are you because your

child is overeating"), and undereating ("How worried are you because your child is

undereating"). The remaining questions probed the child's tendency to be fussy ("How

worried are you because your child is choosy", "How worried are you because your child has

feeding difficulties", "How worried are you because your child is refusing food"). Parents

were given the following response options: "no/did not happen", "not worried", "a bit

worried" and "greatly worry". The two top categories ("a bit worried" and "greatly worry")

were combined to avoid very low frequencies. Children who had at least one measure of any

of the items were included in the analyses (N=12,048). About half (45%) of the included

children had data on all 8 time points and ~85% had data at least 3 time points.

Anthropometric data

Weight and height were measured during clinic visits when the children (N = 4,885) were 11

years old (mean=128.6 months, SD=1.64). Height was measured to the nearest millimeter

with the use of a Harpenden Stadiometer (Holtain Ltd.). Weight was measured with a Tanita

Body Fat Analyzer (Tanita TBF UK Ltd.) to the nearest 50g. BMI was calculated by dividing

weight (in kg) by height squared (in m). Age- and sex-standardized BMI z-scores (zBMI)

were calculated according to UK reference data, indicating the degree to which a child is

heavier (>0) or lighter (<0) than expected according to his/her age and sex(21). We aimed to

relate the trajectories of eating behaviors with zBMI at age 11 years. Children with data on

both eating behavior and zBMI were included in the final stage of the analyses (N=4,884). A

comparison of the distribution of derived trajectories between participants with and without

BMI data at 11 years can be found in eTable 5.

Covariates

The following indicators of socioeconomic status of the family were used: Maternal age at

birth (years) and maternal education status (A-Levels or higher, lower than A-Levels) and

parental occupational status (manual, non-manual labor of the highest earner in the family).

Further birthweight (grams) and gestational age at birth (weeks) of the children were also

used. The indicators of socioeconomic status were treated as potential confounders for the

analyses of zBMI and as predictors of missing data for parent-reported EB data. Details of all

data are available through a fully searchable data dictionary at

www.bristol.ac.uk/alspac/researchers/our-data.

Statistical analyses

Analyses were conducted from October 2017 to May 2018 and included two stages in line

with the classify-analyze framework (22).

First, Latent Class Growth Analysis (LCGA) was used to identify subgroups ("latent

classes") of children who share the same trajectories of eating behaviors (23). In comparison

to Growth Mixture Modelling, an alternative approach to identifying these latent classes,

LCGA constrains the variation within each class to zero, reducing the number of parameters

and simplifying model estimation (23). LCGA was conducted using Full Information

Maximum Likelihood (FIML) (24), incorporating indicators of social class (maternal age,

maternal education, and manual or non-manual labor of the highest earner in the family) as

auxiliary variables to account for the missingness (including attrition) affecting the

longitudinal data, as previously described in ALSPAC (19). FIML assumes data are missing

at random (MAR), once these auxiliary variables are accounted for and therefore children

with at least one measure of eating behavior at any time point. Analyses were stratified by

sex to examine possible effect modification. Stratified results were compared against

unstratified using combined data using Likelihood Ratio Tests. As the number of classes is

not directly estimated, alternative specifications with increasing number of assumed classes

were compared using the following model fit indicators: Akaike Information Criterion (AIC),

Bayesian Information Criterion (BIC), adjusted Bayesian information Criterion (adj BIC),

selecting the lowest values, and entropy, aiming for the highest. In addition to these model fit

indicators, the class size and interpretability of the classes were taken into account as

recommended by Muthén (24). After selection of the best number of classes, estimations

were repeated using 1000 random starts to avoid local maxima.

In the second stage, participants were allocated to their most likely classes according to their

posterior probabilities using the maximum-probability assignment rule (25). These predicted

classes were then included as explanatory variables in regression analyses of zBMI scores at

age 11, which also controlled for the following a priori confounders: maternal age,

gestational age, birthweight, and maternal education at birth. Results are reported in terms of

adjusted regression coefficients (β) for each class in comparison to the first (reference) class.

Since not all children with eating behavior data had data on zBMI, because of attrition

affecting later ages, the characteristics of study participants with/without zBMI were

compared in order to assess their representativeness of the original study membership (eTable

5). LCGA was conducted in MPlus Version 8 (26). Regression analyses were conducted in

Stata 15 (27). All code is available at https://github.com/MoritzHerle/Patterns-of-child-

eating-behaviors-and-later-BMI .

Ethical approval

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee

and the Local Research Ethics Committees. All procedures were performed in accordance

with the ethical standards laid down in the 1964 Declaration of Helsinki and its later

amendments.

Results

Summary statistics of the study population at baseline are listed in Table 1. Eating behaviors

varied at the different time points (Figure 1). Overeating was uncommon, with the majority of

parents reporting that their children never engaged in this behavior (77-85% across the 8 time

points). Being fussy about food was the most common child behavior, especially at 54

months, when a fifth of the children were described as fussy to a worrying extent.

TABLE 1 here

FIGURE 1 here

Eating behavior classes

Comparisons of alternative number of classes for the three LGCA models identified four

classes for the overeating longitudinal data, and six classes each for undereating and fussy

eating according to our pre-specified criteria (eTable 1a-c). Overall, separate models for boys

and girls fit the data better than when analyzed jointly (eTable 2).

Most children were assigned to the "low stable" class of overeating, marked by the absence

of high levels of overeating across time points. Undereating was more heterogeneous; the

most common class was "low transient", characterized by low levels of undereating, which

attenuated completely by age 10. Similarly, the most common class for fussy eating was the

"low transient" group, with increasing numbers of parents reporting fussy eating from 15

months onwards, which decreases again after 62 months. Figure 2a-c illustrates the class

trajectories for overeating, undereating, and fussy eating.

When comparing maternal and gestational age at birth, birthweight, and maternal education

across classes we found that, for overeating, boys and girls in the "early increasing" class had

a higher mean birthweight than those in the "low stable" class; eTable 3). Children in the

"high stable" class of undereating and fussy eating had a lower mean birthweight than their

respective "low stable" classes. In addition, the percentage of mothers with A-Levels or

university degree was lower in the "low stable" class of fussy eating compared to the "high

stable". Child zBMI scores per eating behavior class ranged widely within all classes

(eFigure 1a-b).

FIGURES 2A -2C HERE

Sensitivity analyses

Not all children included in LCGA had zBMI data at 11 years. Trajectory frequencies derived

from all participants were compared to the frequencies among the children who had complete

BMI data at 11 years. Trajectory sizes and distributions were similar (eTable 5).

Association between eating behaviors and zBMI at age 11

Overeating

In comparison to children who were reported to never overeat to a worrying extent ("low

stable"), all other classes were positively associated with greater zBMI at the later age of 11

years: "low transient" (boys: coefficient β=0.26, 95%CI:0.13, 0.39; girls: β=0.32, 95%CI:

0.19, 0.44), "late increasing" (boys: β = 0.94, 95%CI: 0.8, 1.09; girls: β=0.94, 95%CI: 0.82,

1.07) and "early increasing" (boys: β=0.83, 95%CI: 0.65, 1.02; girls: β=1.1, 95%CI: 0.92,

1.28; Table 2).

Undereating

In contrast, undereating classes were associated with lower zBMI. The magnitude of

associations in the "high transient" (boys: β=-0.25, 95%CI:-0.41, -0.08; girls: β=-0.24,

95%CI:-0.38, -0.09) and "high decreasing" (boys: β=-0.27, 95%CI:-0.5, -0.05; girls: β=-0.21,

95%CI:-0.45, 0.03) classes were similar. "Stable high" undereating was most strongly

associated with lower zBMI (boys: β=-0.79, 95%CI:-1.15, -0.42; girls: β=-0.76, -1.06, -0.45;

Table 2).

Fussy Eating

Similarly, fussy eating was associated with lower zBMI, for both boys and girls. "Stable

high" fussy eating was most strongly and negatively associated with zBMI (boys: β=-0.49,

95%CI: -0.68, -0.30; girls: β=-0.35, -0.52, -0.18; Table 2). In contrast to boys, amongst girls

"low transient" and "low increasing" fussy eating were not associated with zBMI at 11 years.

Interactions between class and sex in their effects on zBMI at 11 were not supported for any

of the Eating behaviors. Results from unadjusted regression models are available in eTable

6a-c.

<u>TABLE 2 HERE</u>

Discussion

In this study, differential developmental patterns in eating behaviors across childhood were identified and found to be associated with later child zBMI. This is the first study to address this question, by establishing longitudinal trajectories of eating behaviors during the first ten years of childhood and investigating their association with childhood zBMI at age 11. Results suggested four different trajectories of overeating and six trajectories each for undereating and fussy eating, respectively. Overall, it is notable that the three eating behaviors follow markedly different developmental trajectories. Overeating was found to be generally low, but increased with time, whereas under und fussy eating varied substantially across the observed timeframe. Previous research, on smaller datasets with a lower number of eating behaviors measures have indicated similar patterns of change and stability (18, 28, 29). These differences might be explained by various complex environmental and biological factors. With age, children gain autonomy and have more meals outside the home, which might be associated with the general increase of overeating. On the other hand, in toddlerhood, parents might introduce various and different textures and flavors of foods to their children (30), which they might readily embrace or resist, potentially explaining this early increase in fussy eating.

The majority of children were not described as overeaters by their parents. However, two trajectories (16% of children) were marked by gradual increases in overeating and showed similar positive associations with child zBMI at age 11. Recent longitudinal analysis of dietary data in a UK child cohort highlighted that eating larger portions a few times per week accelerates early childhood growth (31). This tendency to overeat is likely to result in larger portion sizes, which have been suggested to have enduring effects on child weight (32, 33). Of note, is the possible perpetual bi-directional association between overeating and portion size, where one potentially influences the other.

In contrast to overeating, undereating was more common and more heterogeneous. By 15

months, parents reported that children engaged in various levels of undereating, with about

10% of boys and girls reported to undereat at a worrying level. However, undereating

behavior of most children attenuated with time, indicating that parent-perceived undereating

in children under the age of two years may represent a normal pattern of development. Only

2-3% of children engaged in persistent high levels of undereating. This persistent pattern of

undereating was negatively associated with child zBMI at age 11. Parental reports of

undereating might be an indication of satiety sensitivity (34). Previous research has suggested

that children who were attuned to their internal satiety cues ate smaller portions (35), and

grew at a slower rate than their less satiety-responsive siblings (36).

Similarly to undereating, fussy eating behavior was highly heterogeneous in early life. Using

LCGA, we identified a small but substantial group of children (8%) who were persistently

fussy throughout the first ten years of life. These results add to previous studies suggesting

that some fussiness around food is common during childhood, with one third of children

reported to be fussy at some point, but only a small percentage of children remaining highly

fussy eaters across development (37). More persistent fussy eating trajectories were

negatively associated with child zBMI at age 11.

The relationship between food fussiness and weight is complex as fussy children might

undereat certain food groups (e.g., fruits and vegetables) but overeat others (e.g.,

carbohydrates and fats). Previous cross-sectional studies proposed that fussy children ate

fewer vegetables and less fish, but consumed more savory and sweet snack foods at 14

months (38). However, a longitudinal study indicated that persistent fussy eating in childhood

was associated with higher prevalence of underweight in children aged six years (15).

This study supports the prospective association between eating behaviors and weight in

children. Individual differences in weight have consistently been shown to be influenced by

genetic factors (39). The behavioral susceptibility to obesity theory (40) suggests that eating

behaviors might act as a mediator between genetic risk for obesity and exposure to the

current obesogenic environment. Previous studies proposed that increased genetic risk for

obesity is associated with decreased responsiveness to satiety cues, as well as greater

responsiveness to external food cues in ten year old twins (41). Subsequent research has

replicated these findings in Finnish (42), UK (43) and Canadian adults (44). However,

previous studies only included single measures of eating behaviors and it remains unknown

how genetic risk for obesity influences longitudinal trajectories of eating behaviors across

development.

Apart from weight, eating behaviors have been implicated in diet quality and as potential risk

factor for eating disorders. Especially, food fussiness has been associated with poor diet

quality,(45) such as low consumption of vegetables (46). Food fussiness has received

attention in the context of avoidant/restrictive food intake disorder (ARFID) (47). ARFID is a

recently defined diagnosis and little is known about its onset, development, and effect on

health and is characterized by extreme food fussiness affecting growth, weight, and physical

health (48) and that a large proportion of adolescents diagnosed with ARFID were persistent

fussy eaters during childhood (49). More research examining the impact of early food

fussiness and undereating on feeding and eating disorders risk is needed. It is possible that the

persistent fussy and undereating associated with low zBMI in this study may be ARFID

presentations, or risk factors for other eating disorders marked by restrictive eating, such as

anorexia nervosa. Further, child food fussiness has been found to be moderately heritable

(50) and future research is needed to uncover its genetic basis, as well as the role of fussy

eating in neurodevelopmental disorders such as autism spectrum disorder. In addition, the

majority of the research in this field relies on parental report. Parental anxiety could influence

parents' perception and reporting of their child's eating behavior.(51)

Strengths and Limitations

To our knowledge, this is the most comprehensive longitudinal study of child eating

behaviors in a large sample. Data were from a population-based cohort and person-centered

statistical analyses allowed us to clarify the heterogeneity of eating behaviors. Height and

weight were objectively measured during clinic visits. However, measures of eating

behaviors were parent reported and subject to reporting bias. For example parents might be

influenced by their own eating behaviors, their prior experiences with other children and

might be observing their children more closely in early life. As children grow up and enter

school, they will have an increasing numbers of meals outside the family home. Therefore

parents might be less aware of their children's eating behaviors. However, relying on parental

report remains the most commonly used measure of child eating behaviors, given that young

children are not able to report their own behavior reliably, and standardized direct

observational measures are costly and time-consuming, and would be infeasible for large

cohorts such as ALSPAC. One additional limitation is the fact that undereating and

overeating were only measured with one item at each wave, and a more comprehensive

assessment of these eating behaviors would have been desirable. However, in the context of

large-scale data collection efforts, such as ALSPAC, researchers always have to strike a

balance between including the optimal number of items without overwhelming the

participants. Further, the phrasing of the items only enquire how worried parents are about

their children's' eating behavior, and not the frequency of the behaviors themselves. We

implicitly assume that the greater the parental worry, the more pronounced the behavior. The

results however refer to the reporting of the behavior, not the behavior per se. Overall,

previous support for the use parental reports comes from research validating parent reported

child eating against behavioral measures of eating such as eating rate, energy intake at meal,

eating without hunger and caloric compensation (35).

Additionally, analyzing the effect of estimated class membership on an outcome includes

some degree of uncertainty. The values for entropy, which broadly reflects the level of

correct classification, were lower for overeating and undereating than the desired 0.8,

commonly used as cut-off point (51). Classes derived from LCGA are unobserved and hence

class membership is inferred. We used maximum-probability assignment, which allocates

each participant to the class they are most likely to belong to, carrying this class membership

forward to further analyses. This method has been suggested to attenuate the effect of class

on distal outcomes, due to uncertainty in class assignment (52). Hence, effect sizes estimated

from the regression analyses may be conservative.

Conclusions

We identified four trajectories of overeating and six trajectories each of fussy and

undereating in the ALSPAC sample, providing a thorough examination of child Eating

behaviors. EB trajectories were differentially associated with child zBMI, with persistent

behaviors having a stronger effect on BMI. Characterizing the heterogeneity of early life

eating behaviors is an important component to understanding behavioral risk factors for

common conditions, such as obesity.

Acknowledgements:

We are extremely grateful to all the families who took part in this study, the midwives for

their help in recruiting them, and the whole ALSPAC team, which includes interviewers,

computer and laboratory technicians, clerical workers, research scientists, volunteers,

managers, receptionists and nurses.

Funder:

This work was supported by the UK Medical Research Council and the Medical Research

Foundation (ref: MR/R004803/1).

The UK Medical Research Council and Wellcome (Grant ref: 102215/2/13/2) and the

University of Bristol provide core support for ALSPAC. A comprehensive list of grants

funding is available on the ALSPAC website

(http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf).

Prof Bulik acknowledges funding from the Swedish Research Council (VR Dnr: 538-2013-

8864).

The funders were not involved in the design or conduct of the study; collection, management,

analysis, or interpretation of the data; or preparation, review, or approval of the manuscript.

Potential conflict of interest: Bulik reports: Shire (grant recipient, Scientific Advisory

Board member) and Pearson and Walker (author, royalty recipient). All other authors have

indicated they have no conflicts of interest to disclose.

Author Contributions:

MH, BDS, CB, RBW and NM designed the research; MH and BDS performed statistical

analyses; all authors wrote and revised the manuscript for important intellectual content; NM

had primary responsibility for final content. All authors read and approved the final

manuscript.

References

1.   Emmett PM, Hays NP, Taylor CM. Factors Associated with Maternal Worry about Her Young Child Exhibiting Choosy Feeding Behaviour. Int J Env Res Pub He. 2018;15(6).

2.   Emmett PM, Hays NP, Taylor CM. Factors Associated with Maternal Worry about Her Young Child Exhibiting Choosy Feeding Behaviour. Int J Environ Res Public Health. 2018;15(6).

3.   Braet C, Claus L, Goossens L, Moens E, Van Vlierberghe L, Soetens B. Differences in eating style between overweight and normal-weight youngsters. J Health Psychol. 2008;13(6):733-43.

4.   Domoff SE, Miller AL, Kaciroti N, Lumeng JC. Validation of the Children's Eating Behaviour Questionnaire in a low-income preschool-aged sample in the United States. Appetite. 2015;95:415-20.

5.   Hajna S, LeBlanc PJ, Faught BE, Merchant AT, Cairney J, Hay J, et al. Associations between family eating behaviours and body composition measures in peri-adolescents: Results from a community-based study of school-aged children. Can J Public Health. 2014;105(1):E15-E21.

6.   Jansen PW, Roza SJ, Jaddoe VW, Mackenbach JD, Raat H, Hofman A, et al. Children's eating behavior, feeding practices of parents and weight problems in early childhood: results from the population-based Generation R Study. The international journal of behavioral nutrition and physical activity. 2012;9:130.

7.   Webber L, Hill C, Saxton J, Van Jaarsveld CH, Wardle J. Eating behaviour and weight in children. International journal of obesity. 2009;33(1):21-8.

8.   Sanchez U, Weisstaub G, Santos JL, Corvalan C, Uauy R. GOCS cohort: children's eating behavior scores and BMI. Eur J Clin Nutr. 2016.

9.   Svensson V, Lundborg L, Cao Y, Nowicka P, Marcus C, Sobko T. Obesity related eating behaviour patterns in Swedish preschool children and association with age, gender, relative weight and parental weight--factorial validation of the Children's Eating Behaviour Questionnaire. The international journal of behavioral nutrition and physical activity. 2011;8:134.

10.   Cao YT, Svensson V, Marcus C, Zhang J, Zhang JD, Sobko T. Eating behaviour patterns in Chinese children aged 12-18 months and association with relative weight--factorial validation of the Children's Eating Behaviour Questionnaire. The international journal of behavioral nutrition and physical activity. 2012;9:5.

11.   Parkinson KN, Drewett RF, Le Couteur AS, Adamson AJ, T GMSC. Do maternal ratings of appetite in infants predict later Child Eating Behaviour Questionnaire scores and body mass index? Appetite. 2010;54(1):186-90.

12.   van Jaarsveld CHM, Llewellyn CH, Johnson L, Wardle J. Prospective associations between appetitive traits and weight gain in infancy. Am J Clin Nutr. 2011;94(6):1562-7.

13.   Steinsbekk S, Wichstrom L. Predictors of Change in BMI From the Age of 4 to 8. Journal of pediatric psychology. 2015.

14.   Antoniou EE, Roefs A, Kremers SP, Jansen A, Gubbels JS, Sleddens EF, et al. Picky eating and child weight status development: a longitudinal study. J Hum Nutr Diet. 2016;29(3):298-307.

15.   de Barse LM, Tiemeier H, Leermakers ET, Voortman T, Jaddoe VW, Edelson LR, et al. Longitudinal association between preschool fussy eating and body composition at 6 years of age: The Generation R Study. The international journal of behavioral nutrition and physical activity. 2015;12:153.

16.   Mallan KM, Fildes A, Magarey AM, Daniels LA. The Relationship between Number of Fruits, Vegetables, and Noncore Foods Tried at Age 14 Months and Food Preferences, Dietary Intake Patterns, Fussy Eating Behavior, and Weight Status at Age 3.7 Years. J Acad Nutr Diet. 2016;116(4):630-7.

17.   Carruth BR, Skinner JD. Revisiting the picky eater phenomenon: neophobic behaviors of young children. J Am Coll Nutr. 2000;19(6):771-80.

18.   Taylor CM, Steer CD, Hays NP, Emmett PM. Growth and body composition in children who
are picky eaters: a longitudinal view. Eur J Clin Nutr. 2018.
19.   Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile: the
'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. Int J
Epidemiol. 2013;42(1):111-27.
20.   Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Smith GD, et al. Cohort Profile: The
Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. International Journal of
Epidemiology. 2013;42(1):97-110.
21.   Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child
overweight and obesity worldwide: international survey. BMJ. 2000;320(7244):1240-3.
22.   Clogg C. Latent class models: Recent developments and prospects for the future. In:
Arminger G, Clogg C, Sobel M, editors. Handbook of statistical modeling for the social and behavioral
sciences. New York, NY: Plenum Press;; 1995. p. 311 - 59.
23.   Berlin KS, Parra GR, Williams NA. An Introduction to Latent Variable Mixture Modeling (Part
2): Longitudinal Latent Class Growth Analysis and Growth Mixture Models. Journal of pediatric
psychology. 2014;39(2):188-203.
24.   Asparouhov T, Muthen B. Auxiliary Variables in Mixture Modeling: Three-Step Approaches
Using Mplus. Struct Equ Modeling. 2014;21(3):329-41.
25.   Goodman LA. On the Assignment of Individuals to Latent Classes. Sociol Methodol.
2007;37:1-22.
26.   Muthen LK, Muthen B. Mplus User's Guide. Eighth Edition. Los Angeles, CA Muthén &
Muthén; 1997-2017
27.   StataCorp. Stata Statistical Software: Release 15. College Station, TX: StataCorp LLC 2017.
28.   Ashcroft J, Semmler C, Carnell S, van Jaarsveld CHM, Wardle J. Continuity and stability of
eating behaviour traits in children. Eur J Clin Nutr. 2008;62(8):985-90.
29.   Derks IPM, Bolhuis K, Sijbrands EJG, Gaillard R, Hillegers MHJ, Jansen PW. Predictors and
patterns of eating behaviors across childhood: Results from The Generation R study. Appetite.
2019;141:104295.
30.   Northstone K, Emmett P, Nethersole F, Pregnancy ASTALSo, Childhood. The effect of age of
introduction to lumpy solids on foods eaten and reported feeding difficulties at 6 and 15 months. J
Hum Nutr Diet. 2001;14(1):43-54.
31.   Syrad H, Llewellyn CH, Johnson L, Boniface D, Jebb SA, van Jaarsveld CHM, et al. Meal size is
a critical driver of weight gain in early childhood. Sci Rep-Uk. 2016;6.
32.   Huang TT, Howarth NC, Lin BH, Roberts SB, McCrory MA. Energy intake and meal portions:
associations with BMI percentile in U.S. children. Obes Res. 2004;12(11):1875-85.
33.   McConahy KL, Smiciklas-Wright H, Birch LL, Mitchell DC, Picciano MF. Food portions are
positively related to energy intake and body weight in early childhood. J Pediatr-Us.
2002;140(3):340-7.
34.   Carnell S, Wardle J. Measuring behavioural susceptibility to obesity: validation of the child
eating behaviour questionnaire. Appetite. 2007;48(1):104-13.
35.   Syrad H, Johnson L, Wardle J, Llewellyn CH. Appetitive traits and food intake patterns in early
life. Am J Clin Nutr. 2016;103(1):231-5.
36.   van Jaarsveld CHM, Boniface D, Llewellyn CH, Wardle J. Appetite and Growth A Longitudinal
Sibling Analysis. Jama Pediatr. 2014;168(4):345-50.
37.   Cardona Cano S, Tiemeier H, Van Hoeken D, Tharner A, Jaddoe VW, Hofman A, et al.
Trajectories of picky eating during childhood: A general population study. The International journal
of eating disorders. 2015;48(6):570-9.
38.   Tharner A, Jansen PW, Kiefte-de Jong JC, Moll HA, van der Ende J, Jaddoe VW, et al. Toward
an operative diagnosis of fussy/picky eating: a latent profile approach in a population-based cohort.
The international journal of behavioral nutrition and physical activity. 2014;11:14.

39.   Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Felix R, et al. Genetic studies of body mass
index yield new insights for obesity biology. Nature. 2015;518(7538):197-U401.
40.   Llewellyn C, Wardle J. Behavioral susceptibility to obesity: Gene-environment interplay in
the development of weight. Physiology & behavior. 2015;152(Pt B):494-501.
41.   Llewellyn CH, Trzaskowski M, van Jaarsveld CHM, Plomin R, Wardle J. Satiety mechanisms in
genetic risk of obesity. Jama Pediatr. 2014;168(4):338-44.
42.   Konttinen H, Llewellyn C, Wardle J, Silventoinen K, Joensuu A, Mannisto S, et al. Appetitive
traits as behavioural pathways in genetic susceptibility to obesity: a population-based cross-sectional
study. Sci Rep-Uk. 2015;5.
43.   de Lauzon-Guillain B, Clifton EA, Day FR, Clement K, Brage S, Forouhi NG, et al. Mediation
and modification of genetic susceptibility to obesity by eating behaviors. Am J Clin Nutr.
2017;106(4):996-1004.
44.   Jacob R, Drapeau V, Tremblay A, Provencher V, Bouchard C, Perusse L. The role of eating
behavior traits in mediating genetic susceptibility to obesity. Am J Clin Nutr. 2018;108(3):445-52.
45.   Vilela S, Hetherington MM, Oliveira A, Lopes C. Tracking diet variety in childhood and its
association with eating behaviours related to appetite: The generation XXI birth cohort. Appetite.
2018;123:241-8.
46.   de Barse LM, Tiemeier H, Leermakers ETM, Voortman T, Jaddoe VWV, Edelson LR, et al.
Longitudinal association between preschool fussy eating and body composition at 6 years of age:
The Generation R Study. The international journal of behavioral nutrition and physical activity.
2015;12(153):8.
47.   APA APA. Diagnostic and Statistical Manual of Mental Disorders (DSM-V). Arlington, VA: APA
Press; 2013.
48.   Bryant-Waugh R. Avoidant restrictive food intake disorder: An illustrative case example. New
York, N.Y: Wiley; 2013. 1 online resource p.
49.   Fisher MM, Rosen DS, Ornstein RM, Mammel KA, Katzman DK, Rome ES, et al.
Characteristics of avoidant/restrictive food intake disorder in children and adolescents: a "new
disorder" in DSM-5. J Adolesc Health. 2014;55(1):49-52.
50.   Smith AD, Herle M, Fildes A, Cooke L, Steinsbekk S, Llewellyn CH. Food fussiness and food
neophobia share a common etiology in early childhood. J Child Psychol Psyc. 2017;58(2):189-96.
51.   Ram N, Grimm KJ. Growth Mixture Modeling: A Method for Identifying Differences in
Longitudinal Change Among Unobserved Groups. Int J Behav Dev. 2009;33(6):565-76.
52.   Bray BC, Lanza ST, Tan X. Eliminating Bias in Classify-Analyze Approaches for Latent Class
Analysis. Struct Equ Modeling. 2015;22(1):1-11.

Table 1: Summary statistics of the baseline characteristics of the study population;

Baseline characteristics		
	N available	**Mean (SD) or N (%)**
Sex (%boys)	12048	Boys: 6208 (52)
Gestational age at birth (weeks)	12048	39.45 (1.86)
Birthweight (grams)	11902	345 (546)
Maternal age (years)	12048	28.31 (4.86)
Maternal A-Levels or higher	11375	4158 (37)
Parental non-manual labor profession	9366	7558 (81)
zBMI ^{a} of children at age 11 (kg/m^2)	4885	0.60 (1.14)

ALSPAC Study

Table 2: Estimated regression coefficients (β) for assigned eating behavior class on standardized BMI at age 11, separately for boys and girls

Overeating

Outcome: age and sex adjusted BMI aged 11[1], adjusted for covariates[2]

	Boys			**Girls**		
	2286			2596		
N						
Class	**β**	**95% CI[3]**		**β**	**95% CI**	
1 low stable	base			Base		
2 low transient	0.26	0.13	0.39	0.32	0.19	0.44
3 late increasing	0.94	0.8	1.09	0.94	0.82	1.07
4 early increasing	0.83	0.65	1.02	1.1	0.92	1.28

Test for (Sex * Class) interaction $F_{(3, 4869)} = 1.10$, $p=0.35$

Undereating

Outcome: age and sex adjusted BMI aged 11, adjusted for covariates[2]

	Boys			**Girls**		
N	2285			2595		
Class	**β**	**95% CI**		**β**	**95% CI**	
1 low stable	base			base		
2 low transient	-0.11	-0.24	0.01	-0.13	-0.24	-0.01
3 low decreasing	-0.17	-0.31	-0.03	-0.19	-0.32	-0.07
4 high transient	-0.25	-0.41	-0.08	-0.24	-0.38	-0.09
5 high decreasing	-0.27	-0.5	-0.05	-0.21	-0.45	0.03
6 high stable	-0.79	-1.15	-0.42	-0.76	-1.06	-0.45

Test for (Sex * Class) interaction $F_{(5, 4863)} = 0.12$, $p = 0.98$

Fussy Eating

Outcome: age and sex adjusted BMI aged 11, adjusted for covariates[2]

	Boys			**Girls**		
N	2287			2597		
Class	**β**	**95% CI**		**β**	**95% CI**	
1 low stable	base			base		
2 low transient	-0.21	-0.36	-0.05	0.01	-0.13	0.15
3 low increasing	-0.25	-0.39	-0.11	-0.01	-0.13	0.11
4 high decreasing	-0.31	-0.48	-0.15	-0.31	-0.45	-0.17
5 low increasing	-0.34	-0.50	-0.17	-0.26	-0.41	-0.11
6 high stable	-0.49	-0.68	-0.30	-0.35	-0.52	-0.18

Test for (Sex * Class) interaction $F(5, 4867) = 2.01$, $p = 0.07$

[1] Age and sex standardized score in reference to the UK population (21)

[2] Estimates adjusted for: maternal age at birth, gestational age, birthweight and maternal

education [3] CI: Confidence Intervals

Figure 1: Prevalence of eating behaviours across the eight assessment waves

Figure 2a: Trajectories of parental reports of overeating behaviors from 15 to
116 months for boys and girls (6186 boys, 5817 girls)

Caption: The y-axis shows the probability of scoring in the highest category of overeating ("great
worry") at each of the eight time points. Trajectories for boys are in dashed lines. Trajectories for girls
are in solid lines. The legend shows the name of the class for boys (Bs) and girls (Gs), followed by
their percentages in brackets.

Figures 2b: Trajectories of parental reports of undereating behaviors from 15 to 116 months for boys and girls (6189 boys, 5817 girls)

Caption: The y-axis shows the probability of scoring in the highest category of undereating ("a bit worried") at each of the eight time points. Trajectories for boys are in dashed lines. Trajectories for girls are in solid lines. The legend shows the name of the class for boys (Bs) and girls (Gs), followed by their percentages in brackets.

Figures 2c: Trajectories of parental reports of fussy eating behaviors from 15 to 116 months for boys and girls (6208 boys, 5840 girls)

Caption: The y-axis shows the probability of scoring in the highest category of the fussy eating items ("a bit worried"") at each of the eight time points. Trajectories for boys are in dashed lines. Trajectories for girls are in solid lines. The legend shows the name of the class for boys (Bs) and girls (Gs), followed by their percentages in brackets.