Efficient Prediction of Vitamin B Deficiencies via

Machine-learning Using Routine Blood Test Results in Patients

With Intense Psychiatric Episode

Hidetaka Tamune[1)2)3)5)*], Jumpei Ukita[3)4)5)], Yu Hamamoto[1)2)], Hiroko Tanaka[1)2)], Kenji

Narushima[1)], Naoki Yamamoto[1)]

1) Department of Neuropsychiatry, Tokyo Metropolitan Tama Medical Center, Tokyo,

Japan

2) Department of Neuropsychiatry, Graduate School of Medicine, The University of

Tokyo, Tokyo, Japan

3) Mental Health Research Course, Faculty of Medicine, The University of Tokyo,

Tokyo, Japan

4) Department of Physiology, Graduate School of Medicine, The University of Tokyo,

Tokyo, Japan

5) H. Tamune and JU contributed equally to this work

* Correspondence:

Hidetaka Tamune, M.D., tamune-tky@umin.ac.jp

Abstract: 294 words

Main text: 1878 words + 4 Tables + 4 Figures + 24 references

Keywords: Machine Learning; Random Forest Classifier; Vitamin B Deficiency; Folic

Acid; Early Diagnosis; Decision support techniques or decision making.

Abstract

Background: Vitamin B deficiency is common worldwide and may lead to psychiatric

symptoms; however, vitamin B deficiency epidemiology in patients with intense

psychiatric episode has rarely been examined. Moreover, vitamin deficiency testing is

costly and time-consuming. It hampered to effectively rule out vitamin

deficiency-induced intense psychiatric symptoms. In this study, we aimed to clarify the

epidemiology of these deficiencies and efficiently predict them using machine-learning

models from patient characteristics and routine blood test results that can be obtained

within one hour.

Methods: We reviewed 497 consecutive patients deemed to be at imminent risk of

seriously harming themselves or others over 2 years. Machine-learning models were

trained to predict each deficiency from age, sex, and 29 routine blood test results.

Results: We found that 112 (22.5%), 80 (16.1%), and 72 (14.5%) patients had vitamin

B_1, vitamin B_{12}, and folate (vitamin B_9) deficiency, respectively. Also, the

machine-learning models well generalized to predict the deficiency in the future unseen

data; areas under the receiver operating characteristic curves for the validation dataset

(i.e. dataset not used for training the models) were 0.716, 0.599, and 0.796, respectively.

The Gini importance of these vitamins provided further evidence of a relationship

between these vitamins and the complete blood count, while also indicating a hitherto

rarely considered, potential association between these vitamins and alkaline phosphatase

(ALP) or thyroid stimulating hormone (TSH).

Discussion: This study demonstrates that machine-learning can efficiently predict some

vitamin deficiencies in patients with active psychiatric symptoms, based on the largest

cohort to date with intense psychiatric episode. The prediction method may expedite

risk stratification and clinical decision-making regarding whether replacement therapy

should be prescribed. Further research includes validating its external generalizability in

other clinical situations and clarify whether interventions based on this method can

improve patient care and cost-effectiveness.

Vitamin deficiency predicted by ML

1. Introduction

Vitamin B deficiency is common worldwide and may lead to psychiatric symptoms[1–4]. For example, meta-analyses have shown that patients with schizophrenia or first-episode psychosis have lower folate (vitamin B_9) levels than their healthy counterparts[4,5]. Moreover, vitamin therapy can effectively alleviate symptoms in a subgroup of patients with schizophrenia[3,6–8]. However, the epidemiology of vitamin B deficiency in patients with active mental symptoms requiring immediate hospitalization has rarely been examined.

In a psychiatric emergency, psychiatrists should promptly distinguish treatable patients with altered mental status due to a physical disease from patients with an authentic mental disorder (international statistical classification of diseases and related health problems-10, ICD-10 code: F2-9). However, vitamin deficiency testing is very costly (around 60 dollars for each measurement of vitamin B_1 (vitB$_1$), vitamin B_{12} (vitB$_{12}$), or folate in the U.S.; 15–25 dollars for each test in Japan) and usually requires at least two days. Therefore, an efficient, cost-effective method of predicting vitamin B deficiency is needed.

Although several studies have applied machine-learning to the prediction of diagnosis or treatment outcomes[9–11], no study using machine-learning has focused on vitamin B deficiencies. We herein explore whether vitB$_1$, vitB$_{12}$, and folate deficiencies can be predicted using a machine-learning classifier from patient characteristics and routine blood test results obtained within one hour based on a large cohort of patients requiring urgent psychiatric hospitalization.

Vitamin deficiency predicted by ML

2. Methods

2.1. Medical chart review

We reviewed consecutive patients admitted to the Department of Neuropsychiatry at Tokyo Metropolitan Tama Medical Center between September 2015 and August 2017 under the urgent involuntary hospitalization law, which requires the immediate psychiatric hospitalization of patients at imminent risk of seriously harming themselves or others. The necessity of hospitalization was judged by designated mental health specialists. The patient characteristics, ICD-10 codes, and laboratory data were gathered retrospectively.

Since the reference ranges for $vitB_1$, $vitB_{12}$, and folate are 70–180 nmol/L (30–77 ng/mL), 180–914 ng/L, and > 4.0 µg/L, respectively[12], a deficiency of the nutrients was defined as < 30 ng/mL, < 180 ng/L, and < 4.0 µg/L, respectively, unless otherwise stated.

2.2. Random forest classifier and statistics

A random forest classifier was trained to predict the deficiency of each substance from age, sex, and 29 routine blood variables (described in the Result section with values). The random forest classifier was trained using the dataset populated in the period from September 2015 to December 2016 (the "Training set"). First, we optimized the hyperparameters of the classifier by selecting the best combination of hyperparameters that maximized the "5-fold cross validation" accuracy, among many combinations within appropriate ranges. The cross-validation accuracy was computed as follows; in one session, the classifier was trained using 80% of the training set and evaluated on the withheld 20% of the training set. This session was performed five times so that every data would be withheld once. The accuracies were finally averaged across sessions to yield the cross-validation accuracy. By incorporating this process, the classifier was generalized to unseen data (Graphical method is shown in **Figure 1**).

Using the optimized hyperparameters, the classifiers were then validated using

data collected from January 2017 through August 2017 (the "Validation set"). We report

the classification performance on the validation set in the results section unless

otherwise stated. We quantified the sensitivity, specificity, and accuracy (defined as the

average of the sensitivity and the specificity on the optimal operating point) using

receiver operating characteristic curves (ROCs). We also quantified the 95% confidence

interval of the accuracy using 1000-times bootstrapping.

When investigating the Gini importance and the partial dependency[13], we

retrained the classifiers using all datasets. All data analyses were performed using

Python (2.7.10) with the Scikit-learn package (0.19.0) and R (3.4.2) with the edarf

package (1.1.1).

2.3. Ethical considerations

Informed consent was obtained from participants using an opt-out form on the

website. The study protocol was approved by the Research Ethics Committee, Tokyo

Metropolitan Tama Medical Center (Approval number: 28-8). The study complied with

the Declaration of Helsinki and the STROBE statement.

3. Results

3.1. Eligible patients

During the 2-year study period, 497 consecutive patients (496 were Asian) were enrolled. The mean age (standard deviation, SD) was 42.3 ($\pm$15.4) years, and 228 patients (45.9%) were women. F2 (Schizophrenia, schizotypal, delusional, and other non-mood psychotic disorders) was diagnosed in over 60% of the patients. The ICD-10 codes of the patients and the number of deficiencies at several cut-off values for $vitB_1$, $vitB_{12}$, and folate are shown in **Table 1**. According to the predefined cut-off values[12], 112 (22.5%), 80 (16.1%), and 72 (14.5%) patients exhibited a deficiency of $vitB_1$ (<30 ng/mL), $vitB_{12}$ (<180 ng/L), and folate (<4.0 μg/L), respectively. Vitamin B deficiencies in sub-groups are shown in **Table 2**. A summary of the full dataset is shown in **Table 3**. Detailed information (sub-datasets) is shown in **Supplementary Table 1**, **2, and 3** online. Histograms of $vitB_1$, $vitB_{12}$, and folate values are shown in **Figure 2 A-C**.

3.2. Prediction via machine-learning using routine blood test results

A random forest classifier was trained to predict the deficiency of each substance from patient characteristics and routine blood test results. The classifier was trained using the dataset gathered in the period from September 2015 to December 2016 (the "Training set", n = 373), which was then validated from January 2017 through August 2017 (the "Validation set", n = 124).

The area under the ROCs (AUCs) for the validation set were 0.716, 0.599, and 0.796, for $vitB_1$, $vitB_{12}$, and folate, respectively (**Figure 2 D-F** and **Table 4**). With some operative points on the ROC, the sensitivity, specificity and accuracy for the validation set were calculated (**Table 4**. See also **Supplementary Table 4** for training set and **Supplementary Table 5** for different operating points).

When the prediction performances were compared between the classifiers trained using the dataset from the F2 population and the classifiers trained using the

dataset from the other population, the AUC was not statistically different (DeLong's

test), except in the case of vitB$_1$ (see **Supplementary Table 6**).

Figure 3 shows the Gini importance (a–c) and partial dependency plots (d–f)

for the eight most important variables for each substance. The results provided further

evidence of a relationship between the vitamin B levels and complete blood count while

also indicating the hitherto rarely considered, potential association between these

vitamins and alkaline phosphatase (ALP) or thyroid stimulating hormone (TSH).

3.3. Robustness verification

We verified the robustness of the results by two independent means. First, we

used different cut-off values to define the deficiency[14–16]. Although the AUC for the

validation set, shown in **Supplementary Table 7**, tended to be higher when strict

cut-off values were used, the obtained AUCs were not statistically significant ($p > 0.05$,

DeLong's test with Bonferroni correction).

Second, we trained and evaluated random forest classifiers using a dataset split

in a different way; the classifier was trained using the dataset collected in the period

from the 31st of January, 2016 to August 2017, which was then validated with data

gathered from September 2015 to the 31st of January, 2016. Note that the sample sizes

of the training and validation sets were equal to those in the original setting. The AUCs

for the validation set were 0.771, 0.621, and 0.745 for vitB$_1$, vitB$_{12}$, and folate,

respectively; none were statistically different from the AUC trained using the original

setting (DeLong's test), further demonstrating the robustness of the performance.

Vitamin deficiency predicted by ML

4. Discussion

4.1. Relevance of the present study

Based on the largest cohort to date of patients at imminent risk of seriously harming themselves or others, this study indicated that deficiency of certain vitamins can be predicted in an efficient manner via machine-learning using routine blood test results. Given the large number of patients with vitamin B deficiencies, empirical therapy might be acceptable; however, risk stratification is preferred for personalized medicine and shared decision-making. The prediction method presented here may expedite clinical decision-making as to whether vitamins should be prescribed to a patient (Graphical abstract is shown in **Figure 4**).

Remarkably, the AUC for folate deficiency was 0.796. Folate features the potential to maintain neuronal integrity and is one of the homocysteine-reducing B-vitamins[5]; homocysteine has been linked to the etiology of schizophrenia[17], and vitamin B supplements have been reported to reduce psychiatric symptoms significantly in patients with schizophrenia[7]. As our study does not present longitudinal results, an intervention effect of folate supplementation in the cohort remains to be clarified.

4.2. Trade-off of interpretability and generalizability using machine-learning

Compared to the AUC of folate, AUCs of $vitB_1$ and $vitB_{12}$ were relatively low. Using other parameters that were not incorporated into this model or using other models including deep neural networks might increase the accuracy of prediction.

However, interpretability and completeness of machine-learning classifiers are subject to trade-off[17]. Although completeness and generalizability are desirable, interpretability is also indispensable, especially in clinical settings, since it provides meaningful and trustworthy findings for clinical physicians as well as new biological insights[18]. In this study we chose random forest classifiers since they provide expressive and interpretable data, with sufficient accuracy.

Vitamin deficiency predicted by ML

4.3. Biological mechanism suggestion

Using the random forest classifiers, as shown in **Figure 3**, we identified several items related to complete blood count as top hits. Notably, our classifier was blind to any biological knowledge, including the well-established association between anemia and vitamin B deficiency, including folate[19]. The results provide further evidence of a relationship between vitamin B levels and the complete blood count and support the use of machine-learning to investigate novel, underlying biological mechanisms[20].

ALP and its metabolites indicate the vitamin B_6 status[21]; low vitB$_{12}$ is potentially associated with low ALP[22]. More generally, ALP may have a close and complicated relationship with the overall vitamin B group. Autoimmune disorders, especially thyroid disease, are commonly associated with pernicious anaemia[23], but there has been no established hypothesis regarding the causal relationships between thyroid disease and vitamin B deficiencies. The potential association between the levels of these vitamins and ALP or TSH awaits further study, both investigations of populations and basic research[24].

4.4. Limitations

This study is subject to several limitations. First, the findings of this single-center retrospective study may have limited generalizability. Second, the patients' long-term prognosis was not investigated due to administrative restrictions; the extent to which this method can expedite clinical decision-making is therefore unclear. Further, we did not investigate the relationship between serological values and the need for intervention. The lack of data for vitamin B deficiency in the Japanese general population hampered the comparison between the experimental cohort and their counterparts who lacked psychiatric symptoms. Establishing appropriate reference values and an assessment method requires further investigation. Finally, we did not

assess the predictive value of other nutritional impairments, including vitamin B_6 and

homocysteine deficiency, which were previously shown to have a close link with

psychiatric symptoms[3,5]; however, our study provides fundamental data on nutritional

impairment based on the largest cohort of patients with intense psychiatric episode ever

assembled for this purpose and presents a potential framework for predicting nutritional

impairment using machine-learning.

4.5. Conclusion

The present report is, to the best of our knowledge, the first to demonstrate that

machine-learning can efficiently predict nutritional impairment. Further research is

needed to validate the external generalizability of the findings in other clinical situations

and clarify whether interventions based on this method can improve patient care and

cost-effectiveness.

Vitamin deficiency predicted by ML

5. Contribution to the Field Statement

Vitamin B deficiency is common worldwide and may lead to psychiatric symptoms; however, vitamin B deficiency epidemiology in patients with intense psychiatric symptoms has rarely been examined. Moreover, vitamin deficiency testing is costly and time-consuming. Based on the largest cohort to date of patients at imminent risk of seriously harming themselves or others, this study demonstrated that the deficiency of certain vitamins can be predicted in an efficient manner via machine-learning models from patient characteristics and routine blood test results obtained within one hour.

In detail, among the 497 patients investigated (over 60% was diagnosed with schizophrenia or related psychotic disorders), 22.5%, 16.1%, and 14.5% patients had a deficiency of vitamin B_1, B_{12}, and folate, respectively, by direct measurement. Also, the machine-learning models well generalized to predict the deficiency in unseen datasets; areas under the receiver operating characteristic curves for the validation dataset were 0.716, 0.599, and 0.796, respectively. The prediction method presented in this study may expedite risk stratification and clinical decision-making regarding whether replacement therapy should be prescribed. The results also provided further evidence for a well-known relationship between these vitamins and the complete blood count and supported the application of machine-learning to investigate novel, underlying biological mechanisms.

6. Acknowledgements

We thank Mr. James Robert Valera for his assistance in editing this manuscript and all the staff for their care of the patients and their contributions to this study.

7. Author Contributions Statement

H. Tamune has full access to all data and takes responsibility for the integrity of the data. H. Tamune, JU, KN, and NY conceived the study. H. Tamune, YH, and H. Tanaka collected the data. JU performed the statistical analyses. H. Tamune and JU drafted the first version of the manuscript. All authors critically revised the manuscript for intellectual content and approved the final version.

8. Data Availability Statements

The datasets and source code utilized in the current study are available from the corresponding author upon reasonable request.

9. Conflict of Interest Statement

The authors declare no conflict of interest, except for a scholarship grant awarded to JU from Takeda Science Foundation and Masayoshi Son Foundation.

<u>References</u>

1. Harper C. Thiamine (vitamin B1) deficiency and associated brain damage is still

common throughout the world and prevention is simple and safe! *Eur J Neurol.* (2006)

13: 1078–1082.

2. Reynolds E. Vitamin B12, folic acid, and the nervous system. *Lancet Neurol.* (2006)

5: 949–960.

3. Arai M, Yuzawa H, Nohara I, Ohnishi T, Obata N, Iwayama Y *et al.* Enhanced

carbonyl stress in a subpopulation of schizophrenia. *Arch Gen Psychiatry* (2010) 67:

589–597.

4. Cao B, Wang DF, Xu MY, Liu YQ, Yan LL, Wang JY *et al.* Lower folate levels in

schizophrenia: A meta-analysis. *Psychiatry Res.* (2016) 245: 1–7.

5. Firth J, Carney R, Stubbs B, Teasdale SB, Vancampfort D, Ward PB, *et al.* Nutritional

deficiencies and clinical correlates in first-episode psychosis: A systematic review and

meta-analysis. *Schizophr Bull.* (2018) 44: 1275–1292.

6. Levine J, Stahl Z, Sela BA, Ruderman V, Shumaico O, Babushkin I *et al.*

Homocysteine-reducing strategies improve symptoms in chronic schizophrenic patients

with hyperhomocysteinemia. *Biol Psychiatry* (2006) 60: 265–269.

7. Firth J, Stubbs B, Sarris J, Rosenbaum S, Teasdale S, Berk M *et al.* The effects of

vitamin and mineral supplementation on symptoms of schizophrenia: A systematic

review and meta-analysis. *Psychol Med.* (2017) 47: 1515–1527.

8. Itokawa M, Miyashita M, Arai M, Dan T, Takahashi K, Tokunaga T *et al.*

Pyridoxamine: A novel treatment for schizophrenia with enhanced carbonyl stress.

Psychiatry Clin Neurosci. (2018) 72: 35–44.

9. Koutsouleris N, Kahn RS, Chekroud AM, Leucht S, Falkai P, Wobrock T *et al.*

Multisite prediction of 4-week and 52-week treatment outcomes in patients with

first-episode psychosis: A machine learning approach. *Lancet Psychiatry* (2016) 3:

935–946.

Vitamin deficiency predicted by ML

10. Mechelli A, Lin A, Wood S, McGorry P, Amminger P, Tognin S *et al.* Using clinical

information to make individualized prognostic predictions in people at ultra-high risk

for psychosis. *Schizophr Res.* (2017) 184: 32-38.

11. Vieira S, Pinaya WH, Mechelli A. Using deep learning to investigate the

neuroimaging correlates of psychiatric and neurological disorders: Methods and

applications. *Neurosci Biobehav Rev.* (2017) 74: 58–75.

12. Mayo Foundation for Medical Education and Research, Rochester Test Catalog.

https://www.mayomedicallaboratories.com/test-catalog/ (2018).

13. Friedman, JH. Greedy function approximation: A gradient boosting machine. *Ann.*

Stat. (2001) 29: 1189–1232.

14. Sasaki T, Yukizane T, Atsuta H, Ishikawa H, Yoshiike T, Takeuchi T *et al.* A case of

thiamine deficiency with psychotic symptoms: Blood concentration of thiamine and

response to therapy. *Seishin Shinkeigaku Zasshi* (2010) 112: 97–110.

15. Clarke R, Refsum H, Birks J, Evans JG, Johnston C, Sherliker P *et al.* Screening for

vitamin B-12 and folate deficiency in older persons. *Am J Clin Nutr.* (2003) **77**:

1241–1247.

16. Goff DC, Bottiglieri T, Arning E, Shih V, Freudenreich O, Evins AE *et al.* Folate,

homocysteine, and negative symptoms in schizophrenia. *Am J Psychiatry* (2004) 161:

1705–1708.

17. Muntjewerff JW, Kahn RS, Blom HJ, den Heijer M. Homocysteine,

methylenetetrahydrofolate reductase and risk of schizophrenia: A meta-analysis. *Mol*

Psychiatry (2006) 11: 143–149.

18. Gilpin LH, Bau D, Yuan BZ, Bajwa A, Specter M, Kagal L. Explaining

explanations: An overview of interpretability of machine learning. In *2018 IEEE 5th*

International Conference on Data Science and Advanced Analytics (DSAA): 80–89.

19. Evans TC, Jehle D. The red blood cell distribution width. *J Emerg Med.* (1991) 9:

71–74.

Vitamin deficiency predicted by ML

20. So HC, Chau CK, Chiu WT, Ho KS, Lo CP, Yim SH *et al.* Analysis of genome-wide

association data highlights candidates for drug repositioning in psychiatry. *Nat Neurosci.*

(2017) 20: 1342–1349.

21. Ueland PM, Ulvik A, Rios-Avila L, Midttun Ø, Gregory JF. Direct and functional

biomarkers of vitamin B6 status. *Annu Rev Nutr*. (2015) 35: 33–70.

22. Carmel R, Lau KH, Baylink DJ, Saxena S, Singer FR. Cobalamin and

osteoblast-specific proteins. *N Engl J Med.* (1977) 319: 70–75.

23. Stabler SP. Vitamin B12 deficiency. *N Engl J Med.* (2013) 368: 149–160.

24. Zheng Y, Cantley LC. Toward a better understanding of folate metabolism in health

and disease. *J Exp Med.* (2018) 216: 253–266.

Vitamin deficiency predicted by ML

Supporting Material List

Supplementary Table 1 (related to Table 1). Divided patient distribution data (n = 497)

Supplementary Table 2 (related to Table 2). Divided data of vitamin B deficiencies in

sub-groups

Supplementary Table 3 (related to Table 3). Divided dataset of age, sex, and 29

parameters

Supplementary Table 4 (related to Table 4). Summary of sensitivity, specificity, and

accuracy for the training set

Supplementary Table 5 (related to Table 4). Sensitivities and specificities at other

operating points

Supplementary Table 6. Subgroup analyses

Supplementary Table 7. AUC with different cut-off values

Legends

Figure 1: Graphical illustration of method of machine-learning

Figure 2: Histogram and ROC curves of each vitamin B value

(A-C) The histograms for vitamin B_1, vitamin B_{12}, and folate (vitamin B_9). Their medians (1^{st}–3^{rd} quartile) are 35 (30–42) ng/mL, 285 (206–431) ng/L, and 7.2 (4.9–10.8) µg/L, respectively.

(D-F) ROC curves for vitamin B_1, vitamin B_{12}, and folate. Operating points used in **Table 4** and **Supplementary Table 5** are depicted in blue.

Abbreviations: Vit B_1, vitamin B_1; Vit B_{12}, vitamin B_{12}.

Figure 3: Gini importance and partial dependence plots of vitamin B deficiencies

The Gini importance (A-C) and partial dependency plots of the probability of deficiency (D-F) are shown for the eight most important variables for vitamin B_1, vitamin B_{12}, and folate (vitamin B_9). Combined with these, this machine-learning classifier without hypothesis also provided further evidence of a relationship between vitamin B levels and the complete blood count while also indicating a potential association between these vitamins and alkaline phosphatase (ALP) or thyroid-stimulating hormone (TSH).

Abbreviations: Vit B_1, vitamin B_1; Vit B_{12}, vitamin B_{12}; Hb, hemoglobin; Hct, hematocrit; WBC, white blood cell count; CK, creatine kinase; RDW.CV, red blood cell distribution width-coefficient variation; Plt, platelet; ALT, alanine transaminase; Lym, lymphocyte fraction; Cre, creatinine; Neu, neutrocyte fraction; γGTP, γ-glutamyltransferase; MCV, mean corpuscular volume; glu, plasma glucose.

Figure 4: Graphical abstract

Vitamin deficiency predicted by ML

386 **Table 1. Patient distribution data (n = 497)**

	ICD-10 code										VitB$_1$			VitB$_{12}$			Folate		
	F0	F1	F2	F3	F4	F5	F6	F7	F8	F9	<20	<28	<30*	<150	<180*	<200	<3.0	<4.0*	<5.0
N	28	21	300	58	16	0	29	20	24	1	15	81	112	37	80	107	29	72	134
%	5.6	4.2	60.4	11.7	3.2	0.0	5.8	4.0	4.8	0.2	3.0	16.3	22.5	7.4	16.1	21.5	5.8	14.5	27.0

Asterisks show the predefined cut-off values for vitamin B$_1$, vitamin B$_{12}$, and folate

(vitamin B$_9$) based on a reference[12]; different cut-off values based on other papers[14–16]

are also presented for further investigation.

ICD-10 codes. F0, Mental disorders due to known physiological conditions; F1, Mental

and behavioral disorders due to psychoactive substance use; F2, Schizophrenia,

schizotypal, delusional, and other non-mood psychotic disorders; F3, Mood disorders;

F4, Anxiety, dissociative, stress-related, somatoform, and other non-psychotic mental

disorders; F5, Behavioral syndromes associated with physiological disturbances and

physical factors; F6, Disorders of adult personality and behavior; F7, Intellectual

disabilities; F8, Pervasive and specific developmental disorders; F9, Behavioral and

emotional disorders with onset usually occurring in childhood and adolescence.

Vitamin deficiency predicted by ML

402 **Table 2. Vitamin B deficiencies in sub-groups**

	F0	F1	F2	F3	F4	F6	F7	F8	F9
$vitB_1 < 30$	9 (32%)	4 (19%)	70 (23%)	11 (19%)	3 (19%)	7 (24%)	5 (25%)	3 (13%)	0
$vitB_{12} < 180$	5 (18%)	4 (19%)	53 (18%)	7 (12%)	3 (19%)	1 (3%)	4 (20%)	3 (13%)	0
Folate < 4.0	5 (18%)	7 (33%)	38 (13%)	6(10%)	5 (31%)	3 (10%)	4 (20%)	4 (17%)	0

405   Abbreviations; see **Table 1**.

Vitamin deficiency predicted by ML

Table 3. Summary of full dataset of age, sex, and 29 parameters for machine-learning

Parameters	Units	Mean	SD					
Age	years	42.3	15.4	UN	mg/dL	12.9	6.7
Sex	Woman	n = 228		Cre	mg/dL	0.7	0.2
				T.bil	mg/dL	0.7	0.4
WBC	$\times 10^3/\mu L$	8.2	2.8	Na	mmol/L	139	3
Hb	g/dL	13.7	1.7	Cl	mmol/L	105	4
Hct	%	40.3	4.5	K	mmol/L	3.7	0.4
MCV	fL	89	6.6	cor.Ca	mg/dL	9.1	0.5
Plt	$\times 10^4/\mu L$	24.9	6.3	CK	IU/L	514	1230
RDW.CV	%	13.5	1.3	AST	IU/L	31	34
Neu	%	70	11	ALT	IU/L	27	24
Lym	%	23	10	LDH	IU/L	239	91
Mono	%	6	2	ALP	IU/L	224	81
Eo	%	1	2	γGTP	IU/L	37	63
Baso	%	0	0	Glu	mg/dL	112	40
TP	g/dL	7.2	0.6	CRP	mg/dL	0.4	0.9
Alb	g/dL	4.4	0.4	TSH	μIU/mL	1.7	2.4

Two patients lacked age data (no photo ID was available), and one patient lacked biochemistry data (inappropriate sample processing). For machine-learning, the missing values were replaced using the mean.

Abbreviations: WBC, white blood cell count; Hb, hemoglobin; Hct, hematocrit; MCV, mean corpuscular volume; RDW.CV, red blood cell distribution width-coefficient variation; Plt, platelet; Neu, neutrocyte fraction; Lym, lymphocyte fraction; Mono, monocyte fraction; Eo, eosinocyte fraction; Baso, basocyte fraction; TP, total protein; Alb, albumin; UN, urea nitrogen; Cre, creatinine; T.bil, total bilirubin; Na, sodium; Cl,

Vitamin deficiency predicted by ML

421   chloride; K, potassium; cor.Ca, corrected calcium; CK, creatine kinase; AST, aspartate

422   transaminase; ALT, alanine transaminase; LDH, lactate dehydrogenase; ALP, alkaline

423   phosphatase; γGTP, γ-glutamyltransferase; Glu, plasma glucose; CRP, C-reactive

424   protein; TSH, thyroid-stimulating hormone.

Vitamin deficiency predicted by ML

Table 4. Summary of AUC, sensitivity, specificity, and accuracy for the validation set

	vitB$_1$	vitB$_{12}$	Folate
AUC	0.716	0.599	0.796
Sensitivity	0.594	0.316	0.667
Specificity	0.783	0.943	0.917
Accuracy	0.688 [0.597–0.787]	0.629 [0.523–0.746]	0.792 [0.665–0.909]

Generalization performance of the classifiers was evaluated using AUC of the validation set. Sensitivity, specificity, and accuracy of the classification at the optimal operating points that maximized accuracy on the receiver operating characteristic curve of the validation set are also shown (see also **Figure 2 D-F**). Accuracy was defined as the average of the sensitivity and specificity. Square brackets indicate the 95% CI. Note that the 95% CI of each accuracy does not include 0.5, which demonstrates statistical significance. For further information, see **Figure 2** and **Supplementary Table 5**.

Abbreviations: AUC, area under the receiver operating characteristic curve; CI, confidence interval.

Vitamin deficiency predicted by ML

Figure 1

Figure 2

Figure 3

Figure 4