Transmission Dynamics of and Insights from the 2018-2019 Measles Outbreak in New York City: A Modeling Study

Wan Yang (wy2202@columbia.edu)

Department of Epidemiology, Mailman School of Public Health, Columbia University, NY, NY

Abstract:

In 2018-2019, New York City experienced the largest measles outbreak in the US in nearly three decades. To identify key factors contributing to this outbreak to aid future public health interventions, here we developed a model-inference system to infer the transmission dynamics of measles in the affected community, based on incidence data. Our results indicate that delayed vaccination of young children aged 1-4 years enabled the initial spread of measles and that increased infectious contact among this age group, likely via gatherings intended to expose unvaccinated children (i.e. "measles parties"), further aggravated the outbreak and led to widespread of measles beyond this age group. We found that around half of infants were susceptible to measles by age 1 (the age-limit to receive the first vaccine dose in the US); as such, infants experienced a large number of infections during the outbreak. We showed that without the implemented vaccination campaigns, the outbreak severity including numbers of infections and hospitalizations would be 10 times higher and predominantly affect infants and children under 4. These results suggest that recommending the first vaccine dose before age 1 and the second dose before age 4 could allow pro-vaccine parents to vaccinate and protect infants and young children more effectively, should high level of vaccine hesitancy persist. In addition, enhanced public health education is needed to reduce activities that unnecessarily expose children to measles and other infections.

Introduction

Measles is highly contagious and severe viral disease. Thanks to a highly effective vaccine and high coverage of vaccination, endemic transmission of measles—i.e. continuous transmission for more than 12 months—in the US was declared eliminated in 2000. However, due to vaccine hesitancy and declining vaccination rate, in recent years there have been an increasing number of large outbreaks following introduction of measles infection (*1*). Due to long-term fluctuations in vaccine coverage and infection history, population susceptibility could vary substantially by age group. This susceptibility disparity by age can further interact with age-specific social connectivity (i.e., contact rate) to shape the epidemic trajectory. As such, understanding these detailed population characteristics as well as their impact on transmission dynamics in the recent outbreaks is important for devising timely and effective intervention strategies.

In the fall of 2018, several New York City (NYC) residents acquired measles while traveling abroad and subsequently led to the largest measles outbreak in the US in nearly three decades. The first case of this outbreak developed a rash on Sep 30, 2018 and as of Aug 6, 2019 (at the time of this writing), there have been 642 confirmed cases, largely occurring in an Orthodox Jewish community (*2, 3*). To contain the outbreak, the NYC Department of Health and Mental Hygiene (DOHMH) launched extensive vaccination campaigns and, on Apr 9, 2019, ordered mandatary vaccination of all individuals living, working or going to school in the affected zip codes. As a result, over 32 thousand individuals under 19 years were vaccinated with the measles, mumps, and rubella (MMR) vaccine during Oct 2018 – July 2019 and the outbreak subsided (*3, 4*).

In this study, we model the transmission dynamics of this measles outbreak in the

affected Orthodox Jewish community in NYC from Oct 1, 2018 to July 31, 2019, months with

more than one measles cases reported. Using an age-structured model-inference framework, we

are able to estimate key epidemiological features including the initial susceptibilities in five

different age groups (i.e., <1, 1-4, 5-17, 18-49, and 50+ years) and the basic reproductive number

R_0, infer key factors contributing to the spread of measles, estimate the proportions of infection

attributable to each age group, as well as assess the impact of vaccination campaigns. We also

discuss the implications of our findings to current measles vaccination policies.

Results

Overview of the measles outbreak and model fit

The measles outbreak started on Sep 30, 2018, when a young child developed a rash. It evolved

relatively slowly in the first three months; however, the outbreak took off quickly in early 2019,

peaked in April after the city declaring a public health emergence, and recorded a total of 642

cases by July 31, 2019. As shown in Fig. 1, age-grouped incidence, estimated based on health

reports/alerts (*3, 5*), peaked first in Mar 2019 among 1-4 year-olds—the age group with the

largest number of infection (275 cases or 42.8% of the total, as of Aug 6, 2018), followed by <1

year-olds (100 cases or 15.6%) and 5-17 year-olds (138 cases or 21.5%) in Apr 2019, and 18+

year-olds (129 cases or 20.1%) in May 2019.

As shown in Fig. 2 and Table S2, our model-filter system was able to recreate the overall

incidence curve during Oct 2018 – July 2019, estimate the overall age distribution of measles

cases, as well as recreate the estimated age-grouped incidence curves for all age groups. Note

that while our model (Eqn 1) divided 18+ year-olds into two subgroups (i.e. 18-49 and 50+

years) given their differences in contact rates and interactions with other age-groups, we present

the combined results here because data were only available for the entire age group. The

estimated reporting rate was around 90% throughout the study period and slightly lower in Apr

2019, at the peak of the outbreak [89.1%, 95% credible interval (CI): 79.4, 99.2%; Fig. S2].

Inference of key epidemiological characteristics

The model-filter system estimated that, at the beginning of the outbreak (i.e. Sep 2018),

susceptibility was the highest in infants, at approximately 53.2% (95% CI: 49.0, 57.5%; Fig.

3A). This was expected, because maternal immunity wanes within 3 to 9 months after birth and,

as a result, by their first birthday—the age eligible to receive the first dose of MMR vaccine in

the US—almost all infants have lost their maternal immunity and are susceptible to measles.

Young children aged 1-4 years had the second highest susceptibility; approximately 24.9% (95%

CI: 20.4, 29.7%) were susceptible. In comparison, susceptibility was lower among both 5-17

year-olds (6.0%, 95% CI: 4.1, 7.9%) and 18+ year-olds 6.0% (95% CI: 4.4, 7.6%; Fig. 3). These

estimates were consistent with the observation that all cases recorded in Oct 2018 were children

ranging from 11 months to 4 years (*2*). Sensitivity analysis on assumptions related to the

vaccination campaigns showed that estimated susceptibilities were slightly higher for 5-18 year-

olds (7.5%, 95% CI: 5.1, 9.9%) and 18-49 year-olds (7.9%, 95% CI: 5.2, 9.9%) if more vaccine

doses were given to the Jewish Orthodox community or 5-18 year-olds; however, the estimates

were in general consistent with the baseline scenario (Table S2).

The initial vaccination campaigns launched promptly afterwards lowered the

susceptibility to 40.3% (95% CI: 36.4, 44.0%) in infants and 13.9% (95%: 9.7, 18.7%) in 1-4

year-olds, by the end of Dec 2018 (Fig. 3). These efforts along with other transmission and

infection controls (*2*) appeared to effectively contain the outbreak at the time. The effective

reproductive number (R_e) is a key epidemiological parameter reflecting the potential of an

infection to cause an epidemic in a partially immune population; an epidemic is possible when

R_e>1. The estimated R_e was 1.5 (95% CI: 0.7, 2.9) in Oct 2018 and dropped to round 1 (95% CI:

0.7, 1.5) in Dec 2018 (Fig. 4B).

The outbreak, however, took off again in early 2019 (Fig. 1). The estimated R_e increased

and remained above 1 in the first three months of 2019 (Fig. 4B). In a perfectly mixed model, R_e

is computed as the product of the basic reproductive number R_0 and population susceptibility. In

particular, the basic reproductive number R_0 measures the transmissibility of an infection in a

fully susceptible population; for measles, while often reported in the range of 12-18, R_0 could

vary from 1.4 to 770 (*6*). In this study, we estimated that R_0 was approximately 7 during the

entire outbreak (Fig. 4A). Population susceptibility, the other factor for R_e, were to increase in a

close population (i.e., without migration), could only do so slowly as infants lose maternal

immunity. Thus, these two factors alone could not explain the sudden large increase in R_e (Fig.

4B). In this study, we utilized an age-structured model that enables more detailed analysis of the

transmission dynamics. Indeed, our model-filter system detected an increase in contact rate

among 1-4 year-olds during Jan-March 2019 (Fig. 4D); this increased contact rate along with the

high susceptibility among 1-4 year-olds appeared to raise R_e above unity (Eqn 5 in Methods) and

contribute to the re-surge of measles in early 2019. This finding was also consistent with reports

of parents hosting "measles parties" to exposure unvaccinated children at the time (*7*). In

contrast, estimated contact rates were relatively stable for other age groups (e.g., Fig. 4 E and F

for 5-17 and 18-49 year-olds, respectively). As a result, infections increased quickly among 1-4

year-olds (Figures 1 and 4D), reaching a peak of around 80 cases in March. Meanwhile,

infections also increased in other age groups including 5-17 and 18+ year-olds despite their low

overall susceptibilities, due to interactions between age-groups (Fig. S2) and the high contact

rates in these groups (Fig. 4 E and F).

The outbreak began to decline in Apr 2019, following more stringent public health

interventions (*3, 4*). In particular, the model estimated that, thanks to extensive vaccination

campaigns, susceptibility was reduced to 22.8% (95% CI: 19.3, 26.0%) in <1 year-olds, 4.3%

(95% CI: 0.5, 8.9%) in 1-4 year-olds, and 2.4% (95% CI: 0.4, 4.4%) in 5-17 year-olds at the end

of May. Consequently, the effective reproductive number R_e dropped below 1 from April

onwards.

Who acquired infection from whom?

Table 1 shows the estimated proportions of infections caused by each of the five age groups

based on the estimated Who-Acquires-Infection-From-Whom (WAIFW) contact matrix (Eqn 2

in Methods). Children aged 1-4 years not only had the largest number of infections (42.8%), but

also appeared to cause the largest number of infections in other age groups. Tallied over the

entire study period (Oct 2018-July 2019), an estimated 51.6% (95% CI: 39.3, 63.1%) of the total

cases were infected by 1-4 year-olds, compared to 25.2% (95% CI: 15.8, 35.5%) by 5-17 year-

olds, 17.7% (95% CI: 10.4, 26.5%) by 18-49 year-olds, 4.5% (95% CI: 3.0, 6.4%) by <1 year-

olds, and 1.0% (95% CI: 0.4, 1.8%) by 50+ year-olds. In particular, this age group caused

around half of the infections in infants (44.6% or 45 cases) as well as the largest proportions of

inter-group transmission to other age-groups (ranging from 12.9% to 5-17 year-olds to 40.1% to

50+ year-olds; Table 1).

Estimated impact of vaccination campaigns

Figure 5 shows the estimated outbreak outcomes had there be no vaccination campaigns.

Without the vaccination campaigns, the model estimated that the outbreak could continue to the

end of 2019 and infect a total of 6196 (95% CI: 5, 8478) people by then or 6078 (95% CI: 5,

8433) during the observed outbreak period (i.e. Oct 2018 – July 2019), compared to 642 cases

reported as of Aug 6, 2019. In addition, these infections would largely occur in infants under 1

and young children aged 1-4 years. During the observed outbreak period, there would be 1015

(95%: 0, 1430) infections in infants and 3052 (95% CI: 2, 4096) infections in 1-4 year-olds,

more than 10 times of the reported numbers in these two age groups (i.e. 100 and 275,

respectively, as of Aug 6, 2019; Table 2). Children aged 5-17 years would have the third largest

number of infections, with 1107 (95% CI: 1, 1692) cases, 8 times of the reported number (i.e.

138 as of Aug 6, 2019).

Measles can cause severe diseases. According to data by Apr 24, 2019, among 390

individuals with measles, 29 were hospitalized, of which six needed intensive care (*8*). Assuming

the same ratios among all cases, without the vaccination campaigns, during the observed

outbreak period there would be 452 (95% CI: 0, 628) hospitalizations, including 94 (95% CI: 0,

130) needing intensive care, and the majority would be in young children under 4 (Table 2).

Discussion

Using a model-filter inference system, we have reconstructed in detail the transmission dynamics

of the measles outbreak in an Orthodox Jewish community in NYC during Oct 2018 – July 2019.

We have estimated the population characteristics (e.g. age-specific susceptibilities) and

epidemiological parameters (e.g. reproductive numbers) as well as subtle changes in key

parameters (e.g. contact rates) that are critical to the transmission of measles. Using model

simulation and the posterior estimates from the model-inference system, we are also able to

estimate the impact of vaccination campaigns implemented during the outbreak, including

numbers of infections and hospitalizations averted, for each age group. These latter findings echo

those from previous studies (*9-11*) and again highlight the severity of measles disease, should

there be no effective infection and transmission controls (in particular, vaccination).

Our analyses estimate that around a quarter of young children aged 1-4 in the affected

community were susceptible at the onset of the outbreak, likely due to delayed vaccination.

Indeed, 94% (101/108) of the early infections in children were unvaccinated (*12*). In contrast,

vaccination rate remained high in older children 5-17 years, with an estimated 94% immune to

measles. This difference may be due to better compliance with vaccination regulation at school

entry, or a result of vaccination campaigns in response to previous outbreaks (e.g., a large

outbreak occurred in the same community in 2013 (*13*)). Nevertheless, the large number of

unvaccinated children under 4 was sufficient to cause many infections in late 2018,

predominantly in the same age group (Fig. 1 and references (*2, 12, 14*)). This observation

highlights the importance of vaccination compliance with both MMR vaccine doses, especially

given the long lag between the two vaccine doses. In addition, recommending the second MMR

dose earlier than the currently scheduled age 4-6 years could allow pro-vaccine parents to fully

vaccinate their children sooner and reduce the number of susceptible children overall.

Our study also reveals intricate interplays of population dynamics and measles

transmission. While the high susceptibility in children under 4 was likely responsible for the

early spread of measles, our estimates suggest that the second and more severe part of the

outbreak in 2019 was likely due to increased infectious contact among this age group, likely

facilitated by parents hosting "measles parties" that intentionally bring unvaccinated children

together and expose them to those sick with measles (*7*). As shown in Fig. 4, the increase in

infectious contact interacting with the high susceptibility in 1-4 year-olds was able to raise the

effective reproductive number R_e to above unity—the threshold for an epidemic to occur—and

aggravate the outbreak in 2019, despite earlier public health efforts that had reduced R_e to below

1 in late 2018. Similar disease-related gatherings have been noted in previous measles outbreaks

(*15*) as well as other disease outbreaks (*16*). These activities create further challenges for the

control of measles spread and stress the need for enhanced public health education.

In addition, the intensified measles outbreak not only affected children with delayed

vaccination, but also a large number of infants under 1, who were too young to receive their first

dose of MMR vaccine in the US. At least 100 infants under 1 were infected with measles during

the 10-month outbreak period, despite extensive infection and transmission control efforts

including vaccinating infants 6 months or older and post-exposure prophylaxis with immune

globulin given to those under 6 months (*2, 14*). This was largely a result of the high

susceptibility in infants. Our model-inference system estimates that about half of infants were

susceptible by age 1 and that nearly half of the 100 infant cases were infected by 1-4 year-olds

(Table 1). In addition, our simulations suggest that, without the vaccination campaigns, the

number of infections in infants would be over 10 times higher than observed (Table 2). These

findings demonstrate the rippling effects of vaccine hesitancy beyond the risk posted to age-

eligible children with delayed vaccination. These findings also suggest that administration of the

first dose of routine MMR vaccine earlier than the current 1 year age-limit in the US may be

necessary to protect infants should high level of vaccine hesitancy persist. Of note, the World

Health Organization (WHO) recommends administering the first dose of measles vaccine at 9-12

months of age for routine vaccination programs and as early as 6 months for settings such as

during an outbreak (*17*).

Our model simulations, consistent with many previous studies (*11, 15, 18*), demonstrate

the significant public health impact of vaccination in controlling measles outbreaks. Without the

implemented vaccination campaigns, the severity of the measles outbreak—including number of

infections, hospitalizations, and severe infections needing intensive care—would be about 10

times worse than observed (Table 2). These estimates, however, did not include the long-term

health impacts on affected individuals, particularly young children (*10, 19, 20*), nor the

enormous economic burdens (*13, 21, 22*).

We note several limitations of our study. First, we did not explicitly model the impact

of public health interventions other than the vaccination campaigns, due to a lack of data. During

the outbreak, such efforts included prescreening patients prior to presence for treatment, post-

exposure prophylaxis, and closing schools out of vaccination compliance (*2, 12, 14, 23, 24*). Of

note, however, here the estimated basic reproductive number R_0 was around 7, lower than the 12-

18 range based on epidemics in the pre-vaccine era (*25*); and the estimated infectious period was

around 4 days (Fig. 4C), at the lower end of the commonly used range of 4-6 days (*26*). The

lower R_0 and shorter infectious period could be a result of the aforementioned public health

interventions. Second, there were uncertainties in the accuracy of case reporting. Because the

incidence data used here were published on Aug 6, 2016, a later revision of case reports, we used

a relatively high but broad prior range for the reporting rate (i.e. 80-100%). In addition, our

model-inference system explicitly accounted for observational errors (Eqn 7a and 7b). Third,

due to a lack of contact data and for simplicity, we set all terms related to group-1 (i.e. <1 year-

olds) in the WAIFW matrix to the same as the contact rate within the group (Eqn 2). The model

formulation may have led to under-estimation of the proportions of infection in <1 year-olds

attributable to the same age group and/or 1-4 year-olds, given the likely more frequent contact

within the same age group and with similar ages (i.e. 1-4 years) due to more shared settings such

as daycares and pediatric hospitals. Lastly, there were uncertainties regarding the settings of

vaccination campaigns. Nevertheless, sensitivity analysis showed that our main estimates were

robust to a wide range of assumptions (Table S2).

In summary, using a comprehensive model-inference system, we have reconstructed

transmission dynamics of the recent measles outbreak in NYC in great detail. Our estimates

highlight the importance of vaccination in protecting children as well as public health education

to reduce activities that unnecessarily expose children to risk of measles infection. Further, in

light of the persistent vaccine hesitancy, revising current vaccination recommendations may

allow pro-vaccine parents to vaccinate and protect their children more effectively.

Materials and Methods

The measles outbreak occurred predominantly among members of the Orthodox Jewish

community in Williamsburg and Borough Park, two neighborhoods in located in Brooklyn,

NYC. As such, we focused on modeling the outbreak in this subpopulation. Estimated based on

the Jewish Community Study of New York (*27*), approximately N=165,970 Orthodox Jews live

in these two NYC neighborhoods, of which 4552 (2.7%), 18,208 (11%), 59,176 (35.7%), 60,445

(36.4%), and 23,589 (14.2%) are <1, 1-4, 5-17, 18-49, and 50+ years, respectively.

Estimating monthly incidence by age group

Monthly measles incidence aggregated over all ages from Sep 2018 to Aug 2019 and the age

distribution of case-patients over the entire outbreak were published on the NYC DOHMH

website (*3*). Of note, 1 case was reported in Sep 2018 (i.e., the initial case) and none were

reported in Aug 2019 at the time of this writing. In addition, the numbers of reported cases were

subsequently revised by the DOHMH (often adjusted upwards, presumably from retrospective

case identification) and, as such, varied over time. To estimate the monthly incidence for each

age group, we used the age distribution of cases reported in earlier health reports/alerts (*5*) to

apportion the total incidence for each month. For months without age information, we used

estimates either from the preceding month or the following month back-calculated from the

overall age distribution.

Transmission model

The transmission model used here was similar to described in our previous study (*28*). As

illustrated in Fig. S1, the model represents the susceptible-exposed-infectious-recovered (SEIR)

disease dynamics with five age groups (i.e. <1, 1-4, 5-17, 18-49, and 50+ years) to account for

population differences by age group (e.g., susceptibility and contact rate), routine two-dose

vaccination at ages 1 and 5, and immunization during the vaccination campaigns per Eqn 1:

281
$$
\begin{cases}
\dfrac{dS_i}{dt} = -\left(\dfrac{S_i^{m_1}}{N_i}\right)\sum_{j=1}^{5}\beta_{ij}I_j^{m_2} + k(1-v)N\mathbf{1}_{i=1} + \dfrac{M}{180}\mathbf{1}_{i=1} + l_{i-1}S_{i-1}(1-v_{i-1}) - l_iS_i - V_i(t) \\[2ex]
\dfrac{dE_i}{dt} = \left(\dfrac{S_i^{m_1}}{N_i}\right)\sum_{j=1}^{5}\beta_{ij}I_j^{m_2} - \dfrac{E_i}{Z} + \alpha_i(t) + l_{i-1}E_{i-1} - l_iE_i \\[2ex]
\dfrac{dI_i}{dt} = \dfrac{E_i}{Z} - \dfrac{I_i}{D} + l_{i-1}I_{i-1} - l_iI_i \\[2ex]
\dfrac{dR_i}{dt} = \dfrac{I_i}{D} + l_{i-1}R_{i-1} - l_iR_i + l_{i-1}S_{i-1}v_{i-1} + V_i(t) \\[2ex]
\dfrac{dM}{dt} = kvN - \dfrac{M}{180}
\end{cases}
$$

for *i*=1,..,5 (Eqn 1)

where S_i, E_i, I_i, R_i and N_i are, respectively, the numbers of susceptible, exposed (i.e. latently

infected), infectious, recovered (and/or immunized) people and population size in the *i*-th age

group; *M* is the number of infants with maternal immunity, which decays exponentially with a

mean duration of 180 days; *t* is time in days. $V_i(t)$ is the number of people in group-*i* immunized

by the vaccination campaigns on day-*t* (described in detail in the next section). The exponents

m_1 and m_2 describe the level of inhomogeneous mixing (*29, 30*); and $m_1=m_2=1$ represents

homogeneous mixing. *Z* and *D* are the latent and infectious period, respectively.

To model the different contact rates within and between age groups, we used 7

parameters for the Who-Acquires-Infection-From-Whom (WAIFW) matrix as follows:

293
$$\boldsymbol{\beta} = \begin{bmatrix} \beta_1 & \beta_1 & \beta_1 & \beta_1 & \beta_1 \\ \beta_1 & \beta_2 & \beta_6 & \beta_7 & \beta_1 \\ \beta_1 & \beta_6 & \beta_3 & \beta_7 & \beta_1 \\ \beta_1 & \beta_7 & \beta_7 & \beta_4 & \beta_1 \\ \beta_1 & \beta_1 & \beta_1 & \beta_1 & \beta_5 \end{bmatrix} \quad \text{(Eqn 2)}$$

294 where β_1 to β_5 represent within-group contact for the five age groups and β_6 and β_7 represent

295 mixing between siblings and child-parent, respectively. For simplicity, we set all interactions

296 with group 1 (i.e. <1 year) or group 5 (50+ years) to β_1, the lowest contact rate. For group-3 (5-

297 17 years), to capture the varying contact rate following school schedules, we adjusted β_3 for each

298 date per the school calendar in NYC as:

299
$$\beta_3(t) = \frac{\beta_3}{b_{Term}} \cdot [1 + b_1 Term(t)] \quad \text{(Eqn 3)}$$

300 where b_1 is the amplitude of school term-time forcing; $Term(t)$ is set to 1 for school days and -1

301 for non-school days; and b_{Term} is the yearly average of $1+b_1 Term(t)$ (*31*).

303   The basic reproductive number R_0, defined as the average number of secondary infections

304 caused by a primary case-patient in a *naïve* population, reflects the transmissibility of an

305 infection. In an age-structured model, R_0 is computed as:

306
$$R_0 = \text{eigen}_{max}(\boldsymbol{n\beta}D) \quad \text{(Eqn 4)}$$

where $\text{eigen}_{max}(\cdot)$ denotes the function giving the largest eigenvalue of a matrix, and $\boldsymbol{n}$ is a

diagonal matrix with elements $n_i=N_i/\sum N_i$ ($i=1, \ldots, 5$ here), i.e. the fraction of population in

group-i. Based on this relationship between R_0 and the $\boldsymbol{\beta}$ matrix, we reparametrized the model to

include R_0 as a model parameter by setting β_1 to 1 and estimating the relative magnitude of β_2–β_6,

all scaled to R_0. In the current mass vaccination era, most people are immune via vaccination. To

reflect the potential of an infection to cause an epidemic in a partially susceptible population, the

effective reproductive number, R_e, accounts for population susceptibility and is computed as:

$$R_e = \text{eigen}_{\max}(\boldsymbol{s\beta}D) \qquad \text{(Eqn 5)}$$

where $\boldsymbol{s}$ is a diagonal matrix with elements $s_i=S_i/N_i$ (i=1, …, 5 here), i.e. the susceptibility in

group-i.

To model the demographic processes, k is the birth rate (2.7 per 1000 person-year here

(*27*)); v is the immunity level in mothers, approximated by the susceptibility of the child-bearing

age-group (i.e., 18-49 year-olds); N is total population size; and $\mathbf{1}_{i=1}$ is an indicator function, with

value 1 for group-1 (<1 year-olds) and 0 for all other groups. Thus, the term $k(1-v)\,\mathbf{1}_{i=1}$ (1[st] line in

Eqn 1) is the number of susceptible newborns and kvN (last line in Eqn 1) is the number of

newborns with maternal immunity. The term l_i is the rate of aging for group-i (i.e. the inverse of

the sojourn time in each age group) with l_0 set to 0 and l_5 set to the death rate. The term v_i (for

i=1 and 2) is the vaccination rate for the two doses of vaccine (i.e., v_i=0 for i=0, 3, and 4). In this

study, we set v_1 (1[st] dose) to 0.9 times the immunity level of group-2 (i.e., assuming a 90%

vaccine efficacy) and v_2 (2[nd] dose) to 0.7 for days before May 2019; for days afterwards, we used

0.8 for v_1 and 0.9 for v_2, corresponding to a 1-(1-0.8)(1-0.9)=98% overall vaccination rate.

Based on NYC health reports/alerts (*2, 12, 14*), we seeded the model, via the parameter

$\alpha_i(t)$, with 3 cases in group-2 (i.e. 1-4 year-olds)—one each with rash onset on Sep 30, Oct 15,

and Oct 30, 2018, respectively—and 1 case each in group-3 (5-17 year-olds) and group-4 (18-49

year-olds), both during the winter recess (from 12/24/18 to 1/1/19).

Modeling the vaccination campaigns

To contain the outbreak, the NYC DOHMH conducted extensive vaccination campaigns and

administered 31,790 doses of MMR vaccine to children under 19 years in Williamsburg and

Borough Park by July 2019 (*3*). However, information on the age and immune status of

vaccinees was not reported. In this study, per the health reports/alters (*3, 12*), we assumed there

were two phases of vaccination campaign: 1) Oct 2018 – Feb 2019, during which 7000 children,

90% Orthodox Jewish (15% were <1 year, 65% 1-4 years, and 20% 5-17 years) were vaccinated;

and 2) Mar – July 2019, during which 24,790 children, 60% Orthodox Jewish (10%, 40%, and

50%, respectively, were <1, 1-4, and 5-17 years) were vaccinated. For reference, Orthodox Jews

made up for approximately 30% of the total population in the two affected neighborhoods.

For <1 year-olds, immunization could fail due to residual maternal immunity; thus, we

assumed a 85% immunization success rate for group-1. For those above 1 year, some vaccinees

might have received 1 or 2 doses of vaccine and higher the population susceptibility, the less

likely a vaccinee had been immune before the additional vaccine dose. As such, we assumed the

immunization success rate was twice the group-specific susceptibility for 1-4 year-olds and three

times that for 5-17 year-olds or at a minimum of 25% and a maximum of 75%. We further

assumed a 10-day delay in vaccine effect and computed the daily number of individuals

vaccinated per a gamma distribution (mean=30 days and standard deviation=21 days for Phase-1

and mean=56 and standard deviation=15 days for Phase-2 such that it peaked ~1 week after Apr

9, 2019 when the city implemented a vaccination mandate). The estimated numbers matched

with the reports (e.g., 1740 doses by our model vs ~1600 doses given to children under 5 as

reported (*32*) and 1142 doses by our model vs ~1000 doses given in March 2019 as reported (*7,*

358 *33*)). These daily numbers were then included in the transmission model [i.e. $V_i(t)$ in Eqn 1].

359 Sensitivity to model assumptions were tested as described below.

361 *Estimation of model state variables and parameters*

362 To estimate the model state variables (i.e., S_i, E_i, I_i, R_i, and M) and parameters (β_2 to β_7, R_0, b_1, D,

363 Z, m_1, and m_2), we fitted the model to the reported monthly overall incidence and the estimated

364 monthly age-grouped incidence using a particle filter (*34*). Briefly, we first initialized a suite of

365 model realizations (termed "particles", N=10,000 here) using Latin Hypercube sampling (*35*)

366 from the prior distribution of state variables and parameters (Table S1). The particle filter then

367 sequentially incorporated the monthly incidence to the model via repeated prediction-update

368 cycles. In each cycle (i.e., each month here), the particles were stochastically integrated forward

369 in time for a month per the model (i.e., Eqn 1; this generates the prediction). To update the model

370 state including all model variables and parameters, at the end of each month, the model-

371 estimated incidence was aggregated for the month, adjusted by the reporting rate for that month

372 (estimated simultaneously by the filter), and used to compute the likelihood of each particle

373 (described below). The posterior of model state was then computed using Bayes' rule (*34, 36*)

374 and the particles resampled and updated—those with high posterior probabilities were retained

375 and those with very low posterior probabilities discarded.

377   To allow for a wider observational variance than, e.g., the Poisson process, we

378 heuristically modeled the observations using a multivariate Gaussian distribution (i.e. the

379 likelihood function):

380        $$Y_m | r, C_m \sim \mathcal{N}(r C_m, \Sigma) \quad \text{(Eqn 6)}$$

where Y_m is the vector of monthly incidence reported for month-m, including the monthly

incidence for individual age-groups (i.e., <1, 1-4, 5-17 and 18+ years; note that 18-49 and 50+

year-olds were combined due to a lack of data for these two groups separately) and all ages

combined. Correspondingly, C_m is the vector of monthly incidence estimated by the model; and r

is the reporting rate and, for simplicity, assumed the same for all ages. Σ is the covariance

matrix, with the off-diagonal terms set to 0. To account for uncertainties in the estimated age-

grouped incidence, the variance ($\Sigma_{ii}, i = 1, ..., 4$) for each of the 4 aforementioned age-groups

was heuristically computed as:

$$\Sigma_{ii,m} = 100 + \frac{\left(\sum_m^{m-2} Y_m/3\right)^2}{3} \text{ (Eqn 7a)}$$

That is, the observational variance is proportional to the average incidence in the preceding two

months (if available) and the current month, plus a baseline constant. For the overall incidence

with detailed data, a smaller variance was used:

$$\Sigma_{ii,m} = 100 + \frac{\left(\sum_m^{m-2} Y_m/3\right)^2}{5} \text{ (Eqn 7b)}$$

As there were great uncertainties in the susceptibilities of the younger age groups, we

tested prior ranges from 10-40% for 1-4 year-olds, 5-20% for 5-17 year-olds, and 5-15% for 18-

49 year-olds. For the basic reproductive number R_0, we tested prior values ranging from 5 to 12

(note these values were lower than the oft-reported 12-18 range (*6, 25*)). To optimize the model-

filter system, we parsed these wide ranges into smaller segments and tested all combinations by

permutation (5040 in total; see specific prior ranges in Table S1). To account for model

stochasticity, we ran the model-filter system 5 times for each prior combination and 10 times for

the final prior select. We then selected the optimal priors based on the model-goodness-of-fit to

the data (minimal root-mean-square-error and maximal correlation and likelihood) over the

period of Oct 2018 – July 2019 as well as accuracy of the one-step-ahead predictions (recall that

the particle filtering process comprises sequential prediction-update cycles) for the period of Oct

2018 – March 2019 (i.e. before the emergency vaccination mandate). We pooled all 10 final runs

(10,000 particles each run and 100,000 model realizations in total) to compute the posterior

estimates (e.g., mean and 95% CI).

Sensitivity analysis on vaccination campaigns settings

To test the sensitivity of model results to assumptions on vaccination campaign settings, we

tested the model-filter system using the following alternative scenarios:

1) For the 2nd phase (Mar-July 2019), 90% (vs. 60% in the baseline scenario) of the

vaccine doses were given to members of the Orthodox Jewish community;

2) For the 2nd phase, the age distribution of vaccinees was the same as the 1st phase (i.e.

15%, 65%, and 20%, respectively, for <1, 1-4, and 5-17 year-olds vs. 10%, 40%, and 50% for

the three groups in the baseline scenario);

3) For the 2nd phase, 90% of the vaccine doses were given to Orthodox Jewish and the

age distribution of vaccinees was the same as the 1st phase (i.e. 15%, 65%, and 20% for <1, 1-4,

and 5-17 year-olds, respectively).

Because population susceptibility would be affected by the number of individuals

immunized by the vaccination campaigns, we tested susceptibility ranges 10-40% for 1-4 year-

olds, 5-20% for 5-17 year-olds, and 5-15% for 18-49 year-olds, divided in to small segments as

for the baseline scenario (140 different combinations for each alternative scenario; Table S1).

For simplicity, we used the same optimal prior ranges for the parameters under the baseline

scenario in this sensitivity analysis.

Evaluating the impact of vaccination campaigns

To estimate the impact of vaccination campaigns, we generated model-simulated

counterfactuals—i.e., outbreak outcomes should there be no vaccination campaigns

implemented—using the posterior mean estimates of group-specific initial population

susceptibilities and model parameters for each month. We ran the model (10,000 realizations)

stochastically up to the end of 2019 to test how long the outbreak could last without intervention;

for months after Aug 2019, parameters estimated at the end of July 2019 were used.

Acknowledgements

This study was supported by the National Institute of Allergy and Infectious Diseases

(1R01AI145883-01). I thank all public health personnel and individuals involved in combating

the measles outbreak and New York City Department of Health and Mental Hygiene for making

the incidence data publicly available. I also thank Columbia University Mailman School of

Public Health for access to high performance computing.

Conflict of interest: None.

References:

1. V. K. Phadke, R. A. Bednarczyk, D. A. Salmon, S. B. Omer, Association Between Vaccine

Refusal and Vaccine-Preventable Diseases in the United States A Review of Measles and

Pertussis. *Jama-J Am Med Assoc* **315**, 1149-1158 (2016).

2.   New York City Department of Health and Mental Hygiene, ALERT # 39: Update on

Measles Outbreak in New York City in the Orthodox Jewish Community. (2018).

https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2018/alert39-measles-

outbreak.pdf.

3.   New York City Department of Health and Mental Hygiene, Measles. (2019).

https://www1.nyc.gov/site/doh/health/health-topics/measles.page.

4.   New York City Department of Health and Mental Hygiene, Order of the Commissioner.

(2019). 4/9/19. https://www1.nyc.gov/assets/doh/downloads/pdf/press/2019/emergency-

orders-measles.pdf.

5.   New York City Department of Health and Mental Hygiene, Health Alert Network. (2019).

https://www1.nyc.gov/site/doh/providers/resources/health-alert-network.page.

6.   F. M. Guerra, S. Bolotin, G. Lim, J. Heffernan, S. L. Deeks, Y. Li, N. S. Crowcroft, The

basic reproduction number (R0) of measles: a systematic review. *Lancet Infect Dis* **17**, e420-

e428 (2017).

7.   S. Scutti, New York City declares a public health emergency amid Brooklyn measles

outbreak. (2019). 4/9/19. https://www.cnn.com/2019/04/09/health/measles-new-york-

emergency-bn/index.html.

8.   New York City Department of Health and Mental Hygiene, ALERT # 9: Citywide

Recommendations during the Ongoing Measles Outbreak in New York City. (2019).

https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2019/recommendations-during-

measles-outbreak.pdf.

9. M. J. Mina, B. T. Grenfell, C. J. E. Metcalf, Response to Comment on "Long-term measles-

induced immunomodulation increases overall childhood infectious disease mortality".

Science **365**, (2019).

10. M. J. Mina, C. J. E. Metcalf, R. L. de Swart, A. D. M. E. Osterhaus, B. T. Grenfell, Long-

term measles-induced immunomodulation increases overall childhood infectious disease

mortality. *Science* **348**, 694-699 (2015).

11. P. A. Gastanaduy, S. Funk, P. Paul, L. Tatham, N. Fisher, J. Budd, B. Fowler, S. de Fijter,

M. DiOrio, G. S. Wallace, B. Grenfell, Impact of Public Health Responses During a Measles

Outbreak in an Amish Community in Ohio: Modeling the Dynamics of Transmission. *Am J*

Epidemiol **187**, 2002-2010 (2018).

12. New York City Department of Health and Mental Hygiene, ALERT # 2: Update on Measles

Outbreak in New York City in the Orthodox Jewish Community. (2019).

https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2019/update-on-measles-

outbreak-in-nyc-in-the-orthodox-jewish-community.pdf.

13. J. B. Rosen, R. J. Arciuolo, A. M. Khawja, J. Fu, F. R. Giancotti, J. R. Zucker, Public Health

Consequences of a 2013 Measles Outbreak in New York City. *JAMA Pediatr* **172**, 811-817

(2018).

14. New York City Department of Health and Mental Hygiene, ALERT # 38: Measles Outbreak

in New York City in the Orthodox Jewish Community. (2018).

https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2018/alert38-measles-

outbreak.pdf.

15. P. A. Gastanaduy, E. Banerjee, C. DeBolt, P. Bravo-Alcantara, S. A. Samad, D. Pastor, P. A.

Rota, M. Patel, N. S. Crowcroft, D. N. Durrheim, Public health responses during measles

outbreaks in elimination settings: Strategies and challenges. *Human vaccines &*

immunotherapeutics **14**, 2222-2238 (2018).

16. Centers for Disease Control and Prevention, Chickenpox (Varicella): Transmssion. (2018).

December 31, 2018. https://www.cdc.gov/chickenpox/about/transmission.html.

17. W. H. Organization, Measles vaccines: WHO position paper, April 2017–

Recommendations. *Vaccine*, (2017).

18. J. Lessler, C. J. E. Metcalf, F. T. Cutts, B. T. Grenfell, Impact on Epidemic Measles of

Vaccination Campaigns Triggered by Disease Outbreaks or Serosurveys: A Modeling Study.

Plos Medicine **13**, (2016).

19. P. E. Christensen, H. Schmidt, H. O. Bang, V. Andersen, B. Jordal, O. Jensen, An epidemic

of measles in southern Greenland, 1951; measles in virgin soil. III. Measles and tuberculosis.

Acta Med Scand **144**, 450-454 (1953).

20. P. E. Christensen, H. Schmidt, H. O. Bang, V. Andersen, B. Jordal, O. Jensen, An epidemic

of measles in southern Greenland, 1951; measles in virgin soil. II. The epidemic proper.

Acta Med Scand **144**, 430-449 (1953).

21. S. Y. Chen, S. Anderson, P. K. Kutty, F. Lugo, M. McDonald, P. A. Rota, I. R. Ortega-

Sanchez, K. Komatsu, G. L. Armstrong, R. Sunenshine, J. F. Seward, Health Care-

Associated Measles Outbreak in the United States After an Importation: Challenges and

Economic Impact. *Journal of Infectious Diseases* **203**, 1517-1525 (2011).

22. G. H. Dayan, I. R. Ortega-Sanchez, C. W. LeBaron, M. P. Quinlisk, I. M. R. Team, The cost

of containing one case of measles: The economic impact on the public health infrastructure -

Iowa, 2004. *Pediatrics* **116**, E1-E4 (2005).

23. T. Pager, Measles Outbreak: Yeshiva's Preschool Program Is Closed by New York City

Health Officials. (2019). 4/15/2019.

https://www.nytimes.com/2019/04/15/nyregion/measles-nyc-yeshiva-closing.html.

24. A. Sanders, NYC health officials close two more Williamsburg yeshivas for failure to show

immunization records amid measles outbreak. (2019). 6/13/2019.

https://www.nydailynews.com/news/politics/ny-city-closes-williamsburg-brooklyn-

yeshivas-measles-outbreak-20190613-jogozcbe65ejtalweq255lwkka-story.html.

25. R. M. Anderson, R. M. May, *Infectious diseases of humans: dynamics and control.* (Oxford

University Press, Oxford, 1991).

26. M. J. Keeling, B. T. Grenfell, Disease extinction and community size: Modeling the

persistence of measles. *Science* **275**, 65-67 (1997).

27. P. Beck, S. M. Cohen, J. B. Ukeles, R. Miller, Jewish Community Study of New York:

2011, Geographic Profile. revised ed. *New York City: UJA-Federation of New York,* (2013).

28. W. Yang, J. Li, J. Shaman, Characteristics of measles epidemics in China (1951-2004) and

implications for elimination: A case study of three key locations. *Plos Comput Biol* **15**,

e1006806 (2019).

29. B. F. Finkenstädt, B. T. Grenfell, Time series modelling of childhood diseases: a dynamical

systems approach. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* **49**,

187-205 (2000).

30. W.-m. Liu, H. W. Hethcote, S. A. Levin, Dynamical behavior of epidemiological models

with nonlinear incidence rates. *Journal of mathematical biology* **25**, 359-380 (1987).

31. M. J. Keeling, P. Rohani, in *Modeling Infectious Diseases in Humans and Animals*.

(Princeton University Press, 2008), chap. 5, pp. 155.

32. New York City Department of Health and Mental Hygiene, Health Department Reports

Eleven New Cases of Measles in Brooklyn's Orthodox Jewish Community, Urges On Time

Vaccination for All Children, Especially Before Traveling to Israel and Other countries

Experiencing Measles Outbreaks. (2018).

https://www1.nyc.gov/site/doh/about/press/pr2018/pr091-18.page.

33. New York City Department of Health and Mental Hygiene, Measles Outbreak in Orthodox

Jewish Community of Brooklyn Continues to Grow — Health Department Urges Parents to

Vaccinate Their Children. (2019). 2/28/19.

https://www1.nyc.gov/site/doh/about/press/pr2019/measles-outbreak-now-at-121-

cases.page.

34. M. S. Arulampalam, S. Maskell, N. Gordon, T. Clapp, A tutorial on particle filters for online

nonlinear/non-Gaussian Bayesian tracking. *IEEE Trans. Signal Process.* **50**, 174-188 (2002).

35. M. D. McKay, R. J. Beckman, W. J. Conover, Comparison of three methods for selecting

values of input variables in the analysis of output from a computer code. *Technometrics* **21**,

239-245 (1979).

36. W. Yang, A. Karspeck, J. Shaman, Comparison of filtering methods for the modeling and

retrospective forecasting of influenza epidemics. *PLoS Comput Biol* **10**, e1003583 (2014).

Figure Captions:

Fig. 1. Monthly incidence for all ages and by age group. The solid line (y-axis on the right) shows monthly incidence for all ages, reported as of Aug 6, 2019. For comparison, bars (y-axis on the left) show monthly incidence for <1 (blue), 1-4 (orange), 5-17 (grey) and 18+ (yellow) year-olds, respectively, estimated based on health reports.

Fig. 2. Model fit. Box plots show estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F). Results are pooled over all 10 model-filter runs (each with 10,000 and in total 100,000 model realizations). Horizontal thick lines show the median of model estimates; box edges show the 25th and 75th percentiles; whiskers show the 2.5th and 97.5th percentiles; and dots show outliers. Stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Fig. 3. Estimated changes in population susceptibility. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) for <1 year-olds (A), 1-4 year-olds (B), 5-17 year-olds (C) and 18+ year-olds (D), respectively, at the end of each month from Sep 2018 to July 2019. The initial susceptibilities, estimated at the end of Sep 2018, were computed by adding the total numbers of individuals immunized by the vaccination campaigns in Oct 2018 to the posterior estimates at the end of Oct 2018. For comparison, the grey bars (y-axis on the right) show estimated numbers of individuals immunized during the vaccination campaigns; note that the vaccination campaigns targeted individuals under 19 years and thus is not shown for 18+ year-olds.

Fig. 4. Estimates of key model parameters: (A) the basic reproductive number, (B) the effective reproductive number, (C) infectious period, (D) relative contact rate among 1-4 year-olds, (E) relative contact rate among 5-17 year-olds, and (F) relative contact rate among 18-49 year-olds. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) made at the end of each month from Oct 2018 to July 2019. For comparison, the grey bars (y-axis on the left) show monthly incidence for all ages (A-C) or the related age groups (D-F).

Fig. 5. Estimated impact of vaccination campaigns. Box plots show simulated estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F), should there be no vaccination campaigns. Results are pooled over 10,000 model simulations. Horizontal thick lines show the median of model estimates; box edges show the 25th and 75th percentiles; whiskers show the 2.5th and 97.5th percentiles; and dots show outliers. For comparison, stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Table Captions:

Table 1. Estimated proportion of infections caused by each age-group. Rows show the receiving (i.e. infectee) age groups and columns show the source of infection (i.e. infector age group). The numbers are the mean (and 95% CI) estimates in percentage. For instance, for <1 year-olds (3rd row), on average 16.3% of cases were infected by the same age group, 44.6% by 1-4 year-olds, 20.9% by 5-17 year-olds, 15.2% by 18-49 year-olds, and 3% by 50+ year-olds.

Table 2. Estimated impact of vaccination campaigns during Oct 2018 – July 2019. Columns 2-4 show the total numbers of infections (2nd column), hospitalizations (3rd column), and individuals in intensive care unit (ICU) for different age groups (rows 3 to 6) and overall (last row), should there be no vaccination campaigns. Columns 5-7 show the number of infections, hospitalizations, and ICU cases averted by the vaccination campaigns, compared to data reported as of Aug 6, 2019. Numbers are the mean (and 95% confidence intervals) of 10,000 simulations.

TABLES

Table 1. Estimated proportion of infections caused by each age-group. Rows show the receiving (i.e. infectee) age groups and columns show the source of infection (i.e. infector age group). The numbers are the mean (and 95% CI) estimates in percentage. For instance, for <1 year-olds (3[rd] row), on average 16.3% of cases were infected by the same age group, 44.6% by 1-4 year-olds, 20.9% by 5-17 year-olds, 15.2% by 18-49 year-olds, and 3% by 50+ year-olds.

Infectee age groups	Infector age groups				
	<1 year	1-4 years	5-17 years	18-49 years	50+ years
<1 year	16.3 (12.3, 21.0)	44.6 (35.5, 53.5)	20.9 (13.9, 28.4)	15.2 (9.6, 21.6)	3.0 (1.4, 5.0)
1-4 years	1.8 (1.0, 3.0)	85.8 (77.0, 91.9)	7 (3.2, 12.9)	5 (2.2, 9.3)	0.3 (0.1, 0.6)
5-17 years	1.5 (0.9, 2.4)	12.9 (6.9, 21.3)	80.9 (70.4, 88.4)	4.4 (2, 8)	0.3 (0.1, 0.5)
18-49 yeas	2.4 (1.5, 3.7)	18.6 (10.9, 28.0)	9.1 (4.8, 14.9)	69.5 (56.2, 80.8)	0.4 (0.2, 0.8)
50+ years	15.5 (11.6, 19.9)	40.1 (31.1, 49.3)	19.7 (13, 27)	15.6 (9.7, 22.4)	9.1 (3.8, 16.3)

Table 2. Estimated impact of vaccination campaigns during Oct 2018 – July 2019. Columns 2-4 show the total numbers of infections (2nd column), hospitalizations (3rd column), and individuals in intensive care unit (ICU) for different age groups (rows 3 to 6) and overall (last row), should there be no vaccination campaigns. Columns 5-7 show the number of infections, hospitalizations, and ICU cases averted by the vaccination campaigns, compared to data reported as of Aug 6, 2019. Numbers are the mean (and 95% confidence intervals) of 10,000 simulations.

Age group	No. if without vaccination campaigns			No. averted by vaccination campaigns		
	Infection	Hospitalization	ICU	Infections	Hospitalization	ICU
<1	1015 (0, 1430)	75 (0, 106)	16 (0, 22)	937 (0, 1330)	70 (0, 99)	14 (0, 20)
4-5	3052 (3, 4096)	227 (0, 305)	47 (0, 63)	2837 (0, 3821)	211 (0, 284)	44 (0, 59)
5-17	1107 (1, 1692)	82 (0, 126)	17 (0, 26)	999 (0, 1554)	74 (0, 116)	15 (0, 24)
18+	904 (1, 1343)	67 (0, 100)	14 (0, 21)	803 (0, 1214)	60 (0, 90)	12 (0, 19)
All	6078 (5, 8443)	452 (0, 628)	94 (0, 130)	5576 (0, 7801)	415 (0, 580)	86 (0, 120)

FIGURES

Fig. 1. Monthly incidence for all ages and by age group. The solid line (y-axis on the right) shows monthly incidence for all ages, reported as of Aug 6, 2019. For comparison, bars (y-axis on the left) show monthly incidence for <1 (blue), 1-4 (orange), 5-17 (grey) and 18+ (yellow) year-olds, respectively, estimated based on health reports.

Fig. 2. Model fit. Box plots show estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F). Results are pooled over all 10 model-filter runs (each with 10,000 and in total 100,000 model realizations). Horizontal thick lines show the median of model estimates; box edges show the 25th and 75th percentiles; whiskers show the 2.5th and 97.5th percentiles; and dots show outliers. Stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Fig. 3. Estimated changes in population susceptibility. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) for <1 year-olds (A), 1-4 year-olds (B), 5-17 year-olds (C) and 18+ year-olds (D), respectively, at the end of each month from Sep 2018 to July 2019. The initial susceptibilities, estimated at the end of Sep 2018, were computed by adding the total numbers of individuals immunized by the vaccination campaigns in Oct 2018 to the posterior estimates at the end of Oct 2018. For comparison, the grey bars (y-axis on the right) show estimated numbers of individuals immunized during the vaccination campaigns; note that the vaccination campaigns targeted individuals under 19 years and thus is not shown for 18+ year-olds.

Fig. 4. Estimates of key model parameters: (A) the basic reproductive number, (B) the effective reproductive number, (C) infectious period, (D) relative contact rate among 1-4 year-olds, (E) relative contact rate among 5-17 year-olds, and (F) relative contact rate among 18-49 year-olds. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) made at the end of each month from Oct 2018 to July 2019. For comparison, the grey bars (y-axis on the left) show monthly incidence for all ages (A-C) or the related age groups (D-F).

Fig. 5. Estimated impact of vaccination campaigns. Box plots show simulated estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F), should there be no vaccination campaigns. Results are pooled over 10,000 model simulations. Horizontal thick lines show the median of model estimates; box edges show the 25th and 75th percentiles; whiskers show the 2.5th and 97.5th percentiles; and dots show outliers. For comparison, stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Supplementary Materials

Table S1. Main model parameters and prior ranges tested. In total, we tested 5040 combinations of prior ranges. Each combination was used as the lower and upper bounds of Latin Hypercube sampling. The optimal prior used in the final model-filter runs are bolded if multiple ranges were tested.

Parameter	Symbol/Equation	Ranges tested
Initial susceptibility in <1 year-olds	$S_1(t=0)$; Eqn 1	Based on susceptibility in 18-49 year-olds (i.e., the mothers)
Initial susceptibility in 1-4 year-olds	$S_2(t=0)$; Eqn 1	[5, 15], [10, 20], [15, 25], **[20, 30]**, [25, 35], [30, 40], [35, 45]%
Initial susceptibility in 5-17 year-olds	$S_3(t=0)$; Eqn 1	**[4, 8]**, [5, 10], [5, 15], [10, 20], [15, 25]%
Initial susceptibility in 18-49 year-olds	$S_4(t=0)$; Eqn 1	**[4, 8]**, [5, 10], [5, 15], [10, 20]%
Initial susceptibility in 50+ year-olds	$S_5(t=0)$; Eqn 1	[4, 8]%
Initial number of infants with maternal immunity	$M(t=0)$; Eqn 1	Based on susceptibility in 18-49 year-olds (i.e., the mothers)
Latent period	Z; Eqn 1	[7, 9] days
Infectious period	D; Eqn 1	[2, 6] days
Mixing parameter for the susceptibles	m_1; Eqn 1	**1** (perfect mixing), [0.95, 1], [0.9, 0.95]
Mixing parameter for the infectious	m_2; Eqn 1	**1** (perfect mixing), [0.95, 1], [0.9, 0.95]
Relative contact rate among <1 year-olds	β_1; Eqn 2	Set to 1
Relative contact rate among 1-4 year-olds	β_2; Eqn 2	[3, 30]
Relative contact rate among 5-17 year-olds	β_3; Eqn 2	[25, 50]
Relative contact rate among 18-49 year-olds	β_4; Eqn 2	[20, 40]
Relative contact rate among 50+ year-olds	β_5; Eqn 2	[1, 5]
Relative contact rate between 1-4 and 5-17 year-olds (sibling interactions)	β_6; Eqn 2	[1, 5]
Contact rate between 18-49 and 1-4 or 5-17 year-olds (parent-child interactions)	β_7; Eqn 2	[1, 5]
Amplitude of school term-time forcing	b_1; Eqn 3	[0.25, 0.75], **[0.5, 1]**
Basic reproductive number	R_0; Eqn 4	**[5, 10]**, [7, 12]
Reporting rate	r; Eqn 6	[80, 100]%

Table S2. Comparison of model estimates on initial susceptibility and model performance under different assumptions on vaccination campaigns. The baseline setting is as reported in the main text and alternative settings 1 to 3 are as described in the section "*Sensitivity analysis on vaccination campaigns settings.*" The results are summarized by pooling all 10 model-filter runs (10,000 particles each run and 100,000 model realizations in total). The numbers are the mean and, for the susceptibilities, 95% credible intervals in the parentheses. The initial susceptibilities, estimated at the end of Sep 2018, were computed by adding the total numbers of individuals immunized by the vaccination campaigns in Oct 2018 to the posterior estimates at the end of Oct 2018.

| | Age group | Model Settings on Vaccination Campaigns | | | |
		Baseline	Alternative 1	Alternative 2	Alternative 3
Estimated initial susceptibility (%) at end of Sep 2018	<1 year	53.2 (49, 57.5)	54.2 (50, 58.4)	54.2 (50, 58.5)	54.2 (50, 58.5)
	1-4 years	24.9 (20.4, 29.7)	24.9 (20.4, 29.7)	24.9 (20.4, 29.7)	29.9 (25.4, 34.7)
	5-17 years	6.0 (4.1, 7.9)	7.5 (5.1, 9.9)	7.5 (5.1, 9.9)	6.0 (4.1, 7.9)
	18-49 years	6.0 (4.1, 7.9)	7.5 (5.2, 9.9)	7.5 (5.2, 9.9)	7.5 (5.2, 9.9)
	50+ years	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)
Log-likelihood		-255.25	-265.52	-255.25	-257.42
Relative error of total number of cases over the outbreak	<1 year	0.39%	-11.96%	-11.56%	-17.91%
	1-4 years	-0.04%	-7.80%	-6.90%	-8.07%
	5-17 years	-4.13%	14.38%	0.33%	1.84%
	18+ years	-4.70%	7.24%	5.97%	7.95%
	All ages	-1.79%	-0.65%	-3.48%	-4.25%
Root-mean-square-error (RMSE), over Oct 2018 – July 2019	<1 year	1.74	1.86	1.80	2.55
	1-4 years	8.72	8.13	8.17	7.70
	5-17 years	3.91	5.27	3.00	5.10
	18+ years	5.08	4.62	4.30	4.25
	All ages	7.31	7.52	5.88	6.29
Correlation, over Oct 2018 – July 2019	<1 year	0.99	1.00	1.00	0.99
	1-4 years	0.95	0.96	0.96	0.97
	5-17 years	0.99	0.97	0.99	0.97
	18+ years	0.97	0.97	0.97	0.97
	All ages	0.99	0.99	1.00	1.00
1-step-head prediction RMSE, over Oct 2018 – Mar 2019	<1 year	4.37	4.46	4.40	4.60
	1-4 years	17.13	17.34	17.35	25.12
	5-17 years	8.27	8.39	8.37	8.50
	18+ years	4.34	4.93	4.93	5.06
	All ages	27.79	28.23	28.16	35.02

Fig. S1. Schematic of the measles transmission model. Measles transmission model follows the susceptible (S), exposed (E) and latently infected, infectious (I), and recovered (R) and/or immunized SEIR dynamics and includes 5 age-groups as indicated by the subscripts (i.e., <1, 1-4, 5-17, 18-49, and 50+ year-olds, respectively) and a group (M) for infants with maternal immunity. Black solid arrows show the disease-related processes; grey solid arrows show the demographic processes including birth (horizontal), aging (vertical), and death (tilted). Black dashed arrows show processes related to the routine 2-dose measles vaccination where susceptible individuals are vaccinated at ages 1 and 5 and move to the respective immune groups. Red dotted arrows show processes related to vaccination of susceptible individuals under 18 during the vaccination campaigns.

Fig. S2. Estimates of model parameters not listed in Fig. 4: (A) amplitude of school term-time forcing, (B) latent period, (C) reporting rate, (D) relative contact rate among 50+ year-olds, (E) relative contact rate between 1-4 and 5-17 year-olds (i.e. sibling interactions), and (F) relative contact rate between 18-49 and 1-4 or 5-17 year-olds (i.e. parent-child interactions). Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) made at the end of each month from Oct 2018 to July 2019. For comparison, the grey bars (y-axis on the left) show monthly incidence for all ages. Note that m_1 and m_2 are not shown as both optimal priors are the value 1 (Table S1).