Transmission Dynamics of and Insights from the 2018-2019 Measles Outbreak in

New York City: A Modeling Study

Wan Yang

Department of Epidemiology, Mailman School of Public Health, Columbia University, NY, NY

Correspondence to: wy2202@columbia.edu

Abstract

Background

In 2018-2019, New York City experienced the largest measles outbreak in the US in nearly three

decades. The objectives of this study are to identify key population and epidemiological

characteristics that may have contributed to this outbreak and gain insights to aid future public

health interventions.

Methods and Findings

We developed a model-inference system to infer the transmission dynamics of measles in the

affected community and assess the impact of vaccination campaigns implemented during the

outbreak, based on monthly age-grouped and overall incidence data reported from Oct 2018 to

July 2019. The model-inference system accounted for differential mixing among 5 age-groups

(<1, 1-4, 5-17, 18-49, and 50+ years), routine two-dose vaccination, vaccination campaigns, and

observational errors including underreporting. However, it did not account for other public

health interventions such as post-exposure prophylaxis and school closure.

Approximately 25% of 1-4 year-olds were susceptible when the outbreak began,

enabling measles to spread predominantly in this age group in the initial months. Early

vaccination campaigns were able to lower this susceptibility and contain measles spread by Dec

2018. However, an increase in infectious contact among 1-4 year-olds, likely facilitated by

parents hosting "measles parties" to expose unvaccinated children, appeared to rekindle and

aggravate the outbreak in early 2019. Consequently, 1-4 year-olds not only had the largest

number of infections (275/642 or 42.8%, as of Aug 6, 2019), but also appeared to cause the

largest number of infections in other age groups, including around half of the 100 infections in

infants.

The model estimated that, without the implemented vaccination campaigns, there

would be 6078 (95% CI: 5, 8433) infections and 452 (95% CI: 0, 628) hospitalizations, including

94 (95% CI: 0, 130) needing intensive care, during the observed outbreak period. In addition, it

would largely affect infants under 1 too young to receive the first dose of vaccine and young

children under 4, with over 10 times more infections and hospitalizations in both age groups

than reported.

Conclusions

These findings highlight the importance of vaccination in protecting children from measles and

public health education to reduce activities that unnecessarily expose children to the disease.

Measles is highly contagious and severe viral disease. Thanks to a highly effective vaccine and

high coverage of vaccination, endemic transmission of measles—i.e. continuous transmission

for more than 12 months—in the US was declared eliminated in 2000. However, due to vaccine

hesitancy and declining vaccination rate, in recent years there have been an increasing number

of large outbreaks following introduction of measles infection [1]. Due to long-term fluctuations

in vaccine coverage and infection history, population susceptibility could vary substantially by

age group. This susceptibility disparity by age can further interact with age-specific social

connectivity (i.e., contact rate) to shape the epidemic trajectory. As such, understanding these

detailed population characteristics as well as their impact on transmission dynamics in the

recent outbreaks is important for devising timely and effective intervention strategies.

In the fall of 2018, several New York City (NYC) residents acquired measles while

traveling abroad and subsequently led to the largest measles outbreak in the US in nearly three

decades. The first case of this outbreak developed a rash on Sep 30, 2018 and as of Aug 6, 2019

(at the time of this writing), there have been 642 confirmed cases, largely occurring in the

Orthodox Jewish community [2,3]. To contain the outbreak, the NYC Department of Health and

Mental Hygiene (DOHMH) launched extensive vaccination campaigns and, on Apr 9, 2019,

ordered mandatary vaccination of all individuals living, working or going to school in the

affected zip codes. As a result, over 32 thousand individuals under 19 years were vaccinated

with the measles, mumps, and rubella (MMR) vaccine during Oct 2018 – July 2019 and the

outbreak subsided [3,4].

In this study, we modeled the transmission dynamics of this measles outbreak in the

affected Orthodox Jewish community in NYC from Oct 1, 2018 to July 31, 2019, months with

more than one measles cases reported. Using an age-structured model-inference framework,

we were able to estimate key epidemiological features including the initial susceptibilities in

five different age groups (i.e., <1, 1-4, 5-17, 18-49, and 50+ years) and basic reproductive

number R_0, infer key factors contributing to the spread of measles, estimate the proportions of

infection attributable to each age group, as well as assess the impact of vaccination campaigns.

METHODS

The measles outbreak occurred predominantly among members of the Orthodox Jewish

community in Williamsburg and Borough Park, two neighborhoods in located in Brooklyn, NYC.

As such, we focused on modeling the outbreak in this subpopulation. Estimated based on the

Jewish Community Study of New York [5], approximately N=165,970 Orthodox Jews live in

these two NYC neighborhoods, of which 4552 (2.7%), 18,208 (11%), 59,176 (35.7%), 60,445

(36.4%), and 23,589 (14.2%) are <1, 1-4, 5-17, 18-49, and 50+ years, respectively.

Estimating monthly incidence by age group

Monthly measles incidence aggregated over all ages from Sep 2018 to Aug 2019 and the age

distribution of case-patients over the entire outbreak were published on the NYC DOHMH

website [3]. Of note, 1 case was reported in Sep 2018 (i.e., the initial case) and none were

reported in Aug 2019 at the time of this writing. In addition, the numbers of reported cases

were subsequently revised by the DOHMH (often adjusted upwards, presumably from

retrospective case identification) and, as such, varied over time. To estimate the monthly

incidence for each age group, we used the age distribution of cases reported in earlier health

reports/alerts [6] to apportion the total incidence for each month. For months without age

information, we used estimates either from the preceding month or the following month back-

calculated from the overall age distribution.

Transmission model

The transmission model used here was similar to described in our previous study [7]. As

illustrated in Figure 1, the model represents the susceptible-exposed-infectious-recovered

(SEIR) disease dynamics with five age groups (i.e. <1, 1-4, 5-17, 18-49, and 50+ years) to account

for population differences by age group (e.g., susceptibility and contact rate), routine two-dose

vaccination at ages 1 and 5, and immunization during the vaccination campaigns per Eqn 1:

100
$$
\begin{cases}
\dfrac{dS_i}{dt} = -\left(\dfrac{S_i^{m_1}}{N_i}\right)\sum_{j=1}^{5}\beta_{ij} I_j^{m_2} + k(1-v)N\mathbf{1}_{i=1} + \dfrac{M}{180}\mathbf{1}_{i=1} + l_{i-1}S_{i-1}(1-v_{i-1}) - l_i S_i - V_i(t) \\[2ex]
\dfrac{dE_i}{dt} = \left(\dfrac{S_i^{m_1}}{N_i}\right)\sum_{j=1}^{5}\beta_{ij} I_j^{m_2} - \dfrac{E_i}{Z} + \alpha_i(t) + l_{i-1}E_{i-1} - l_i E_i \\[2ex]
\dfrac{dI_i}{dt} = \dfrac{E_i}{Z} - \dfrac{I_i}{D} + l_{i-1}I_{i-1} - l_i I_i \\[2ex]
\dfrac{dR_i}{dt} = \dfrac{I_i}{D} + l_{i-1}R_{i-1} - l_i R_i + l_{i-1}S_{i-1}v_{i-1} + V_i(t) \\[2ex]
\dfrac{dM}{dt} = kvN - \dfrac{M}{180}
\end{cases}
$$

for *i*=1,..,5 (Eqn 1)

where S_i, E_i, I_i, R_i and N_i are, respectively, the numbers of susceptible, exposed (i.e. latently

infected), infectious, recovered (and/or immunized) people and population size in the *i*-th age

group; *M* is the number of infants with maternal immunity, which decays exponentially with a

mean duration of 180 days; t is time in days. $V_i(t)$ is the number of people in group-i immunized

by the vaccination campaigns on day-t (described in detail in the next section). The exponents

m_1 and m_2 describe the level of inhomogeneous mixing [8,9]; and $m_1=m_2=1$ represents

homogeneous mixing. Z and D are the latent and infectious period, respectively.

To model the different contact rates within and between age groups, we used 7

parameters for the Who-Acquires-Infection-From-Whom (WAIFW) matrix as follows:

112
$$\beta = \begin{bmatrix} \beta_1 & \beta_1 & \beta_1 & \beta_1 & \beta_1 \\ \beta_1 & \beta_2 & \beta_6 & \beta_7 & \beta_1 \\ \beta_1 & \beta_6 & \beta_3 & \beta_7 & \beta_1 \\ \beta_1 & \beta_7 & \beta_7 & \beta_4 & \beta_1 \\ \beta_1 & \beta_1 & \beta_1 & \beta_1 & \beta_5 \end{bmatrix} \text{ (Eqn 2)}$$

113   where β_1 to β_5 represent within-group contact for the five age groups and β_6 and β_7 represent

114   mixing between siblings and child-parent, respectively. For simplicity, we set all interactions

115   with group 1 (i.e. <1 year) or group 5 (50+ years) to β_1, the lowest contact rate. For group-3 (5-

116   17 years), to capture the varying contact rate following school schedules, we adjusted β_3 for

117   each date per the school calendar in NYC as:

118
$$\beta_3(t) = \frac{\beta_3}{b_{Term}} \cdot [1 + b_1 Term(t)] \quad \text{(Eqn 3)}$$

where b_1 is the amplitude of school term-time forcing; $Term(t)$ is set to 1 for school days and -1

for non-school days; and b_{Term} is the yearly average of $1+b_1 Term(t)$ [10].

The basic reproductive number R_0, defined as the average number of secondary

infections caused by a primary case-patient in a *naïve* population, reflects the transmissibility of

an infection. In an age-structured model, R_0 is computed as:

$$R_0 = \text{eigen}_{\max}(\boldsymbol{n\beta}D) \qquad \text{(Eqn 4)}$$

where $\text{eigen}_{\max}(\cdot)$ denotes the function giving the largest eigenvalue of a matrix, and $\boldsymbol{n}$ is a

diagonal matrix with elements $n_i=N_i/\sum N_i$ (i=1, …, 5 here), i.e. the fraction of population in group-

i. Based on this relationship between R_0 and the $\boldsymbol{\beta}$ matrix, we reparametrized the model to

include R_0 as a model parameter by setting β_1 to 1 and estimating the relative magnitude of β_2–

β_6, all scaled to R_0. In the current mass vaccination era, most people are immune via

vaccination. To reflect the potential of an infection to cause an epidemic in a partially

susceptible population, the effective reproductive number, R_e, accounts for population

susceptibility and is computed as:

$$R_e = \text{eigen}_{\max}(\boldsymbol{s\beta}D) \qquad \text{(Eqn 5)}$$

where $\boldsymbol{s}$ is a diagonal matrix with elements $s_i=S_i/N_i$ (i=1, …, 5 here), i.e. the susceptibility in

group-i.

To model the demographic processes, k is the birth rate (2.7 per 1000 person-year here

[5]); v is the immunity level in mothers, approximated by the susceptibility of the child-bearing

age-group (i.e., 18-49 year-olds); N is total population size; and $\mathbf{1}_{i=1}$ is an indicator function, with

value 1 for group-1 (<1 year-olds) and 0 for all other groups. Thus, the term $k(1-v)\,\mathbf{1}_{i=1}$ (1st line in

Eqn 1) is the number of susceptible newborns and kvN (last line in Eqn 1) is the number of

newborns with maternal immunity. The term l_i is the rate of aging for group-i (i.e. the inverse

of the sojourn time in each age group) with l_0 set to 0 and l_5 set to the death rate. The term v_i

(for i=1 and 2) is the vaccination rate for the two doses of vaccine (i.e., v_i=0 for i=0, 3, and 4). In

this study, we set v_1 (1st dose) to 0.9 times the immunity level of group-2 (i.e., assuming a 90%

vaccine efficacy) and v_2 (2^{nd} dose) to 0.7 for days before May 2019; for days afterwards, we

used 0.8 for v_1 and 0.9 for v_2, corresponding to a 1-(1-0.8)(1-0.9)=98% overall vaccination rate.

Based on NYC health reports/alerts [2,11,12], we seeded the model, via the parameter

$\alpha_i(t)$, with 3 cases in group-2 (i.e. 1-4 year-olds)—one each with rash onset on Sep 30, Oct 15,

and Oct 30, 2018, respectively—and 1 case each in group-3 (5-17 year-olds) and group-4 (18-49

year-olds), both during the winter recess (from 12/24/18 to 1/1/19).

Modeling the vaccination campaigns

To contain the outbreak, the NYC DOHMH conducted extensive vaccination campaigns and

administered 31,790 doses of MMR vaccine to children under 19 years in Williamsburg and

Borough Park by July 2019 [3]. However, information on the age and immune status of

vaccinees was not reported. In this study, per the health reports/alters [3,12], we assumed

there were two phases of vaccination campaign: 1) Oct 2018 – Feb 2019, during which 7000

children, 90% Orthodox Jewish (15% were <1 year, 65% 1-4 years, and 20% 5-17 years) were

vaccinated; and 2) Mar – July 2019, during which 24,790 children, 60% Orthodox Jewish (10%,

40%, and 50%, respectively, were <1, 1-4, and 5-17 years) were vaccinated. For reference,

Orthodox Jews made up for approximately 30% of the total population in the two affected

neighborhoods.

For <1 year-olds, immunization could fail due to residual maternal immunity; thus, we

assumed a 85% immunization success rate for group-1. For those above 1 year, some vaccinees

might have received 1 or 2 doses of vaccine and higher the population susceptibility, the less

likely a vaccinee had been immune before the additional vaccine dose. As such, we assumed

the immunization success rate was twice the group-specific susceptibility for 1-4 year-olds and

three times that for 5-17 year-olds or at a minimum of 25% and a maximum of 75%. We further

assumed a 10-day delay in vaccine effect and computed the daily number of individuals

vaccinated per a gamma distribution (mean=30 days and standard deviation=21 days for Phase-

1 and mean=56 and standard deviation=15 days for Phase-2 such that it peaked ~1 week after

Apr 9, 2019 when the city implemented a vaccination mandate). The estimated numbers

matched with the reports (e.g., 1740 doses by our model vs ~1600 doses given to children

under 5 as reported [13] and 1142 doses by our model vs ~1000 doses given in March 2019 as

reported [14,15]). These daily numbers were then included in the transmission model [i.e. $V_i(t)$

in Eqn 1]. Sensitivity to model assumptions were tested as described below.

Estimation of model state variables and parameters

To estimate the model state variables (i.e., S_i, E_i, I_i, R_i, and M) and parameters (β_2 to β_7, R_0, b_1, D,

Z, m_1, and m_2), we fitted the model to the reported monthly overall incidence and the

estimated monthly age-grouped incidence using a particle filter [16]. Briefly, we first initialized

a suite of model realizations (termed "particles", N=10,000 here) using Latin Hypercube

sampling [17] from the prior distribution of state variables and parameters (Table 1). The

particle filter then sequentially incorporated the monthly incidence to the model via repeated

prediction-update cycles. In each cycle (i.e., each month here), the particles were stochastically

integrated forward in time for a month per the model (i.e., Eqn 1; this generates the

prediction). To update the model state including all model variables and parameters, at the end

of each month, the model-estimated incidence was aggregated for the month, adjusted by the

reporting rate for that month (estimated simultaneously by the filter), and used to compute the

likelihood of each particle (described below). The posterior of model state was then computed

using Bayes' rule [16,18] and the particles resampled and updated—those with high posterior

probabilities were retained and those with very low posterior probabilities discarded.

To allow for a wider observational variance than, e.g., the Poisson process, we

heuristically modeled the observations using a multivariate Gaussian distribution (i.e. the

likelihood function):

$$\boldsymbol{Y_m}|r, \boldsymbol{C_m} \sim \mathcal{N}(r\boldsymbol{C_m}, \boldsymbol{\Sigma}) \quad \text{(Eqn 6)}$$

where $\boldsymbol{Y_m}$ is the vector of monthly incidence reported for month-m, including the monthly

incidence for individual age-groups (i.e., <1, 1-4, 5-17 and 18+ years; note that 18-49 and 50+

year-olds were combined due to a lack of data for these two groups separately) and all ages

combined. Correspondingly, $\boldsymbol{C_m}$ is the vector of monthly incidence estimated by the model; and

r is the reporting rate and, for simplicity, assumed the same for all ages. $\boldsymbol{\Sigma}$ is the covariance

matrix, with the off-diagonal terms set to 0. To account for uncertainties in the estimated age-

grouped incidence, the variance ($\Sigma_{ii}, i = 1, \dots, 4$) for each of the 4 aforementioned age-groups

was heuristically computed as:

$$\Sigma_{ii,m} = 100 + \frac{\left(\sum_{m}^{m-2} Y_m/3\right)^2}{3} \quad \text{(Eqn 7a)}$$

That is, the observational variance is proportional to the average incidence in the preceding

two months (if available) and the current month, plus a baseline constant. For the overall

incidence with detailed data, a smaller variance was used:

$$\Sigma_{ii,m} = 100 + \frac{(\sum_{m}^{m-2} Y_m/3)^2}{5} \text{ (Eqn 7b)}$$

As there were great uncertainties in the susceptibilities of the younger age groups, we

tested prior ranges from 10-40% for 1-4 year-olds, 5-20% for 5-17 year-olds, and 5-15% for 18-

49 year-olds. For the basic reproductive number R_0, we tested prior values ranging from 5 to 12

(note these values were lower than the oft-reported 12-18 range [19,20]). To optimize the

model-filter system, we parsed these wide ranges into smaller segments and tested all

combinations by permutation (5040 in total; see specific prior ranges in Table 1). To account for

model stochasticity, we ran the model-filter system 5 times for each prior combination and 10

times for the final prior select. We then selected the optimal priors based on the model-

goodness-of-fit to the data (minimal root-mean-square-error and maximal correlation and

likelihood) over the period of Oct 2018 – July 2019 as well as accuracy of the one-step-ahead

predictions (recall that the particle filtering process comprises sequential prediction-update

cycles) for the period of Oct 2018 – March 2019 (i.e. before the emergency vaccination

mandate). We pooled all 10 final runs (10,000 particles each run and 100,000 model

realizations in total) to compute the posterior estimates [e.g., mean and 95% credible intervals

(CI)].

Sensitivity analysis on vaccination campaigns settings

To test the sensitivity of model results to assumptions on vaccination campaign settings, we

tested the model-filter system using the following alternative scenarios:

1) For the 2nd phase (Mar-July 2019), 90% (vs. 60% in the baseline scenario) of the

vaccine doses were given to members of the Orthodox Jewish community;

2) For the 2nd phase, the age distribution of vaccinees was the same as the 1st phase (i.e.

15%, 65%, and 20%, respectively, for <1, 1-4, and 5-17 year-olds vs. 10%, 40%, and 50% for the

three groups in the baseline scenario);

3) For the 2nd phase, 90% of the vaccine doses were given to Orthodox Jewish and the

age distribution of vaccinees was the same as the 1st phase (i.e. 15%, 65%, and 20% for <1, 1-4,

and 5-17 year-olds, respectively).

Because population susceptibility would be affected by the number of individuals

immunized by the vaccination campaigns, we tested susceptibility ranges 10-40% for 1-4 year-

olds, 5-20% for 5-17 year-olds, and 5-15% for 18-49 year-olds, divided in to small segments as

for the baseline scenario (140 different combinations for each alternative scenario; Table 1).

For simplicity, we used the same optimal prior ranges for the parameters under the baseline

scenario in this sensitivity analysis.

Evaluating the impact of vaccination campaigns

To estimate the impact of vaccination campaigns, we generated model-simulated

counterfactuals—i.e., outbreak outcomes should there be no vaccination campaigns

implemented—using the posterior mean estimates of group-specific initial population

susceptibilities and model parameters for each month. We ran the model (10,000 realizations)

stochastically up to the end of 2019 to test how long the outbreak could last without

intervention; for months after Aug 2019, parameters estimated at the end of July 2019 were

used.

RESTULTS

Overview of the measles outbreak and model fit

The measles outbreak started on Sep 30, 2018, when a young child developed a rash. It evolved

relatively slowly in the first three months; however, the outbreak took off quickly in early 2019,

peaked in April after the city declaring a public health emergence, and recorded a total of 642

cases by July 31, 2019. As shown in Figure 2, age-grouped incidence, estimated based on health

reports/alerts [3,6], peaked first in Mar 2019 among 1-4 year-olds—the age group with the

largest number of infection (275 cases or 42.8% of the total, as of Aug 6, 2018), followed by <1

year-olds (100 cases or 15.6%) and 5-17 year-olds (138 cases or 21.5%) in Apr 2019, and 18+

year-olds (129 cases or 20.1%) in May 2019.

The model-filter system was able to recreate the overall incidence curve during Oct

2018 – July 2019, estimate the overall age distribution of measles cases, as well as recreate the

estimated age-grouped incidence curves for all age groups (Figure 3 and Table S1). Note that

while our model (Eqn 1) divided 18+ year-olds into two subgroups (i.e. 18-49 and 50+ years)

given their differences in contact rates and interactions with other age-groups, we present the

combined results here because data were only available for the entire age group. The

estimated reporting rate was around 90% throughout the study period and slightly lower in Apr

2019, at the peak of the outbreak (89.1%, 95% CI: 79.4, 99.2%; Figure S1).

Inference of key epidemiological characteristics

The model-filter system estimated that, at the beginning of the outbreak (i.e. Sep 2018),

susceptibility was the highest in infants, at approximately 53.2% (95% CI: 49.0, 57.5%; Figure

4A). This was expected, because maternal immunity wanes within 3 to 9 months after birth

and, as a result, by their first birthday—the age eligible to receive the first dose of MMR vaccine

in the US—almost all infants have lost their maternal immunity and are susceptible to measles.

Young children aged 1-4 years had the second highest susceptibility; approximately 24.9% (95%

CI: 20.4, 29.7%) were susceptible. In comparison, susceptibility was lower among both 5-17

year-olds (6.0%, 95% CI: 4.1, 7.9%) and 18+ year-olds 6.0% (95% CI: 4.4, 7.6%; Figure 4). These

estimates were consistent with the observation that all cases recorded in Oct 2018 were

children ranging from 11 months to 4 years [2]. Sensitivity analysis on assumptions related to

the vaccination campaigns showed that estimated susceptibilities were slightly higher for 5-18

year-olds (7.5%, 95% CI: 5.1, 9.9%) and 18-49 year-olds (7.9%, 95% CI: 5.2, 9.9%) if more vaccine

doses were given to the Jewish Orthodox community or 5-18 year-olds; however, the estimates

were in general consistent with the baseline scenario (Table S1).

The initial vaccination campaigns launched promptly afterwards lowered the

susceptibility to 40.3% (95% CI: 36.4, 44.0%) in infants and 13.9% (95%: 9.7, 18.7%) in 1-4 year-

olds, by the end of Dec 2018 (Figure 4). These efforts along with other transmission and

infection controls [2] appeared to effectively contain the outbreak at the time. The effective

reproductive number (R_e) is a key epidemiological parameter reflecting the potential of an

infection to cause an epidemic in a partially immune population; an epidemic is possible when

R_e>1. The estimated R_e was 1.5 (95% CI: 0.7, 2.9) in Oct 2018 and dropped to round 1 (95% CI:

0.7, 1.5) in Dec 2018 (Figure 5B).

The outbreak, however, took off again in early 2019 (Figure 2). The estimated R_e

increased and remained above 1 in the first three months of 2019 (Figure 5B). In a perfectly

mixed model, R_e is computed as the product of the basic reproductive number R_0 and

population susceptibility. In particular, the basic reproductive number R_0 measures the

transmissibility of an infection in a fully susceptible population; for measles, while often

reported in the range of 12-18, R_0 could vary from 1.4 to 770 [19]. In this study, we estimated

that R_0 was approximately 7 during the entire outbreak (Figure 5A). Population susceptibility,

the other factor for R_e, were to increase in a close population (i.e., without migration), could

only do so slowly as infants lose maternal immunity. Thus, these two factors alone could not

explain the sudden large increase in R_e (Figure 5B). In this study, we utilized an age-structured

model that enables more detailed analysis of the transmission dynamics. Indeed, our model-

filter system detected an increase in contact rate among 1-4 year-olds during Jan-March 2019

(Figure 5D); this increased contact rate along with the high susceptibility among 1-4 year-olds

appeared to raise R_e above unity (Eqn 5) and contribute to the re-surge of measles in early

2019. This finding was also consistent with reports of parents hosting "measles parties" to

exposure unvaccinated children at the time [14]. In contrast, estimated contact rates were

relatively stable for other age groups (e.g., Figure 5 E and F for 5-17 and 18-49 year-olds,

respectively). As a result, infections increased quickly among 1-4 year-olds (Figures 2 and 5D),

reaching a peak of around 80 cases in March. Meanwhile, infections also increased in other age

groups including 5-17 and 18+ year-olds despite their low overall susceptibilities, due to

interactions between age-groups (Figure S1) and the high contact rates in these groups (Figure

5 E and F).

The outbreak began to decline in Apr 2019, following more stringent public health

interventions [3,4]. In particular, the model estimated that, thanks to extensive vaccination

campaigns, susceptibility was reduced to 22.8% (95% CI: 19.3, 26.0%) in <1 year-olds, 4.3%

(95% CI: 0.5, 8.9%) in 1-4 year-olds, and 2.4% (95% CI: 0.4, 4.4%) in 5-17 year-olds at the end of

May. Consequently, the effective reproductive number R_e dropped below 1 from April onwards.

Who acquired infection from whom?

Table 2 shows the estimated proportions of infections caused by each of the five age groups

based on the estimated WAIFW contact matrix (Eqn 2). Children aged 1-4 years not only had

the largest number of infections (42.8%), but also appeared to cause the largest number of

infections in other age groups. Tallied over the entire study period (Oct 2018-July 2019), an

estimated 51.6% (95% CI: 39.3, 63.1%) of the total cases were infected by 1-4 year-olds,

compared to 25.2% (95% CI: 15.8, 35.5%) by 5-17 year-olds, 17.7% (95% CI: 10.4, 26.5%) by 18-

49 year-olds, 4.5% (95% CI: 3.0, 6.4%) by <1 year-olds, and 1.0% (95% CI: 0.4, 1.8%) by 50+ year-

olds. In particular, this age group caused around half of the infections in infants (44.6% or 45

cases) as well as the largest proportions of inter-group transmission to other age-groups

(ranging from 12.9% to 5-17 year-olds to 40.1% to 50+ year-olds; Table 2).

Estimated impact of vaccination campaigns

Figure 6 shows the estimated outbreak outcomes had there be no vaccination campaigns.

Without the vaccination campaigns, the model estimated that the outbreak could continue to

the end of 2019 and infect a total of 6196 (95% CI: 5, 8478) people by then or 6078 (95% CI: 5,

8433) during the observed outbreak period (i.e. Oct 2018 – July 2019), compared to 642 cases

reported as of Aug 6, 2019. In addition, these infections would largely occur in infants under 1

and young children aged 1-4 years. During the observed outbreak period, there would be 1015

(95%: 0, 1430) infections in infants and 3052 (95% CI: 2, 4096) infections in 1-4 year-olds, more

than 10 times of the reported numbers in these two age groups (i.e. 100 and 275, respectively,

as of Aug 6, 2019; Table 3). Children aged 5-17 years would have the third largest number of

infections, with 1107 (95% CI: 1, 1692) cases, 8 times of the reported number (i.e. 138 as of Aug

6, 2019).

Measles can cause severe diseases. According to data by Apr 24, 2019, among 390

individuals with measles, 29 were hospitalized, of which six needed intensive care [21].

Assuming the same ratios among all cases, without the vaccination campaigns, during the

observed outbreak period there would be 452 (95% CI: 0, 628) hospitalizations, including 94

(95% CI: 0, 130) needing intensive care, and the majority would be in young children under 4

(Table 3).

DISUCSSION

Using a model-filter inference system, we have reconstructed in detail the transmission

dynamics of the measles outbreak in an Orthodox Jewish community in NYC during Oct 2018 –

July 2019. We have estimated the population characteristics (e.g. age-specific susceptibilities)

and epidemiological parameters (e.g. reproductive numbers) as well as subtle changes in key

parameters (e.g. contact rates) that are critical to the transmission of measles. Using model

simulation and the posterior estimates from the model-inference system, we are also able to

estimate the impact of vaccination campaigns implemented during the outbreak, including

numbers of infections and hospitalizations averted, for each age group. These latter findings

echo those from previous studies [22-24] and again highlight the severity of measles disease,

should there be no effective infection and transmission controls (in particular, vaccination).

Our analyses estimate that around a quarter of young children aged 1-4 in the affected

community were susceptible at the onset of the outbreak, likely due to delayed vaccination.

Indeed, 94% (101/108) of the early infections in children were unvaccinated [12]. In contrast,

vaccination rate remained high in older children 5-17 years, with an estimated 94% immune to

measles. This difference may be due to better compliance with vaccination regulation at school

entry, or a result of vaccination campaigns in response to previous outbreaks (e.g., a large

outbreak occurred in the same community in 2013 [25]). Nevertheless, the large number of

unvaccinated children under 4 was sufficient to cause many infections in late 2018,

predominantly in the same age group (Figure 2 and references [2,11,12]). This observation

highlights the importance of vaccination compliance with both MMR vaccine doses, especially

given the long lag between the two vaccine doses.

Our study also reveals intricate interplays of population dynamics and measles

transmission. While the high susceptibility in children under 4 was likely responsible for the

early spread of measles, our estimates suggest that the second and more severe part of the

outbreak in 2019 was likely due to increased infectious contact among this age group, likely

facilitated by parents hosting "measles parties" that intentionally bring unvaccinated children

together and expose them to those sick with measles [14]. As shown in Figure 5, the increase in

infectious contact interacting with the high susceptibility in 1-4 year-olds was able to raise the

effective reproductive number R_e to above unity—the threshold for an epidemic to occur—and

aggravate the outbreak in 2019, despite earlier public health efforts that had reduced R_e to

below 1 in late 2018. Similar disease-related gatherings have been noted in previous measles

outbreaks [26] as well as other disease outbreaks [27]. These activities create further

challenges for the control of measles spread and stress the need for enhanced public health

education.

In addition, the intensified measles outbreak not only affected children with delayed

vaccination, but also a large number of infants under 1, who were too young to receive their

first dose of MMR vaccine in the US. At least 100 infants under 1 were infected with measles

during the 10-month outbreak period, despite extensive infection and transmission control

efforts including vaccinating infants 6 months or older and post-exposure prophylaxis with

immune globulin given to those under 6 months [2,11]. This was largely a result of the high

susceptibility in infants. Our model-inference system estimates that about half of infants were

susceptible by age 1 and that nearly half of the 100 infant cases were infected by 1-4 year-olds

(Table 2). In addition, our simulations suggest that, without the vaccination campaigns, the

number of infections in infants would be over 10 times higher than observed (Table 3). These

findings demonstrate the rippling effects of vaccine hesitancy beyond the risk posted to age-

eligible children with delayed vaccination. These findings also suggest that administration of

the first dose of routine MMR vaccine earlier than the current 1 year age-limit in the US may be

necessary to protect infants should the high level of vaccine hesitancy persist.

Our model simulations, consistent with many previous studies [24,26,28], demonstrate

the significant public health impact of vaccination in controlling measles outbreaks. Without the

implemented vaccination campaigns, the severity of the measles outbreak—including number

of infections, hospitalizations, and severe infections needing intensive care—would be about 10

times worse than observed (Table 3). These estimates, however, did not include the long-term

health impacts on affected individuals, particularly young children [23,29], nor the enormous

economic burdens [25,30,31].

We note several limitations of our study. First, we did not explicitly model the impact

of public health interventions other than the vaccination campaigns, due to a lack of data.

During the outbreak, such efforts included prescreening patients prior to presence for

treatment, post-exposure prophylaxis, and closing schools out of vaccination compliance

[2,11,12,32,33]. Of note, however, here the estimated basic reproductive number R_0 was

around 7, lower than the 12-18 range based on epidemics in the pre-vaccine era [20]; and the

estimated infectious period was around 4 days (Figure 5C), at the lower end of the commonly

used range of 4-6 days [34]. The lower R_0 and shorter infectious period could be a result of the

aforementioned public health interventions. Second, there were uncertainties in the accuracy

of case reporting. Because the incidence data used here were published on Aug 6, 2016, a later

revision of case reports, we used a relatively high but broad prior range for the reporting rate

(i.e. 80-100%). In addition, our model-inference system explicitly accounted for observational

errors (Eqn 7a and 7b). Third, due to a lack of contact data and for simplicity, we set all terms

related to group-1 (i.e. <1 year-olds) in the WAIFW matrix to the same as the contact rate

within the group (Eqn 2). The model formulation may have led to under-estimation of the

proportions of infection in <1 year-olds attributable to the same age group and/or 1-4 year-

olds, given the likely more frequent contact within the same age group and with similar ages

(i.e. 1-4 years) due to more shared settings such as daycares and pediatric hospitals. Lastly,

there were uncertainties regarding the settings of vaccination campaigns. Nevertheless,

sensitivity analysis showed that our main estimates were robust to a wide range of assumptions

(Table S1).

In summary, using a comprehensive model-inference system, we have reconstructed

transmission dynamics of the recent measles outbreak in NYC in great detail. Our estimates

highlight the importance of vaccination in protecting children as well as public health education

to reduce activities that unnecessarily expose children to risk of measles infection.

Acknowledgements

I thank all public health personnel and individuals involved in combating the measles outbreak

and New York City Department of Health and Mental Hygiene for making the incidence data

publicly available. I also thank Columbia University Mailman School of Public Health for access

to high performance computing.

Finding source: This study was supported by the National Institute of Allergy and Infectious

Diseases (1R01AI145883-01).

Conflict of interest: None.

References:
1. Phadke VK, Bednarczyk RA, Salmon DA, Omer SB (2016) Association between vaccine refusal
and vaccine-preventable diseases in the United States a review of measles and
pertussis. Jama-J Am Med Assoc 315: 1149-1158.
2. New York City Department of Health and Mental Hygiene. (2018) Alert # 39: Update on
measles outbreak in New York City in the Orthodox Jewish community.
https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2018/alert39-measles-
outbreak.pdf
3. New York City Department of Health and Mental Hygiene. (2019) Measles.
https://www1.nyc.gov/site/doh/health/health-topics/measles.page
4. New York City Department of Health and Mental Hygiene. (2019) Order of the commissioner.
https://www1.nyc.gov/assets/doh/downloads/pdf/press/2019/emergency-orders-
measles.pdf
5. Beck P, Cohen SM, Ukeles JB, Miller R (2013) Jewish community study of New York: 2011,
geographic profile. Revised ed. New York City: UJA-Federation of New York.
6. New York City Department of Health and Mental Hygiene. (2019) Health alert network.
https://www1.nyc.gov/site/doh/providers/resources/health-alert-network.page
7. Yang W, Li J, Shaman J (2019) Characteristics of measles epidemics in China (1951-2004) and
implications for elimination: A case study of three key locations. Plos Comput Biol 15:
e1006806.
8. Finkenstädt BF, Grenfell BT (2000) Time series modelling of childhood diseases: A dynamical
systems approach. Journal of the Royal Statistical Society: Series C (Applied Statistics)
49: 187-205.

9. Liu W-M, Hethcote HW, Levin SA (1987) Dynamical behavior of epidemiological models with
nonlinear incidence rates. Journal of mathematical biology 25: 359-380.
10. Keeling MJ, Rohani P (2008) Temporally forced models. Modeling infectious diseases in
humans and animals. 1st ed: Princeton University Press. pp. 155.
11. New York City Department of Health and Mental Hygiene. (2018) Alert # 38: Measles
outbreak in New York City in the Orthodox Jewish community.
https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2018/alert38-measles-
outbreak.pdf
12. New York City Department of Health and Mental Hygiene. (2019) Alert # 2: Update on
measles outbreak in New York City in the Orthodox Jewish community.
https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2019/update-on-measles-
outbreak-in-nyc-in-the-orthodox-jewish-community.pdf
13. New York City Department of Health and Mental Hygiene. (2018) Health department
reports eleven new cases of measles in brooklyn's Orthodox Jewish community, urges
on time vaccination for all children, especially before traveling to israel and other
countries experiencing measles outbreaks.
https://www1.nyc.gov/site/doh/about/press/pr2018/pr091-18.page
14. Scutti S. (2019) New York City declares a public health emergency amid Brooklyn measles
outbreak. https://www.cnn.com/2019/04/09/health/measles-new-york-emergency-
bn/index.html
15. New York City Department of Health and Mental Hygiene. (2019) Measles outbreak in
Orthodox Jewish community of Brooklyn continues to grow — health department urges
parents to vaccinate their children.
https://www1.nyc.gov/site/doh/about/press/pr2019/measles-outbreak-now-at-121-
cases.page
16. Arulampalam MS, Maskell S, Gordon N, Clapp T (2002) A tutorial on particle filters for online
nonlinear/non-Gaussian Bayesian tracking. IEEE Trans Signal Process 50: 174-188.
17. Mckay MD, Beckman RJ, Conover WJ (1979) Comparison of three methods for selecting
values of input variables in the analysis of output from a computer code. Technometrics
21: 239-245.
18. Yang W, Karspeck A, Shaman J (2014) Comparison of filtering methods for the modeling and
retrospective forecasting of influenza epidemics. PLoS Comput Biol 10: e1003583.
19. Guerra FM, Bolotin S, Lim G, Heffernan J, Deeks SL, et al. (2017) The basic reproduction
number (R0) of measles: A systematic review. Lancet Infect Dis 17: e420-e428.
20. Anderson RM, May RM (1991) Infectious diseases of humans: Dynamics and control.
Oxford: Oxford University Press.
21. New York City Department of Health and Mental Hygiene. (2019) Alert # 9: Citywide
recommendations during the ongoing measles outbreak in New York City.
https://www1.nyc.gov/assets/doh/downloads/pdf/han/alert/2019/recommendations-
during-measles-outbreak.pdf
22. Mina MJ, Grenfell BT, Metcalf CJE (2019) Response to comment on "long-term measles-
induced immunomodulation increases overall childhood infectious disease mortality".
Science 365.

23. Mina MJ, Metcalf CJE, De Swart RL, Osterhaus ADME, Grenfell BT (2015) Long-term
measles-induced immunomodulation increases overall childhood infectious disease
mortality. Science 348: 694-699.
24. Gastanaduy PA, Funk S, Paul P, Tatham L, Fisher N, et al. (2018) Impact of public health
responses during a measles outbreak in an Amish community in Ohio: Modeling the
dynamics of transmission. Am J Epidemiol 187: 2002-2010.
25. Rosen JB, Arciuolo RJ, Khawja AM, Fu J, Giancotti FR, et al. (2018) Public health
consequences of a 2013 measles outbreak in New York City. JAMA Pediatr 172: 811-817.
26. Gastanaduy PA, Banerjee E, Debolt C, Bravo-Alcantara P, Samad SA, et al. (2018) Public
health responses during measles outbreaks in elimination settings: Strategies and
challenges. Human vaccines & immunotherapeutics 14: 2222-2238.
27. Centers for Disease Control and Prevention. (2018) Chickenpox (varicella): Transmssion.
https://www.cdc.gov/chickenpox/about/transmission.html
28. Lessler J, Metcalf CJE, Cutts FT, Grenfell BT (2016) Impact on epidemic measles of
vaccination campaigns triggered by disease outbreaks or serosurveys: A modeling study.
Plos Medicine 13.
29. Christensen PE, Schmidt H (1953) An epidemic of measles in southern greenland 1951;
measles in virgin soil. Iv. The significance of specific prophylaxis. Acta Med Scand 145:
126-142.
30. Chen SY, Anderson S, Kutty PK, Lugo F, Mcdonald M, et al. (2011) Health care-associated
measles outbreak in the United States after an importation: Challenges and economic
impact. Journal of Infectious Diseases 203: 1517-1525.
31. Dayan GH, Ortega-Sanchez IR, Lebaron CW, Quinlisk MP, Team IMR (2005) The cost of
containing one case of measles: The economic impact on the public health
infrastructure - iowa, 2004. Pediatrics 116: E1-E4.
32. Pager T. (2019) Measles outbreak: Yeshiva's preschool program is closed by New York City
health officials. https://www.nytimes.com/2019/04/15/nyregion/measles-nyc-yeshiva-
closing.html
33. Sanders A. (2019) NYC health officials close two more Williamsburg yeshivas for failure to
show immunization records amid measles outbreak.
https://www.nydailynews.com/news/politics/ny-city-closes-williamsburg-brooklyn-
yeshivas-measles-outbreak-20190613-jogozcbe65ejtalweq255lwkka-story.html
34. Keeling MJ, Grenfell BT (1997) Disease extinction and community size: Modeling the
persistence of measles. Science 275: 65-67.
564

TABLES

Table 1. Main model parameters and prior ranges tested. In total, we tested 5040 combinations of prior ranges. Each combination was used as the lower and upper bounds of Latin Hypercube sampling. The optimal prior used in the final model-filter runs are bolded if multiple ranges were tested.

Parameter	Symbol/Equation	Ranges tested
Initial susceptibility in <1 year-olds	$S_1(t=0)$; Eqn 1	Based on susceptibility in 18-49 year-olds (i.e., the mothers)
Initial susceptibility in 1-4 year-olds	$S_2(t=0)$; Eqn 1	[5, 15], [10, 20], [15, 25], **[20, 30]**, [25, 35], [30, 40], [35, 45]%
Initial susceptibility in 5-17 year-olds	$S_3(t=0)$; Eqn 1	**[4, 8]**, [5, 10], [5, 15], [10, 20], [15, 25]%
Initial susceptibility in 18-49 year-olds	$S_4(t=0)$; Eqn 1	**[4, 8]**, [5, 10], [5, 15], [10, 20]%
Initial susceptibility in 50+ year-olds	$S_5(t=0)$; Eqn 1	[4, 8]%
Initial number of infants with maternal immunity	$M(t=0)$; Eqn 1	Based on susceptibility in 18-49 year-olds (i.e., the mothers)
Latent period	Z; Eqn 1	[7, 9] days
Infectious period	D; Eqn 1	[2, 6] days
Mixing parameter for the susceptibles	m_1; Eqn 1	**1** (perfect mixing), [0.95, 1], [0.9, 0.95]
Mixing parameter for the infectious	m_2; Eqn 1	**1** (perfect mixing), [0.95, 1], [0.9, 0.95]
Relative contact rate among <1 year-olds	β_1; Eqn 2	Set to 1
Relative contact rate among 1-4 year-olds	β_2; Eqn 2	[3, 30]
Relative contact rate among 5-17 year-olds	β_3; Eqn 2	[25, 50]
Relative contact rate among 18-49 year-olds	β_4; Eqn 2	[20, 40]
Relative contact rate among 50+ year-olds	β_5; Eqn 2	[1, 5]
Relative contact rate between 1-4 and 5-17 year-olds (sibling interactions)	β_6; Eqn 2	[1, 5]
Contact rate between 18-49 and 1-4 or 5-17 year-olds (parent-child interactions)	β_7; Eqn 2	[1, 5]
Amplitude of school term-time forcing	b_1; Eqn 3	[0.25, 0.75], **[0.5, 1]**
Basic reproductive number	R_0; Eqn 4	**[5, 10]**, [7, 12]
Reporting rate	r; Eqn 6	[80, 100]%

Table 2. Estimated proportion of infections caused by each age-group. Rows show the receiving (i.e. infectee) age groups and columns show the source of infection (i.e. infector age group). The numbers are the mean (and 95% CI) estimates in percentage. For instance, for <1 year-olds (3[rd] row), on average 16.3% of cases were infected by the same age group, 44.6% by 1-4 year-olds, 20.9% by 5-17 year-olds, 15.2% by 18-49 year-olds, and 3% by 50+ year-olds.

Infectee age groups	Infector age groups				
	<1 year	1-4 years	5-17 years	18-49 years	50+ years
<1 year	16.3 (12.3, 21.0)	44.6 (35.5, 53.5)	20.9 (13.9, 28.4)	15.2 (9.6, 21.6)	3.0 (1.4, 5.0)
1-4 years	1.8 (1.0, 3.0)	85.8 (77.0, 91.9)	7 (3.2, 12.9)	5 (2.2, 9.3)	0.3 (0.1, 0.6)
5-17 years	1.5 (0.9, 2.4)	12.9 (6.9, 21.3)	80.9 (70.4, 88.4)	4.4 (2, 8)	0.3 (0.1, 0.5)
18-49 yeas	2.4 (1.5, 3.7)	18.6 (10.9, 28.0)	9.1 (4.8, 14.9)	69.5 (56.2, 80.8)	0.4 (0.2, 0.8)
50+ years	15.5 (11.6, 19.9)	40.1 (31.1, 49.3)	19.7 (13, 27)	15.6 (9.7, 22.4)	9.1 (3.8, 16.3)

Table 3. Estimated impact of vaccination campaigns during Oct 2018 – July 2019. Columns 2-4 show the total numbers of infections (2nd column), hospitalizations (3rd column), and individuals in intensive care unit (ICU) for different age groups (rows 3 to 6) and overall (last row), should there be no vaccination campaigns. Columns 5-7 show the number of infections, hospitalizations, and ICU cases averted by the vaccination campaigns, compared to data reported as of Aug 6, 2019. Numbers are the mean (and 95% confidence intervals) of 10,000 simulations.

Age group	No. if without vaccination campaigns			No. averted by vaccination campaigns		
	Infection	Hospitalization	ICU	Infections	Hospitalization	ICU
<1	1015 (0, 1430)	75 (0, 106)	16 (0, 22)	937 (0, 1330)	70 (0, 99)	14 (0, 20)
4-5	3052 (3, 4096)	227 (0, 305)	47 (0, 63)	2837 (0, 3821)	211 (0, 284)	44 (0, 59)
5-17	1107 (1, 1692)	82 (0, 126)	17 (0, 26)	999 (0, 1554)	74 (0, 116)	15 (0, 24)
18+	904 (1, 1343)	67 (0, 100)	14 (0, 21)	803 (0, 1214)	60 (0, 90)	12 (0, 19)
All	6078 (5, 8443)	452 (0, 628)	94 (0, 130)	5576 (0, 7801)	415 (0, 580)	86 (0, 120)

FIGURES

Figure 1. Schematic of the measles transmission model. Measles transmission model follows the susceptible (S), exposed (E) and latently infected, infectious (I), and recovered (R) and/or immunized SEIR dynamics and includes 5 age-groups as indicated by the subscripts (i.e., <1, 1-4, 5-17, 18-49, and 50+ year-olds, respectively) and a group (M) for infants with maternal immunity. Black solid arrows show the disease-related processes; grey solid arrows show the demographic processes including birth (horizontal), aging (vertical), and death (tilted). Black dashed arrows show processes related to the routine 2-dose measles vaccination where susceptible individuals are vaccinated at ages 1 and 5 and move to the respective immune groups. Red dotted arrows show processes related to vaccination of susceptible individuals under 18 during the vaccination campaigns.

Figure 2. Monthly incidence for all ages and by age group. The solid line (y-axis on the right) shows monthly incidence for all ages, reported as of Aug 6, 2019. For comparison, bars (y-axis on the left) show monthly incidence for <1 (blue), 1-4 (orange), 5-17 (grey) and 18+ (yellow) year-olds, respectively, estimated based on health reports.

Figure 3. Model fit. Box plots show estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F). Results are pooled over all 10 model-filter runs (each with 10,000 and in total 100,000 model realizations). Horizontal thick lines show the median of model estimates; box edges show the 25[th] and 75[th] percentiles; whiskers show the 2.5[th] and 97.5[th] percentiles; and dots show outliers. Stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Figure 4. Estimated changes in population susceptibility. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) for <1 year-olds (A), 1-4 year-olds (B), 5-17 year-olds (C) and 18+ year-olds (D), respectively, at the end of each month from Sep 2018 to July 2019. The initial susceptibilities, estimated at the end of Sep 2018, were computed by adding the total numbers of individuals immunized by the vaccination campaigns in Oct 2018 to the posterior estimates at the end of Oct 2018. For comparison, the grey bars (y-axis on the right) show estimated numbers of individuals immunized during the vaccination campaigns; note that the vaccination campaigns targeted individuals under 19 years and thus is not shown for 18+ year-olds.

Figure 5. Estimates of key model parameters: (A) the basic reproductive number, (B) the effective reproductive number, (C) infectious period, (D) relative contact rate among 1-4 year-olds, (E) relative contact rate among 5-17 year-olds, and (F) relative contact rate among 18-49 year-olds. Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) made at the end of each month from Oct 2018 to July 2019. For comparison, the grey bars (y-axis on the left) show monthly incidence for all ages (A-C) or the related age groups (D-F).

Figure 6. Estimated impact of vaccination campaigns. Box plots show simulated estimates of monthly incidence for all ages (A), percentage of cases reported in each age group (B), and monthly incidence for <1 year-olds (C), 1-4 year-olds (D), 5-17 year-olds (E), and 18+ year-olds (F), should there be no vaccination campaigns. Results are pooled over 10,000 model simulations. Horizontal thick lines show the median of model estimates; box edges show the 25[th] and 75[th] percentiles; whiskers show the 2.5[th] and 97.5[th] percentiles; and dots show outliers. For comparison, stars (*) in A and B show monthly incidence for all ages and the age distribution, reported as of Aug 6, 2019; crosses (x) in C-F show age-grouped monthly incidence estimated from health reports.

Supplementary Material

Table S1. Comparison of model estimates on initial susceptibility and model performance under different assumptions on vaccination campaigns. The baseline setting is as reported in the main text and alternative settings 1 to 3 are as described in the section "*Sensitivity analysis on vaccination campaigns settings.*" The results are summarized by pooling all 10 model-filter runs (10,000 particles each run and 100,000 model realizations in total). The numbers are the mean and, for the susceptibilities, 95% credible intervals in the parentheses. The initial susceptibilities, estimated at the end of Sep 2018, were computed by adding the total numbers of individuals immunized by the vaccination campaigns in Oct 2018 to the posterior estimates at the end of Oct 2018.

		Model Settings on Vaccination Campaigns			
	Age group	Baseline	Alternative 1	Alternative 2	Alternative 3
Estimated initial susceptibility (%) at end of Sep 2018	<1 year	53.2 (49, 57.5)	54.2 (50, 58.4)	54.2 (50, 58.5)	54.2 (50, 58.5)
	1-4 years	24.9 (20.4, 29.7)	24.9 (20.4, 29.7)	24.9 (20.4, 29.7)	29.9 (25.4, 34.7)
	5-17 years	6.0 (4.1, 7.9)	7.5 (5.1, 9.9)	7.5 (5.1, 9.9)	6.0 (4.1, 7.9)
	18-49 years	6.0 (4.1, 7.9)	7.5 (5.2, 9.9)	7.5 (5.2, 9.9)	7.5 (5.2, 9.9)
	50+ years	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)	6.0 (4.1, 7.9)
Log-likelihood		-255.25	-265.52	-255.25	-257.42
Relative error of total number of cases over the outbreak	<1 year	0.39%	-11.96%	-11.56%	-17.91%
	1-4 years	-0.04%	-7.80%	-6.90%	-8.07%
	5-17 years	-4.13%	14.38%	0.33%	1.84%
	18+ years	-4.70%	7.24%	5.97%	7.95%
	All ages	-1.79%	-0.65%	-3.48%	-4.25%
Root-mean-square-error (RMSE), over Oct 2018 – July 2019	<1 year	1.74	1.86	1.80	2.55
	1-4 years	8.72	8.13	8.17	7.70
	5-17 years	3.91	5.27	3.00	5.10
	18+ years	5.08	4.62	4.30	4.25
	All ages	7.31	7.52	5.88	6.29
Correlation, over Oct 2018 – July 2019	<1 year	0.99	1.00	1.00	0.99
	1-4 years	0.95	0.96	0.96	0.97
	5-17 years	0.99	0.97	0.99	0.97
	18+ years	0.97	0.97	0.97	0.97
	All ages	0.99	0.99	1.00	1.00
1-step-head prediction RMSE, over Oct 2018 – Mar 2019	<1 year	4.37	4.46	4.40	4.60
	1-4 years	17.13	17.34	17.35	25.12
	5-17 years	8.27	8.39	8.37	8.50
	18+ years	4.34	4.93	4.93	5.06
	All ages	27.79	28.23	28.16	35.02

Figure S1. Estimates of model parameters not listed in Figure 5: (A) amplitude of school term-time forcing, (B) latent period, (C) reporting rate, (D) relative contact rate among 50+ year-olds, (E) relative contact rate between 1-4 and 5-17 year-olds (i.e. sibling interactions), and (F) relative contact rate between 18-49 and 1-4 or 5-17 year-olds (i.e. parent-child interactions). Red lines and surrounding areas (y-axis on the left) show the mean and 95% credible intervals of estimates pooled over all 10 model-filter runs (100,000 model realizations in total) made at the end of each month from Oct 2018 to July 2019. For comparison, the grey bars (y-axis on the left) show monthly incidence for all ages. Note that m_1 and m_2 are not shown as both optimal priors are the value 1 (Table 1).