

**Title: Relationship between early childhood non-parental childcare and diet, physical activity, sedentary behaviour, and sleep: A systematic review of longitudinal studies**

Silvia Costa<sup>1,2</sup>, Sara E. Benjamin Neelon,<sup>1,3</sup> Eleanor Winpenny<sup>1</sup>, Veronica Phillips<sup>4</sup>, Jean Adams<sup>1</sup>

1 UKCRC Centre for Diet and Activity Research, MRC Epidemiology Unit, School of Clinical Medicine, University of Cambridge, United Kingdom

2 School of Sport, Exercise and Health Sciences, Loughborough University, United Kingdom

3 Department of Health, Behavior and Society, Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, United States

4 Medical Library, School of Clinical Medicine, University of Cambridge, Cambridge, United Kingdom

**Corresponding author:** Silvia Costa

**Address:** School of Sport, Exercise and Health Sciences, Loughborough University, Epinal Way, Loughborough, LE11 3TU, United Kingdom

**Email:** [s.costa@lboro.ac.uk](mailto:s.costa@lboro.ac.uk)

**Telephone:** +44 (0)1509 226 304

1 **ABSTRACT**

2 **Background:** The rising prevalence of childhood obesity is a global public health concern. Evidence  
3 suggests that exposure to non-parental childcare before age six years is associated with increased  
4 risk of obesity, diet, and activity behaviours (physical activity, sedentary behaviour, and sleep).

5 However, findings are inconsistent and mostly from cross-sectional studies, making it difficult to  
6 identify the direction of causation in associations. This review identified and synthesised the published  
7 research on longitudinal associations between non-parental childcare during early childhood, diet, and  
8 activity behaviours.

9 **Methods:** Seven databases were searched using a predefined search strategy. Results were  
10 independently double-screened through title/abstract and full-text stages according to predefined  
11 criteria. Included studies were tabulated, and evaluated for risk of bias using the Nutrition Evidence  
12 Library Bias Assessment Tool.

13 **Results:** Of 18793 references screened, 13 studies met eligibility criteria and were included in the  
14 review. Eight studies reported on diet and seven studies reported on activity behaviour outcomes  
15 (three on physical activity, three on sedentary behaviour, and one on sleep). These studies included  
16 results on 89 tested childcare:outcome associations. Of 63 associations testing diet outcomes, 37  
17 (59%) were null, and the remainder showed inconsistent patterns. There was an indication of a  
18 potential benefit of Head Start providers (vs other care, including parental) on dietary behaviours. Of  
19 26 associations testing activity behaviour outcomes, 22 (85%) were null, and the remainder were  
20 inconsistent. Most studies (92%) did not use (or did not report using) valid and reliable outcome  
21 measures, and outcome assessors were not blinded (or it was unclear if they were blinded) to  
22 children's exposure status (77%).

23 **Conclusions:** The scarce available literature indicates little and mixed evidence of a longitudinal  
24 association between exposure to non-parental childcare before age six years and diet or activity  
25 behaviours. This reflects a paucity of research, rather than clear evidence of no effect. There is an  
26 urgent need for studies investigating the longitudinal associations of non-parental childcare on diet  
27 and activity behaviours to assess potential lasting effects and mechanisms. Studies should assess  
28 whether and how associations vary by provider and child sub-groups, as well as differences by  
29 intensity and duration of care.

30

31 **Keywords:** Early childhood; Early care and education; Dietary behaviours; Activity behaviours;  
32 Obesity risk factors

33

34

## 35 **BACKGROUND**

36 High rates of childhood obesity are a worldwide concern (1, 2). Globally in 2016, approximately 40.6  
37 million (6%) children under 5 years of age were overweight or obese (1), an increase of over one third  
38 from 30.4 million in 2000 (1). Obesity during childhood is associated with increased risk of both  
39 obesity and a range of other conditions later in life, including low self-esteem, high blood pressure,  
40 insulin resistance, coronary heart disease and stroke (3-5) . The early years (<6 years of age) have  
41 been repeatedly highlighted as a critical period for the development and prevention of obesity (4, 6,  
42 7), as well as the establishment of related healthy habits, such as healthy diet, physical activity, and  
43 sleep patterns (8, 9). Several individual, inter-personal and environmental factors influence the  
44 development of childhood obesity (10). Because they affect large numbers of children, environmental  
45 factors such as childcare settings represent potential targets for obesity prevention (11, 12).

46 An increasing number of children now attend non-parental childcare prior to 6 years of age, and many  
47 spend large proportions of their week days in such care (13, 14). A report by the United Nations'  
48 Children's Emergency Fund shows that roughly 80% of 3–6 year olds and 25% of 0–3 year olds in  
49 developed countries spend time in some form of childcare (14). A growing body of research, including  
50 a number of systematic reviews, suggests that attendance at childcare is associated with increased  
51 adiposity or risk of obesity in children (15-17). However, the available evidence is inconsistent (15, 16,  
52 18, 19), and may partly depend on different aspects of the childcare received, such as the type (i.e.,  
53 informal or formal care) or intensity (e.g., number of hours per week).

54 The pathways through which non-parental childcare might affect obesity are poorly understood (20,  
55 21). Different types and characteristics of childcare settings may have different influences on the  
56 development of obesity-related risk factors, such as diet and activity behaviours (including physical  
57 activity, sedentary behaviour, and sleep) (12, 22-25). Evidence suggests that some types of non-  
58 parental childcare (e.g., grandparents or Head Start in the US) and staff behaviours (e.g., giving non-  
59 food rewards and allowing children to self-serve) are associated with diet patterns and behaviours  
60 (24, 26). Similarly, different types (e.g., home-based versus centre-based settings) and features (e.g.,

61 staff behaviours like playing with children) of childcare are associated with physical activity (27-30)  
62 and sedentary behaviour (23, 30, 31) in young children. There is also some evidence that attending  
63 some types of childcare is associated with problematic sleeping patterns in young children (22, 25,  
64 32). However, the direction of these associations is mixed, and associations are not consistently  
65 found in all population sub-groups or studies. Additionally, the vast majority of the current evidence  
66 comes from cross-sectional studies, which makes it difficult to determine the direction of causation.  
67 The aim of this review was to systematically gather and synthesise the published research on the  
68 longitudinal relationship between non-parental childcare in the early years and diet, physical activity,  
69 sedentary behaviour, and sleep. We focused exclusively on longitudinal studies to increase  
70 confidence that any association we find might be causal.

71

## 72 **METHODS**

73 This review was part of a larger programme of systematic reviews (including obesity and stress  
74 outcomes alongside the diet and activity behaviour outcomes reported here) that was registered with  
75 the PROSPERO database (registration number CRD42015027233) (33), and reported in line with the  
76 Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) recommendations  
77 (34). The protocol for the overall programme of systematic reviews has been published elsewhere  
78 (35).

79

### 80 **Search Strategy**

81 Seven electronic bibliographic databases were searched in January 2016, using a predefined search  
82 strategy: MEDLINE, EMBASE, PsycINFO, Web of Science, Scopus, Applied Social Sciences Index  
83 and Abstracts (ASSIA), and the Scientific Electronic Library Online (SciELO). Searches were  
84 restricted to human subjects, but there were no restrictions placed on publication date or language.  
85 The search strategy was based on the key themes of relevance to the overall review as described in  
86 the protocol – childcare, adiposity and body mass, physical activity, sedentary behaviour, sleep, diet,  
87 and stress – and informed by search strategies of relevant previous systematic reviews (36). An  
88 experienced university librarian (VP) reviewed the search strategy, adapted it for different databases,  
89 and ran the searches. An example of the search strategy used for the MEDLINE and Embase

90 databases can be found in Supplement 1. Results were merged from the different databases and  
91 managed using EndNote® software. The searches were re-run at the end of May 2017.

92

### 93 **Study Selection and eligibility criteria**

94 After removal of duplicates, records were screened in two phases using a pre-piloted procedure (Box  
95 1). In phase one, title and abstracts were screened by two reviewers working independently against  
96 the five phase one eligibility criteria described in Figure 1. The full texts of all studies identified by  
97 either reviewer as potentially eligible were retrieved. In phase two, full texts were screened by two  
98 reviewers working independently against the seven phase two eligibility criteria described in Figure 1.  
99 In cases of uncertainty or discrepancy between reviewers, we consulted a third reviewer and  
100 consensus was achieved by discussion.

101

102 Figure 1 – Eligibility criteria

103

104 The number of papers included and excluded at each stage of the review process can be seen in the  
105 PRISMA diagram presented in Figure 2. The diagram shows overall number of references screened  
106 and excluded (by reason) instead of number of references by outcome of interest, because eligibility  
107 at full-text screening (phase two) was done for all six outcomes of interest in the registered protocol  
108 for the programme of systematic reviews (obesity, stress, diet, physical activity, sedentary behaviour  
109 and sleep) (33). Concurrently, the final number of included studies are presented for behaviour  
110 outcomes only that were the focus of the current review – diet, physical activity, sedentary behaviour,  
111 and sleep.

112

113 Figure 2 – PRISMA diagram.

114

115 Details of and justification for each eligibility criteria are described in full in the protocol (35). Studies  
116 were included where participants were children aged <6 years and not in primary school at first  
117 assessment, and living in middle- and high-income countries as defined by the World Bank (37). Only  
118 observational longitudinal study designs, including case-control, prospective, and retrospective  
119 designs, were included. The exposure of interest was non-parental childcare where there was

120 between-child variation in exposure, for example by timing of attendance (i.e., age when care started  
121 and stopped), intensity (i.e., full- or part-time care), duration (i.e., years of childcare), types (i.e.,  
122 formal or informal; private or public), or simply attendance versus non-attendance. Studies were  
123 included where outcomes were objectively assessed or proxy/self-reported measures of diet, physical  
124 activity, sedentary behaviour, or sleep. Studies were excluded if they were not published in peer-  
125 reviewed journals.

126 When the team could not resolve issues of whether eligibility criteria were met, study authors were  
127 contacted via email for clarification. If authors did not reply by the end of the data extraction stage,  
128 studies were excluded from the review. Conference abstracts, masters and doctoral theses were  
129 excluded, as these do not necessarily go through a formal peer-review process. Nevertheless, the  
130 authors of any potentially relevant records of these types of references were contacted via email to  
131 determine if peer-reviewed journal articles had resulted, and screened articles identified through this  
132 route according to the phase two process above.

133

#### 134 **Data Extraction and Management**

135 A standardised and pre-piloted form was used to extract data from included studies for assessment of  
136 study quality and evidence synthesis. This captured information about study setting and population,  
137 exposure and outcome variables, statistical analyses and results, as stated in the protocol (35). The  
138 first author extracted these data into an Excel® database, and a second author (JA/SBN)  
139 independently checked the extracted information against the full-texts of included studies.

140

#### 141 **Data Synthesis**

142 Key information was tabulated (e.g., sample characteristics, exposure, and outcome measures) for  
143 each study, grouped by outcome variable, and performed a narrative synthesis of the included  
144 studies. Because of heterogeneity in exposure and outcome variables, it was not appropriate to  
145 perform a meta-analysis. This also meant that it was not possible, as originally planned, to perform a  
146 quantitative synthesis of differences in effect between different types and features of childcare,  
147 different outcomes, high- and middle-income countries, ages at exposure, and socio-demographic  
148 sub-groups (e.g., by ethnicity). Instead, and because of the sometimes large number of relevant

149 exposure and outcome variables used in included studies, all individual relevant associations reported  
150 in the included studies were included here.

151

## 152 **Quality Assessment**

153 An adaptation of the United States Department of Agriculture's Nutrition Evidence Library Bias  
154 Assessment Tool (NEL-BAT) (38) was used to assess risk of bias in included studies. This tool  
155 assesses risk of selection, performance, detection, and attrition bias. For observational studies,  
156 responses to each of the 13 questions are scored 0–2 (possible range of scores: 0–26), where lower  
157 scores indicate lower risk of bias. SC and JA independently assessed all included studies for risk of  
158 bias, and disagreements in scores were resolved by discussion.

159

## 160 **RESULTS**

161 The literature search identified 47 529 articles. After de-duplication, 18 793 articles underwent  
162 title/abstract screening, and the full texts of 175 articles were reviewed. Thirteen studies (39-51) met  
163 all of the eligibility criteria, and were included in the review. Of these, eight studies reported on diet  
164 (39, 45-51), three reported on physical activity (39, 41, 44), three on sedentary behaviour (41-43), and  
165 one on sleep (40) outcomes. Some studies reported on more than one of diet, physical activity,  
166 sedentary behaviour and sleep.

167

### 168 **Summary of Included Studies**

169 A detailed description of each study's characteristics can be seen in Table 1. Most included studies  
170 were from high-income countries, with seven originating from the United States (39, 40, 42, 43), one  
171 from Australia (41), one from New Zealand (44), and one from the UK (48). Samples were generally  
172 balanced with relation to children's gender (although three studies did not report gender composition  
173 of the sample) (47, 48, 51), but varied greatly both in size (between 34 and 18 050 subjects) and  
174 ethnic composition (between 0% and 87% white, with one study not reporting race/ethnicity or country  
175 of birth (41), and five providing information only for country of birth) (45, 46, 48, 49, 51).

176

177 [Table 1– See additional file 1]

178

179 All studies assessed exposure to non-parental childcare between birth and 5 years. Measurement of  
180 exposure was by proxy-report from a parent or primary caregiver in all studies. Nine studies described  
181 childcare exposure in simple categorical terms (e.g., centre-based preschool or Head Start centre  
182 versus other/mixed care, including parents) (39-41, 45, 46, 48-51). One study assessed duration of  
183 exposure (centre-based preschool for at least 2 years versus other/mixed care, including parents)  
184 (39). Five studies assessed intensity of non-parental childcare (e.g., average number of hours in  
185 childcare between 24-36 months) (42-44, 47, 48).

186 Age at outcome assessment varied from 1-12 months to 51 years. For diet outcomes, all studies used  
187 proxy-report by a mother or main caregiver (39, 45-51). For physical activity outcomes, most studies  
188 used proxy- or self-report (39, 41, 44), with the exception of average accelerometer counts/minute in  
189 one case (44). All studies assessing sedentary behaviour outcomes (e.g., hours/day of television (TV)  
190 viewing) using proxy-report by a parent. For sleep, quantitative outcomes (e.g., nap durations) were  
191 measured objectively by an accelerometer, whereas qualitative outcomes (e.g., difficulty going to bed  
192 or falling asleep) were subjectively measured by parent-report (40).

193 No two studies used the same outcome variables. All diet studies presented outcomes as categorical  
194 variables, with three studies investigating breastfeeding related outcomes (e.g., breastfeeding  $\geq 4$   
195 months) (47, 48, 51), four studies investigating consumption of healthy and unhealthy foods (e.g.  
196 consuming sweetened beverages, salty snacks, or fruits/vegetables) (39, 45, 46, 50), and one study  
197 investigating between-meal eating (49). Two studies presented physical activity outcomes as  
198 categorical variables (e.g., high, medium, and low physical activity level versus sedentary) (39, 41),  
199 whereas one study used continuous variables (e.g., average accelerometer counts per minute). All  
200 sedentary behaviour variables were categorical (e.g.,  $>4$  versus  $\leq 4$  hours/day of TV viewing), and two  
201 out of three studies used TV viewing as a proxy for sedentary behaviour. There was a wide range of  
202 sleep variables, from number and duration of naps to variables relating to the quality of sleep.

203 Four studies investigated only one exposure and one outcome variable (41-43, 47). The remainder  
204 explored several outcome (40) or exposure variables (39, 44). Thus, the eight included studies  
205 reported on 63 associations between non-parental childcare and diet outcomes, nine associations  
206 with physical activity outcomes, three associations with sedentary behaviour outcomes, and 15  
207 associations with sleep outcomes. Nine studies employed analytical strategies that accounted for  
208 potential confounding effects of other variables (i.e., multivariable regression models) (39, 41, 45-51),



209 8 of which investigated diet outcomes, whereas the remainder used simple statistical tests which did  
210 not account for potential confounding factors (40, 42-44). In the majority of cases, this was because  
211 the association between childcare and physical activity, sedentary behaviour or sleep was not the  
212 main focus on the paper.

213

## 214 **Synthesis of Findings**

215 Table 2 presents detailed results for all relevant associations explored in each study.

216

### 217 *Diet*

218 Eight studies evaluated the longitudinal relationship between non-parental childcare during early  
219 childhood and diet outcomes (39, 45-51). Results were highly mixed. Lee et al (46) reported that  
220 children who attended Head Start settings at 4 years of age showed significantly higher frequency of  
221 healthy eating patterns at 5-6 years of age than those attending other settings (all  $p < 0.05$ ), except  
222 Pre-Kindergarten. Conversely, no differences in frequency of unhealthy eating patterns were found  
223 between the groups (46). Another study assessing attendance at Head Start (39) found that children  
224 who attended Head Start or other centre-based childcare at 4 years of age (irrespective of length of  
225 exposure) were more likely to report frequent fruit consumption than those in other/mixed care  
226 (including parental care) at age 5-6 years. In this study, children who attended other (non-Head Start)  
227 centre-base childcare were also less likely to regularly consume soda at age 5-6 years than those in  
228 other/mixed care (all  $p < 0.05$ ). However, other centre-base childcare was also associated with higher  
229 likelihood of regular consumption of chips ( $p < 0.05$ ). There was no difference in the likelihood of  
230 regular consumption of fast-food, candy, and chips consumption, and frequent consumption of  
231 vegetables between those attending Head Start or other centre-based childcare (irrespective of length  
232 of exposure) versus other/mixed care (including parental care).

233 Similarly, Wasser et al (50) also reported mixed findings. They found that children in any non-  
234 maternal childcare had higher odds of consuming whole fruit (Odds Ratio (OR): 1.15,  $p < 0.05$ ), and  
235 juice (any childcare OR: 1.51; Grandparents OR: 1.91,  $p < 0.05$ ) than those in maternal care. But there  
236 was no association between childcare (overall or by type) and consumption of five other food and  
237 drinks including vegetables and salty snacks. Camara et al (45) also reported mostly null associations  
238 between childcare and the two dietary patterns investigated, except a higher adherence to a

239 processed/fast-food pattern at 2-5 years of age in those being cared for at home by someone other  
240 than the mother compared to those cared for by their mother ( $B= 0.70$  ( $SE:0.14$ ),  $p<0.001$ ) at 2-3  
241 years of age. Sata et al's (49) reported more frequent between meal eating before dinner at age 6  
242 years in those cared for by grandparents and nursery/kindergarten than those cared for by mothers,  
243 as well as between meal eating  $\geq 3$  times per day those cared for by grandparents versus those cared  
244 for by mothers at 3 years of age. However, no other associations were found with any care at 12 or  
245 22 years of age.

246 Three studies investigated breastfeeding outcomes (47, 48, 51), showing mixed results. Pearce et al  
247 (48) reported lower likelihood of breastfeeding for  $\geq 4$  months in children attending informal compared  
248 to parental care (independent of attending full- or part-time, lone parenthood, or area of deprivation),  
249 but mixed results for those attending formal care. For example, in the analyses stratified by family  
250 structure, children living in single parent families receiving formal care were more likely to be  
251 breastfed for  $\geq 4$  months (Risk ratio (RR)= 1.65) than those being cared for by parents, but the reverse  
252 was true for children living in couple families (RR= 0.79, all  $p<0.05$ ). While Weille et al (51) reported a  
253 higher risk of changing from mostly breastfed to mostly or solely formula-fed in those attending  
254 childcare compared to those cared for at home (RR= 2.05 to 2.50,  $p<0.05$ ), Levy et al (47) found an  
255 increased risk of earlier cessation of breastfeeding in children who used pacifier and did not attend  
256 any childcare and those who attended 15 days of childcare between 0-6 months of age versus those  
257 not attending childcare and not using pacifier.

258

259 Overall, the eight included studies tested 63 associations between non-parental childcare exposures  
260 and diet outcomes. Of these, 37 (59%) were null, 10 (16%) indicated significant beneficial effects of  
261 non-parental care on dietary behaviours, seven (11%) indicated significant detrimental effects of non-  
262 parental care on dietary behaviours, one (2%) found significant association with a diet behaviour that  
263 is of unsure detrimental effect (between-meal snacking) (49), and eight (13%) found mixed results, As  
264 an example of the latter, when investigating associations by maternal education, Pearce et al. (48)  
265 reported that those in informal care had significantly lower odds of being breastfed  $\geq 4$  months versus  
266 those in parental care, but only when parents had GCSE D-G or lower education. At the same time,  
267 those in formal care also had significantly lower odds of being breastfed  $\geq 4$  months versus those in  
268 parental care, but only when parents had degree-level education (48).

269

270 *Physical Activity*

271 Three studies evaluated the longitudinal relationship between childcare during early childhood and  
272 physical activity outcomes (39, 41, 44). Results were highly mixed. Belfield & Kelly (39) found that  
273 children who attended Head Start at 4 years old had significantly lower physical activity levels in  
274 kindergarten than those who received parental care. However, there was no difference in physical  
275 activity between those attending other centre-based care versus parental care, irrespective of length  
276 of exposure to such care. Conversely, D'Onise et al (41) reported that those attending Kindergarten  
277 Union preschool between 2 and 4 years old were more likely to be in the high physical activity level  
278 group (versus sedentary group) at around age 45 years than those who did not attend this preschool.  
279 In the only study looking at intensity of childcare use, Taylor et al (44) found no significant  
280 associations between weekly hours of childcare attendance at 3 or 4 years old and objectively  
281 measured physical activity 1 or 2 years later.

282 Overall, the six included studies tested nine associations between non-parental childcare exposures  
283 and physical activity outcomes. Of these, seven (78%) were null (39, 44), whereas two (22%) found  
284 significant differences but in competing directions, with one showing childcare to be associated with  
285 more, and one with less physical activity (39, 41).

286

287 *Sedentary Behaviour*

288 Three studies evaluated the longitudinal relationship between childcare during early childhood and  
289 sedentary behaviour outcomes (41-43), including one study that conceptualised sedentary behaviour  
290 as the absence of physical activity (also reported on above) (41). As noted, D'Onise et al (41)  
291 reported that those who attended Kindergarten Union preschool between ages 2 and 4 years were  
292 less likely to be in the sedentary group (versus the high physical activity level group) at around 45  
293 years, than those who did not attend this preschool. The remaining two studies (42, 43) found no  
294 significant associations between number of hours per week of childcare at 24-36 months and 3-5  
295 years and subsequent daily hour of television viewing at 36 months and 6-12 years respectively.  
296 Thus, three associations between non-parental childcare and sedentary behaviour were tested in  
297 included studies. Two (67%) were null (42, 43), and one (33%) showed a significant association  
298 between childcare and lower risk of sedentary behaviour (41).

299

300 *Sleep*

301 The only study investigating the longitudinal relationship between childcare and sleep outcomes  
302 yielded mixed results (40). Cairns and Harsh (40) reported that those attending all day preschool or  
303 day care at age 5 years (versus a primary/secondary caregiver) transitioned to earlier sleep onset and  
304 wake up time on week days in the first months of preschool. The authors are clear that the health  
305 implications of these differences are unknown. There were no differences between the groups in any  
306 other variables (e.g., difficulty in going to bed and nocturnal sleep duration on week days).  
307 Overall, 15 associations between non-parental childcare and sleep outcomes were tested in this  
308 single included study. The majority (n=13) were null, with only two showing significant results. The  
309 health implications of these are unknown.

310

311 [Table 2 – See additional file 2]

312

### 313 **Quality Evaluation**

314 Risk of bias scores ranged from 1-12 out of a total maximum of 26 (with lower scores indicating lower  
315 risk of bias) (see Table 3). The most common sources of bias were not reporting or using valid and  
316 reliable outcome measures (NEL-BAT question 13; 12 studies), and outcome assessors not blinded  
317 (or not clear whether they were blinded) to the intervention or exposure status of participants (NEL-  
318 BAT question 12; 10 studies). There was low risk of bias throughout in terms of inclusion and  
319 exclusion criteria (question 2), recruitment strategy (question 2), accounting for variations in the  
320 execution of the study from the proposed protocol or research plan (question 7), and length of follow-  
321 up across study groups (question 14).

322

323 [Table 3 – See additional file 3]

324

## 325 **DISCUSSION**

### 326 **Summary of findings**

327 To our knowledge, this is the first systematic review to investigate the longitudinal relationship  
328 between non-parental childcare before age 6 years and diet, physical activity, sedentary behaviour

329 and sleep. Overall, the evidence base is very limited with only 13 studies meeting eligibility criteria. In  
330 total, eight studies reported on diet outcomes, three on physical activity, three on sedentary behaviour  
331 and one on sleep. Included studies varied widely in terms of definition and measurement of both  
332 exposure and outcomes, and lacked in-depth exploration of different aspects of childcare that may  
333 influence any relationship with the outcomes studied. The available, limited, longitudinal literature  
334 suggests that attending certain types of non-parental childcare (particularly informal providers) might  
335 be related to less breastfeeding, but the evidence regarding other dietary outcomes is mixed, and  
336 sometimes contradictory. Moreover, the data reviewed suggests that attending non-parental childcare  
337 is unrelated to physical activity, sedentary behaviour or sleep outcomes. Included studies were of  
338 mixed quality with most (92%) not reporting using valid and reliable outcome measures, and few  
339 (23%) including blinding of outcome assessors to participants' exposure status.

340

#### 341 **Strengths and limitations of studies included in the review**

342 The measurement of exposure to childcare in included studies was highly variable. Some studies  
343 focused on one particular type of childcare provider (e.g., attending Kindergarten Union) (41) versus a  
344 reference group that was an amalgam of all other types (39-41). Other studies included only the  
345 number of hours per week in non-parental childcare (42-44). Only four studies explored differences  
346 between the type of childcare provider (45, 48-50), but no studies performed detailed analyses  
347 exploring differences by duration, intensity, and timing of childcare. Thus, we were unable to explore  
348 differing effects according to these characteristics of childcare, as originally intended.

349 Similarly, there was substantial heterogeneity in how outcomes were assessed in included studies,  
350 which did not allow for direct comparisons. Apart from Cairns & Harsh's study (40), no study reported  
351 validity or reliability of the methods used for measuring outcomes. Concurrently, the common use of  
352 proxy-report measures of diet, physical activity, sedentary and sleep behaviour outcomes increases  
353 the risk of measurement error and bias (e.g., overestimating physical activity and underestimating  
354 sedentary behaviour).

355 Only seven (54%) studies used an adequate analytical framework that accounted for the potential  
356 complexity of the relationship between non-parental childcare and outcomes (physical activity in both  
357 cases) (39, 41, 45, 48-50), by including and statistically adjusting for potential confounding and  
358 mediating variables. Thus, the evidence base may be substantially compromised by uncontrolled

359 confounding by factors such as family and socioeconomic characteristics. Furthermore, apart from  
360 Pearce et al's (48) study focusing on breastfeeding, no other study explored variations in any  
361 relationships between exposures and outcomes according to contextual factors, such as maternal  
362 education or socioeconomic status. Thus, we were not able to report on these, as originally proposed  
363 in the protocol (33). Failure to adjust for confounding variables is often a result of the association  
364 between childcare and health behaviours not being the primary focus of the study. Greater attention  
365 to these associations as primary aims of studies is required to increase the strength of available  
366 evidence.

367

### 368 **Strengths and limitations of the review**

369 This systematic review has several strengths. A large number and variety of databases were  
370 searched using a comprehensive search strategy designed in partnership with an experienced  
371 librarian (VP), without limits on date of publication or language. Independent double screening was  
372 used at both abstract and full-text screening stages, and a third reviewer was included to resolve any  
373 inconsistencies in these processes, reducing risk of researcher bias. In cases of remaining  
374 uncertainty, study authors were contacted. The risk of bias assessment (NEL-BAT) was also  
375 performed independently in duplicate. This review focused on longitudinal studies because these  
376 provide a better indication of causality between exposure and outcomes than cross-sectional studies  
377 (52).

378 However, there are a number of limitations that need highlighting. The low number of studies for each  
379 of the outcomes did not allow us to present a summary of findings table as advised by the Cochrane  
380 Library Handbook, nor to perform a meta-analysis as planned (33). Heterogeneity in the study  
381 designs, definition of exposure and outcomes, and the methods and measurement tools used also  
382 made comparisons difficult; therefore, data were narratively synthesized and described. The results  
383 cannot be generalised to middle-income countries, as all studies were located in high-income  
384 countries. Furthermore, it was not possible to determine if associations persisted or emerged later,  
385 because only two articles and four (4%) associations had outcomes that were measured after  
386 childhood (41, 49); the remaining 11 articles and 85 (96%) associations assessed outcomes that were  
387 measured during childhood only (0-12 years).

388

### 389 **Interpretation of findings**

390 Overall, there were substantial null results with a few scattered and mostly inconsistent statistically  
391 significant associations between non-parental childcare and diet, physical activity, sedentary  
392 behaviour and sleep outcomes. There was an indication that attending Head Start settings might be  
393 associated with positive dietary behaviours compared to other/mixed care (including parental care)  
394 (39, 46). However, this evidence comes from only from two studies (39, 46), and was not seen across  
395 all dietary outcomes studied (e.g., there was an association with more frequent fruit but not vegetable  
396 consumption) (39), or in relation to all other childcare types (e.g., no significant difference in healthy  
397 eating patterns between Head Start and state-funded pre-kindergarten programmes). Few consistent  
398 findings were found for physical activity, sedentary behaviour or sleep.

399 Whilst cross-sectional studies generally find more evidence of an unhealthy effect of childcare on diet  
400 and activity behaviours (22, 23, 26, 28, 30, 53, 54), this does not appear to be reflected in the limited  
401 available longitudinal data. It is possible that any cross-sectional relationship does not persist  
402 longitudinally and, hence, that there is no long-term impact of childcare on diet and activity  
403 behaviours. This would suggest that identified longitudinal associations between childcare and  
404 adiposity occurs via other mechanisms, such as stress. Alternatively, and maybe more likely, the  
405 quality and quantity of the longitudinal evidence available on the relationship between childcare and  
406 diet and activity behaviours is not strong enough to draw conclusions on the presence or nature of  
407 any relationship.

408 Most included studies measured outcomes in childhood, up to age 12 years, only (39, 40, 42-44). It is  
409 possible that any effects of childcare on diet and activity behaviours emerge later in life – particularly  
410 when children start to develop into more independent adolescents and adults. Although not enough to  
411 corroborate it, the significant associations found in D'Onise et al's study (41), where physical activity  
412 level was assessed during mid-adulthood, support the plausibility of that hypothesis at least for  
413 activity behaviours.

414 The wide range of different outcome and exposure measures used in the included studies indicates  
415 poor theorisation and conceptualisation of any potential association. In general, there is limited  
416 evidence of shared understanding of exactly what aspect of childcare is expected to be associated  
417 with exactly what aspect of diet or type of activity behaviour, what the direction of such associations  
418 is, and why. Furthermore, authors rarely addressed the many dimensions that can vary in the

419 exposure to non-parental childcare in terms of provider, timing, duration and intensity (particularly in  
420 relation to the activity behaviours outcomes). Greater conceptual clarity in these areas may help drive  
421 stronger longitudinal investigations. Clearer disentanglement of all of the potential dimensions in  
422 which exposure to childcare may vary will help identify if there are more and less healthful ways in  
423 which children can receive childcare.

424 Although it was not possible to perform meta-analyses or meta-regressions, there is no obvious  
425 indication that results were related to study size, whether outcomes were considered as continuous  
426 and categorical variables, whether outcomes were measured using subjective or objective methods,  
427 and whether studies were prospective or retrospective. However, the very small number of studies  
428 included for each outcome makes it difficult to draw conclusions on these issues.

429

#### 430 **Implications for policy, practice and research**

431 Although overall there was little evidence of a longitudinal relationship between childcare and diet and  
432 physical activity, sedentary behaviour and sleep, this more likely represents a current absence of high  
433 quality evidence, rather than good evidence of absence of an effect. Given this, it is difficult to draw  
434 any firm implications for policy and practice. Nevertheless, and given that there is some evidence of  
435 an association between childcare and adiposity,(15-17) it would be prudent for those regulating and  
436 providing childcare to continue to consider how they can provide a healthful environment for the  
437 children in their care.

438 The small number of studies included in the current review highlights the need for more longitudinal  
439 studies investigating the relationship between non-parental childcare and diet, physical activity,  
440 sedentary behaviour and sleep. These studies should employ valid and reliable measures of both  
441 exposure and outcomes; analytical frameworks that recognise the potential complexity of the  
442 relationship between exposure and outcome, and account for known and possible confounding and  
443 mediating factors (e.g., socioeconomic status and maternal employment). Additionally, studies should  
444 also perform more detailed investigations to explore potential differences in the effect of childcare  
445 according to the type of provider, duration, intensity and timing of childcare. This would help in  
446 clarifying whether specific patterns of exposure to non-parental childcare have a more or less  
447 healthful impact on children's diet or activity behaviours.



448 The majority of studies included in this review assessed outcomes only during childhood (0-12 years),  
449 with only two studies assessing outcomes during adulthood (meal eating at age 22 years, and  
450 physical activity level at ages 45-51 years) (41, 49). There is a need for more studies examining long  
451 term relationships, to assess whether relationships between childcare and diet, physical activity,  
452 sedentary behaviour, or sleep emerge and continue into adolescence and adulthood. Existent birth  
453 cohorts may be useful in this respect, although they may impose limitations in the detail and methods  
454 of assessment of childcare exposure.

455 There is a lack of studies in middle-income countries, as well as consideration of differences in effect  
456 by ethnicity and socioeconomic status. Studies located in middle-income countries and investigating  
457 interactions with ethnicity and socioeconomic status would allow us to assess whether context  
458 influences the relationship between childcare and activity behaviours and hence whether targeted  
459 interventions may be justified. Indeed, the only included study which explored this aspect found that  
460 informal childcare was consistently associated with higher risk of not breastfeeding for at least 4  
461 months across two indicators of socioeconomic status; but there were inconsistent, mostly null,  
462 associations with formal childcare (48).

463

## 464 **CONCLUSIONS**

465 This review provides the first systematic summary of studies examining the longitudinal relationship  
466 between non-parental childcare and diet, physical activity, sedentary behaviour and sleep. Results  
467 were dominated by null findings with little consistent evidence that non-parental childcare was with  
468 any of the outcomes of interest. However, the available evidence is limited, highly heterogeneous in  
469 the definition and measurement of non-parental childcare, diet and activity behaviours, and lacks an  
470 in-depth exploration of different aspects of childcare that may influence this relationship, such as the  
471 type, duration or intensity. Further work is required to clearly conceptualise proposed pathways linking  
472 childcare with diet and activity behaviours, and to determine whether and what aspects of childcare  
473 might discourage physical activity, and promote sedentary behaviour, less healthful diet and sleep  
474 patterns. This would, in turn, help identify potential targets for interventions, policies, or regulations to  
475 help childcare settings provide healthful environments for the children in their care.

476

## 477 **DECLARATIONS**

478 **Ethics approval and consent to participate:** Not applicable.

479 **Consent for publication:** Not applicable.

480 **Availability of data and material:** Not applicable.

481 **Competing interests:** The authors declare that they have no competing interests.

482 **Funding:** The work was undertaken by the Centre for Diet and Activity Research (CEDAR), a UKCRC  
483 Public Health Research Centre of Excellence. Funding from the British Heart Foundation, Cancer  
484 Research UK, Economic and Social Research Council, Medical Research Council, the National  
485 Institute for Health Research, and the Wellcome Trust, under the auspices of the UK Clinical  
486 Research Collaboration, is gratefully acknowledged.

487 **Authors' contributions:** SC, SBN, and JA devised the systematic review, performed the risk of bias  
488 assessment, discussed and resolved disagreements on inclusion of studies at both title/abstract and  
489 full-text screening stages. VP advised on the design of the search strategy, conducted the database  
490 searches, and provided the search results for the review. SC lead on the merger and de-duplicating of  
491 the results database, and the drafting and submission of the manuscript. All authors performed  
492 screening at both stages, reviewed several drafts of the manuscript, and agreed on the final submitted  
493 version of the manuscript

494 **Acknowledgements:** The authors thank Dr. Sarah Gonzalez-Nahm for her invaluable help in the  
495 double-screening process of this review, and Kai Schulze and Amy Yau for their help with translating  
496 and screening of non-English full-texts.

497

498

## 499 **REFERENCES**

- 500 1. United Nations Children's Fund, World Health Organization, World Bank Group. Key findings of the  
501 Joint Child Malnutrition Estimates (2017 edition). 2017.
- 502 2. Onis M, Blössner M, Borghi E. Global prevalence and trends of overweight and obesity among  
503 preschool children. *Am J Clin Nutr.* 2010;92(5):1257-64.
- 504 3. Reilly JJ, Methven E, McDowell ZC, Hacking B, Alexander D, Stewart L, et al. Health  
505 consequences of obesity. *Arch Dis Child.* 2003;88(9):748-52.
- 506 4. Lakshman R, Elks CE, Ong KK. Childhood obesity. *Circulation.* 2012;126:1770-9.

- 507 5. Lawlor DA, Leon DA. Association of body mass index and obesity measured in early childhood  
508 with risk of coronary heart disease and stroke in middle age: findings from the aberdeen children of  
509 the 1950s prospective cohort study. *Circulation*. 2005;111(15):1891-6.
- 510 6. Cameron N, Demerath EW. Critical periods in human growth and their relationship to diseases of  
511 aging. *Am J Phys Anthropol*. 2002;119(Suppl 35):159–84.
- 512 7. Dietz WH. Critical periods in childhood for the development of obesity. *Am J Clin Nutr*.  
513 1994;59(5):955-9.
- 514 8. Gaffney KF, Kitsantas P, Brito A, Castello J. Baby Steps in the Prevention of Childhood Obesity:  
515 IOM Guidelines for Pediatric Practice. *J Pediatr Nurs*. 2014;29(2):108-13.
- 516 9. Jones RA, Hinkley T, Okely AD, Salmon J. Tracking physical activity and sedentary behavior in  
517 childhood: a systematic review. *Am J Prev Med*. 2013;44(6):651-8.
- 518 10. Davison KK, Birch LL. Childhood overweight: a contextual model and recommendations for future  
519 research. *Obes Rev*. 2001;2(3):159-71.
- 520 11. Leal C, Chaix B. The influence of geographic life environments on cardiometabolic risk factors: a  
521 systematic review, a methodological assessment and a research agenda. *Obes Rev*.  
522 2011;12(3):217-30.
- 523 12. Larson N, Ward DS, Neelon SB, Story M. What role can child-care settings play in obesity  
524 prevention? A review of the evidence and call for research efforts. *J Am Diet Assoc*.  
525 2011;111(9):1343-62.
- 526 13. Laughlin L. Who's minding the kids? Child care arrangements: Spring 2011. Washington, DC,  
527 Bureau C; 2011 2013.
- 528 14. United Nations Children's Fund. The child care transition, Innocenti Report Card 8. Florence:  
529 Innocenti Research Centre; 2008. Contract No.: 8.
- 530 15. Black L, Matvienko-Sikar K, Kearney PM. The association between childcare arrangements and  
531 risk of overweight and obesity in childhood: a systematic review. *Obes Rev*. 2017;Epub ahead of  
532 print.
- 533 16. Alberdi G, McNamara AE, Lindsay KL, Scully HA, Horan MH, Gibney ER, et al. The association  
534 between childcare and risk of childhood overweight and obesity in children aged 5 years and  
535 under: a systematic review. *Eur J Pediatr*. 2016;175(10):1277-94.

- 536 17.Swyden K, Sisson SB, Lora K, Castle S, Copeland KA. Association of childcare arrangement with  
537 overweight and obesity in preschool-aged children: a narrative review of literature. *Int J Obes.*  
538 2016;41:1.
- 539 18.Costa S, Adams J, Gonzalez-Nahm S, Benjamin Neelon SE. Childcare in Infancy and Later  
540 Obesity: a Narrative Review of Longitudinal Studies. *Curr Pediatr Rep.* 2017.
- 541 19.Swyden K, Sisson SB, Lora K, Castle S, Copeland KA. Association of childcare arrangement with  
542 overweight and obesity in preschool-aged children: a narrative review of literature. *Int J Obes.*  
543 2017;41(1):1-12.
- 544 20.Lumeng JC, Taveras EM, Birch L, Yanovski SZ. Prevention of Obesity in Infancy and Early  
545 Childhood: A National Institutes of Health Workshop. *JAMA Pediatr.* 2015; 169(5):484-90.
- 546 21.Monasta L, Batty GD, Cattaneo A, Lutje V, Ronfani L, Van Lenthe FJ, et al. Early-life determinants  
547 of overweight and obesity: a review of systematic reviews. *Obes Rev.* 2010;11(10):695-708.
- 548 22.Iwata S, Iwata O, Iemura A, Iwasaki M, Matsuishi T. Determinants of sleep patterns in healthy  
549 Japanese 5-year-old children. *International journal of developmental neuroscience : the official*  
550 *journal of the International Society for Developmental Neuroscience.* 2011;29(1):57-62.
- 551 23.Vanderloo LM. Screen-viewing among preschoolers in childcare: a systematic review. *BMC*  
552 *pediatrics.* 2014;14:205.
- 553 24.Ward S, Belanger M, Donovan D, Carrier N. Systematic review of the relationship between  
554 childcare educators' practices and preschoolers' physical activity and eating behaviours. *Obes*  
555 *Rev.* 2015.
- 556 25.Fukuda K, Sakashita Y. Sleeping pattern of kindergartners and nursery school children: function of  
557 daytime nap. *Percept Mot Skills.* 2002;94(1):219-28.
- 558 26.Kim J, Peterson KE. Association of infant child care with infant feeding practices and weight gain  
559 among US infants. *Arch Pediatr Adolesc Med.* 2008;162(7):627-33.
- 560 27.Tandon PS, Saelens BE, Christakis DA. Active play opportunities at child care. *Pediatrics.*  
561 2015;135(6):e1425-31.
- 562 28.Bingham DD, Costa S, Hinkley T, Shire KA, Clemes SA, Barber SE. Physical Activity During the  
563 Early Years: A Systematic Review of Correlates and Determinants. *Am J Prev Med.*  
564 2016;51(3):384-402.

- 565 29. Vanderloo LM, Tucker P, Johnson AM, Burke SM, Irwin JD. Environmental influences on  
566 preschoolers' physical activity levels in various early-learning facilities. *Res Q Exerc Sport*.  
567 2015;86(4):360-70.
- 568 30. Cerin E, Baranowski T, Barnett A, Butte N, Hughes S, Lee RE, et al. Places where preschoolers  
569 are (in)active: an observational study on Latino preschoolers and their parents using objective  
570 measures. *The international journal of behavioral nutrition and physical activity*. 2016;13(1):29.
- 571 31. Peden ME, Jones R, Costa S, Ellis Y, Okely AD. Relationship between children's physical activity,  
572 sedentary behavior, and childcare environments: A cross sectional study. *Prev Med Rep*.  
573 2017;6:171-6.
- 574 32. Plancoulaine S, Lioret S, Regnault N, Heude B, Charles MA. Gender-specific factors associated  
575 with shorter sleep duration at age 3 years. *Journal of sleep research*. 2015;24(6):610-20.
- 576 33. Costa S, Adams J, Benjamin Neelon SE. The relationship between childcare and adiposity, body  
577 mass, and obesity-related risk factors: a systematic review of longitudinal studies. PROSPERO;  
578 2015. Report No.: CRD42015027233 Contract No.: CRD42015027233.
- 579 34. Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items  
580 for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation.  
581 *BMJ*. 2015;349.
- 582 35. Costa S, Adams J, Phillips V, Benjamin Neelon SE. The relationship between childcare and  
583 adiposity, body mass and obesity-related risk factors: protocol for a systematic review of  
584 longitudinal studies. *Syst Rev*. 2016;5:141.
- 585 36. Lakshman R, Mazarello Paes V, Hesketh K, O'Malley C, Moore H, Ong K, et al. Protocol for  
586 systematic reviews of determinants/correlates of obesity-related dietary and physical activity  
587 behaviors in young children (preschool 0 to 6 years): evidence mapping and syntheses. *Syst Rev*.  
588 2013;2.
- 589 37. The World Bank. World Bank Country and Lending Groups – World Bank Data Help Desk  
590 Washington, DC, USA2017 [Available from:  
591 [https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-](https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups)  
592 [lending-groups](https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups).  
593 38. Obbagy JE, Lyon JMG, Spahn JM. USDA Nutrition Evidence Library Systematic Reviews: using  
594 food and nutrition research to inform nutrition programs and policies. *The Digest*. 2014;49(2):1-7.

- 595 39. Belfield CR, Kelly IR. Early education and health outcomes of a 2001 U.S. Birth Cohort. *Econ Hum*  
596 *Biol.* 2013;11(3):310-25.
- 597 40. Cairns A, Harsh J. Changes in Sleep Duration, Timing, and Quality as Children Transition to  
598 Kindergarten. *Behav Sleep Med.* 2014;12(6):507-16.
- 599 41. D'Onise K, Lynch JW, McDermott RA, Esterman A. The beneficial effects of preschool attendance  
600 on adult cardiovascular disease risk. *Aust N Z J Public Health.* 2011;35(3):278-83.
- 601 42. Lumeng JC, Gannon K, Appugliese D, Cabral HJ, Zuckerman B. Preschool child care and risk of  
602 overweight in 6- to 12-year-old children. *Int J Obes (2005).* 2005;29(1):60-6.
- 603 43. Lumeng JC, Rahnama S, Appugliese D, Kaciroti N, Bradley RH. Television exposure and  
604 overweight risk in preschoolers. *Arch Pediatr Adolesc Med.* 2006;160(4):417-22.
- 605 44. Taylor RW, Murdoch L, Carter P, Gerrard DF, Williams SM, Taylor BJ. Longitudinal study of  
606 physical activity and inactivity in preschoolers: the FLAME study. *Medicine and science in sports*  
607 *and exercise.* 2009;41(1):96-102.
- 608 45. Camara S, de Lauzon-Guillain B, Heude B, Charles MA, Botton J, Planoulaine S, et al.  
609 Multidimensionality of the relationship between social status and dietary patterns in early  
610 childhood: longitudinal results from the French EDEN mother-child cohort. *Int J Behav Nutr Phys*  
611 *Activ.* 2015;12:122.
- 612 46. Lee R, Zhai F, Han WJ, Brooks-Gunn J, Waldfogel J. "Head Start and Children's Nutrition, Weight,  
613 and Health Care Receipt". *Early childhood research quarterly.* 2013;28(4).
- 614 47. Levy SM, Slager SL, Warren JJ, Levy BT, Nowak AJ. Associations of pacifier use, digit sucking,  
615 and child care attendance with cessation of breastfeeding. *J Fam Pract.* 2002;51(5):465.
- 616 48. Pearce A, Li L, Abbas J, Ferguson B, Graham H, Law C. Childcare use and inequalities in  
617 breastfeeding: findings from the UK Millennium Cohort Study. *Arch Dis Child.* 2012;97(1):39-42.
- 618 49. Sata M, Yamagishi K, Sairenchi T, Ikeda A, Irie F, Watanabe H, et al. Impact of Caregiver Type for  
619 3-Year-Old Children on Subsequent Between-Meal Eating Habits and Being Overweight From  
620 Childhood to Adulthood: A 20-Year Follow-up of the Ibaraki Children's Cohort (IBACHIL) Study. *J*  
621 *Epidemiol.* 2015;25(9):600-7.
- 622 50. Wasser HM, Thompson AL, Siega-Riz AM, Adair LS, Hodges EA, Bentley ME. Who's feeding  
623 baby? Non-maternal involvement in feeding and its association with dietary intakes among infants  
624 and toddlers. *Appetite.* 2013;71:7-15.

- 625 51. Weile B, Rubin DH, Krasilnikoff PA, Kuo HS, Jekel JF. Infant feeding patterns during the first year  
626 of life in Denmark: factors associated with the discontinuation of breast-feeding. *J Clin Epidemiol.*  
627 1990;43(12):1305-11.
- 628 52. Rajulton F. The fundamentals of longitudinal research: an overview. *Can Stud Popul.*  
629 2001;28(2):169-85.
- 630 53. Kagamimori S, Yamagami T, Sokejima S, Numata N, Handa K, Nanri S, et al. The relationship  
631 between lifestyle, social characteristics and obesity in 3-year-old Japanese children. *Child Care*  
632 *Health Dev.* 2008;25(3):235-48.
- 633 54. Nevarez MD, Rifas-Shiman SL, Kleinman KP, Gillman MW, Taveras EM. Associations of Early Life  
634 Risk Factors With Infant Sleep Duration. *Acad Pediatr.* 2010;10(3):187-93.
- 635

Table 1 - Description of included studies.

Study (date)	Location	Sample size	Study design	Sex	Ethnicity/country of birth	Outcome
Belfield & Kelly (2013)	USA	6550	Prospective longitudinal cohort study (Early Childhood Longitudinal Survey – Birth Cohort)	<u>Preschool</u> : 4124 (50.6%) girls, 4026 (49.4%) boys <u>Kindergarten</u> : 3301 (50.4%) girls, 3249 (49.6%) boys	<u>Preschool</u> : 1231 (15.1%) Black, 1157 (14.2%) Hispanic, 864 (10.6%) Asian, 1410 (17.3%) Other non-White, 3488 (42.8%) White <u>Kindergarten</u> : 1009 (15.4%) Black, 943 (14.4%) Hispanic, 766 (11.7%) Asian, 1212 (18.5%) Other non-White, 2620 (40.0%) White	Physical activity, Diet
Cairns & Harsh (2014)	USA	34	Prospective longitudinal study	15 (44.1%) girls, 19 (55.9%) boys	62% White, 32% Black, 6% Other	Sleep
D'Onise et al (2011)	Australia	1063	Prospective longitudinal cohort study (North West Adelaide Health Study)	580 (54.6%) girls, 483 (45.4%) boys	Not reported	Physical activity, Sedentary behaviour
Lumeng et al (2005)	USA	1244	Longitudinal panel survey study	630 (50.6%) girls, 614 (49.4%) boys	488 (39.2%) Black, 69 (5.5%) Hispanic, 623 (50.1%) White, 64 (5.1%) Other	Sedentary behaviour
Lumeng et al (2006)	USA	1016	Prospective longitudinal study	498 (49.0%) girls, 518 (51.0%) boys	841 (82.8%) White, 175 (17.2%) not White	Sedentary behaviour
Camara et al (2015)	France	974	Prospective longitudinal cohort study (EDEN - Etude des Déterminants pré et post natus du développement et de la santé de l'ENfant)	454 (46.6%) girls, 520 (53.4%) boys	France birth (ethnic composition not presented)	Diet
Levy et al (2002)	USA	1387	Prospective longitudinal cohort study (Iowa Fluoride Study)	Not reported	95% White, 5% Other	Diet
Pearce et al (2012)	UK	18050	Prospective longitudinal cohort study (Millennium Cohort Study)	Not reported	UK birth (ethnic composition not presented)	Diet



Sata et al (2015)	Japan	4281	Prospective longitudinal cohort study (Ibaraki Children's Cohort (IBACHIL) Study)	2042 (47.7%) girls, 2239 (52.3%) boys	Japan birth (ethnic composition not presented)	Diet
Wasser et al (2013)	USA	210	Prospective longitudinal study	116 (53.5%) girls, 101 (46.5%) boys	African-American	Diet
Lee et al (2013)	USA	2150	Prospective longitudinal cohort study (Early Childhood Longitudinal Study-Birth Cohort (ECLS-B))	49% girls, 51% boys (at birth)	USA birth (ethnic composition not presented)	Diet
Weile et al (1990)	Denmark	500	Prospective longitudinal study	Not reported	Denmark birth (ethnic composition not presented)	Diet
Taylor et al (2009)	New Zealand	<u>3 years:</u> 238 <u>4 years:</u> 216 <u>5 years:</u> 204	Prospective longitudinal cohort study (The Family Lifestyle, Activity, Movement, and Eating study)	<u>3 years:</u> 107 (43.9%) girls, 137 (56.1%) boys. <u>4 years:</u> 104 (43.9%) girls, 133 (56.1%) boys. <u>5 years:</u> 99 (44%) girls, 126 (56%) boys	<u>Baseline:</u> 87% Caucasian, 10.8% Maori, 3.7% Pacific Islanders	Physical activity

Legend: BMI – body mass index; SD – standard deviation. \* Numbers are approximate, calculated from the percentages and total preschool and kindergarten sample sizes presented in article's appendix table A2.

Table 2 – Results of included studies.

Study	Exposure	Age at childcare exposure	Outcome	Age at outcome	Analysis	Adjustment	Results (most adjusted model)
Belfield & Kelly (2013)	Center-based preschool	4 y	Low activity level	5-6 y	Multivariable probit regression models	Child's race/ethnicity, gender, age, and number of siblings, twin (yes/no), maternal employment, education, and marital status, health insurance status, father non-resident, household income, geographic region, and prior health at 24 mo (general health status, and indicators of asthma, gastroenteritis, respiratory condition, and ear infection)	AME=0.124 (SE: 0.120), p>0.05
	Center-based preschool	4 y	Screened low activity	5-6 y	Multivariable probit regression models		AME=0.056 (SE: 0.084), p>0.05
	Center-based preschool	4 y	Regular fast food consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=-0.15 (SE: 0.055), p>0.05
	Center-based preschool	4 y	Regular Soda consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=-0.117 (SE: 0.056), p<0.05
	Center-based preschool	4 y	Regular Candy consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=0.031 (SE: 0.053), p>0.05
	Center-based preschool	4 y	Regular Chips consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=0.113 (SE: 0.054), p<0.05
	Center-based preschool	4 y	Infrequent Vegetable consumption (vs. frequent)	5-6 y	Multivariable probit regression models		AME=0.049 (SE: 0.058), p>0.05
	Center-based preschool	4 y	Infrequent Fruit consumption (vs. frequent)	5-6 y	Multivariable probit regression models		AME=-0.120 (SE: 0.060), p<0.05
	Center-based preschool for 2 y	4 y	Low activity level	5-6 y	Multivariable probit regression models		AME=0.007 (SE: 0.173), p>0.05
	Center-based preschool for 2 y	4 y	Screened low activity	5-6 y	Multivariable probit regression models		AME=0.058 (SE: 0.126), p>0.05
	Center-based preschool for 2 y	4 y	Regular fast food consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=0.022 (SE: 0.085), p>0.05
	Center-based preschool for 2 y	4 y	Regular Soda consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=-0.260 (SE: 0.083), p<0.01
	Center-based preschool for 2 y	4 y	Regular Candy consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=-0.031 (SE: 0.080), p>0.05
	Center-based preschool for 2 y	4 y	Regular Chips consumption (vs. not)	5-6 y	Multivariable probit regression models		AME=0.024 (SE: 0.081), p>0.05
	Center-based preschool for 2 y	4 y	Infrequent Vegetable consumption (vs. frequent)	5-6 y	Multivariable probit regression models		AME=-0.113 (SE: 0.090), p>0.05
Center-based preschool for 2 y	4 y	Infrequent Fruit consumption (vs. frequent)	5-6 y	Multivariable probit regression models	AME=-0.231 (SE: 0.093), p<0.01		

Head Start	4 y	Low activity level	5-6 y	Multivariable probit regression models	AME=0.313 (SE: 0.142), p<0.05
Head Start	4 y	Screened low activity	5-6 y	Multivariable probit regression models	AME=0.128 (SE: 0.112), p>0.05
Head Start	4 y	Regular fast food consumption (vs. not)	5-6 y	Multivariable probit regression models	AME=0.050 (SE: 0.077), p>0.05
Head Start	4 y	Regular Soda consumption (vs. not)	5-6 y	Multivariable probit regression models	AME=-0.065 (SE: 0.081), p>0.05
Head Start	4 y	Regular Candy consumption (vs. not)	5-6 y	Multivariable probit regression models	AME=-0.108 (SE: 0.073), p>0.05
Head Start	4 y	Regular Chips consumption (vs. not)	5-6 y	Multivariable probit regression models	AME=0.040 (SE: 0.074), p>0.05
Head Start	4 y	Infrequent Vegetable consumption (vs. frequent)	5-6 y	Multivariable probit regression models	AME=-0.067 (SE: 0.083), p>0.05
Head Start	4 y	Infrequent Fruit consumption (vs. frequent)	5-6 y	Multivariable probit regression models	AME=-0.266 (SE: 0.085), p<0.01

Cairns & Harsh (2014)	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Total sleep duration weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA	None reported	Group: not significant Time: F(2,64)=5.2, p=0.008, $\eta^2=0.14$ Group by time: not significant
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Nocturnal sleep duration weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not significant Time: not significant Group by time: not significant
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Sleep onset weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: F(1,32)=5.8, p=0.022, $\eta^2=0.15$ Time: F(2,64)=40.9, p<.001, $\eta^2=0.56$ Group by Time: F(2,64)=6.1, p=0.004, $\eta^2=0.16$
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Sleep onset time weekend	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: F(2,64)=6.9, p<0.01, $\eta^2=0.18$ Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Wake up time weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: F(1,32)=14.9, p=0.001, $\eta^2=0.32$ Time: F(2,64)=81.6, p<0.001, $\eta^2=0.72$ Group by time: F(2,64)=17.5, p<0.001, $\eta^2=0.35$
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Wake up time weekend	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: F(2,64)=4.4, p<0.05, $\eta^2=0.12$ Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Sleep efficiency weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not significant Time: F(2,64)=3.5, p<0.05, $\eta^2=0.10$ Group by time: not significant

	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Nap duration weekday	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: F(2,55)=20.46, p<0.001, η <sup>2</sup> =0.436 Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Nap duration weekend	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time= not significant Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Number of weekdays with a nap	5 y <sup>b</sup>	T-test (Summer vs 2 weeks after start of kindergarten)		Group: not reported Time: T(13)=3.4, p=0.005 Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Number of weekday naps	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not significant Time: not significant Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Number of weekend naps	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not significant Time: not significant Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Caregivers rating children as having less difficulty in going to bed	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: F(2,55)=20.46, p<0.001, η <sup>2</sup> =0.436 Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Caregivers rating children as having less difficulty falling asleep	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: F(2,42)=3.9, p=0.03, η <sup>2</sup> =0.16 Group by time: not reported
	All day preschool/ daycare (vs primary/secondary caregiver)	5 y <sup>a</sup>	Caregivers ratings of returning to wakefulness in the morning	5 y <sup>b</sup>	Group by assessment mixed model ANOVA		Group: not reported Time: not reported Group by time: F(2,42) = 6.3, p=0.004, η <sup>2</sup> =0.23
Camara et al (2015)	Childcare arrangement	2-3 y	Processed, fast-foods at 2, 3 and 5 y dietary pattern	2 y, 3 y, 5 y	Multivariable linear regression	Child's age, gender, recruitment centre, season when the food frequency questionnaire was completed household disadvantage composite index, older sibling at home (2 y), maternal age at delivery, education level, and current/ past occupation, working time, and unemployed/student when child aged 2 y	At home, cared for by mother: Reference At home, not cared for by mother: B= 0.70 (SE: 0.14), p<0.001 Crèche/pre-school: B= -0.03 (SE: 0.13), p>0.05 At nanny's home: B=0.13 (SE: 0.13), p>0.05
	Childcare arrangement	2-3 y	Guidelines at 2, 3 and 5 y dietary pattern	2 y, 3 y, 5 y	Multivariable linear regression	Child's age, gender, recruitment centre, season when the food frequency questionnaire was completed household disadvantage composite index, older sibling at home (2 y), maternal age at delivery, education level, and current/ past occupation, working time, and unemployed/student when child aged 2 y	At home, cared for by mother: Reference At home, not cared for by mother: B= 0.01 (SE:0.15), p>0.05 Crèche/pre-school: B= 0.08 (SE: 0.13), p>0.05 At nanny's home: B= 0.10 (SE: 0.13), p>0.05
D'Onise et al (2011)	Attended Kindergarten Union preschool (vs not attended)	2-5 y	PA level	Preschool mean: 45.3y No Preschool mean= 51.1 y	Multinomial logistic regression	Age, gender, child socioeconomic position, adult height, educational attainment, and adult income	Sedentary: Reference Low PA: RRR= 1.24 (95%CI: 0.89-1.74) Moderate PA: RRR=1.26 (95%CI: 0.87-1.81) High PA: RRR=1.99 (95%CI: 1.19-3.35)
Lee et al (2013)	Type of childcare arrangement on a regular basis - Head Start vs. not Head Start	4 y	Frequency of having healthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions	Child's variables (e.g., gender, ethnicity, multiple birth, prematurity, breastfeeding and number of siblings at 9 mo,	M= 2.21 (SE:0.74), p<0.01

Type of childcare arrangement on a regular basis - Head Start vs. not Head Start	4 y	Frequency of having unhealthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions	pre-treatment outcomes at 2y); <u>Maternal variables</u> (e.g., married at birth (yes/no), pre-pregnancy age and body mass index, depression at 9 mo, health status and employment at 2 y, foreign born); Parenting behaviours/ home environments (e.g., KIDI at 9 mo, having sleep routine, weekday watching TV, and indoor and outdoor activities at 2 y);	M=0.63 (SE:0.57), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. Pre-Kindergarten	4 y	Frequency of having healthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 1.26 (SE: 1.33), p>0.05
Type of childcare arrangement on a regular basis - Head Start vs. Pre-Kindergarten	4 y	Frequency of having unhealthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 0.36 (SE:0.97), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. Other centre-based	4 y	Frequency of having healthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions	<u>Family variables</u> (e.g., parent's education at birth, parental occupation and family income at 9 mo, living in urban area, region of country, and number of times receiving Special Supplemental Nutrition Program for Women, Infants, and Children, food stamps, and Temporary Assistance for Needy Families by 2 y)	M= 2.35 (SE: 1.14), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. Other centre-based	4 y	Frequency of having unhealthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 0.80 (SE: 0.78), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. Other non-parental	4 y	Frequency of having healthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 2.74 (SE: 1.32), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. Other non-parental	4 y	Frequency of having unhealthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 0.77 (SE:0.98), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. parental	4 y	Frequency of having healthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 2.07 (SE: 1.01), p<0.05
Type of childcare arrangement on a regular basis - Head Start vs. parental	4 y	Frequency of having unhealthy eating patterns (times/week)	5-6 y	Propensity-score weighted regressions		M= 0.47 (SE: 0.77), p<0.05

Levy et al (2002)	Number of days in childcare between 0-6 mo of age	6 weeks, 3 mo, 6 mo (referring to preceding time period)	Time until cessation of breastfeeding	6 weeks, 3 mo, 6 mo	Cox proportional hazard regression	Pacifier use, digit sucking, maternal and paternal age and education, family income, breastfeeding plans, maternal smoking, infant's gender, and infant antibiotic use.	No pacifier use, or digit sucking, or childcare: Reference No pacifier use, does digit sucking, no childcare days: p≥0.05 No pacifier use, does digit sucking, 15 childcare days: HR=1.41 (95%CI: 1.02-1.96), p<0.05 No pacifier use, does digit sucking, 30 childcare days: p≥0.05 No pacifier use, does digit sucking, 60 childcare days: p≥0.05 Pacifier use, no digit sucking, no childcare: HR=1.67 (95%CI: not reported#), p<0.05 Pacifier use, no digit sucking, 15 days childcare: p≥0.05 Pacifier use, no digit sucking, 30 days childcare: significant # Pacifier use, no digit sucking, 60 days childcare: borderline significant # Pacifier use and digit sucking, no childcare: HR=1.88 (95%CI: 1.36-2.62), p<0.05 Pacifier use and digit sucking, 15 childcare days: HR=1.52 (95%CI: 1.03-2.25), p<0.05 Pacifier use and digit sucking, 30 childcare days: p≥0.05 Pacifier use and digit sucking, 60 childcare days: not significant
Lumeng et al (2005)	Center-based childcare attendance intensity (None vs 15h/week vs ≥15h/week)	3-5 y	>4 h/day of TV viewing (yes vs no)	6-12 y	Turkey's test	None	Not significant, p=0.27
Lumeng et al (2006)	Average number of hours in non-parental childcare	24-36 mo	TV viewing (<2h/day vs ≥2h/day)	36 mo	T-test	None	Not significant, p=0.58
Pearce et al (2012)	Overall childcare type	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression	Mother's ethnicity, parity, age at first live birth, and whether the mother returned to work before the infant was age 4 mo	Parent: Reference Informal: RR= 0.51 (95%CI: 0.43–0.59), p<0.05 Formal: RR= 0.84 (95%CI: 0.72–0.99), p<0.05
	Childcare type by intensity	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression		Parent: Reference Informal Part-time: RR= 0.54 (95%CI: 0.45-0.63), p<0.05 Informal Full-time: RR= 0.42 (95%CI: 0.28-0.64), p<0.05 Formal Part-time: RR= 1.01 (95%CI: 0.82-1.24), p≥0.05 Formal Full-time: RR= 0.68 (95%CI: 0.51-0.92), p<0.05

Childcare type by National Statistics Socio-economic Classification	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression	<p><u>Routine and Manual</u>            Parent: Reference            Informal: RR= 0.47 (95%CI: 0.34-0.66), p&lt;0.05            Formal: RR= 0.54 (95%CI: 0.21-1.36), p≥0.05</p> <p><u>Intermediate</u>            Parent: Reference            Informal: RR= 0.50 (95%CI: 0.37-0.67), p&lt;0.05            Formal: RR= 0.84 (95%CI: 0.57-1.23), p≥0.05</p> <p><u>Managerial and Professional</u>            Parent: Reference            Informal: RR= 0.50 (95%CI: 0.39-0.65), p&lt;0.05            Formal: RR= 0.76 (95%CI: 0.62-0.94), p&lt;0.05</p>
Childcare type by Maternal Education	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression	<p><u>None – GCSE D–G</u>            Parent: Reference            Informal: RR= 0.44 (95%CI: 0.27-0.71), p&lt;0.05            Formal: RR= 1.00 (95%CI: 0.44-2.28), p≥0.05</p> <p><u>GCSE A–C, A Levels, Diploma</u>            Parent: Reference            Informal: RR= 0.47 (95% CI: 0.37-0.59), p≥0.05            Formal: RR= 0.83 (95%CI: 0.64-1.08), p≥0.05</p> <p><u>Degree</u>            Parent: Reference            Informal: RR= 0.82 (95%CI: 0.64-1.06), p≥0.05            Formal: RR= 0.71 (95%CI: 0.58-0.86), p&lt;0.05</p>
Childcare type by Lone Parenthood	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression	<p><u>Lone parent</u>            Parent: Reference            Informal: RR= 0.40 (95%CI: 0.25-0.65), p&lt;0.05            Formal: RR= 1.65 (95%CI: 1.04-2.63), p&lt;0.05</p> <p><u>Couple family</u>            Parent: Reference            Informal: RR= 0.53 (95%CI: 0.44-0.63), p&lt;0.05            Formal: RR= 0.79 (95%CI: 0.66-0.94), p&lt;0.05</p>

Childcare type by Area Deprivation	<4 to 9 mo	Breastfeeding for ≥4 mo	9 mo	Poisson regression		<p><u>5 (Most deprived)</u>            Parent: Reference            Informal: RR= 0.72 (95%CI: 0.53-0.97), p&lt;0.05            Formal: RR= 0.63 (95% CI: 0.28-1.39), p≥0.05</p> <p><u>4</u>            Parent: Reference            Informal: RR= 0.54 (95%CI: 0.36-0.81), p&lt;0.05            Formal: RR= 1.12 (95% CI: 0.73-1.72), p≥0.05</p> <p><u>3</u>            Parent: Reference            Informal: RR= 0.51 (95%CI: 0.33-0.80), p&lt;0.05            Formal: RR= 1.27 (95% CI: 0.86-1.85), p≥0.05</p> <p><u>2</u>            Parent: Reference            Informal: RR= 0.37 (95%CI: 0.21-0.65), p&lt;0.05            Formal: RR= 0.71 (95% CI: 0.47-1.06), p≥0.05</p> <p><u>1 (Least deprived)</u>            Parent: Reference            Informal: RR= 0.48 (95%CI: 0.26-0.88), p&lt;0.05            Formal: RR= 0.64 (95% CI: 0.42-1.00), p≥0.05</p>	
Sata et al (2015)	Main daytime caregiver	3 y	Between-meal eating before dinner	6 y	Logistic regression models, stratified by gender	Baseline types of feeding, wake-up time, time of sleep, physical activity, playing outside, living with brothers or sisters, picky eating, and father's employment.	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 2.1 (95%CI: 1.4-3.1), p&lt;0.001            Nursery school / kindergarten staff: OR=1.6 (95%CI: 1.1-2.4), p&lt;0.05</p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 2.5 (95%CI: 1.7-3.8), p&lt;0.001            Nursery school / kindergarten staff: OR=1.6 (95%CI: 1.1-2.4), p&lt;0.05</p>
	Main daytime caregiver	3 y	Between-meal eating ≥3 times/day	6 y	Logistic regression models, stratified by gender	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 3.2 (95%CI: 1.3-7.7), p&lt;0.05            Nursery school / kindergarten staff: OR=1.9 (95%CI: 0.7-5.4), p≥0.05</p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 2.7 (95%CI: 1.1-6.7), p&lt;0.05            Nursery school / kindergarten staff: OR=2.3 (95%CI: 0.9-6.3), p≥0.05</p>	



Main daytime caregiver	3 y	Between-meal eating before bedtime $\geq 3$ times/week	6 y	Logistic regression models, stratified by gender	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 1.5 (95%CI: 0.8-2.7), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=1.1 (95% CI: 0.6-2.0), <math>p \geq 0.05</math></p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 1.4 (95%CI: 0.7-2.5), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=1.6 (95% CI: 0.8-3.0), <math>p \geq 0.05</math></p>
Main daytime caregiver	3 y	Between-meal eating before dinner	12 y	Logistic regression models, stratified by gender	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 1.3 (95%CI: 0.9-1.8), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=1.0 (95%CI: 0.7-1.5), <math>p \geq 0.05</math></p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 1.9 (95%CI: 1.3-2.8), <math>p &lt; 0.01</math>            Nursery school / kindergarten staff: OR=1.7 (95%CI: 1.1-2.5), <math>p &lt; 0.05</math></p>
Main daytime caregiver	3 y	Between-meal eating $\geq 5$ times/week	12 y	Logistic regression models, stratified by gender	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 1.0 (95%CI: 0.7-1.4), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=1.2 (95%CI: 0.8-1.7), <math>p \geq 0.05</math></p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 0.9 (95%CI: 0.6-1.3), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=0.9 (95%CI: 0.6-1.3), <math>p \geq 0.05</math></p>
Main daytime caregiver	3 y	Between-meal eating before bedtime $\geq 3$ times/week	12 y	Logistic regression models, stratified by gender	<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 1.5 (95%CI: 0.9-2.5), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=0.7 (95%CI: 0.4-1.3), <math>p \geq 0.05</math></p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 1.1 (95%CI: 0.6-2.2), <math>p \geq 0.05</math>            Nursery school / kindergarten staff: OR=1.2 (95%CI: 0.6-2.5), <math>p \geq 0.05</math></p>

	Main daytime caregiver	3 y	Between-meal eating before dinner	22 y	Logistic regression models, stratified by gender		<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 0.9 (95%CI: 0.6-1.5), p≥0.05            Nursery school / kindergarten staff: OR=1.2 (95%CI: 0.8-1.9), p≥0.05</p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 1.2 (95%CI: 0.7-2.0), p≥0.05            Nursery school / kindergarten staff: OR=0.9 (95%CI: 0.5-1.5), p≥0.05</p>
	Main daytime caregiver	3 y	Between-meal eating ≥5 times/week	22 y	Logistic regression models, stratified by gender		<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 0.9 (95%CI: 0.5-1.5)            Nursery school / kindergarten staff: OR=1.0 (95%CI: 0.6-1.6)</p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 0.8 (95%CI: 0.5-1.3)            Nursery school / kindergarten staff: OR=1.1 (95%CI: 0.7-1.8)</p>
	Main daytime caregiver	3 y	Between-meal eating before bedtime ≥3 times/week	22 y	Logistic regression models, stratified by gender		<p><u>Boys:</u>            Mothers: Reference            Grandparents: OR= 1.0 (95%CI: 0.6-1.7), p≥0.05            Nursery school / kindergarten staff: OR=0.6 (95%CI: 0.3-1.1), p≥0.05</p> <p><u>Girls:</u>            Mothers: Reference            Grandparents: OR= 1.3 (95%CI: 0.6-2.6), p≥0.05            Nursery school / kindergarten staff: OR=1.3 (95%CI: 0.6-2.8), p≥0.05</p>
Taylor et al (2009)	Number of hours per week childcare attendance	3 y, 4 y, 5 y	Total active time (minutes/day)	3 y, 4 y, 5 y	Random coefficient regression	None reported	Not significant, p=0.069–0.806
	Number of hours per week childcare attendance	3 y, 4 y, 5 y	Average accelerometer counts (counts/minute)	3 y, 4 y, 5 y	Random coefficient regression	None reported	Not significant, p=0.069–0.806
Wasser et al (2013)	Any non-maternal caregiver use	6-18 mo	Consuming any whole fruit	6-18 mo	Random-effects logistic regression	Maternal age, employment, depression, any maternal college, and marital status	None: Reference Any: OR= 1.51 (95%CI: 1.03-2.23), p<0.05
	Any non-maternal caregiver use	6-18 mo	Consuming any vegetable	6-18 mo	Random-effects logistic regression		None: Reference Any: OR= 1.25 (95%CI: 0.79-1.99), p≥0.05
	Any non-maternal caregiver use	6-18 mo	Consuming any juice	6-18 mo	Random-effects logistic regression		None: Reference Any: OR= 1.64 (95%CI: 1.01-2.67), p<0.05

Any non-maternal caregiver use	6-18 mo	Consuming any fried potatoes	6-18 mo	Random-effects logistic regression	None: Reference Any: OR= 0.82 (95%CI: 0.46-1.43), $p \geq 0.05$
Any non-maternal caregiver use	6-18 mo	Consuming any desserts and sweets	6-18 mo	Random-effects logistic regression	None: Reference Any: OR= 1.20 (95%CI: 0.77-1.86), $p \geq 0.05$
Any non-maternal caregiver use	6-18 mo	Consuming any sweetened beverages	6-18 mo	Random-effects logistic regression	None: Reference Any: OR= 1.17 (95%CI: 0.65-2.12), $p \geq 0.05$
Any non-maternal caregiver use	6-18 mo	Consuming any salty snacks	6-18 mo	Random-effects logistic regression	None: Reference Any: OR= 1.45 (95%CI: 0.67-3.12), $p \geq 0.05$
Type of non-maternal caregiver use	6-18 mo	Consuming any whole fruit	6-18 mo	Random-effects logistic regression	None: Reference Father: OR= 1.12 (95%CI: 0.64 -1.97), $p \geq 0.05$ Grandmother: OR= 0.92 (95%CI: 0.57-1.5), $p \geq 0.05$ Licensed provider: OR= 1.55 (95%CI: 0.93-2.59), $p \geq 0.05$
Type of non-maternal caregiver use	6-18 mo	Consuming any vegetable	6-18 mo	Random-effects logistic regression	None: Reference Father: OR= 0.93 (95%CI: 0.48-1.8), $p \geq 0.05$ Grandmother: OR= 0.89 (95%CI: 0.5-1.59), $p \geq 0.05$ Licensed provider: OR= 0.96 (95%CI: 0.52-1.79), $p \geq 0.05$
Type of non-maternal caregiver use	6-18 mo	Consuming any juice	6-18 mo	Random-effects logistic regression	None: Reference Father: OR= 0.83 (95%CI: 0.42-1.64), $p \geq 0.05$ Grandmother: OR= 1.97 (95%CI: 1.02-3.81), $p < 0.05$ Licensed provider: OR= 1.2 (95%CI: 0.61-2.34), $p \geq 0.05$
Type of non-maternal caregiver use	6-18 mo	Consuming any fried potatoes	6-18 mo	Random-effects logistic regression	None: Reference Father: OR= 1.13 (95%CI: 0.48-2.69), $p \geq 0.05$ Grandmother: OR= 0.97 (95%CI: 0.48-1.96), $p \geq 0.05$ Licensed provider: OR= 0.75 (95%CI: 0.38-1.48), $p \geq 0.05$
Type of non-maternal caregiver use	6-18 mo	Consuming any desserts and sweets	6-18 mo	Random-effects logistic regression	None: Reference Father: OR= 0.85 (95%CI: 0.44-1.67), $p \geq 0.05$ Grandmother: OR= 0.74 (95%CI: 0.42-1.28), $p \geq 0.05$ Licensed provider: OR= 1.30 (95%CI: 0.75-2.26), $p \geq 0.05$

	Type of non-maternal caregiver use	6-18 mo	Consuming any sweetened beverages	6-18 mo	Random-effects logistic regression		None: Reference Father: OR= 1.71 (95%CI: 0.71-4.11), p≥0.05 Grandmother: OR= 0.97 (95%CI: 0.46-2.05), p≥0.05 Licensed provider: OR= 1.28 (95%CI: 0.63-2.62), p≥0.05
	Type of non-maternal caregiver use	6-18 mo	Consuming any salty snacks	6-18 mo	Random-effects logistic regression		None: Reference Father: OR= 2.06 (95%CI: 0.66-6.39), p≥0.05 Grandmother: OR= 1.03 (95%CI: 0.40-2.69), p≥0.05 Licensed provider: OR= 0.71 (95%CI: 0.28-1.79), p≥0.05
Weile et al (1990)	Attending daycare (vs. cared for at home)	1-12 mo	Changing from feeding categories 1/2 to categories 3/4/5*	1-12 mo	Cox proportional hazards model	Other children in family and socioeconomic status	RR= 2.08 (95%CI: 1.43-3.01), p<0.05
	Attending daycare (vs. cared for at home)	1-12 mo	Changing from feeding categories 1/2/3 to categories 4/5*	1-12 mo	Cox proportional hazards model	Other children in family and socioeconomic status	RR= 2.05 (95%CI: 1.39-3.02), p<0.05
	Attending daycare (vs. cared for at home)	1-12 mo	Changing from feeding categories 1/2/3/4 to category 5*	1-12 mo	Cox proportional hazards model	Other children in family and socioeconomic status	RR= 2.50 (95%CI: 1.66-3.78), p<0.05

Legend: AME - average marginal effects; CI - confidence interval; h - hours; m - months; OR- odds ratio; PA - physical activity; RR - relative risk; RRR - relative risk ratio; SD - standard deviation; SE - robust standard errors; TV – television; y - years. weeks before start of kindergarten;<sup>b</sup> 2 weeks, 1 month after start of kindergarten; \* Categories: (1) 100% breast-fed, (2) breast-fed>formula-fed, (3) breast-fed=formula-fed, (4) breast-fed<formula-fed, and (5) 100% formula-fed. # Estimates and significance figures taken from text only, as article did not present tables and we could not obtain these from publishers or authors.

Table 3 – Results of the Nutrition Evidence Library Bias Assessment Tool risk of bias evaluation.

NEL-BAT question	Belfield & Kelly (2013)	Cairns & Harsh (2014)	Camara et al (2015)	D'Onise et al (2011)	Lee et al (2013)	Levy et al (2002)	Lumeng et al (2005)	Lumeng et al (2006)	Pearce et al (2012)	Stata et al (2015)	Taylor et al (2009)	Wasser et al (2013)	Weile et al (1990)	Total score by question
1. Were the inclusion/exclusion criteria similar across study groups?	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2. Was the strategy for recruiting or allocating participants similar across study groups?	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5. Was there an attempt to balance the allocation between the study groups or match the study groups (e.g., through stratification, matching, propensity scores)?	0	2	1	0	0	1	2	0	1	2	0	1	0	10
6. Was distribution of health status, demographics, and other critical confounding factors similar across study groups at baseline? If not, does the analysis control for baseline differences between groups?	0	1	0	0	0	1	2	2	1	2	0	1	1	11
7. Did the investigators account for important variations in the execution of the study from the proposed protocol or research plan?	0	0	0	0	0	0	1	0	0	0	1	0	0	2
8. Was adherence to the study protocol similar across study groups?	0	0	1	0	1	1	0	0	1	1	0	1	1	7
9. Did the investigators account for the impact of unintended/unplanned concurrent interventions or exposures that were differentially experienced by study groups and might bias results?	0	0	1	0	1	1	0	0	1	1	0	1	0	6
12. Were outcome assessors blinded to the intervention or exposure status of participants?	0	1	1	1	2	2	0	0	1	2	1	1	2	14
13. Were valid and reliable measures used consistently across all study groups to assess inclusion/exclusion criteria, interventions/exposures, outcomes, participant health benefits and harms, and confounding?	1	0	2	2	2	2	2	2	1	2	2	2	2	22

14. Was the length of follow-up similar across study groups?	0	0	0	2	0	0	0	0	0	0	0	0	2
15. In cases of high or differential loss to follow-up, was the impact assessed (e.g., through sensitivity analysis or other adjustment method)?	0	1	2	1	0	1	1	1	0	2	0	0	9
16. Were other sources of bias taken into account in the design and/or analysis of the study (e.g., through matching, stratification, interaction terms, multivariate analysis, or other statistical adjustment such as instrumental variables)?	0	2	0	0	0	0	2	2	0	0	2	0	8
17. Were the statistical methods used to assess the primary outcomes adequate?	0	0	0	0	0	0	2	2	0	0	2	0	6
<b>Total study score:</b>	1	7	8	6	6	9	12	9	6	12	8	7	6