

Psychiatric comorbid disorders of cognition

Original Research

Psychiatric comorbid disorders of cognition: A machine learning approach using 1,175 UK Biobank participants

Chenlu Li, Delia A. Gheorghe, John E.J. Gallacher, Sarah Bauermeister

Department of Psychiatry and on behalf of Dementias Platform UK, Warneford Hospital,
University of Oxford, Oxford, OX3 7JX

Corresponding author: Sarah Bauermeister sarah.bauermeister@psych.ox.ac.uk

Psychiatric comorbid disorders of cognition

Abstract

Background

Conceptualising comorbidity is complex and the term is used variously. Here, it is the coexistence of two or more diagnoses which might be defined as ‘chronic’ and, although they may be pathologically related, they may also act independently¹. Of interest here is the comorbidity of common psychiatric disorders and impaired cognition.

Objectives

To examine whether anxiety and/or depression are important longitudinal predictors of cognitive change.

Methods

UK Biobank participants used at three time points (n= 502,664): baseline, 1st follow-up (n= 20,257) and 1st imaging study (n=40,199). Participants with no missing data were 1,175 participants aged 40 to 70 years, 41% female. Machine learning (ML) was applied and the main outcome measure of reaction time intraindividual variability (cognition) was used.

Findings

Using the area under the Receiver Operating Characteristic (ROC) curve, the anxiety model achieves the best performance with an Area Under the Curve (AUC) of 0.68, followed by the depression model with an AUC of 0.63. The cardiovascular and diabetes model, and the covariates model have weaker performance in predicting cognition, with an AUC of 0.60 and 0.56, respectively.

Conclusions

Outcomes suggest psychiatric disorders are more important comorbidities of long-term cognitive change than diabetes and cardiovascular disease, and demographic factors. Findings suggest that

Psychiatric comorbid disorders of cognition

psychiatric disorders (anxiety and depression) may have a deleterious effect on long-term cognition and should be considered as an important comorbid disorder of cognitive decline.

Clinical implications

Important predictive effects of poor mental health on longitudinal cognitive decline should be considered in secondary and also primary care.

Psychiatric comorbid disorders of cognition

Summary Box

What is already known about this subject? 3-4 bullet points

- Poor mental health is associated with cognitive deficits.
- One in four older adults experience a decline in affective state with increasing age.
- ML approaches have certain advantages in identifying patterns of information useful for the prediction of an outcome.

What are the new findings? 3-4 bullet points

- Psychiatric disorders are important comorbid disorders of long-term cognitive change.
- Machine-learning methods such as sequence learning based methods are able to offer non-parametric joint modelling, allow for multiplicity of factors and provide prediction models that are more robust and accurate for longitudinal data
- The outcome of the RNN analysis found that anxiety and depression were stronger predictors of change IIV over time than either cardiovascular disease and diabetes or the covariate variables.

How might it impact on clinical practice in the foreseeable future?

The important predictive effect of mental health on longitudinal cognition should be noted and, its comorbidity relationship with other conditions such as cardiovascular disease likewise to be considered in primary care and other clinical settings

Psychiatric comorbid disorders of cognition

Background

Conceptualising comorbidity is complex and the term is used variously. Here, we refer to it as the coexistence of two or more diagnoses which might be defined as ‘chronic’ and, although they may be pathologically related to each other, they may also act independently¹. Of interest here is the comorbidity of common psychiatric disorders and impaired cognition.

Psychiatric disorders such as anxiety and depression (poor mental health) are associated with cognitive deficits², with higher baseline levels of depressive symptoms predicting a steeper decline in delayed recall and global cognition at follow-up³, and longitudinal slowing in processing speed⁴. Furthermore, at least one in four older adults experience a gradual decline in affective state⁵ with increasing age, suggesting that along with age-related cognitive decline, there could be a comorbid relationship of unclear temporality⁶. The challenge is to measure this relationship in the most effective way. A cognitive measure which is associated with both longitudinal cognitive decline and poor mental health is reaction time intraindividual variability (RT IIV)^{2,7}. RT IIV can be defined as trial-to-trial within-person variability in reaction time across individual trials within a single task for an individual. It is indicative of neurobiological disturbance, an increase of which is associated older age, mild cognitive impairment, dementia and mortality⁸.

Machine learning (ML) approaches have certain advantages in identifying patterns of information useful for the prediction of an outcome, even when complex high-dimensional interactions exist⁹. Unlike hypothesis-driven models, ML approaches are refrained from assumptions on data distributions, especially one of normally distributed independence of attributes or linear data behaviour which are often failed to be met by large registries incorporating biological data. In addition, ML approaches have shown certain advantages in examining potential predictors simultaneously in an unbiased manner. Although ML methods have been used successfully to

Psychiatric comorbid disorders of cognition

develop risk-prediction schemes in other areas of medicine, applications to psychiatric comorbid disorders have so far relied on small samples and thin predictor sets, failing to realize the full potential of the methods. With this in consideration, this study applies an ML technique on UK Biobank data to longitudinally examine whether anxiety/depression are important predictors of cognitive change.

The UK Biobank is a large population-based prospective cohort study of 502,664 participants. Invitations to participate in the UK Biobank study were sent to 9.2 million community-dwelling persons in the UK who were registered with the UK National Health Service (NHS) aged between 37 and 73 years. A response rate of 5.5% was recorded. Assessments took place at 22 centres across the UK where participants completed undertook comprehensive mental health, cognitive, lifestyle, biomedical and physical assessments. Mental health and cognitive assessments were completed on a touchscreen computer¹⁰. Here we use a ML approach to investigate whether the presence of anxiety and depressive symptoms in UK Biobank leads to greater cognitive change measured from baseline to follow-up.

Objective

To examine whether anxiety and/or depression are important longitudinal predictors of cognitive change.

Psychiatric comorbid disorders of cognition

Methods

Governance

Ethical approval was granted to Biobank from the Research Ethics Committee - REC reference 11/NW/0382¹⁰. All analyses were conducted using data from UK Biobank application 15008.

Population sample

The study considered for inclusion, the baseline sample of 502,664 UK Biobank participants aged between 37 and 73 years. For the analysis, participants with mental health and cognitive data at three time periods (recruitment, 5 year follow-up, and the 1st imaging sub-study) were used. For the purposes of this analysis the imaging sub-study data represents approximately a 10 year follow-up period from recruitment. Participants who chose to respond, 'I don't know' or 'I do not wish to answer' to items within either of the scales (depression and neuroticism) items at the three time periods were excluded, respectively (leaving 373,210 participants at baseline, 129,468 at first assessment and 3,501 at imaging sub-study). Casewise deletion was applied to all demographic variables (2,326 were excluded). All analyses were conducted listwise and the number of participants with no missing data across all measurement variables of interest were $n = 1,175$.

Assessment

Details of the UK Biobank assessment procedures may be found elsewhere¹⁰. Cognitive performance was assessed using a 'stop-go' task where RT IIV was used as a sensitive measure of cognitive change with greater RT IIV over time indicating decline in performance⁷. Mental health was assessed using two items from the Patient Health Questionnaire nine-item scale (PHQ-9)¹¹ to measure depression (PHQ-2: 'Over the last 2 weeks, how often have you been bothered by little interest or pleasure in doing things?'; 'Over the last 2 weeks, how often have you been bothered by feeling down,

Psychiatric comorbid disorders of cognition

depressed, or hopeless?')¹² and the 12-item Eysenck Personality Questionnaire-Revised (EPQ-R) Neuroticism scale as a proxy measure of anxiety.^{13 14} The other measures relevant to this analysis included self-reported cardiovascular disease (CVD: including heart attack, angina and stroke) and diabetes, and observed body mass index (BMI) as indicators of wider co-morbidity^{15 16}. Self-reported smoking, alcohol consumption, and fruit and vegetable consumption were used as indicators of lifestyle. Self-reported education, employment, ethnicity, and household income were used as socioeconomic indicators.

Analytic strategy

Data were modelled as follows. Both age and RT IIV were modelled as continuous variables. The RT IIV was calculated as the standard deviation of each participant's RTs over the test trials¹⁷. Participants with only one valid score at baseline or follow-up were omitted. The IIVs showed log-normal distributions and were natural log transformed. Depressed mood items were modelled as categorical variables (Not at all as 1, Several days as 2, More than half the days as 3 and Nearly every day as 4), and anxiety items were modelled as binary variables (yes/no). CVD and diabetes were modelled as binary indicators (present/absent). Gender was modelled as binary indicator (male/female). BMI was treated as a 3 level factor (normal (<25), overweight (25-29.9), obese (>29.9)). Lifestyle data were largely considered as binary indicators (smoker/non-smoker, <5 portions of fruit and vegetable per day / 5 or more portions per day), however alcohol was treated as a continuous variable (g/day). Socioeconomic data were considered as binary indicators (employed/not employed, college degree/no degree, white ethnicity/other) excepting income which was modelled as a 5 level factor (<£18,000, £18,000 -£30,999, £31,000 - £51,999; £52,000 - £100,000, >£100,000).

Psychiatric comorbid disorders of cognition

Machine learning model

A recurrent neural network (RNN) is a class of artificial neural networks that is often utilized as an effective prediction tool for longitudinal biomedical data. By exhibiting temporal dynamic behaviour for time sequences, it allows for temporal dependencies between measurements. As a sequence learning based method, it is able to offer non-parametric joint modelling of longitudinal data^{18 19}. Long short-term memory (LSTM) is an RNN architecture developed to deal with the exploding and vanishing gradient problem during backpropagation through time. It is able to effectively capture long-term temporal dependencies by employing a constant error carousel (memory cell) which maintains its state over arbitrary time intervals, and three nonlinear gating units (input, forget and output) which regulate the information flow into and out of the cell.^{18 20-23}. This architecture is applied to longitudinally model the prediction abilities of psychiatric comorbidity disorders on cognition via sequence-to-sequence learning.

The dataset was partitioned into two non-overlapping subsets: 75% of the within-class subjects for training and 25% for testing. Data were rescaled [0 and 1] to meet the default hyperbolic tangent activation function of the LSTM model. The Adam version of stochastic gradient descent algorithm was applied to tune the weights of the network. This optimisation algorithm combines the advantages of Adaptive Gradient Algorithm (AdaGrad) and Root Mean Square Propagation (RMSProp), thus being effective in handling sparse gradients on non-stationary problems. Furthermore, it compares favourably to other stochastic optimization methods in practice²⁴. Hyperparameters are optimised on the training set, e.g., the model was set to fit 1000 training epochs with a batch size set to the number of available training subjects. Mean absolute error (MAE) and Area under the Receiver Operating Characteristic (ROC) Curve (AUC) were used to assess model performance, respectively. Permutation tests were applied to obtain *p*-values of AUC. To evaluate whether depression and anxiety are important comorbid disorders of cognition and, to compare the

Psychiatric comorbid disorders of cognition

prediction abilities of diabetes and cardiovascular disease the model was run systematically to include the following as features:

- I. Covariates only as reference model (age; BMI; lifestyle factors; gender; socioeconomic factors)
- II. Neuroticism scale + reference model
- III. Depression scale + reference model
- IV. Diabetes and cardiovascular disease + reference model

Exposure (outcome) was cognition (RT IIV). Given that depression mood disorder and anxiety have previously been considered to be comorbid disorders of impaired cognition, we expected predictions based on comorbid mental health disorders to outperform predictions based only on covariate variables.

A complete data analysis approach was used and all analyses were conducted on the Dementias Platform UK Data Portal ²⁵ in Python 3.6.7 and MATLAB R2019.

Findings

Complete data were available for 1,175 Participants (Table 1). The sample was aged between 40 and 70 years at baseline, 41% female, and relatively healthy with few participants showing comorbidities and few reporting unhealthy behaviour such as smoking (5%).

The outcome of the RNN analysis found that anxiety and depression were stronger predictors of change IIV over time than either cardiovascular disease and diabetes or the covariate variables, as measured by the ROC curve (Figure 1 and Table 2). The AUC curve was 0.68 ($p < 0.0001$) for anxiety, followed by the depression model with an AUC of 0.63 ($p < 0.0001$). The cardiovascular and diabetes

Psychiatric comorbid disorders of cognition

model and the demographics model had weaker performance with an AUC of 0.60 ($p = 0.0036$) and 0.56 ($p = 0.1126$), respectively. The anxiety model prospectively identified 63.24% of patients who eventually reached high IIV (i.e., sensitivity), and 53.35% of patients who maintained low IIV (i.e., specificity). Correspondingly, the anxiety model had a positive predictive value (PPV) of 52.72%, and a negative predictive value (NPV) of 63.68%. The depression model achieved slightly smaller yet still significant ($p < 0.0001$) predictive performance in comparison with the other models. The RNNs results suggest that the anxiety and depression model outperformed the reference model (Model I – covariates only). In general, the neuroticism and depression scales helped improve the accuracy in capturing the pattern of IIV, and therefore may be more important predictors of cognitive change over cardiovascular disease and diabetes. The results also suggest increased risk of multiple mental health disorders occurring together, suggesting depression and anxiety are potential comorbid disorders of cognition.

Discussion

A longitudinal analysis of the comorbid predictors of mental health disorders on cognitive change was conducted using ML in a relatively healthy population sample. Evidence was found suggesting that anxiety and depression are important comorbid disorders of cognition but when cardiovascular disease, diabetes and other covariate predictors were included in the model these were not as important on long-term effect. This suggests that poor mental health has a significant deleterious effect on long-term cognition, and may be considered an important comorbid disorder of cognitive decline. The contribution of the present investigation is thereby three-fold.

First, this study used the community-based sample from UK Biobank with a general population large enough for identification of valid effects which might be implied across clinical and research settings.

Psychiatric comorbid disorders of cognition

Clinical-based samples have been criticised for inconsistency in diagnoses when patients are treated over time in a variety of facilities or at different times in the same facility, or for failure to report all the comorbid diagnoses of a patient²⁶. In comparison, community-based samples are based on a system in which a practical number of criteria are rated, or collected and assessed on every patient, providing unbiased prevalence or incidence rates of comorbidity, or unbiased estimates of risk factors for comorbidity²⁶.

Second, this study investigates the psychiatric comorbid disorders of cognition in a comprehensive ML design which was applied to examine if psychiatric disorders as measured by self-report scales are important predictors of impaired cognition. Unlike hypothesis-driven statistical techniques, the sequence learning based method is able to offer non-parametric joint modelling, allow for multiplicity of factors and provide prediction models that are more robust and accurate for longitudinal data¹⁸.

Third, this study is conducted longitudinally. Disease progression modelling using longitudinal data may provide clinicians with improved tools for diagnosis and monitoring of diseases. We apply the widely used LSTM network to capture the long-term temporal dependencies among measurements without making parametric assumptions about cognitive trajectories. To our knowledge there are no studies which have employed the ML techniques presented here to investigate the comorbidity of mental health disorders and longitudinal cognitive decline using cardiovascular disease and diabetes as comorbidity covariates. Utilising ML methodologies for clinical data such as psychiatric outcomes may provide additional information which is not possible with traditional statistical methodologies. In future research, approaches used here may include imputation on missing data in other longitudinal cohorts. For instance, by pre-processing data interpolation using mean or forward imputation, or utilizing possible correlations between missing values' patterns and the target²⁷.

Psychiatric comorbid disorders of cognition

There are general limitations to our work which should be noted. Participants in UK Biobank are selected from a general healthy population and outcomes may not reflect implications for those with clinically diagnosed psychiatric disorders of anxiety and depression. Here, also, measures of psychological state at point of assessment have been used to measure poor mental health (anxiety and depression). A further limitation is that only one cognitive measure has been used but findings would benefit from application across measures of wider cognition, e.g., memory and executive function.

Clinical Implications

The implication of this work is that the important predictive effect of mental health on longitudinal cognition should be noted,²⁸ and its comorbid relationship with other conditions such as cardiovascular disease likewise to be considered in primary care and other clinical settings.

Psychiatric comorbid disorders of cognition

Acknowledgements

This is a DPUK supported project with all analyses conducted on the DPUK Data Portal, constituting part 1 of DPUK Application 0132.

The Medical Research Council supports DPUK through grant MR/L023784/2.

CL was funded by DPUK to complete this analysis.

Authors' Contributions

CL, JG and SB conceptualised the idea. CL analysed the data, CL, DG, JG and SB interpreted the data, CL wrote the 1st draft of the manuscript. DG and SB commented, edited and proofread the manuscript. All authors read and approved the final manuscript.

Psychiatric comorbid disorders of cognition

References

1. Kraemer HC. Statistical issues in assessing comorbidity. *Stat Med* 1995;14(8):721-33. [published Online First: 1995/04/30]
2. Bauermeister S, Bunce D. Poorer mental health is associated with cognitive deficits in old age. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2015;22(1):95-105. doi: 10.1080/13825585.2014.893554 [published Online First: 2014/03/13]
3. Panza F, Frisardi V, Capurso C, et al. Late-life depression, mild cognitive impairment, and dementia: possible continuum? *Am J Geriatr Psychiatry* 2010;18(2):98-116. doi: 10.1097/JGP.0b013e3181b0fa13 [published Online First: 2010/01/28]
4. Bunce D, Batterham PJ, Christensen H, et al. Causal associations between depression symptoms and cognition in a community-based cohort of older adults. *Am J Geriatr Psychiatry* 2014;22(12):1583-91. doi: 10.1016/j.jagp.2014.01.004 [published Online First: 2014/02/08]
5. Arve S, Tilvis RS, Lehtonen A, et al. Coexistence of lowered mood and cognitive impairment of elderly people in five birth cohorts. *Aging (Milano)* 1999;11(2):90-5. [published Online First: 1999/07/01]
6. John A, Patel U, Rusted J, et al. Affective problems and decline in cognitive state in older adults: a systematic review and meta-analysis. *Psychol Med* 2019;49(3):353-65. doi: 10.1017/S0033291718001137 [published Online First: 2018/05/25]
7. Bunce D, Bauermeister S. Intraindividual Reaction Time Variability, Attention, and Age-Related Outcomes. Oxford Research Encyclopedia of Psychology: Oxford University Press 2019.
8. Bunce D, Tzur M, Ramchurn A, et al. Mental health and cognitive function in adults aged 18 to 92 years. *J Gerontol B Psychol Sci Soc Sci* 2008;63(2):P67-74. doi: 10.1093/geronb/63.2.p67 [published Online First: 2008/04/29]
9. Chekroud AM, Zotti RJ, Shehzad Z, et al. Cross-trial prediction of treatment outcome in depression: a machine learning approach. *The Lancet Psychiatry* 2016;3(3):243-50.
10. Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine* 2015;12(3):e1001779.
11. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16(9):606-13. doi: 10.1046/j.1525-1497.2001.016009606.x [published Online First: 2001/09/15]
12. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care* 2003;41(11):1284-92. doi: 10.1097/01.MLR.0000093487.78664.3C [published Online First: 2003/10/30]
13. Eysenck SBG, Eysenck HJ, Barrett P. A Revised Version of the Psychoticism Scale. *Personality and Individual Differences* 1985;6(1):21-29. doi: Doi 10.1016/0191-8869(85)90026-1
14. Ormel J, Jeronimus BF, Kotov R, et al. Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clin Psychol Rev* 2013;33(5):686-97. doi: 10.1016/j.cpr.2013.04.003 [published Online First: 2013/05/25]
15. Hassing LB, Hofer SM, Nilsson SE, et al. Comorbid type 2 diabetes mellitus and hypertension exacerbates cognitive decline: evidence from a longitudinal study. *Age Ageing* 2004;33(4):355-61. doi: 10.1093/ageing/afh100 [published Online First: 2004/05/12]
16. Biessels GJ, Deary IJ, Ryan CM. Cognition and diabetes: a lifespan perspective. *Lancet Neurol* 2008;7(2):184-90. doi: 10.1016/S1474-4422(08)70021-8 [published Online First: 2008/01/22]
17. Dykiert D, Der G, Starr JM, et al. Age differences in intra-individual variability in simple and choice reaction time: systematic review and meta-analysis. *PLoS One* 2012;7(10):e45759.
18. Ghazi MM, Nielsen M, Pai A, et al. Robust training of recurrent neural networks to handle missing data for disease progression modeling. *arXiv preprint arXiv:180805500* 2018
19. Pearlmutt BA. Learning state space trajectories in recurrent neural networks. *Neural Computation* 1989;1(2):263-69.

Psychiatric comorbid disorders of cognition

20. Gers FA, Schmidhuber E. LSTM recurrent networks learn simple context-free and context-sensitive languages. *IEEE Transactions on Neural Networks* 2001;12(6):1333-40.
21. Gers FA, Schmidhuber J, Cummins F. Learning to forget: Continual prediction with LSTM. 1999
22. Greff K, Srivastava RK, Koutník J, et al. LSTM: A search space odyssey. *IEEE transactions on neural networks and learning systems* 2017;28(10):2222-32.
23. Hochreiter S, Schmidhuber J. Long short-term memory. *Neural computation* 1997;9(8):1735-80.
24. Kingma DP, Ba J. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980* 2014
25. Bauermeister S, Orton C, Thompson S, et al. The Dementias Platform UK (DPUK) Data Portal. *BioRxiv Preprint* doi: 10.1101/582155
26. Cloninger CR. Implications of comorbidity for the classification of mental disorders: the need for a psychobiology of coherence. *Psychiatric diagnosis and classification* 2002;79:106.
27. Lipton ZC, Kale DC, Wetzell R. Modeling missing data in clinical time series with RNNs. *Machine Learning for Healthcare* 2016(arXiv:1606.04130v5)
28. Baruch N, Burgess J, Pillai M, et al. Treatment for depression comorbid with dementia. *Evid Based Mental Health* 2019;22:167-171.

Psychiatric comorbid disorders of cognition

Table 1 Descriptive statistics

	Baseline		First Assessment			Imaging			
	%	IIV mean	IIV SD	%	IIV mean	IIV SD	%	IIV mean	IIV SD
Panel A – Neuroticism EPQ-R Scale									
“Does your mood often go up and down?”									
Yes	33%	72.18	53.38	30%	71.10	47.07	28%	78.64	54.66
No	67%	72.38	47.54	70%	73.00	48.50	72%	74.98	52.08
“Do you ever feel just miserable for no reason?”									
Yes	33%	68.89	46.54	30%	70.34	41.65	27%	78.13	51.82
No	67%	74.03	50.90	70%	73.31	50.54	73%	75.22	53.21
“Are you an irritable person?”									
Yes	25%	72.80	52.87	24%	73.43	52.06	22%	72.75	51.38
No	75%	72.15	48.39	76%	72.11	46.76	78%	76.96	53.23
“Are your feelings easily hurt?”									
Yes	48%	75.40	50.37	46%	73.53	51.28	43%	75.19	48.15
No	52%	69.40	48.60	54%	71.50	45.19	57%	76.64	56.13
“Do you often feel fed-up?”									
Yes	28%	70.08	47.66	28%	69.53	46.24	23%	78.87	53.15
No	72%	73.18	50.23	72%	73.53	48.72	77%	75.16	52.73
“Would you call yourself a nervous person?”									
Yes	18%	77.87	58.06	17%	72.94	52.84	15%	76.48	54.22
No	82%	71.11	47.41	83%	72.32	47.07	85%	75.93	52.60
“Are you a worrier?”									
Yes	45%	71.83	46.39	46%	73.84	50.85	42%	74.81	50.13
No	55%	72.71	51.99	54%	71.24	45.60	58%	76.91	54.75
“Would you call yourself tense or highly strung?”									
Yes	10%	74.25	48.17	10%	80.28	60.56	8%	74.99	48.79
No	90%	72.09	49.70	90%	71.57	46.46	92%	76.11	53.20
“Do you worry too long after an embarrassing experience?”									
Yes	45%	69.91	40.66	46%	72.36	51.05	47%	73.12	47.35
No	55%	74.27	55.66	54%	72.48	45.40	53%	78.57	57.14
“Do you suffer from nerves?”									
Yes	16%	75.18	53.87	14%	71.05	50.58	12%	67.83	44.36
No	84%	71.78	48.68	86%	72.65	47.66	88%	77.10	53.77
“Do you often feel lonely?”									
Yes	12%	73.67	54.61	12%	70.26	36.67	11%	74.46	50.43

Psychiatric comorbid disorders of cognition

No	88%	72.12	48.80	88%	72.73	49.46	89%	76.21	53.13
“Are you often troubled by feelings of guilt?”									
Yes	23%	68.99	36.86	22%	73.61	51.99	24%	76.89	50.80
No	77%	73.33	52.78	78%	72.08	46.89	76%	75.74	53.47
Panel B – PHQ-2 Depression Scale									
“frequency of depressed mood”									
Not at all	83%	72.53	51.25	86%	73.47	49.39	87%	76.29	53.42
Several days	15%	71.72	40.85	12%	65.35	39.48	11%	72.94	47.49
More than	1%	72.45	43.13	1%	79.28	38.60	1%	89.04	60.27
Nearly	1%	60.70	24.81	1%	60.88	28.68	1%	54.53	36.12
“frequency of unenthusiasm / disinterest”									
Not at all	87%	71.99	50.41	89%	72.99	48.92	90%	76.37	53.27
Several days	11%	74.07	43.29	9%	70.45	42.47	9%	70.05	47.67
More than	1%	65.79	35.19	1%	55.48	25.83	1%	89.56	59.17
Nearly	1%	91.88	53.72	1%	58.10	34.68	0%	101.1	40.76
Panel C – Other Comorbid Disorders									
Heart attack									
Yes	1%	62.04	33.94	1%	75.11	55.66	2%	65.52	51.29
No	99%	72.42	49.66	99%	72.39	47.97	98%	76.22	52.86
Angina									
Yes	1%	67.15	34.69	1%	72.81	44.78	1%	62.47	27.99
No	99%	72.37	49.67	99%	72.42	48.12	99%	76.18	53.04
Stroke									
Yes	1%	79.49	38.78	1%	89.73	51.71	1%	71.77	45.78
No	99%	72.26	49.61	99%	72.23	48.01	99%	76.06	52.90
High blood pressure									
Yes	18%	73.25	43.89	22%	73.51	53.56	22%	77.21	48.83
No	82%	72.11	50.67	78%	72.12	46.43	78%	75.68	53.92
Diabetes									
Yes	2%	64.91	23.10	2%	69.54	43.77	3%	99.29	51.22
No	98%	72.47	49.92	98%	72.50	48.18	97%	75.32	51.74
Panel D – Covariates									
Gender									
Male	59%	72.32	48.20	59%	71.28	48.09	59%	75.22	54.45
Female	41%	72.31	51.38	41%	74.05	48.02	41%	77.14	50.48
Education									
College	55%	69.50	45.23	55%	72.84	48.99	58%	73.04	51.96
Other	45%	75.77	54.17	44%	71.92	46.93	42%	80.06	53.77
BMI									
Normal	41%	73.14	54.74	42%	74.02	49.15	45%	77.98	57.45
Overweight	44%	73.14	48.19	42%	73.23	48.69	40%	74.43	49.64
Obese	15%	67.72	36.90	15%	65.81	42.69	15%	74.32	46.10
Household Income									
<£18,000	10%	85.17	74.41	12%	76.31	42.56	11%	86.90	65.49

Psychiatric comorbid disorders of cognition

(£18,000,£30,99	23%	76.47	44.82	26%	74.29	47.63	27%	79.02	47.50
(£31,000,£51,99	30%	72.42	47.12	30%	73.08	50.67	30%	69.38	45.53
(£52,000,£100,0	28%	68.76	47.75	23%	73.69	52.21	23%	79.33	60.52
>£100,000	10%	59.03	35.00	9%	55.66	29.50	9%	68.16	49.72
Smoking status									
No	95%	72.85	50.20	96%	72.28	48.24	97%	75.86	52.53
Yes	5%	62.77	34.29	4%	76.20	43.39	3%	81.25	62.33
Employment status									
Employed	67%	68.33	45.79	51%	68.48	46.10	42%	73.02	49.65
Other	33%	80.48	55.60	49%	76.60	49.75	58%	78.22	54.98
Ethnicity									
White	98%	72.37	49.44	98%	72.42	47.64	98%	75.62	52.47
Other	2%	69.32	55.05	2%	72.66	68.22	2%	94.84	66.43
Vegetable and fruit intake									
≥ 5 unit per	76%	72.39	49.47	74%	72.72	47.39	79%	75.30	50.92
< 5 unit per	24%	72.06	49.76	26%	71.59	49.99	21%	78.64	59.32
					Mean				
Age	56			60			63		
Alcohol Consumption (g/day)	18.33			16.39			16.37		

Table 2 Model performance for comorbid longitudinal predictors of IIV

	Anxiety	Depression	Cardiovascular & Diabetes	Covariates
AUC	0.68 (<0.0001 ^{***})	0.63 (<0.0001 ^{***})	0.60 (0.0036 ^{**})	0.56 (0.1126)
Sensitivity	63.24%	59.72%	55.05%	54.90%
Specificity	53.35%	52.94%	55.45%	50.99%
PPV	52.72%	52.08%	49.27%	47.06%
NPV	63.68%	61.42%	60.09%	59.72%
$P(AUC_{Anxiety} - AUC_{Depression})$			(0.1056)	
$P(AUC_{Anxiety} - AUC_{Cardiovascular \& Diabetes})$			(0.0014 ^{**})	
$P(AUC_{Anxiety} - AUC_{Covariates})$			<0.0001 ^{***}	

AUC = area under receiver characteristics curve. PPV = positive predictive value. NPV = negative predictive value. The P-values are in the parentheses. Significance at the 0.1%, 1%, and the 5% level is indicated by ^{***}, ^{**}, and ^{*}, respectively

Psychiatric comorbid disorders of cognition

Psychiatric comorbid disorders of cognition




Figure. ROCs for RNN Models I to IV

ROC

