Inferring the role of the microbiome on survival in patients treated with immune checkpoint inhibitors: causal modeling, timing, and classes of concomitant medications

Daniel Spakowicz[1,2][§][*], Rebecca Hoyd[1][§], Mitchell Muniak[1], Marium Husain[1], James S.

Bassett[1], Lei Wang[2], Gabriel Tinoco[1], Sandip H. Patel[1], Jarred Burkart[1], Abdul Miah[1], Mingjia

Li[3], Andrew Johns[1], Madison Grogan[1], David P. Carbone[1], Claire F. Verschraegen[1], Kari L.

Kendra[1], Gregory A. Otterson[1], Lang Li[2], Carolyn J. Presley[1], and Dwight H. Owen[1]

[1] Division of Medical Oncology, Department of Internal Medicine, Ohio State University,

Columbus, OH, USA

[2] Department of Biomedical Informatics, Ohio State University, Columbus, OH, USA

[3] Division of Hospital Medicine, Department of Internal Medicine, The Ohio State

University College of Medicine

[§] These authors contributed equally

*To whom correspondence should be addressed: Daniel.Spakowicz@osumc.edu

ABSTRACT

The microbiome has been shown to affect the response to Immune Checkpoint Inhibitors (ICIs) in a small number of cancers. Here, we sought to more broadly survey cancers to identify those in which the microbiome will play a role using retrospective analyes. We created a causal model for the relationship between medications, the microbiome and ICI response and used it to guide the abstraction of electronic health records of 690 patients who received ICI therapy for advanced cancer. Medications associated with changes to the microbiome including antibiotics, corticosteroids, proton pump inhibitors, histamine receptor blockers, non-steroid anti-inflammatories and statins were abstracted. We tested the effect of medication timing on overall survival (OS) and evaluated the robustness of medication effects in each cancer. Finally, we compared the size of the effect observed for antibiotics classes to taxa correlated with ICI response and a literature review of culture-based antibiotic susceptibilities. Of the medications assessed, only antibiotics and corticosteroids significantly associated with lower OS. The hazard ratios (HRs) for antibiotics and corticosteroids were highest near the start of ICI treatment but remained significant when given prior to ICI. Antibiotics and corticosteroids remained significantly associated with OS even when controlling for multiple factors such as Eastern Cooperative Oncology Group performance status and stage. When grouping antibiotics by class, β-lactams showed the strongest association with OS across all tested cancers. The timing and strength of these effects after controlling for confounding factors are consistent with role for the microbiome in response to ICIs.

KEYWORDS (6)

Microbiome, immune checkpoint inhibitors, antibiotics, corticosteroids, cancer, immunotherapy

BACKGROUND

Treatment with Immune Checkpoint Inhibitors (ICIs) has improved patient outcomes across a wide variety of cancers. Not all patients respond to these drugs and there is a need to identify biomarkers of outcomes. Three recent studies have shown that microbes are associated with response and overall survival (OS) in renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC) and melanoma (1–3). The microbiome may be a key player in response to ICI therapy and a potential biomarker of treatment response.

The microbiome is known to interact with the immune system, but how it affects response to ICIs is not understood. The effectiveness of ICI treatment relies on active T-cell infiltration of a tumor; microbes have been associated with increased Tumor Infiltrating Lymphocytes in an IL12-depended manner (4). However, other immune cells dampen response to ICIs such as myeloid-derived suppressor cells and FOXP3 & CD4+CD25+ T-regulatory cells. Another, systemic form of immune repression is characterized by the production of prostaglandins (5–7).

Several medications commonly used during routine oncologic care and ICI treatment can influence inflammation pathways and/or the microbiome. Corticosteroids (CS) affect both of the aforementioned T-cell subtypes and the prostaglandin-related inflammatory pathways (8). Additionally, antibiotics (ABx) have a direct effect on the microbiome by killing or halting the growth of bacteria. Proton pump inhibitors (PPIs), histamine 2 blockers (H2Bs), non-steroid anti-inflammatory drugs (NSAIDs), and CS have also been associated with changes in the microbiome but, in contrast to antibiotics, this mechanism is indirect (9). PPIs, by inhibiting gastric acid secretion, alter the pH of the gut and change the number and types of bacteria that pass through the stomach (10). Notably, if the taxa enriched by the PPI-induced pH change are also important for response to ICIs, then PPI treatment during ICI may influence clinical

outcomes. The effect of other medications on clinical response may be challenging to interpret

given that the effects may influence both the microbiome and ICI response.

In order to disentangle these complex interactions, we created a model of the

relationship between patient characteristics, medications that affect the microbiome,

inflammation, and survival. Second, we performed a retrospective analysis of patients who

received ICI therapy for advanced cancer between 2011 and 2017 including medications with

known effects on either the microbiome or its pathway toward affecting ICI response. Third, we

estimated the association for each medication with OS. Fourth, we analyzed the effects of

medications longitudinally, in order to decouple confounding variables at different time points.

Fifth, we controlled for variables that broadly describe differences in baseline statuses (e.g.

Eastern Cooperative Oncology Group performance status (PS)) of individuals who received

concomitant medications and those who did not. Sixth, we compared the associations across

several cancers, for which the medications are prescribed in subtly different ways that can be

leveraged to gain further insight into the causal effects. Finally, we related these results to the

microbes shown to be enriched or depleted in individuals who respond to ICIs. The combination

of these strategies gives layers of support to defining the role of the microbiome in the context

ICI treatment of cancer.

METHODS

Causal Model

We performed a literature review of the relationship between the microbiome and

response to ICIs and medications that affect the microbiome (**Figure 1**, references in **Figure**

S1). From these references, a causal model was then constructed such that the nodes

correspond to observable endogenous variables (V_i), as a subset of a set of U exogenous and

unobserved variables that affect the relationship between the microbiome and OS in patients

treated with ICIs. Directed edges denote a relationship between variables when the following

conditions are met: (1) there is a reported relationship between variables in which both variables

were either observed or defined by intervention, and (2) the relationship cannot be explained

through using an existing path. For example, Gopalakrishnan et al reported a correlation

between the microbiome and ICI response (1). This relationship exists in the graph as mediated

by the nodes Microbiome → T-cell Mediated Inflammation → ICI Response, therefore no edge

is drawn directly from Microbiome → ICI Response. The resulting directed acyclic graph was

constructed using the igraph and dagitty packages in R (11,12).

Retrospective data collection

We identified patients with advanced cancer treated between 2011 and 2017 at the Ohio

State University Comprehensive Cancer Center/Arthur G. James Cancer Hospital (OSUCCC-

James) who received at least one dose of ICIs as part of an IRB approved retrospective study

(OSU-2016C0070, OSU-2017C0063). Patient data were collected and housed in REDCap (13).

Medication timing, dose and names were collected from the electronic medical record

information warehouse and validated by manual chart review.

Medication history curation

ABx and CS data were retrieved from the information warehouse within 180 days of ICI

start. All medications matching a comprehensive list of steroid generic and brand names were

collected with dates and routes of administration. Medications were filtered to those confirmed

to be administered and the results checked against a manually-curated subset of the records.

Survival Analysis

Overall survival (OS) was reported in days from the initiation of ICI to the date of death

or last follow-up. All univariate and multivariate analyses were conducted using the survminer

package in R (14,15). Univariate analyses used Kaplan-Meier survival curves with log-rank

tests. Multivariate analyses used Cox-Proportional Hazards models, defining the hazard

function for each patient k as:

$$h_k(t) = h_0(t)e^{\{\sum_{t=1}^{n} \beta_1 A + \beta_2 S + \beta_3 B + \beta_4 E + \beta_5 G\}}$$

Where h(t) is the hazard function at time t = 1 to n, A is a binary indicator of antibiotic

use (+/- 28 days from start of ICIs), S is a binary indicator of corticosteroid use (+/- 28 days from

start of ICIs), B is BMI, E is the Eastern Cooperative Oncology Group performance status (PS)

score (1-5), and G is age. We constructed the models using the survival package and evaluated

model fits using a likelihood ratio tests in R (16–18).

Timing analysis

A 30-day sliding window was used to evaluate the effect of medication timing on the

association with OS. Patients prescribed medications within the window were compared to a

cohort of individuals who were not prescribed those medications within 180 days before or after

the start of ICI treatment. Kaplan-Meier survival curves were used to estimate a hazard ratio

(HR) of association with each treatment window, incremented by single-days, e.g. prescribed

180-150 days before ICI start vs no prescribed medications, and then prescribed 179-149 days

before ICI start vs no prescribed medications. HRs and confidence intervals were calculated in

the survival package and plotted with ggplot2 in R (17–19).

Antibiotics and Corticosteroids classes

ABx and CS were collapsed into categories by DrugBank v5.0 accession numbers (20).

HRs were estimated for medication class and cancer combinations if the total sample set

included at least 20 individuals. Cox Proportional Hazards models for the effects of ABx and CS

class were used to allow for simultaneous estimation of the effects of more than one class,

when applicable. Plots showing prescriptions of more than one class were created with the

UpSetR package in R (21).

Regularized Cox Regression

Regularized Cox survival models for each cancer were implemented in the glmnet

package in R. We optimized the regularization parameter by coordinate descent via 10-fold

cross-validation and then tested the robustness of the parameter selection and resulting

covariates with 1000 bootstrap replicates of different random samples of the dataset (22,23).

Reproducibility

Scripts to reproduce all figures and analyses can be found at

https://github.com/spakowiczlab/co-med-io.

RESULTS

Causal Model

The relationships between clinical variables, medications, the microbiome, ICI response

and OS are strongly interconnected. Our literature review to predict their causal relationships

(**Figure 1**) led to several hypotheses observable within retrospective data. First, medications

that directly affect the microbiome will alter OS via effects on T-cell mediated inflammation (i.e.

ABx → Microbiome → T-cell mediated inflammation → ICI Response → OS). Second, CS and

ABx may have additive effects on ICIs through a collider effect on T-cell mediated inflammation

(i.e. ABx → Microbiome → T-cell mediated inflammation ← CS). Finally, the clinical variables of

stage, BMI, age, and medications such as CS, confound the relationship between the

microbiome and ICI response, mediated by Prostaglandin-based inflammation (which itself is a

collider), and therefore must be controlled for in order to infer the role of the microbiome on OS

(**Figure 1**).

Patient Characteristics

Retrospective analysis of electronic medical records from 2011 to 2017 at the OSUCCC-

James identified 690 patients treated with ICIs (**Table 1**). Most (76.6%) had a PS of 0 or 1. The

most common diagnoses were melanoma (28.5%) and non-small cell lung cancer (NSCLC)

(23.4%). Cancers represented by fewer than 20 patients were categorized as "Other" (23.4%).

The majority of patients (90%) had metastatic disease. ICI treatments included nivolumab in

52.8% of patients, ipilimumab in 18.0% and pembrolizumab in 15.1%.

Microbiome and inflammation-related concomitant medication use

Among the medications included in the causal model, ABx, CS, PPIs, H2Bs, statins and

NSAIDs were identified in this cohort. ABx were prescribed in 35% of patients within 28 days of

the start of ICIs (Table 1). The most commonly prescribed ABx were β-lactams (**Figure S1**). CS

were prescribed in 40% patients within 28 days of the start of ICIs. The most commonly

prescribed CS were dexamethasone and prednisone (**Figure S2**). PPIs were prescribed in 37%

of patients. Some patients received a single medication and no others during the study period,

however, more frequently patients received several medications, e.g. CS with PPI and ABx,

consistent with prophylaxis for developing an ulcer or pneumonia (**Figure S3**). The analysis

strategy first tested for an association of a medication with OS without controlling for

confounding effects of other medications and then further explored those medications with

strong associations.

Across all cancer types, patients who were prescribed ABx within 28 days of the start of

ICIs showed decreased OS (**Figure 2A**). This was also true of patients prescribed CS (**Figure**

2B), but not of patients prescribed other medications (**Figure 2C**). ABx showed a strong

negative correlation with OS in RCC, NSCLC, melanoma, and bladder cancer. CS showed a

strong negative correlation with OS in NSCLC, melanoma and other cancers. While other

medications were not significantly associated with OS across all cancers, several showed

significant associations with specific cancers. For example, H2Bs and NSAIDs associated with

decreased OS in sarcoma and NSCLC, respectively. On the other hand, PPIs and Statins

positively associated with OS in sarcoma. However, we observed the strongest associations for

ABx and CS, and therefore followed these medications in further analyses.

Timing of medication use

Next we focused on the timing of ABx and CS prescriptions and their associations with

OS in each cancer, using a 30-day sliding window (see Methods). ABx showed a greater HR

than CS over nearly the entire period, and both were negatively associated with OS (**Figure**

3A). ABx treatment showed the highest HR more than 100 days before the start of ICIs with a

second, lesser peak around day 0. CS showed a single, strong peak at day 0. We therefore

focused the timing analyses around ICI day 0 to capture the largest HR for both ABx and CS

and examined the effects across cancers and drug subclasses.

Antibiotics and Corticosteroids classes

The effect of ABx on treatment response in different cancer types was not consistently

associated with ABx class (**Figure 3B**). For example, β-lactams showed the highest HR in

melanoma, but vancomycin (oral) had the highest HR in head and neck squamous cell

carcinoma (HNSC). In addition, the overall effect of ABx was sometimes associated with a

single ABx subclass and sometimes distributed over many. Additionally, NSCLC strongly

associated with fluoroquinolone ABx, and this effect was stronger than the effect observed for

all ABx combined. By contrast, the combined effect of all ABx in melanoma was much stronger

than any individual ABx class. In fact, tetracycline was positively associated with OS in

melanoma patients, despite the overall effect of ABx being negatively associated. On the other

hand, the effects of CS classes on different cancers was more consistent (**Figure 3C**), though

these comparisons were often limited by the sample size. Particularly with small sample sizes,

confounding effects of patients receiving multiple drugs, e.g. ABx and CS, may dominate

associations with OS. We therefore used combined models of ABx and CS to examine the

effects of each.

Combined modeling of ABx and CS, controlling for covariates

Models containing both ABx and CS showed that both are significantly associated with OS. A Kaplan-Meier curve stratifying patients by ABx, CS, or both, showed nearly identical intermediate effects of either ABx or CS, and an additive combined effect (**Figure 4A**). We next sought to control for confounding covariates using a Cox Proportional Hazards model. Including PS, BMI, sex, stage, and age in the model confirmed that ABx and CS remained highly significant, as were PS, BMI and age (**Figure 4B**). This suggests that ABx and CS are affecting OS in the context of ICI therapy by a mechanism other than that which is captured by PS, BMI or age, and is consistent with the microbiome parent to T-cell inflammation and child of ABx (**Figure 1**).

In order to estimate the effects of ABx and CS within each cancer, we applied a method that (1) allowed different covariates to be included in each cancer, commensurate with the different clinical features of each cancer, and (2) removed uninformative variables, increasing the power for those cancers with smaller numbers of patients in this dataset. In addition, we repeated the analysis with different random samplings of the data in order to estimate the robustness of the variable selection. We found ABx to consistently and significantly associate with OS in bladder cancer, melanoma and RCC, but not in HNSC, NSCLC, or sarcoma. The HR was above 1 in each of the cancers where ABx was a consistently-selected covariate. Melanoma was notable in that all variables were consistently selected, with ABx showing the highest HRs.

The relationship between ABx, OS, and the microbiome

The bacterial taxa that showed the strongest enrichment in responders or non-responders to ICIs were selected from the literature and combined into a phylogenetic tree (**Figure 5**) (1–3). The taxa spanned several phyla and few ranks were consistently enriched in either responders or non-responders. For example, Firmicutes was found to be enriched in

responders (1), but within the phylum are several taxa found to be enriched in non-responders

(2,3). An exception to this was Bacteroidetes, which was found to be enriched in non-

responders and each of the four species in the phylum were also enriched in non-responders

(1,3). We performed a literature review of ABx susceptibilities for each of these taxa to estimate

whether the size of the HR of the ABx would relate to the taxa for which it is active. For

example, an ABx that target only bacteria enriched in non-responders may be beneficial

because it may shift the community toward those taxa enriched in responders. On the other

hand, if the overall diversity of the microbiome is important, broad-spectrum ABx may have

higher HRs than narrow-spectrum.

The ABx class with the largest HR across all cancers was the β-lactams. Within this

group category are the cephalosporins, which have a relatively narrow spectrum of activity and

a unique pattern relative to other ABx classes. The cephalosporins are ineffective against the

Bacteroidetes, found to be enriched in non-responders, but so were ABx such as vancomycin

and sulfamethoxazole-trimethoprim (SXT). However, unlike vancomycin and SXT,

cephalosporins effectively target *A. muciniphila*, which was shown to causally modify response

to ICIs. Cephalosporins are also ineffective against several Firmicutes, similar to clindamycin,

macrolides and metronidazole (**Figure 5**).

DISCUSSION

The effects of medications or other variables are difficult to parse in a dynamic setting

such as during treatment for cancer. We used a variety of methods to show that ABx and CS

are significantly associated with decreased OS in several cancer types and that these results

are consistent with mediation via the gut microbiome.

The association of CS with ICI response and OS remains controversial. Our observed

association is consistent with other observations of decreased OS in NSCLC (8). However,

Riccuiti et al showed no effect of CS on OS in NSCLC when given on the same day as ICI start,

when the CS was prescribed for reasons other than "cancer-related palliative indications" (24).

Our records lack some variables needed to replicate those results, however our results are

consistent with aspects those findings. For example, dexamethasone treatment showed a

strong negative association with OS across several cancer types, consistent with its use for

brain metastases and anorexia, which are all indicators of poor clinical outcome. On the other

hand, several of our analyses demonstrated associations between CS and OS that may not be

due to selecting a sub-cohort with a poor prognosis. Our first causal strategy, the time analysis,

showed similar results when restricting CS medications to a single day, but a larger effect when

a wider time window was used (**Table 2**). Our second causal strategy, controlling for covariates,

cannot be directly compared because our dataset did not include central nervous system

metastases. However, when we control for metastatic stage and PS, the CS association

remains. Our third causal strategy, comparisons between cancers, shows that the CS

association with OS is observed in cancers for which brain metastases are not common, such

as RCC, and for specific CS not typically used for brain metastases, such as

methylprednisolone in HNSC. This suggests that understanding the association between CS

and the response to ICIs may require more granular assessment of CS types (i.e. rather than

collapsing to 10 mg prednisone equivalent) and cancers.

We applied the same logical framework to ABx treatment to demonstrate an effect on

OS. Unlike CS, the majority of studies have found an association between ABx use and ICI

response, independent of the time window (**Table 2**). First, our longitudinal analysis showed a

global maximum HR well before the start of ICIs, consistent with the ABx effects persisting for

long periods. Given this result, it is unlikely that acute illnesses drive the association between

ABx and OS. Second, when controlling for illness-related covariates that report on the overall

health status of the individual, e.g. PS, the effect of ABx remained significant. Third, the

associations of ABx and OS were observed across cancer types (e.g. patients with bladder

cancer versus melanoma). A larger fraction of bladder cancer patients were treated with ABx

than any other cancer (56%), consistent with their use for urinary tract or as prophylaxis for

invasive urologic procedures. On the other hand, melanoma patients treated with ABx were the

smallest fraction of any cancer (25%), consistent with this population being less likely to

undergo procedures in which prophylactic ABx are used. It is reasonable to suspect that

melanoma patients treated with ABx are therefore more compromised than those not treated

with ABx. However, an effect of ABx remains, even for bladder cancer. Although it remains

probable that the cohorts who receive ABx are different from those who did not in ways that

have not been controlled for in analyses, these three analyses add confidence to the

association of ABx with OS in the context of ICIs.

We next related the strength of the association of ABx classes with OS and the microbes

that those ABx classes affect. The β-lactam ABx were shown to have the strongest association

with OS across cancer types. The literature review of antibiotic susceptibilities showed that this

diverse class is effective against the Gram-positive phylum Firmicutes. The literature review of

the bacterial taxa associated with response to ICIs, showed that the Firmicutes are enriched in

responders to ICIs. Moreover, β-lactams are not consistently effective against members of the

phylum Bacteroidetes, which was found to be enriched in non-responders. This suggests that

the β-lactams may show the strongest signal across all cancers in our dataset because they

disrupt the microbiome in such a way that they reduce response to ICIs by depleting the

Firmicutes more so than the Bacteroidetes.

The association between ABx prescriptions and OS that we observe is consistent with

direct measurements of the microbiome and response to ICIs (1–3). However, there is no

consensus for which taxa are enriched in the responders to ICIs (**Figure 5**). For example, there

is causal evidence for *Akkermansia muciniphila* increasing response to ICIs, however, it was not

among the most enriched in the other datasets (1–3). Nonetheless, agreement can be observed

at broader taxonomic characterizations. Non-responders are enriched for taxa in the phylum

Bacteroidetes and this is consistent with the strength of association we see for different classes

of ABx (1,3). Narrow spectrum β-lactams (e.g. cephalosporins), which show the strongest

association with OS, are effective against Firmicutes (enriched in responders) but less so

against Bacteroidetes (enriched in non-responders).

The results presented here contrast with several assumptions gathered from the

literature and described by the causal model (**Figure 1**). First, we found that ABx and CS are

the only medications significantly associated with OS, despite the inclusion of several

medications associated with changes to the microbiome (**Figure 2**). This may be due to the

types of changes incurred (e.g. PPIs may not significantly change the abundances of those taxa

linked to ICI response) or the strength of the effect amid the noise in the data. However, the

other two hypotheses were borne out by the analyses.

The CS and ABx medications showed an additive effect on OS, consistent with a collider

interaction in the model (**Figure 4A**). Also, there was an effect of ABx after controlling for many

covariates, consistent with its direct effect on the microbiome and the microbiome playing a role

in ICIs (**Figure 4B**). This result was consistent with the relationship between the strength of the

ABx signal and the bacterial taxa susceptible to that ABx (**Figure 5**).

Limitations

A key challenge in this and other retrospective analyses is inferring causal relationships

in non-randomized cohorts. For example, patients who receive medications such as antibiotics

may be quite different from those who do not. However, it is difficult to imagine an ethical trial

that could randomize treatment with ABx in this setting. Therefore, retrospective analyses may

be the best option until direct measurements of the microbiome are widely available. We used a

variety of methods to show that ABx and CS are significantly associated with decreased OS

across a variety of cancers and that these results are consistent with a role for the gut

microbiome.

Our study remains limited by being unable to account for important factors known to

affect OS in the context of ICI treatment. For example, the complete ABx history of patients --

much longer than the windows reported here -- are very likely of consequence. Several groups

have studied the recovery of microbiome diversity following ABx exposure and results show

reasonable recovery 90 days later (25,26). However, multiple courses of ABx prevented such a

recovery; i.e. diversity returned to baseline after one treatment with ABx, but not after a second

ABx treatment within 60 days (27). It is therefore possible that individuals who show extreme

effects of ABx treatment are beyond the time scale of this study. Without baseline microbiome

diversity measures we are unable to capture such information. Similarly, estimating the effects

of ABx on communities from data on microbes in isolation is, at best, approximate. A better

understanding of how ABx affect complex communities is needed. Other limitations include our

small sample size relative to the heterogeneity in the data. Future directions should capture

variables such as other co-morbidities or the presence of brain metastases, tumor biomarkers

such as tumor mutational burden and PD-1/PDL-1 status, and outcome variables like ICI

response or the number of tumor-infiltrating lymphocytes.

CONCLUSIONS

ABx and CS, but not other medications known to affect the microbiome, are associated

with reduced OS when administered near the start of ICI treatment. Our results show this finding

several cancer types, and for several subclasses of these drugs. These results are consistent

with a role of the microbiome in response to ICIs and identify clinical settings where the

microbiome is likely to play the largest role, namely NSCLC, melanoma, RCC, HNSC, and

bladder cancer. A clear understanding of which microbes are important for ICI responses and in

what cancers will require the collection of microbiome samples across a wide variety of clinical

settings. However, some information can be gathered by indirect means, which identifies the

settings where the microbiome is likely to have the greatest effects. Medications that affect the

microbiome given concomitantly with ICIs provide evidence for where microbes play a role.

Further work is needed to identify which microbes are important and identify solutions to

mitigate these effects and perhaps promote greater response to ICIs.

LIST OF ABBREVIATIONS

ICIs: Immune Checkpoint Inhibitors

OS: Overall Survival

ABx: Antibiotics

CS: Corticosteroids

PS: Eastern Cooperative Oncology Group Performance Status

BMI: Body Mass Index

NSCLC: Non-Small Cell Lung Cancer

RCC: Renal Cell Carcinoma

HNSC: Head and Neck Squamous Cell Carcinoma

DECLARATIONS

Ethic Approval and Consent to Participate

Data from this work were collected under an IRB-approved retrospective study (OSU-

2016C0070, OSU-2017C0063).

Consent for Publication

Not applicable.

Availability of Data and Materials

Scripts to reproduce all figures and analyses can be found at

https://github.com/spakowiczlab/co-med-io.

Competing Interests

DHO has a consulting or advisory role to L&M Policy Research and AstraZeneca. DHO

has research funding from Bristol-Myers Squibb, Palobiofarma, Merch Sharp & Dohme and

Genentech/Roche. DHO received travel, accommodations and/or expenses from AstraZeneca.

DPC is on the advisory board of Abbvie, AMGEN, Ariad, AZ, Biocept, BI, BMS, Celgene, EMD

Serono, Genentech, Gritstone, Helsinn, Inivata, Inovio, Janssen, Loxo Oncology, Merck, MSD,

Pfizer, Roche, Takeda, and Trinity. DPC received honoraria from AstraZeneca and Nexus

Oncology. DPC is a consultant for Abbvie, Agenus, Ariad, Celgene, Genentech, Incyte, Inivata,

Janssen, Kyowa Kirin, Loxo, MSD, Novartis, Pfizer and Takeda.

Funding

This project was supported by The Ohio State University Center for Clinical and

Translational Science grant support (National Center for Advancing Translational Sciences,

Grant 8UL1TR000090-05) and a P30 CCSG award (2 P30 CA016058-42). Daniel Spakowicz is

supported by a Pelotonia New Investigator Award and Carolyn Presley is a Paul Calabresi

Scholar supported by the OSU K12 Training Grant for Clinical Faculty Investigators (5K12

CA133250-09). No funding body was involved in the design of the study or the collection,

analysis, or interpretation of data, or in writing the manuscript.

Author's Contributions

MH, GT, SP, JTB, CFV, KLK, SH, JP, DQ, GAO, MG, CP and DHO collected the data.

RH, MM, JB, LW, LL, DPC, DHO and DS analyzed the data and generated the figures and

tables. DS, RH and DHO wrote the manuscript. All authors read and approved the final

manuscript.

Acknowledgements

This work was performed on the Ohio Supercomputer Center servers (Owens, Pitzer;

Project PA1490 awarded to Daniel Spakowicz). The authors would like to thank Stephanie

Fortier (Division of Medical Oncology, The Ohio State University; ORCID 0000-0002-2380-

2378) for critically reviewing and editing the manuscript.

REFERENCES

1. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, et al. Gut
microbiome modulates response to anti–PD-1 immunotherapy in melanoma patients.
Science. 2018 Jan 5;359(6371):97–103.

2. Matson V, Fessler J, Bao R, Chongsuwat T, Zha Y, Alegre M-L, et al. The commensal
microbiome is associated with anti–PD-1 efficacy in metastatic melanoma patients.
Science. 2018 Jan 5;359(6371):104–8.

3. Routy B, Le Chatelier E, Derosa L, Duong CP, Alou MT, Daillère R, et al. Gut microbiome
influences efficacy of PD-1–based immunotherapy against epithelial tumors. Science.
2018;359(6371):91–97.

4. Leach DR, Krummel MF, Allison JP. Enhancement of Antitumor Immunity by CTLA-4
Blockade. Science. 1996 Mar 22;271(5256):1734–6.

5. Chen P-L, Roh W, Reuben A, Cooper ZA, Spencer CN, Prieto PA, et al. Analysis of
Immune Signatures in Longitudinal Tumor Samples Yields Insight into Biomarkers of
Response and Mechanisms of Resistance to Immune Checkpoint Blockade. Cancer
Discov. 2016;6(8):827–37.

6. Limagne E, Richard C, Thibaudin M, Fumet J-D, Truntzer C, Lagrange A, et al. Tim-
3/galectin-9 pathway and mMDSC control primary and secondary resistances to PD-1
blockade in lung cancer patients. Oncoimmunology. 2019;8(4):e1564505.

7. Huang AC, Orlowski RJ, Xu X, Mick R, George SM, Yan PK, et al. A single dose of
neoadjuvant PD-1 blockade predicts clinical outcomes in resectable melanoma. Nat Med.
2019;25(3):454–61.

8. Arbour KC, Mezquita L, Long N, Rizvi H, Auclin E, Ni A, et al. Impact of Baseline Steroids
on Efficacy of Programmed Cell Death-1 and Programmed Death-Ligand 1 Blockade in
Patients With Non–Small-Cell Lung Cancer. JCO. 2018 Oct;36(28):2872–8.

9. Rogers MAM, Aronoff DM. The influence of non-steroidal anti-inflammatory drugs on the
gut microbiome. Clinical Microbiology and Infection. 2016 Feb 1;22(2):178.e1-178.e9.

10. Imhann F, Bonder MJ, Vich Vila A, Fu J, Mujagic Z, Vork L, et al. Proton pump inhibitors
affect the gut microbiome. Gut. 2016 May;65(5):740–8.

11. Csardi G, Nepusz T. The igraph software package for complex network research.
InterJournal. 2006;Complex Systems:1695.

12. Textor J, Zander B van der. dagitty: Graphical Analysis of Structural Causal Models
[Internet]. 2016. Available from: https://CRAN.R-project.org/package=dagitty

13. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data
capture (REDCap)--a metadata-driven methodology and workflow process for providing
translational research informatics support. J Biomed Inform. 2009 Apr;42(2):377–81.

14. Kassambara A, Kosinski M. survminer: Drawing Survival Curves using "ggplot2" [Internet].
2018. Available from: https://CRAN.R-project.org/package=survminer

15. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2017. Available from:
https://www.R-project.org/

16. Zeileis A, Hothorn T. Diagnostic checking in regression relationships. 2002 [cited 2017
May 8]; Available from: http://idg.pl/mirrors/CRAN/web/packages/lmtest/vignettes/lmtest-
intro.pdf

17. Therneau TM. A Package for Survival Analysis in S [Internet]. 2015. Available from:
https://CRAN.R-project.org/package=survival

18. Terry M. Therneau, Patricia M. Grambsch. Modeling Survival Data: Extending the Cox
Model. New York: Springer; 2000.

19. Wickham H. ggplot2: elegant graphics for data analysis [Internet]. Springer Science &
Business Media; 2009 [cited 2015 Dec 14]. Available from:
https://books.google.com/books?hl=en&lr=&id=bes-
AAAAQBAJ&oi=fnd&pg=PR5&dq=ggplot2&ots=SAc_My0SQR&sig=kxI4eEM08tk2hMmUo
5UgBJQE4AQ

20. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a
major update to the DrugBank database for 2018. Nucleic Acids Res. 2018
04;46(D1):D1074–82.

21. Gehlenborg N. UpSetR: A More Scalable Alternative to Venn and Euler Diagrams for
Visualizing Intersecting Sets [Internet]. 2019. Available from: https://CRAN.R-
project.org/package=UpSetR

22. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. Journal of Statistical Software. 2010;33(1):1–22.

23. Simon N, Friedman J, Hastie T, Tibshirani R. Regularization Paths for Cox's Proportional
Hazards Model via Coordinate Descent. Journal of Statistical Software. 2011;39(5):1–13.

24. Ricciuti B, Dahlberg SE, Adeni A, Sholl LM, Nishino M, Awad MM. Immune Checkpoint
Inhibitor Outcomes for Patients With Non–Small-Cell Lung Cancer Receiving Baseline
Corticosteroids for Palliative Versus Nonpalliative Indications. JCO. 2019 Jun
17;JCO.19.00189.

25. Nandi T, Lee IR, Ghosh T, Ng AHQ, Chng KR, Li C, et al. Gut microbiome recovery after
antibiotic usage is mediated by specific bacterial species. bioRxiv. 2018 Jun 19;350470.

26.   Pérez-Cobas AE, Gosalbes MJ, Friedrichs A, Knecht H, Artacho A, Eismann K, et al. Gut
microbiota disturbance during antibiotic therapy: a multi-omic approach. Gut. 2013 Nov
1;62(11):1591–601.

27.   Dethlefsen L, Relman DA. Incomplete recovery and individualized responses of the human
distal gut microbiota to repeated antibiotic perturbation. Proceedings of the National
Academy of Sciences. 2011 Mar 15;108(Supplement_1):4554–61.

28.   Zhao S, Gao G, Li W, Li X, Zhao C, Jiang T, et al. Antibiotics are associated with
attenuated efficacy of anti-PD-1/PD-L1 therapies in Chinese patients with advanced non-
small cell lung cancer. Lung Cancer. 2019 Apr;130:10–7.

29.   Elkrief A, El Raichani L, Richard C, Messaoudene M, Belkaid W, Malo J, et al. Antibiotics
are associated with decreased progression-free survival of advanced melanoma patients
treated with immune checkpoint inhibitors. Oncoimmunology. 2019;8(4):e1568812.

30.   Chheda ZS, Sharma RK, Jala VR, Luster AD, Haribabu B. Chemoattractant Receptors
BLT1 and CXCR3 Regulate Antitumor Immunity by Facilitating CD8 [+] T Cell Migration into
Tumors. JI. 2016 Sep 1;197(5):2016–26.

31.   Wu T, Yang L, Jiang J, Ni Y, Zhu J, Zheng X, et al. Chronic glucocorticoid treatment
induced circadian clock disorder leads to lipid metabolism and gut microbiota alterations in
rats. Life Sciences. 2018 Jan;192:173–82.

32.   Garant A, Guilbault C, Ekmekjian T, Greenwald Z, Murgoi P, Vuong T. Concomitant use of
corticosteroids and immune checkpoint inhibitors in patients with hematologic or solid
neoplasms: A systematic review. Critical Reviews in Oncology/Hematology. 2017
Dec;120:86–92.

33.   Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and
resilience of the human gut microbiota. Nature. 2012 Sep 13;489(7415):220–30.

34.   González-Amaro R, Sánchez-Madrid F. Drugs, inflammation and cell adhesion receptors.
Expert Opinion on Pharmacotherapy. 2001 Jan;2(1):3–17.

35.   Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, et al. Extensive impact
of non-antibiotic drugs on human gut bacteria. Nature. 2018 Mar;555(7698):623–8.

36.   Fucà G, Galli G, Poggi M, Lo Russo G, Proto C, Imbimbo M, et al. Modulation of peripheral
blood immune cells by early use of steroids and its association with clinical outcomes in
patients with metastatic non-small cell lung cancer treated with immune checkpoint
inhibitors. ESMO Open. 2019 Feb;4(1):e000457.

37.   Derosa L, Hellmann MD, Spaziano M, Halpenny D, Fidelle M, Rizvi H, et al. Negative
association of antibiotics on clinical activity of immune checkpoint inhibitors in patients with
advanced renal cell and non-small-cell lung cancer. Annals of Oncology. 2018 Jun
1;29(6):1437–44.

38.   Hussain M, Javeed A, Ashraf M, Al-Zaubai N, Stewart A, Mukhtar MM. Non-steroidal anti-
inflammatory drugs, tumour immunity and immunotherapy. Pharmacological Research.
2012 Jul;66(1):7–18.

39.   Wang D, DuBois RN. Prostaglandins and cancer. Gut. 2006 Jan;55(1):115–22.

40.   Jackson MA, Goodrich JK, Maxan M-E, Freedberg DE, Abrams JA, Poole AC, et al. Proton
pump inhibitors alter the composition of the gut microbiota. Gut. 2016 May;65(5):749–56.

41.   Boursi B, Mamtani R, Haynes K, Yang Y-X. Recurrent antibiotic exposure may promote
cancer formation – Another step in understanding the role of the human microbiota?
European Journal of Cancer. 2015 Nov;51(17):2655–64.

42.   Lidar M, Giat E, Garelick D, Horowitz Y, Amital H, Steinberg-Silman Y, et al. Rheumatic
manifestations among cancer patients treated with immune checkpoint inhibitors.
Autoimmunity Reviews. 2018 Mar;17(3):284–9.

43.   Leonard J, Marshall JK, Moayyedi P. Systematic Review of the Risk of Enteric Infection in
Patients Taking Acid Suppression. Am J Gastroenterology. 2007 Sep;102(9):2047–56.

44.   Park JH, McMillan DC, Horgan PG, Roxburgh CS. The impact of anti-inflammatory agents
on the outcome of patients with colorectal cancer. Cancer Treatment Reviews. 2014
Feb;40(1):68–77.

45.   Huang EY, Inoue T, Leone VA, Dalal S, Touw K, Wang Y, et al. Using Corticosteroids to
Reshape the Gut Microbiome: Implications for Inflammatory Bowel Diseases. Inflamm
Bowel Dis. 2015 May;21(5):963–72.

46.   Jala VR, Bodduluri SR, Satpathy SR, Chheda Z, Sharma RK, Haribabu B. The yin and
yang of leukotriene B4 mediated inflammation in cancer. Semin Immunol. 2017;33:58–64.

47.   Zitvogel L, Ayyoub M, Routy B, Kroemer G. Microbiome and Anticancer
Immunosurveillance. Cell. 2016 Apr 7;165(2):276–87.

FIGURES AND TABLES

Figure 1. Causal model for the effect of concomitant medications on Immunotherapy

Response and Overall Survival. Numbers along edges refer to references supporting the

connection. Hypothesized dominant pathways are shown in heavily-weighted edges. (1–3,8–

10,27–47)

Table 1. Cohort characteristics

Overall n	690
BMI (mean (SD))	27.79 (6.66)
Staging (%)	
1	1 (0.2)
2	4 (0.7)
3	44 (7.2)
4	548 (90.0)
Unknown	12 (2.0)
Age (mean (SD))	62.25 (13.21)
Sex = Male (%)	403 (58.4)
Cancer (%)	
Bladder Cancer	32 (4.9)
Head and Neck Carcinoma	42 (6.5)
Melanoma	185 (28.5)
Non-Small Cell Lung Cancer	152 (23.4)
Renal Cell Carcinoma	65 (10.0)
Sarcoma	21 (3.2)
Other	152 (23.4)
Immunotherapy (%)	
Atezolizumab	22 (3.2)
Durvalumab	12 (1.7)
Durvalumab + Tremelimumab	6 (0.9)
Ipilimumab	126 (18.3)
Nivolumab	364 (52.8)
Nivolumab + Ipilimumab	37 (5.4)
Nivolumab + Chemotherapy	7 (1.0)
Pembrolizumab	104 (15.1)
Tremelimumab	3 (0.4)
Other	9 (1.3)
ECOG (%)	
0	185 (31.0)
1	272 (45.6)
2	113 (19.0)
> 2	26 (4.4)
ABx within 28 days of ICI (mean (SD))	0.36 (0.48)
CS within 28 days of ICI (mean (SD))	0.40 (0.49)

556 **Figure 2. The effect of medications at the start of ICI treatment across all cancers for A)**

557 **Antibiotics, B) Corticosteroids, and C) other medications.** The cell color indicates the p-

558 value of the Kaplan-Meier curve and the "+" or "-" the direction of the HR, in reference to its

559 association with OS (i.e. a "-" indicates an association with decreased OS, therefore a HR > 1).

562 **Figure 3. Associations of ABx and CS over time and by drug class.** A) Hazard ratios with

563 95% confidence intervals of a Cox Proportional-Hazards model comparing individuals treated

564 with ABx or CS during a 30-day sliding window compared to indivduals who did not receive ABx

565 or CS, respectively. The significance and direction of associations of Cox Proportional Hazards

566 models by (B) ABx or (C) CS class and cancer, using a window 28 days around ICI treatment

567 start.

Figure 4. Combined models for ABx and CS and controlling for covariates. (A) Kaplan-

Meier curves for ABx and CS in combination. B) Cox Proportional Hazards model incorporating

both ABx and CS as well as several covariates. C) Cox-LASSO models for each cancer

showing the hazard ratios estimated for covariates and the number of times the covariate was

included in the model. The regularization parameter was selection by 10-fold cross validation,

and then the robustness was assessed by 1000 bootstrap replicates using different random

samples of the data.

Figure 5. Relating the ABx effect to microbes enriched in responders to ICIs. A

dendrogram of the microbes recently shown to be most enriched in responders (black) or non-

responders (red), are related to known ABx susceptibilities (references for each cell in **Table**

S1). The ABx are ordered by hazard ratio across all cancers (i.e. β-lactams showed the largest

hazard ratio and linezolid the smallest).

SUPPLEMENTARY FIGURES AND TABLES

Figure S1. Causal diagram with references.

Figure S2. Number of medications prescribed across all cancers and the frequency of

multiple medications.

Figure S3. Number of antibiotics, separated by class, prescribed across all cancers and

the frequency of multiple antibiotics.

Figure S4. Number of corticosteroids, separated by class, prescribed across all cancers

and the frequency of multiple antibiotics.

Table S1. References for the antibiotic susceptibilities of bacterial taxa found to be
enriched in responders or non-responders to ICI therapy