

1 **Basic determinants of child growth in sub-Saharan Africa: cross-sectional survey analysis of**
2 **positive deviants in poor households.**

3

4 Dickson A. Amugsi^{1*}

5 damugsi2002@yahoo.com; damugsi@aphrc.org

6 Zacharie T. Dimbuene^{2,3}

7 zacharie.tsala.dimbuene@gmail.com

8

9 1. African Population and Health Research Center, APHRC Campus, P.O Box 10787-00100,
10 Nairobi, Kenya

11 2. Department of Population Sciences and Development, University of Kinshasa, Democratic
12 Republic of the Congo

13 3. Statistics Canada, Microdata Access Division, 100 Tunney's Pasture Driveway, Ottawa,
14 Canada, K1A 0T6

15

16 * Corresponding author: Dr. Dickson A. Amugsi

17 Email: damugsi2002@yahoo.com; damugsi@aphrc.org

18

19

20

21

22

23

24 **Abstract**

25 **Background:** Childhood malnutrition is a significant public health problem confronting
26 countries across the globe. Nonetheless, recent evidence suggests a downward trend in
27 undernutrition among children globally. Despite the progress made at the global level, sub-
28 Saharan Africa did not experience significant improvement in the past decades. The objective of
29 this study was to investigate the basic determinants associated with linear growth among
30 children under 5 years living in poor households.

31 **Methods:** The study used nationally representative cross-sectional survey data from Ghana,
32 Kenya, Democratic Republic of Congo (DRC), Nigeria and Mozambique. The participants are
33 children aged 0–59 months (N=24,264) living in poor households. The DHS obtained
34 information on children through face-to-face interviews with mothers. The height of the
35 children was also measured and used to compute the height-for-age Z-scores (HAZ). In this
36 study, HAZ is categorised into HAZ>-2 standard deviations (SD) (not stunted/better growth) and
37 HAZ<-2 SD (stunted/poor growth).

38 **Results:** A unit change in maternal years of education was associated with increased odds of
39 better growth among children living in poor households in DRC [adjusted odds ratio (aOR)=
40 1.03, 95% CI=1.01,1.07)], Ghana (aOR=1.06, 95% CI=1.01,1.11), Kenya (aOR=1.03, 95% CI= 1.01,
41 1.05) and Nigeria (aOR=1.08, 95%=1.06,1.10). Maternal antenatal attendance of at least four
42 visits was associated positively with better child growth in DRC (aOR=1.32, 95% CI=1.05, 1.67)
43 and Ghana (aOR=1.67, 95% CI=1.19, 2.33). In Ghana, Mozambique and DRC, breastfeeding was
44 associated significantly with the likelihood of better linear growth when only socio-
45 demographic correlates were included in the models but disappeared after the inclusion of
46 child-level covariates. In Nigeria, normal maternal weight was associated with increased odds
47 (aOR=1.24, 95% CI=1.08, 1.43) of positive growth among children living in poor households, so
48 was overweight (aOR=1.51, 95% CI= 1.24, 1.83). In all the countries except Ghana, child
49 biological factors such as sex and age were associated with reduced odds of better growth.

50 **Conclusions:** The socio-demographic factors included in this analysis have the potential to
51 promote linear growth of children under 5 years living in poor households. Interventions aimed

52 at fostering linear growth among children living in poverty should target at enhancing these
53 factors.

54 **Keywords:** child growth, sub-Saharan Africa, positive deviance, cross sectional survey

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74 Introduction

75 Childhood undernutrition is a significant public health problem confronting countries across the
76 globe. However, there is evidence that global trends in child malnutrition have improved over
77 the years. For example, it is estimated that childhood stunting (short stature for age), a critical
78 undernutrition metric, decreased from 39.7% in 1990 to 26.7% in 2010 (1). The trend is
79 predicted to reduce to 22% in 2020 (1). Despite global-level progress in the reduction of the
80 problem during the past decades (2), Africa has not seen much improvement. There was a
81 decline in stunting trends from 40.5% in 1980 to 35.2% in 2000 (3), the trends stagnated at 40%
82 between 1990 and 2010 (1). The level of decline in sub-Saharan Africa (SSA) is not different
83 from the African region as a whole. For example, in SSA, the stunting trends decreased from
84 43% in 2000 to 34% in 2018 (4). The risk factors contributing to the high stunting prevalence in
85 Africa are well documented (5-9).

86

87 The consequences of stunting on the later life of the child are well known. Indeed, there is
88 strong evidence that stunting can have long-term effects on cognitive development, school
89 achievement, economic productivity in adulthood and maternal reproductive outcomes (10-
90 13). Stunting is also a condition that may be very difficult to reverse (10). Given the negative
91 consequences of stunting on child health outcomes, the international community has paid
92 considerable attention to the problem. For instance, the World Health Assembly (WHA)
93 Resolution (2012) set a 40% reduction in the number of stunted children under-5 as one of the
94 six global nutrition targets for 2025 (11, 14). The Sustainable Development Goals (SDGs) (15)
95 also captured this undernutrition metric as a critical developmental target. The above
96 discussion suggests the need for substantial investment in nutrition interventions to address
97 childhood stunting, as averting stunting could produce life-long benefits. However, to achieve
98 this goal, a clear understanding of the broader factors that promote child growth is necessary
99 to provide evidence for the design of effective nutrition interventions. The present study is set
100 out to provide this evidence by focusing on factors that foster child growth rather than risk
101 factors of child growth deficiencies.

102

103 The evidence further suggests a disproportionate burden of stunting among children in low and
104 middle-income countries, which is attributable to poverty, lack of food and high incidence of
105 infectious diseases among others (3, 11, 16, 17). In sub-Saharan Africa (SSA), several countries
106 are confronted with a high prevalence of stunting among children under five years of age (3, 11,
107 17). The problem is notably more severe among children living in poor households (17-19)--they
108 tend to have the highest prevalence of childhood stunting (17). It is the case because poverty
109 creates conditions that favour poor child growth outcomes and prevents affected populations
110 from obtaining adequate access to prevention and care (17). Despite the health challenges
111 facing children living in poverty, some children are living in the same conditions (*positive*
112 *deviants*) or even worse yet have positive growth outcomes comparable to children residing in
113 privilege households anywhere in the world (20-23). The questions this paper intends to
114 address is why some children in poor households are not stunted, although they are faced with
115 similar adversity as those who are stunted are? What are the possible factors that help them to
116 have better growth outcomes? Understanding this will help design programmes to promote the
117 growth of children in impoverished households or environments.

118

119 The concept of *positive deviance* (as referenced above) is based on the premise is that even in
120 places where deprivation is severe and widespread, some families are able to cope and harness
121 scant resources, sufficient to support optimal child development (24, 25). The positive deviance
122 (PD) approach is founded on idea that problems can be overcome using solutions that already
123 exist within the community (26, 27). PD often studies the behaviours and characteristics of
124 individuals who have better health outcomes than their peers who live in the same community
125 (26). The PD approach was used previously to investigate several health-related issues in
126 diverse settings (28-36). In statistical analysis, it is often quantified as those who do not
127 experience a negative outcome of interest compared to those around them with the same
128 resources (26). Using the PD approach can be useful because it studies the 'positive' aspects of
129 an outcome or community instead of the 'negative', and can identify potential points of

130 intervention. The *positive deviants* in the present study are children who live in poor
131 households and yet are not stunted relative to their counterparts who live in the same
132 environment but are stunted. The objective of the present study is to examine the basic factors
133 associated with better growth outcomes among children living in poor households. This
134 resource-focused approach moves away from the dominant risk model approach, where the
135 focus is usually on risk factors of child growth deficiencies. Using the PD approach is to help
136 understand the drivers of better child growth and interventions to promote these drivers in
137 poor households effectively.

138

139 **Methodology**

140 *Data sources and sampling strategy*

141 We analysed the Demographic and Health Surveys (DHS) (37) data from Ghana, Kenya, Nigeria,
142 Mozambique and Democratic Republic of Congo (DRC) (38, 39). The DHS are nationally
143 representative surveys conducted every five years in lower- and middle-income countries, using
144 the same questionnaires to enable comparison across countries (40, 41). The DHS utilises a
145 two-stage sample design. The detail description of the DHS design and sampling strategies can
146 be found elsewhere (42-46). The DHS data collectors then interviewed all eligible study
147 participants in their respective households of each country using standardised questionnaires
148 and interview protocols. In this analysis, we used data of children aged 0-59 months and their
149 mothers aged 15-49 years. The DHS obtained children data through face-to-face interviews
150 with their mothers or caregivers. The length/height of the children was measured using an
151 adjustable measuring board calibrated in millimeters. Recumbent length (lying down on the
152 board) was measured for young children while standing height was measured for older
153 children. The height data were converted into Z-scores based on the 2006 WHO growth
154 standards, taking into account the age and sex of the child (47). The total samples used in the
155 present analysis were: Ghana, n= 1,453; Nigeria, n= 10,378; Kenya, n= 4,967; Mozambique, n=
156 3,487; and DRC, n= 3,979.

157

158 **Ethics statement**

159 The government recognized Ethical Review Committees of the respective countries approved
160 the DHS study protocols. Besides, the ethical clearance was granted by the Institutional Review
161 Board of ICF International, USA before the surveys were conducted. Informed consent was
162 obtained from the mothers of the study children before they were included in the study. The
163 DHS Program permitted the authors to use the data. The data were wholly anonymised, and
164 therefore, the authors did not seek further ethical clearance before their use.

165

166 ***Outcome and predictor variables***

167 *Outcome Variables*

168 We used the child height-for-age Z-scores (HAZ) as the indicator of child linear growth in the
169 analysis. For all datasets, HAZ scores were computed using 2006 WHO growth standards (47).
170 We reclassified the child HAZ into stunted (poor growth) and not stunted (better growth).
171 Children who have HAZ above -2 SD ($HAZ > -2SD$) (47, 48) were considered having a better linear
172 growth and described in this study as positive deviants. Similarly, children who have HAZ below
173 -2 SD ($HAZ < -2$) from the median HAZ of the WHO reference population (47) were considered
174 stunted or having poor growth. It is significant to underscore that DHS data contained all the
175 three indicators of child nutritional status: height-for-age z-scores (HAZ), weight-for-age z-
176 scores (WAZ) and weight-for-height z-scores (WHZ). However, we opted for HAZ because it is a
177 cumulative indicator of a child nutritional status, and therefore more informative and
178 appropriate for use in the positive deviance analysis. The WHZ, on the other hand, reflects
179 more recent processes often associated with acute food shortages and/or illnesses leading to
180 weight loss, while the WAZ lies between HAZ and WHZ. For example, a child who has poor HAZ
181 is also likely to be underweight, so is a child who has poor WHZ.

182

183 We stratified the analysis by household wealth index (WI). The WI has been used severally as an
184 indicator for measuring inequalities associated with health outcomes as well as expenditure

185 and income among households (41, 42, 44, 49). The detail discussion on how the DHS created
186 the WI is well documented (40, 41, 44). In the datasets, the WI is classified into five quintiles:
187 poorest, poor, middle, richer and richest(40, 41, 44). In this paper, we recoded poor and
188 poorest into poor/worse-off households. We combined the poorest and poor households
189 wealth quintiles because the literature suggests that children in these households have similar
190 health outcomes (17-19). We restricted all the analyses to children living in these households.

191 *Analytical framework*

192 The modified UNICEF conceptual framework underpinned our analysis (50, 51). This framework
193 outlines how the various factors/determinants influence child survival, growth and
194 development at different levels. These factors are analysed in terms of immediate, underlying
195 and basic determinants. The immediate determinants are adequate nutrients intake and health,
196 while the underlying determinants are food security, care for children and women, healthcare
197 and a healthy environment (51). The underlying determinants either influence child health
198 directly or through the immediate determinants. The basic determinants, in turn, influence the
199 underlying determinants. In this context, the basic determinants are described as “exogenous”
200 factors, which effects on child nutrition are through their effects on the intervening
201 proximate/underlying determinants. Thus, the underlying determinants are endogenously
202 determined by the exogenous determinants (52). For example, the effect of an exogenous
203 variable such as maternal education on child growth outcomes is through its impact on good
204 child-caring practices, including high utilisation of health care services, among others by
205 educated mothers.

206

207 **Data analysis**

208 The present empirical analysis focused mainly on the basic determinants (i.e. socio-
209 demographic factors). The scientific basis for this type of analysis is well documented (52-55).
210 Apart from the socio-demographic factors, variables such as antenatal care (ANC) and
211 breastfeeding practices, which depend mostly on exogenous public health provisions (52) were
212 included in our empirical models. The significance for the inclusion these two variables is that

213 improvement or otherwise of ANC and breastfeeding practices are more likely to inform
214 policies, programmes and interventions rather than changes in socio-demographic
215 endowments of households (52). For example, the available data suggest that policy,
216 institutional and contextual settings are critical determinants of the prevalence of
217 breastfeeding practices (52, 56). In the analysis, we built two regression models for each of the
218 five countries. In the first model, we included the various socio-demographic factors [maternal
219 body mass index (BMI), education, age, work status, parity, breastfeeding practices, marital
220 status, ANC, the gender of head of household, size of household, total children under five years
221 and place of residence]. We adjusted for child dietary diversity (DD)—the details of how the DD
222 is created can be found elsewhere (38), age and sex in the second and final model. The
223 conceptual framework and the literature guided the selection of the explanatory variables (51).
224 We estimated adjusted odds ratio (aORs) of the effects of the basic determinants on child
225 growth in poor households.

226

227 **Results**

228 *Characteristics of study samples*

229 Tables 1 presents the results of the descriptive analysis. The results showed that Ghana (76%)
230 has the highest number of children with better growth followed by Kenya (68%), while in
231 Mozambique, DRC and Nigeria, the prevalence ranged from 50% to 52%. Regarding dietary
232 diversity intake, Mozambique had the highest prevalence of children who consumed at least
233 four food groups (24%), with DRC (6%) and Nigeria (6%) having the lowest prevalence. Similarly,
234 Mozambique had the highest number of women with normal weight (85%). The prevalence
235 ranged from 68% to 76% in DRC, Ghana, Kenya and Nigeria. For maternal education, Ghana has
236 the highest prevalence (23%) of women who had attained a secondary school education, while
237 Mozambique has the lowest prevalence (1.20%). Higher education was less than 1% among
238 women in poor households across all countries. Regarding antenatal attendance among
239 women, DRC registered the highest prevalence (77%) followed by Ghana (59%), while Nigeria
240 registered the lowest prevalence (19%).

241

242

Table 1: Characteristics of the study samples of the five countries

Variables	DRC		Ghana		Kenya		Mozambique		Nigeria	
	%/mean	SD	%/mean	SD	%/mean	SD	%/mean	SD	%/mean	SD
Child-level covariates										
Height-for-age (HAZ)	50.0		76.0		68.0		52.0		51.0	
DD < 4 food groups	94.0		90.6		88.5		76.4		93.9	
DD >= 4 food groups	6.0		9.4		11.5		23.6		6.1	
Sex of child										
Female	50.4		48.8		50.4		51.3		50.6	
Male	49.6		51.2		49.6		48.7		49.4	
Mother-level covariates										
Body Mass Index (BMI)										
BMI <18.50	15.8		7.4		17.0		7.99		12.4	
BMI = 18.50-24.99	76.3		73.3		68.0		84.6		74.4	
BMI = 25-29.99	6.7		15.7		12.2		6.91		10.5	
BMI >= 30	0.40		3.48		2.83		0.03		2.17	
Education										
No education	30.6		54.0		34.4		52.6		75.3	
Primary education	50.4		22.8		55.6		46.3		17.0	
Secondary education	19.0		23.2		9.43		1.17		7.6	
Higher education	0.05		0.1		0.1		na		0.1	
Working status										
Not working	20.0		16.5		45.9		61.5		35.3	
Is working	79.9		83.3		54.0		38.5		64.4	
Parity	4.44	2.46	4.21	2.3	4.36	2.46	4.37	2.43	4.83	2.75
Is Breastfeeding	73.5		65.0		64.4		68.7		63.9	
Marital status										
Not in union	11.4		11.1		13.7		14.3		3.2	
Married	65.2		69.4		81.2		68.8		95.9	
Cohabiting	23.3		19.5		5.1		16.9		0.1	
Number of antenatal visits >=4	77.2		59.2		31.1		28.3		18.8	

Household-level covariates

Sex of household head

Household head is Female	24.1		18.3		34.4		28.0		6.5	
Household head is Male	75.9		81.7		65.6		72		93.5	
Household size	6.4	2.55	6.67	3.14	6.22	2.35	5.97	2.5	7.63	3.42
Number of children under 5	2.24	0.98	1.98	0.99	2.02	0.87	2.03	0.91	2.5	1.22

Community-level covariates

Urban residence	9.32		11.6		15.2		9.19		8.1	
-----------------	------	--	------	--	------	--	------	--	-----	--

DD= Dietary diversity; DRC= Democratic Republic of Congo; SD= Standard deviation

Multivariable results of the association between socio-demographic factors and better child growth

Tables 2-6 present the results of the association between sociodemographic factors at the child-level, maternal-level, household-level and community-level, and better linear growth among children in five SSA countries. The results showed that a unit change in maternal years of education was associated with increased odds of better linear growth among children in DRC (aOR=1.03, 95% CI=1.01,1.07), Ghana (aOR=1.06, 95% CI=1.01,1.11), Kenya (aOR=1.03, 95% CI=1.01, 1.05) and Nigeria (aOR=1.08, 95%=1.06,1.10). Antenatal attendance of at least four visits was associated significantly with the likelihood of better child growth in DRC (aOR=1.32, 95% CI=1.05, 1.67) and Ghana (aOR=1.67, 95% CI=1.19, 2.33). The association did not reach statistical significance in the remaining three countries. In Kenya, children of mothers who were working and live in poor households had 23% reduced odds of better growth (aOR= 0.77, 95% CI=0.66, 0.91) relative to children of non-working mothers. In Nigeria, Mozambique and DRC, breastfeeding was positively associated with better child growth, but this association disappeared after the child level covariates were included in the model. Urban place of residence was associated with 28% reduced odds of better child growth (aOR=0.72, 95% CI=0.55, 0.95) in Mozambique, and increased odds in Nigeria (aOR=1.58, 95% CI= 1.33, 1.87). In Nigeria, normal maternal weight (BMI) was associated with increased odds (aOR=1.24, 95% CI=1.08, 1.43) of better child growth. Maternal overweight was also associated with increased odds (aOR=1.51, 95% CI= 1.24, 1.83) of better child growth in Nigeria. A unit change in household size was associated with increased odds (aOR=1.05, 95% CI= 1.01, 1.10) of better child growth. Maternal parity reduces the odds of better child growth (aOR=0.95, 95% CI=0.92, 0.98) in Nigeria. In all the countries except Ghana, child biological factors such as sex and age were associated with reduced odds of better child growth.

Table 2: Multivariable analysis of the effects of socio-demographic factors on better linear growth among children living in poor households in DRC

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m ²) = 18.50 - 24.99	0.972 (0.758 - 1.247)	0.979 (0.753 - 1.274)
BMI (kg/m ²) = 25 - 29.99	0.913 (0.604 - 1.379)	0.861 (0.551 - 1.348)
BMI (kg/m ²) >= 30	0.310* (0.0795 - 1.207)	0.301 (0.0558 - 1.620)
Maternal education (in single years)	1.030** (1.001 - 1.060)	1.034** (1.003 - 1.065)
Age of the mother (in years)	0.994 (0.972 - 1.016)	1.012 (0.988 - 1.037)
Working status = Is working	0.840 (0.671 - 1.052)	0.873 (0.684 - 1.113)
Parity	0.995 (0.934 - 1.060)	0.989 (0.923 - 1.061)
Is Breastfeeding = YES	1.379*** (1.110 - 1.712)	0.813* (0.637 - 1.036)
Marital Status = Married	0.880 (0.648 - 1.196)	0.937 (0.677 - 1.297)
Marital Status = Cohabiting	0.986 (0.704 - 1.379)	1.036 (0.731 - 1.469)
Number of antenatal visits = 4+ visits	2.125*** (1.710 - 2.641)	1.321** (1.046 - 1.668)
Household-level covariates		
Head of HH is Male	0.957 (0.759 - 1.206)	0.936 (0.741 - 1.183)
Household size	1.014 (0.965 - 1.065)	1.005 (0.954 - 1.058)
Number of children under 5 years	1.018 (0.904 - 1.147)	1.068 (0.943 - 1.210)
Community-level covariates		
Urban residence = Urban	0.989 (0.740 - 1.322)	0.972 (0.720 - 1.312)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.049 (0.686 - 1.602)
Age of the child (in months)		0.959*** (0.953 - 0.965)
Sex of child = Male		0.773*** (0.639 - 0.935)
Observations	3,979	3,979

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index

*** p<0.01, ** p<0.05, * p<0.1

Table 3: Multivariable analysis of the effects of socio-demographic factors on better linear growth among children living in poor households in Ghana

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m ²) = 18.50 - 24.99	0.944 (0.554 - 1.608)	0.946 (0.554 - 1.617)
BMI (kg/m ²) = 25 - 29.99	1.702 (0.888 - 3.262)	1.727* (0.902 - 3.305)
BMI (kg/m ²) >= 30	2.048 (0.739 - 5.677)	2.183 (0.783 - 6.089)
Maternal education (in single years)	1.059** (1.012 - 1.107)	1.057** (1.010 - 1.106)
Age of the mother (in years)	1.004 (0.966 - 1.044)	1.009 (0.969 - 1.050)
Working status = Is working	0.772 (0.509 - 1.170)	0.798 (0.524 - 1.214)
Parity	1.009 (0.902 - 1.129)	1.007 (0.899 - 1.129)
Is Breastfeeding = YES	1.705*** (1.206 - 2.410)	1.412* (0.975 - 2.045)
Marital Status = Married	1.316 (0.761 - 2.278)	1.300 (0.755 - 2.239)
Marital Status = Cohabiting	0.994 (0.539 - 1.831)	0.950 (0.516 - 1.747)
Number of antenatal visits = 4+ visits	2.004*** (1.464 - 2.743)	1.667*** (1.193 - 2.329)
Household-level covariates		
Head of HH is Male	0.889 (0.549 - 1.440)	0.911 (0.566 - 1.468)
Household size	0.987 (0.928 - 1.050)	0.983 (0.923 - 1.047)
Number of children under 5	0.946 (0.774 - 1.156)	0.942 (0.771 - 1.150)
Community-level covariate		
Urban residence = Urban	1.239 (0.735 - 2.087)	1.224 (0.733 - 2.046)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.281 (0.765 - 2.146)
Age of the child (in months)		0.989* (0.979 - 1.000)
Sex of child = Male		0.850 (0.624 - 1.159)
Observations	1,453	1,453

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index

*** p<0.01, ** p<0.05, * p<0.1

Table 4: Multivariable analysis of the effects of socio-demographic factors on better linear growth among children living in poor households in Kenya

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m ²) = 18.50 - 24.99	0.894 (0.716 - 1.116)	0.896 (0.716 - 1.122)
BMI (kg/m ²) = 25 - 29.99	1.234 (0.910 - 1.674)	1.255 (0.923 - 1.706)
BMI (kg/m ²) >= 30	0.982 (0.580 - 1.663)	0.971 (0.579 - 1.630)
Maternal education (in single years)	1.028** (1.006 - 1.051)	1.029** (1.006 - 1.052)
Age of the mother (in years)	1.026*** (1.006 - 1.046)	1.029*** (1.009 - 1.050)
Working status = Is working	0.760*** (0.646 - 0.896)	0.774*** (0.656 - 0.914)
Parity	0.965 (0.911 - 1.023)	0.962 (0.907 - 1.020)
Is Breastfeeding = YES	1.324*** (1.116 - 1.571)	1.182* (0.978 - 1.429)
Marital Status = Married	0.994 (0.777 - 1.270)	1.019 (0.796 - 1.303)
Marital Status = Cohabiting	0.951 (0.639 - 1.417)	0.967 (0.647 - 1.443)
Number of antenatal visits = 4+ visits	1.288*** (1.084 - 1.531)	1.173* (0.978 - 1.407)
Household-level covariates		
Head of HH is Male	1.041 (0.873 - 1.242)	1.037 (0.868 - 1.239)
Household size	0.965* (0.924 - 1.007)	0.959* (0.919 - 1.002)
Number of children under 5	0.976 (0.878 - 1.084)	0.984 (0.884 - 1.095)
Community-level covariate		
Urban residence = Urban	1.111 (0.900 - 1.373)	1.111 (0.898 - 1.375)
Child-level covariates		
Dietary Diversity (DD) >= 4		0.914 (0.720 - 1.161)
Age of the child (in months)		0.991*** (0.987 - 0.996)
Sex of child = Male		0.717*** (0.615 - 0.836)
Observations	4,967	4,967

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index

*** p<0.01, ** p<0.05, * p<0.1

Table 5: Multivariable analysis of the effects of socio-demographic factors on better linear growth among children living in poor households in Mozambique

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m ²) = 18.50 - 24.99	1.305* (0.968 - 1.760)	1.328* (0.985 - 1.789)
BMI (kg/m ²) = 25.00 - 29.99	1.201 (0.792 - 1.821)	1.225 (0.806 - 1.863)
BMI (kg/m ²) >= 30	1.503 (0.389 - 5.810)	1.528 (0.417 - 5.603)
Maternal education (in single years)	1.030 (0.990 - 1.072)	1.031 (0.990 - 1.073)
Age of the mother (in years)	1.012 (0.994 - 1.029)	1.017* (0.999 - 1.036)
Working status = Is working	0.938 (0.798 - 1.102)	0.936 (0.795 - 1.102)
Parity	0.988 (0.935 - 1.045)	0.993 (0.939 - 1.050)
Is Breastfeeding = YES	1.182* (0.991 - 1.411)	0.968 (0.798 - 1.173)
Marital Status = Married	0.941 (0.715 - 1.237)	0.923 (0.701 - 1.216)
Marital Status = Cohabiting	0.972 (0.710 - 1.330)	0.967 (0.706 - 1.326)
Number of antenatal visits = 4+ visits	1.182* (0.990 - 1.411)	1.001 (0.831 - 1.207)
Household-level covariates		
Head of HH is Male	1.126 (0.912 - 1.390)	1.124 (0.910 - 1.388)
Household size	1.067*** (1.018 - 1.118)	1.053** (1.005 - 1.104)
Number of children under 5	1.006 (0.894 - 1.132)	1.040 (0.923 - 1.171)
Community-level covariate		
Urban residence = Urban	0.709** (0.540 - 0.931)	0.721** (0.550 - 0.947)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.169 (0.968 - 1.413)
Age of the child (in months)		0.985*** (0.980 - 0.990)
Sex of child = Male		0.743*** (0.635 - 0.870)
Observations	3,487	3,487

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index

*** p<0.01, ** p<0.05, * p<0.1

Table 6: Multivariable analysis of the effects of socio-demographic factors on better linear growth among children living in poor households in Nigeria

Variables	Model 1	Model 2
Mother-level covariates		
BMI (kg/m ²) = 18.50 - 24.99	1.231*** (1.073 - 1.413)	1.241*** (1.080 - 1.428)
BMI (kg/m ²) = 25.00 - 29.99	1.484*** (1.228 - 1.794)	1.508*** (1.243 - 1.828)
BMI (kg/m ²) >= 30	1.216 (0.874 - 1.693)	1.221 (0.868 - 1.720)
Maternal education (in single years)	1.072*** (1.052 - 1.092)	1.076*** (1.056 - 1.096)
Age of the mother (in years)	1.012** (1.001 - 1.023)	1.018*** (1.007 - 1.030)
Working status = Is working	1.056 (0.960 - 1.162)	1.081 (0.981 - 1.191)
Parity	0.952*** (0.925 - 0.979)	0.950*** (0.923 - 0.978)
Is Breastfeeding = Yes	1.320*** (1.199 - 1.453)	1.035 (0.933 - 1.149)
Marital Status = Married	0.900 (0.685 - 1.183)	0.923 (0.702 - 1.213)
Marital Status = Cohabiting	1.133 (0.646 - 1.987)	1.107 (0.622 - 1.970)
Number of antenatal visits = 4+ visits	1.354*** (1.201 - 1.525)	1.081 (0.955 - 1.224)
Household-level covariates		
Head of HH is Male	0.955 (0.782 - 1.168)	0.953 (0.778 - 1.168)
Household size	0.991 (0.971 - 1.011)	0.986 (0.966 - 1.007)
Number of children under 5	1.017 (0.964 - 1.072)	1.040 (0.986 - 1.097)
Community-level covariate		
Urban residence = Urban	1.518*** (1.283 - 1.796)	1.575*** (1.325 - 1.872)
Child-level covariates		
Dietary Diversity (DD) >= 4		1.148 (0.941 - 1.401)
Age of the child (in months)		0.981*** (0.978 - 0.984)
Sex of child = Male		0.843*** (0.771 - 0.923)
Observations	10,378	10,378

95% Confidence Intervals (CIs) in parentheses; DD-Dietary diversity; HH-Household; BMI-Body mass index

*** p<0.01, ** p<0.05, * p<0.1

Discussion

The study examined the socio-demographic factors associated with better child growth in poor households in five sub-Saharan African countries. We utilised a *positive deviance* approach as our analytical lens, whereby children who were not stunted though living in poor households were considered *positive deviants*. The findings showed that the effects of socio-demographic factors on child growth vary across countries. Maternal higher years of education had a significant positive effect on better linear growth among children living in poverty in DRC, Ghana, Kenya and Nigeria. This finding suggests that maternal education could mitigate the adverse effects of poverty on children's nutritional status. Thus, education is an essential resource for improving child growth outcomes in the face of adversity. It is possibly the case because our conceptual framework (51) suggests that an exogenous variable such as education has a direct effect on caring practices, adequate dietary intake, utilisation of health care service and healthy environment among others. The proximate factors mentioned above, in turn, have direct effects on positive child growth outcomes. The literature on the effects of mothers education on child-caring practices and the utilisation of health services and the consequential positive effect on child health outcomes abound (57-60). Our study findings are consistent with the existing literature (59, 61). A study using data from three SSA countries showed that a higher level of maternal education was associated with reduced odds of child stunting (61). The literature, together with the present study, though using slightly different analytical approaches, demonstrated the importance of education in improving child growth outcomes.

Further, the results showed that in DRC and Ghana, mothers who attended at least four antenatal visits (ANC) have children with better linear growth outcomes. This could be due to the fact that mothers who attend ANC are likely to receive health and nutrition education, which may have a positive impact on their caring practices, with its consequential effect on better child health outcomes. The preceding explanation is in line with the conceptual framework used in this study, that posits that the proximate factors are pathways through which the exogenous factors influence child growth outcomes (51). Our findings are similar to others by previous researchers. For example, Kuhnt and Vollmer (62) found that having at least

four ANC visits was associated with a reduced risk of stunting in pre-school children. Our findings, together with the literature, suggest that promoting ANC attendance among women can have a beneficial effect not only on the mothers but also their offspring. Therefore, interventions to promote child growth in poor environments should incorporate ANC as a critical intervention package.

Our analysis also illuminated the widely recognised benefits of breastfeeding for improved child health and developmental outcomes (63-65), but only when child-level covariates were not included in the empirical regression models. For instance, in Ghana, Mozambique and DRC, breastfeeding showed a significant positive effect on better childhood growth in the models containing only the socio-demographic factors. However, this significant association disappeared after child-level covariates such as dietary diversity, age and sex were included. This finding may mean that whether breastfeeding will have a positive effect on child growth or not depends to some extent on the inclusion or otherwise of child-level covariates. Therefore, in examining the effects of socio-demographic factors on child linear growth, it is significant to include child-level covariates to avoid presenting misleading estimates (66). The non-significant association between breastfeeding on child growth has previously been documented (60, 66, 67). Indeed, Marquis and colleagues (66) observed an inverse relationship between breastfeeding and child linear growth. They attributed this inverse relationship to what they termed *reverse causality*—that is, the breastfeeding did not lead to poor growth, but poor growth and health led to increased breastfeeding.

Surprisingly, in Mozambique, the widely recognised urban advantage in terms of favourable health outcomes was not observed in the present study. The analysis showed that urban place of residence associated negatively with child linear growth in poor households. The reason for this inverse relationship could be attributed to the precarious conditions under which some of the urban poor live (68). In the literature, both negative and positive effects have been found with the urban place of residence and child growth outcomes (68, 69). Some previous studies

have observed that urban children are usually taller and heavier (69, 70). However, this may not include those children in poor urban settings, as there is evidence that children in these settings tend to have shorter heights than expected (68). This finding may mean that the so-called urban advantage does not benefit the urban poor in Mozambique.

Strengths and limitations of the study

The use of large nationally representative samples provides more robust estimates of observed associations as well as enhancing the generalisability of the findings. The analysis of multi-country data helps to illuminate differences and highlights commonalities in the effects of the determinants on child growth across countries. The outcome variable was objectively measured, thereby reducing potential biases. The novelty of this analysis is the focus on factors that promote better child growth rather than risks factors for child growth deficiencies. A limitation worth mentioning is the cross-sectional nature of the data, which does not lend itself to the establishments of a causal relationship between the predictor and outcome variables. The conclusions in the paper are, therefore interpreted as mere associations between the predictor variables and the outcome variable. The use of quantitative data to investigate the PD approach may be a bit limiting as it may not be possible to explore all PD behaviours quantitatively. This limitation Notwithstanding, PD is a well-established concept and hence makes it possible to explore the approach (PD) using quantitative data.

Conclusions

The study examined the effects of child, maternal, household and community-level socio-demographic factors on better linear growth among children in five SSA countries. The results showed that the effects of socio-demographic factors on child linear growth vary across countries. Maternal education has a positive effect on better growth among children in all countries except Mozambique. Improving maternal education in poor households may have a beneficial effect on child growth outcomes. A higher number of ANC visits has a significant positive effect on better child growth. Interventions to promote linear growth among children living in poverty should incorporate ANC as one of the critical intervention packages.

Declarations

Consent for publication

The study used completely anonymous secondary data in the analysis. Therefore, no consent for publication was required

Availability of data and materials

This study was a re-analysis of existing data that are publicly available from The DHS Program at <http://dhsprogram.com/publications/publication-fr221-dhs-final-reports.cfm>. Data are accessible free of charge upon registration with the Demographic and Health Survey program (The DHS Program). The registration is done on the DHS website indicated above.

Competing Interest

The authors have no competing interests to declare.

Funding

This study did not receive funding from any source.

Authors' Contribution

DAA conceived and designed the study, interpreted the results, wrote the first draft of the manuscript, and contributed to revision of the manuscript. DAA and ZTD analysed the data. ZTD contributed to study design, data interpretation, and critical revision of the manuscript. All authors take responsibility for any issues that might arise from the publication of this manuscript.

Acknowledgements

We wish to express our profound gratitude to The DHS Program, USA for providing us access to the data. We also wish to acknowledge institutions of respective countries that played critical roles in the data collection process.

References

1. de Onis M, Blössner M, Borghi E. Prevalence and trends of stunting among pre-school children, 1990–2020. *Public health nutrition*. 2011;15(1):142-8.
2. de Onis M, Blossner M, Borghi E, Morris R, Frongillo EA. Methodology for estimating regional and global trends of child malnutrition. *International journal of epidemiology*. 2004;33(6):1260-70.
3. de Onis M, Frongillo EA, Blossner M. Is malnutrition declining? An analysis of changes in levels of child malnutrition since 1980. *Bull World Health Organ*. 2000;78(10):1222-33.
4. UNICEF / WHO / World Bank Group. Levels and trends in child malnutrition: Joint Child Malnutrition Estimates 2019 [Available from: <https://www.who.int/nutgrowthdb/jme-2019-key-findings.pdf?ua=1>].
5. Danaei G, Andrews KG, Sudfeld CR, Fink G, McCoy DC, Peet E, et al. Risk Factors for Childhood Stunting in 137 Developing Countries: A Comparative Risk Assessment Analysis at Global, Regional, and Country Levels. *PLoS medicine*. 2016;13(11):e1002164.
6. Mosites E, Dawson-Hahn E, Walson J, Rowhani-Rahbar A, Neuhaus ML. Piecing together the stunting puzzle: a framework for attributable factors of child stunting. *Paediatr Int Child Health*. 2017;37(3):158-65.
7. Prado EL, Yakes Jimenez E, Vosti S, Stewart R, Stewart CP, Somé J, et al. Path analyses of risk factors for linear growth faltering in four prospective cohorts of young children in Ghana, Malawi and Burkina Faso. *BMJ Global Health*. 2019;4(1):e001155.
8. Budge S, Parker AH, Hutchings PT, Garbutt C. Environmental enteric dysfunction and child stunting. *Nutrition Reviews*. 2019;77(4):240-53.
9. Black RE, Allen LH, Bhutta ZA, Caulfield LE, de Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*. 2008;371(9608):243-60.
10. Muller O, Krawinkel M. Malnutrition and Health in Developing Countries *Canadian Medical Association Journal*. 2005;171:279–93.
11. de Onis M, Blossner M, Borghi E. Prevalence and trends of stunting among pre-school children, 1990-2020. *Public health nutrition*. 2012;15(1):142-8.
12. Delisle H. Early nutritional influences on obesity, diabetes and cardiovascular disease risk. *International Workshop, 6-9 June 2004, Montreal University, Quebec, Canada. Maternal and Child Nutrition*. 2005;1(3):128-9.
13. Kar BR, Rao SL, Chandramouli BA. Cognitive development in children with chronic protein energy malnutrition. *Behav Brain Funct*. 2008;4:31-.
14. WHO. Global nutrition targets 2025: stunting policy brief (WHO/NMH/NHD/14.3). Geneva: World Health Organization; 2014.
15. UN. Sustainable Development Goals 2015 [cited 2019 23.08]. Available from: <http://www.un.org/sustainabledevelopment/sustainable-development-goals/>.
16. WHO. World Health Organization global data base on child growth and malnutrition Geneva2011 [cited 2019 23.08]. Available from: <http://www.who.int/nutgrowthdb/database/countries/gha/en/>.
17. de Poel V, Hosseinpoor RA, Jehu-Appiah C, Vega J, Speybroeck N. Malnutrition and the disproportionate burden on the poor: the case of Ghana. *Int J Equity Health*. 2007;6(21).
18. Hong R. Effects of economic inequality on chronic childhood undernutrition in Ghana *Public Health Nutr*. 2005;4(10):372-8.

19. Kismul H, Acharya P, Mapatano MA, Hatloy A. Determinants of childhood stunting in the Democratic Republic of Congo: further analysis of Demographic and Health Survey 2013-14. *BMC Public Health*. 2017;18(1):74.
20. Keino S, Plasqui G, Etyyang G, van den Borne B. Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa *Food Nutr Bull*. 2014;35(2):167-78.
21. Lamontagne JF, Engle PL, Zeitlin MF. Maternal employment, child care, and nutritional status of 12-18-month-old children in Managua, Nicaragua *Social science & medicine* (1982). 1998;46(3):403-14.
22. Amugsi DA, Mittelmark MB, Lartey A. An analysis of socio-demographic patterns in child malnutrition trends using Ghana demographic and health survey data in the period 1993–2008. *BMC Public Health*. 2013;13(1):960.
23. Beeghly M, Tronick E. Early resilience in the context of parent-infant relationships: a social developmental perspective *Current problems in pediatric and adolescent health care*. 2011;41(7):197-201.
24. Zeitlin MF, Ghassemi H, Mansour M, Levine RA, Dillanneva M, Carballo M, et al. Positive deviance in child nutrition: with emphasis on psychosocial and behavioural aspects and implications for development: United Nations University Tokyo. 1990.
25. Sternin M, Sternin J, Marsh D. Rapid, sustained childhood malnutrition alleviation through a “positive deviance” approach in rural Vietnam: preliminary findings. In: Keeley E, Burkhalter BR, Wollinka O, Bashir N, eds. *The hearth nutrition model: applications in Haiti, Vietnam, and Bangladesh, Report of a Technical Meeting at World Relief Corporation, Wheaton, IL, June 19-21, 1996*. Arlington: BASICS, 1997.1997.
26. Long KNG, Gren LH, Rees CA, West JH, Hall PC, Gray B, et al. Determinants of better health: a cross-sectional assessment of positive deviants among women in West Bengal. *BMC Public Health*. 2013;13(1):372.
27. Marsh DR, Schroeder DG. The positive deviance approach to improve health outcomes: experience and evidence from the field: preface. *Food Nutr Bull* 2002;23(suppl 4):5-8.
28. Bolles K, Speraw C, Berggren G, Lafontant JG. Ti Foyer (hearth) community-based nutrition activities informed by the positive deviance approach in Leogane, Haiti: A programmatic description. *Food Nutr Bull*. 2002;23(suppl 4):11-7.
29. EcoYoff. Positive deviance—take 2. *Living and learning newsletter*. 2003 [Available from: <http://ifnc.tufts.edu/pdf/ecoyoff21.pdf>].
30. Sethi V, Kashyap S, Seth V, Agarwal S. Encouraging appropriate infant feeding practices in slums: a positive deviance approach. *Pakistan J Nutr* 2003;2:164-6.
31. Shafique M, Edwards HM, De Beyl CZ, Thavrin BK, Min M, Roca-Feltrer A. Positive deviance as a novel tool in malaria control and elimination: methodology, qualitative assessment and future potential. *Malaria journal*. 2016;15:91.
32. Marsh DR, Schroeder DG, Dearden KA, Sternin J, Sternin M. The power of positive deviance *BMJ* 2004;329:1177-9.
33. Marsh DR, Pachón H, Schroeder DG, Ha TT, Dearden K, Lang TT, et al. Design of a prospective, randomized evaluation of an integrated nutrition program in rural Viet Nam *Food Nutr Bull* 2002;23(suppl 4):36-47.
34. Dearden K, Quan N, Do M, Marsh DR, Schroeder G, Pachón H, et al. What influences health behavior? Learning from caregivers of young children in Vietnam *Food Nutr Bull*. 2002;23(suppl 4):119-29.
35. Baxter R, Taylor N, Kellar I, Lawton R. What methods are used to apply positive deviance within healthcare organisations? A systematic review. *BMJ Quality & Safety*. 2016;25(3):190-201.

36. Ahrari M, Kuttab A, Khamis S, Farahat AA, Darmstadt GL, Marsh DR, et al. Socioeconomic and behavioral factors associated with successful pregnancy outcomes in upper Egypt: a positive deviance inquiry *Food Nutr Bull.* 2002;23:83-8.
37. The DHS Program. Demographic and Health Surveys [Available from: <http://dhsprogram.com/data/available-datasets.cfm>].
38. Amugsi DA, Dimbuene ZT, Kimani-Murage EW, Mberu B, Ezeh AC. Differential effects of dietary diversity and maternal characteristics on linear growth of children aged 6-59 months in sub-Saharan Africa: a multi-country analysis. *Public health nutrition.* 2017;20(6):1029-45.
39. Amugsi DA, Dimbuene ZT, Kyobutungi C. Correlates of the double burden of malnutrition among women: an analysis of cross sectional survey data from sub-Saharan Africa. *BMJ Open.* 2019;9(7):e029545.
40. The DHS Program. Demographic and Health Surveys [cited 2018 06.09]. Available from: https://dhsprogram.com/data/dataset_admin/download-datasets.cfm [
41. The DHS Program. DHS Methodology [Available from: <http://dhsprogram.com/What-We-Do/Survey-Types/DHS-Methodology.cfm>].
42. Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF Macro. Ghana Demographic and Health Survey 2008 Accra, Ghana: GSS, GHS, and ICF Macro; 2009.
43. Ministry of Health and Social Services (MoHSS), Macro International Inc. Namibia Demographic and Health Survey 2006-07 Windhoek, Namibia and Calverton, Maryland, USA: MoHSS and Macro International Inc. ; 2008.
44. Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF International. Ghana Demographic and Health Survey 2014. Rockville, Maryland, USA: GSS, GHS, and ICF International.; 2015.
45. National Population Commission (NPC) [Nigeria], ICF International. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International. . 2014.
46. National Bureau of Statistics-Kenya, ICF International. 2014 KDHS Key Findings. Rockville, Maryland, USA: KNBS and ICF International. 2015.
47. WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards based on length/height, weight and age *Acta paediatrica* (Oslo, Norway : 1992) Supplement. 2006;450:76-85.
48. Faye CM, Fonn S, Levin J, Kimani-Murage E. Analysing child linear growth trajectories among under-5 children in two Nairobi informal settlements. *Public health nutrition.* 2019;22(11):2001-11.
49. The Namibia Ministry of Health and Social Services (MoHSS) and ICF International, International. The Namibia Demographic and Health Survey 2013 Windhoek, Namibia, and Rockville, Maryland, USA: MoHSS and ICF. 2014.
50. UNICEF. The State of World's Children: Causes of Child Malnutrition 1998 [Available from: <https://www.unicef.org/sowc98/fig5.htm>].
51. Engle PL, Menon P, Haddad L. Care and nutrition: concepts and measurements. Washington, DC: UNICEF/International Food Policy Institute. 1997.
52. Zanello G, Srinivasan CS, Shankar B. What Explains Cambodia's Success in Reducing Child Stunting-2000-2014? *PloS one.* 2016;11(9):e0162668.
53. Smith LC, Ruel MT, Ndiaye A. Why is child malnutrition lower in urban than in rural areas? Evidence from 36 developing countries *World Devel.* 2005;33(8):1285-305.
54. Sastry N. What explains rural-urban differentials in child mortality in Brazil? *Social science & medicine* (1982). 1997;44(7):989-1002.
55. Rollins NC, Bhandari N, Hajeebhoy N, Horton S, Lutter CK, Martines JC, et al. Why invest, and what it will take to improve breastfeeding practices? *Lancet.* 2016;387(10017):491-504.
56. Mirmiran P, Mohammadi F, Allahverdian S, Azizi F. Association of educational level and marital status with dietary intake and cardiovascular risk factors in Tehranian adults: Tehran lipid and glucose study (TLGS). *Nutr Res.* 2002;22.

57. Armar-Klemesu M, Ruel MT, Maxwell DG, Levin CE, Morris SS. Poor maternal schooling is the main constraint to good child care practices in Accra. *J Nutr.* 2000;130(6):1597-607.
58. Urke HB, Mittelmark MB, Amugsi DA, Matanda DJ. Resources for nurturing childcare practices in urban and rural settings: Findings from the Colombia 2010 Demographic and Health Survey. *Child Care Health Dev* 2018;44(4):572-82.
59. Amugsi DA, Mittelmark MB, Lartey A, Matanda DJ, Urke HB. Influence of childcare practices on nutritional status of Ghanaian children: a regression analysis of the Ghana Demographic and Health Surveys. *BMJ Open.* 2014;4.
60. Makoka D, Masibo PK. Is there a threshold level of maternal education sufficient to reduce child undernutrition? Evidence from Malawi, Tanzania and Zimbabwe. *BMC Pediatrics.* 2015;15(1):96.
61. Kuhnt J, Vollmer S. Antenatal care services and its implications for vital and health outcomes of children: evidence from 193 surveys in 69 low-income and middle-income countries. *BMJ open.* 2017;7(11):e017122-e.
62. WHO. Continued breastfeeding for healthy growth and development of children 2019 [cited 2019 03.09]. Available from: https://www.who.int/elena/titles/continued_breastfeeding/en/.
63. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: a meta-analysis. *BMC public health.* 2014;14:1267-.
64. Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L, et al. Breastfeeding and child cognitive development: new evidence from a large randomized trial. *Archives of general psychiatry.* 2008;65(5):578-84.
65. Marquis GS, Habicht JP, Lanata CF, Black RE, Rasmussen KM. Association of breastfeeding and stunting in Peruvian toddlers: an example of reverse causality. *International journal of epidemiology.* 1997;26(2):349-56.
66. Mulder-Sibanda M, Sibanda-Mulder FS. Prolonged breastfeeding in Bangladesh: indicators of inadequate feeding practices or mothers' response to children's poor health? *Public Health.* 1999;113(2):65-8.
67. Fotso JC, Madise N, Baschieri A, Cleland J, Zulu E, Mutua MK, et al. Child growth in urban deprived settings: does household poverty status matter? At which stage of child development? *Health & place.* 2012;18(2):375-84.
68. Paciorek CJ, Stevens GA, Finucane MM, Ezzati M, Nutrition Impact Model Study G. Children's height and weight in rural and urban populations in low-income and middle-income countries: a systematic analysis of population-representative data. *Lancet Glob Health.* 2013;1(5):e300-e9.
69. Amugsi DA, Mittelmark MB, Lartey A. Dietary Diversity is a Predictor of Acute Malnutrition in Rural but Not in Urban Settings: Evidence from Ghana *British Journal of Medicine and Medical Research.* 2014;4(25):4310-24.