Secrets of the hospital underbelly: patterns of abundance of antimicrobial resistance genes in

hospital wastewater vary by specific antimicrobial and bacterial family

Meghan R. Perry [1,2,3†], Hannah C. Lepper[1†], Luke McNally[4,5], Bryan A. Wee[1], Patrick Munk[6],

Amanda Warr[7], Barbara Moore[2], Pota Kalima[2], Carol Philip[2], Ana Maria de Roda Husman[8],

Frank M. Aarestrup[6], Mark Woolhouse[1], Bram A.D. van Bunnik[1]

[†]Meghan R. Perry and Hannah C. Lepper contributed equally to this work

Affiliations

1. Usher Institute, University of Edinburgh, Edinburgh, UK

2. Edinburgh Clinical Infection Research Group, NHS Lothian Infection Service,

Edinburgh, UK

3. Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

4. Centre for Synthetic and Systems Biology, School of Biological Sciences, University

of Edinburgh, Edinburgh, UK

5. Institute of Evolutionary Biology, School of Biological Sciences, University of

Edinburgh, Edinburgh, UK

6. National Food Institute, Technical University of Denmark, Kgs. Lyngby 2800,

Denmark

7. Roslin Institute, University of Edinburgh, Edinburgh, UK

8. National Institute for Public Health and the Environment (RIVM), Bilthoven 3721, The

Netherlands

Word count for main text: 3694

Number of figures: 3

Number of tables: 1

Corresponding author:

Dr Meghan Perry

Edinburgh Clinical Infection Research Group

NHS Lothian Infection Service

Regional Infectious Diseases Unit

Western General Hospital

Telford Road

Edinburgh

EH4 2JP

Email: meghan.perry@nhslothian.scot.nhs.uk

Alternate corresponding author:

Dr Bram van Bunnik

Epidemiology Research Group

Ashworth laboratories

Kings Buildings

Charlotte Auerbach Road

Edinburgh,

EH9 3FL

Email: Bram.vanBunnik@ed.ac.uk

ABSTRACT

Background

Hospital wastewater is a major source of antimicrobial resistance (AMR) outflow into the environment. This study uses metagenomics to study how hospital clinical activity impacts antimicrobial resistance genes (ARGs) abundances in hospital wastewater.

Methods

Sewage was collected over a 24-hour period from multiple wastewater collection points representing different specialties within a tertiary hospital site and simultaneously from community sewage works. High throughput shotgun sequencing was performed using Illumina HiSeq4000. ARG abundances were correlated to hospital antimicrobial usage (AMU), data on clinical activity and resistance prevalence in clinical isolates.

Results

Microbiota and ARG composition varied between collection points and overall ARG abundance was higher in hospital wastewater than in community influent. ARG and microbiota compositions were correlated (Procrustes analysis, *P*=0.014). Total antimicrobial usage was not associated with higher ARG abundance in wastewater. However, there was a small positive association between resistance genes and antimicrobial usage matched to ARG phenotype (IRR 1.11, CI 1.06 - 1.16, *P*<0.001). Furthermore, analysing carbapenem and vancomycin resistance separately indicated that counts of ARGs to these antimicrobials were positively associated with their increased usage (carbapenem rate ratio (RR) 1.91, 95% confidence intervals (CI) 1.01 – 3.72, *P*=0.07, and vancomycin RR 10.25, CI 2.32 – 49.10, *P*<0.01). Overall, ARG abundance within hospital wastewater did not reflect resistance patterns in clinical isolates from concurrent hospital inpatients. However, for clinical isolates of the family *Enterococcaceae* and *Staphylococcaceae*, there was a positive relationship with wastewater ARG abundance (odds ratio (OR) 1.62, CI 1.33 – 2.00, *P*<0.001, and OR 1.65, CI 1.21 – 2.30, *P*=0.006 respectively).

Conclusions

We found that the relationship between hospital wastewater ARGs and antimicrobial usage or clinical isolate resistance varies by specific antimicrobial and bacterial family studied. One explanation we consider is that relationships observed from multiple departments within a single hospital site will be detectable only for ARGs against parenteral antimicrobials uniquely

used in the hospital setting. Our work highlights that using metagenomics to identify the full

range of ARGs in hospital wastewater is a useful surveillance tool to monitor hospital ARG

carriage and outflow and guide environmental policy on AMR.

KEYWORDS

antimicrobial resistance; metagenomics; hospital waste water; surveillance; environmental

risk; Resistance dissemination; Antibiotic usage

INTRODUCTION

In response to the antimicrobial resistance (AMR) crisis, a challenge for the research and medical communities is understanding the flow of AMR between different environmental niches (Woolhouse et al. 2015) and deciding where to focus surveillance and interventions to inform effective policies and action (Laxminarayan et al. 2016). There is an increasing interest in the contribution of hospital wastewater to AMR in the environment. Sewage treatment does not completely eradicate antimicrobial resistance genes (ARGs) and thus ARGs can enter the food chain through water and the use of sewage sludge in agriculture (Woolhouse and Ward 2013; Woolhouse et al. 2015). As a complex matrix representing human bodily waste the potential of community sewage as a surveillance tool to monitor the global epidemiology of AMR has recently been explored (Hendriksen et al. 2019; Aarestrup and Woolhouse 2020).

Hospitals are epidemiologically important nodal points for concentrated antimicrobial consumption and are sources of resistant pathogens (Versporten et al. 2018). Secondary care surveillance, guided by national and international policies, is based on passive reporting of phenotypic and molecular laboratory results for specific pathogens or from screening samples on specific high risk patients (Tornimbene et al. 2018; Department of Health and Social Care 2019). These methods do not represent the full impact of antimicrobial use and inpatient activity on AMR carriage within a hospital and thus risk of transmission. Nor do they capture all pertinent ARGs. As hospital wastewater contains inpatient bodily waste we hypothesised that it could be used as a representation of hospital inpatient carriage of AMR and as such may be a useful surveillance tool.

Many previous studies have identified key pathogens and resistant genes in hospital wastewater and attempts have been made to correlate resistance of specific organisms from hospital clinical isolates with hospital wastewater isolates with conflicting results (Tuméo et al. 2008; Drieux et al. 2016; Talebi et al. 2008; Yang et al. 2009; Maheshwari et al. 2016; Santoro, Romao, and Clementino 2012). There is currently a knowledge gap on how resistance in hospital wastewater quantitatively reflects clinical activity within hospitals. By applying the technique of metagenomics (Hendriksen et al. 2019) to obtain a universal view of ARG composition in hospital wastewater in this study we were able to interrogate this relationship in a multi-departmental study.

MATERIALS AND METHODS

Sewage collection and antibiotic residue analysis

Sampling was performed in June 2017 on eight wastewater collection points (CP), representing different clinical departments, identified to capture the effluent from the majority of the Western General Hospital, Edinburgh (Supplementary Figure 1). Using composite sampling machines, 100 mL of wastewater was sampled every 15 minutes over a 24-hour period thus aiming to collect a representative sample of waste from the hospital inpatient population. Simultaneously, a 24-hour time proportional sample was collected at the inflow site to Seafield community sewage works (hereafter "Seafield"), which serves a population equivalent of 760,000 from Edinburgh and the Lothians. Samples were transported from the site on dry ice and stored at -80°C. Antibiotic residue analysis was performed on 1L of composite hospital wastewaters and 1L of domestic sewage using LC-MS/MS as previously described (Berendsen et al. 2015; Hendriksen et al. 2019).

DNA extraction and analysis

DNA was extracted from sewage using the QIAamp Fast DNA Stool mini kit with an optimized protocol as previously described(Knudsen et al. 2016) and sequenced on the HiSeq4000 platform (Illumina) using 2x 150bp paired-end sequencing. The taxonomic origin of paired reads were assigned using Kraken2 (Wood and Salzberg 2014) to the standard database, a database of representative bacterial genomes and a database of known vector sequences, UniVec_Core (downloaded 9th April 2019). Taxonomic assignments were summarized at the genus level using kraken-biom (Dabdoub 2019). One sample, CP2, was heavily contaminated and removed from further analysis. We used KMA version 1.2.12 to assign the paired and singleton reads to a database consisting of ResFinder reference genes (downloaded 5th of September, 2019). The following flags were used: "-mem_mode -ef -1t1 -cge -nf -shm 1 -t 1" [20]. Reads mapping to the human reference genome (GCA_000001405.15) were removed prior to submission to public sequence databases according to the protocol used in the Human Microbiome Project (Human Microbiome Project 2021; Sherry 2011).

Data collection

Data was collected on clinical isolates from the week surrounding the hospital wastewater sampling to represent pathogens in hospital inpatients. All types of clinical isolate were included but duplicate samples from the same patient within a 48-hour period were excluded. Antimicrobial usage was collated from weekly pharmacy issues to each ward over the 3 months prior to sampling and presented as defined daily dose per 100 occupied bed days

(DDD/100OBDs). Pharmacy issues for prescriptions for outpatient use and for theatres were

excluded.

Data analysis

All statistical analysis and plots were produced using R version 3.6.0. The abundance of ARGs

and bacterial genera were calculated as Fragments Per Kilobase of transcript per Million

mapped bacterial reads (FPKM) (Munk et al. 2018) Bray-Curtis dissimilarity matrices were

determined using Hellinger transformation of the FPKM. Resistance genes from the ResFinder

database were grouped into clusters with 90% sequence homology. The top 50 ARGs were

visualised using a heatmap and gene-wise and collection point dendrograms as previously

described (Hendriksen et al. 2019). Procrustes analysis was used to test the association between

the resistome and bacteriome dissimilarities.

Correlation between inpatient activity and ARG abundance

The source of variance in the abundance of ARGs between the collection points was

investigated using a multilevel Poisson model with the dependent variable as counts of ARG

reads at each collection point aggregated at the 90% homology cluster level. We used an offset

term with the log of the average gene-length per cluster in the ResFinder database, multiplied

by the total bacterial reads per collection point. Random effects of collection point, 70%

sequence homology cluster, and observation were included in the model, the latter to model

the over dispersion inherent to count data (Harrison 2014).

In the main model, we accounted for co- and cross-resistance by fitting both a measure of direct

selection for resistance (effect of department-level usage of antimicrobials on ARGs that confer

resistance to those antimicrobials) and indirect selection (effect of total department-level AMU

on ARG abundance). In a second set of three models we tested the association between

resistance genes and antimicrobial usage of three specific antimicrobials of interest chosen to

represent parenteral antimicrobials only used in a hospital setting (carbapenems, vancomycin)

and an antimicrobial widely used in both community and hospital (amoxicillin). We use a

Bonferroni correction on P values of these additional tests to account for increased risk of type

I error. We used all antimicrobial resistance phenotypes suggested for any gene in a 90%

homology cluster from either the ResFinder or STARAMR (National Microbiology Laboratory

2021) databases. The average length of stay per department was also used to assess the role of

clinical activity on sewage resistance abundance in the main model. The fixed effects structure

of the main model was further adjusted using AIC minimising methods, assessing whether any
interaction effect should be included.
198
To assess the relationship between AMR in clinical isolates and ARG abundance in hospital
wastewater a binomial generalised linear mixed effects model was used including random
effects for site, the class of the antimicrobial used to test the isolates, and for the species of the
isolate to control for inter-species heterogeneity. Two fixed effects were estimated for the log
FPKM of all resistance genes in the sewage that had the same resistance phenotype as the
isolates: one for isolates that were urinary or faecal, and a second for all other isolate types,
due to the different dynamics of inpatient bodily waste being represented in the wastewater
system. Using separate binomial regression models, we accounted for heterogeneity between
the taxonomic family of the isolates in the relationship between AMR in clinical isolates and
sewage ARGs. As some families were rarely tested, the sample size was too small for this
heterogeneity to be assessed in a single model. Therefore, the three most frequently isolated
families were assessed (*Enterobacteriaceae, Enterococcaceae,* and *Staphylococcaceae*), with
the log FPKM of phenotypically matched resistance genes as the only model effect. A
Bonferroni correction was used to adjust the P values of the effects of these models to account
for multiple testing. A similar model was used to evaluate the relationship between AMU and
AMR in clinical isolates.
215

216   **Ethics**
217   This study was conducted following approval from NHS Lothian Research and Development
218   committee under the sponsorship of University of Edinburgh. There was no direct patient
219   contact and therefore the study did not require ethical board approval.
220

221   **RESULTS**
222   The hospital departments served by the wastewater collection points differed by pattern of
223   antimicrobial use (Table 1, S2) and resistance in the 181 clinical isolates identified in the week
224   surrounding wastewater sampling (Figure S3).
225

Metagenomics of wastewater
An average read pair count of 38.4 million (range 35.7-39.2 million) was obtained with an
average of 62% (range 52-73%) of reads allocated to bacteria from the seven hospital
wastewater samples and one community sewage sample

(https://www.ebi.ac.uk/ena/data/view/PRJEB34410). An average of 0.25% of reads mapped to

ARGs in the seven hospital wastewater samples versus 0.1% from Seafield (Table S1).

One thousand, one hundred and fifty-four unique bacterial genera were detected across all

samples (range 1151 - 1154 genera per sample, Table S2). The top nineteen genera accounted

for >70% of bacterial abundance in all samples (Figure 1.D). The most predominant genera

were *Pseudomonas* and *Acinetobacter*, mainly environmental species such as *Pseudomonas*

fluorescens, *Acinetobacter johnsonii*, likely representing bacteria usually present in the hospital

pipes. When compared with Seafield, there was a difference in diversity in the hospital samples

with a higher predominance of gut associated bacteria including *Faecalibacterium,*

Bacteroides, Bifidobacterium and *Escherichia*. (Figure 1.B & D).

ARG abundance and composition varied across different hospital collection points and Seafield

(Figures 1.A & C, Figure 2, Figures S4 & S6). Apart from the wastewater collected at CP4

which represents the acute receiving unit with patients directly admitted from the community,

ARG abundance from hospital wastewater was higher than ARG abundance in Seafield (Figure

2, Fig S4). ARG composition was strongly correlated with bacterial genus level composition

(Procrustes, p=0.014) (Supplementary Figure 5).

We detected 502 different resistance genes belonging to ten different antimicrobial classes

(Table S3) but over 65% of the sample resistomes were composed of the 15 most abundant

genes (Figure S6), mainly belonging to the aminoglycoside and macrolide antimicrobial classes

(Figure 1.C). Key ARGs of interest to infection control including *bla*OXA, *bla*IMP and genes

of the *vanA* cluster were identified.

Inpatient activity and ARG abundance

No significant relationships were observed between total antimicrobial usage or length of stay

and the abundance of ARGs in sewage (Fig. 3, Table S5). This result indicates there was no

evidence for indirect selection or for the impact of transmission among hospital patients on

ARG abundance in sewage when all resistance phenotypes were modelled. There was a

significant positive effect of increased phenotypically-matched antimicrobial usage on

resistance gene abundance, indicating support for a small role of direct selection (IRR 1.11, CI

1.06 - 1.16, *P* < 0.001). AIC comparison of fixed effect structures for the model indicated that

no interaction effects improved model fit.

We next analysed data on the association between carbapenem, vancomycin and amoxicillin

usage and ARGs conferring resistance to these specific antimicrobials in 3 separate models

(Fig 3A, Table S5). We found positive associations that were significant between vancomycin

ARGs and vancomycin usage (IRR 10.25, CI 2.32 – 49.10, $P < 0.001$) and showed a trend

towards significance between carbapenem ARG abundance and carbapenem antimicrobial

usage (IRR 1.91, CI 1.01 – 3.72, $P = 0.07$). No evidence for an association between amoxicillin

usage and amoxicillin ARGs was identified. We omitted the observation-level random effect

from vancomycin model due to singular model fits, so overdispersion was not accounted for.

ARG abundance at a class level within hospital wastewater did not reflect resistance patterns

in clinical isolates when all the data was analysed in one model (Fig 3B, Table S6). There was

no difference between the relationship of isolates from urine and faecal samples with ARG

abundance and isolates from other sample types, e.g. skin, which we expect to enter the

wastewater system at different rates via sinks and showers. We next separately modelled the

three most frequently isolated taxonomic families (Fig 3, Table S6). *Enterococcaceae* and

Staphylococcaceae had a significant positive association with the abundance of ARGs

conferring resistance to the same antimicrobial class (OR: 1.62, C.I. 1.32 – 2.00, p < 0.001, and

OR: 1.65, C.I. 1.21 – 2.30, p < 0.01, respectively), but there was no such relationship for

resistance levels in Enterobacteriaceae. At an antimicrobial class level, clinical isolate

resistance did not reflect the antimicrobial usage of that class in the preceding 3 months

(Supplementary Table 4).

Analysis of antibiotic residues reflected the high AMU within the hospital compared to the

community with an average 12-fold increased residue concentration in hospital effluent

(ranging between 4 and 13 μL^{-1}) for the five classes measured (Supplementary Figure 7). Our

residue data only represents the residue levels from the whole hospital and not individual

collection points and thus could not be specifically correlated with ARG abundance.

DISCUSSION

This study identified that hospital AMU impacts ARG abundances in hospital effluent, with

implications upstream for infection control in the hospital and downstream for AMR in the

environment. Overall, the distribution of bacterial genera and ARGs in our hospital wastewater

samples and domestic sewage sample is similar to previously described sewage composition

in European regions (Hendriksen et al. 2019; Buelow et al. 2018).

There was a significant positive relationship between inpatient department-level AMU and the

abundance of antimicrobial resistance phenotype matched ARGs when all data was considered

together. No relationship was found for total department AMU and ARG abundance. This

supports a role of direct selection from antimicrobial usage in overall patterns of ARGs in

hospital waste water, but not for indirect selection. Previous studies have found a relationship

at a country level between antimicrobial residues and ARG abundance in sewage from the

community (Hendriksen et al. 2019). Indeed, our data shows that the hospital antimicrobial

residues are within the minimum selection concentration range for *Escherichia coli* and

ciprofloxacin resistance (Sandegren 2014), although recent work suggests that higher

antimicrobial concentrations are needed to select for resistance in microbial communities such

as sewage (Klümper et al. 2019).

The association between phenotype-matched ARGs and AMU was weak. Sewage captures

resistance acquired in both the community and in the hospital, but drivers of hospital- and

community-acquired resistance differ. For example, amoxicillin is used in both the community

and hospitals, and resistance is widespread in the UK (60% hospital isolates resistant to

amoxicillin or ampicillin in 2019) (European Centre for Disease Prevention and Control 2020),

suggesting patients commonly arrive in hospital with carriage of amoxicillin resistance genes.

The acquisition of vancomycin or carbapenem resistance, on the other hand, is associated with

prior use of these antibiotics in hospital (Vasilakopoulou et al. 2020; Zhao et al. 2020), and

these antibiotics are solely used parenterally in a hospital setting. Factors affecting within-

hospital selection for and transmission of resistance, such as hospital antimicrobial usage, may

play a stronger role in patterns of ARGs of vancomycin and carbapenems in hospital waste

water than the ubiquitously used antibiotic amoxicillin. In support of this theory, we found a

positive relationship between AMU and waste water ARGs for vancomycin and carbapenems,

but not amoxicillin. Where a particular ward or department consumes high levels of

carbapenem or vancomycin then this work demonstrates that there could be high levels of

undetected faecal or urinary carriage of carbapenem and vancomycin resistance genes. This

could warrant more stringent isolation of these patients, in fitting with concerns about

"unsampled transmission chains" in carbapenem-resistant *Enterobacteriaceae* (Cerqueira et al.

2017). In addition, if the 70% renal excretion of unchanged meropenem (Mouton and van den

Anker 1995) selects for resistant organisms in waste water, then procedures for treatment of

the bodily waste of patients on meropenem may need to be reconsidered.

Length of stay did not impact ARG abundance in this dataset, despite prolonged duration of

inpatient stay being a risk factor for carriage and infection with resistant microorganisms in

previous studies (Safdar and Maki 2002; Gupta et al. 2011; Founou, Founou, and Essack 2018).

This appears not to support the theory of transmission of antimicrobial resistant organisms

amongst patients and their local environment, including from the hospital water system (Kotay

et al. 2017), during their inpatient stay. However, as these data were aggregated at the

department-level there were few observations of length of stay, and further research with a

greater sample size is needed to investigate this relationship.

Metagenomics can capture ARGs carried by a wide variety of bacterial genera, which is of

benefit as the majority of ARGs are carried by non-pathogenic commensal bacteria (Sommer,

Dantas, and Church 2009). Although short-read sequencing cannot conclusively resolve

associations between bacteria and ARGs, in our results ARGs are highly correlated with the

bacteria identified at that collection point (Supplementary Fig 7). This can explain why levels

of ARGs for aminoglycosides, tetracyclines and macrolides are higher than levels of

phenotypic resistance in clinical isolates; the composition of bacterial genera within wastewater

may have intrinsic or high levels of resistance to these antimicrobial classes. The potential for

transfer of ARGs within the sewage network onto and between human pathogens has been

demonstrated indicating the benefit of obtaining a universal view of ARGs (Ludden et al.

2017).

No quantitative relationship was observed between clinical isolates and ARG abundance in

hospital wastewater when all data was considered together. In addition there was no

relationship between AMU in the previous three months and resistance in clinical isolates. This

may be because clinical isolates are not representative enough of carriage of resistance in the

inpatient population as there is a low rate of culture positivity. However, when examined

separately, there was a positive relationship between resistance in *Enterococcaceae* or

Staphylococcaceae, but not *Enterobactericeae,* and hospital wastewater ARG abundance. The

literature on these relationships is divided (Tuméo et al. 2008; Talebi et al. 2008; Yang et al.

2009; Ory et al. 2016; Hutinel et al. 2019; Zarfel et al. 2013) and future work on antimicrobial

usage, specific organisms, isolate types and ARG abundance in sewage potentially over a

longer time period is required to interrogate these relationships further (Mladenovic-Antic et
al. 2016; Rogues et al. 2007).

There was a higher abundance of ARGs in all hospital wastewater samples, bar one (CP4)
which represents acute admissions unit, compared to Seafield. The lower abundance in Seafield
could be due to dilution, and a decline in the relative abundance of AMR-gene carrying human
commensal bacteria in the environment of sewerage system (Pehrsson et al. 2016), or possibly
lower exposure to antimicrobial residues in community waste water. Associations between
antimicrobial residues in community waste water and ARGs have been found (Hendriksen et
al. 2019; Ju et al. 2019), and hospital waste water has been previously shown to have higher
antimicrobial residue levels (Booth, Aga, and Wester 2020). Some studies comparing sewage
influent in paired communities with and without a hospital have found minimal effect of a
hospital on community influent (Gouliouris et al. 2019; Buelow et al. 2018). In other work,
comparing resistance in hospital and community waste water has indicated some associations
(Ludden et al. 2017; Rogues et al. 2007; Pehrsson et al. 2016), although not all studies making
this comparison have found evidence for a relationship (Paulshus et al. 2019).

Concern has been raised about the impact of hospital wastewater on urban influent and effluent
and specific water treatments for hospital wastewater have been called for. This work
highlights that physicians could consider prescribing environmentally degradable
antimicrobials such as beta-lactams over antimicrobials which have persistent residues across
environmental niches e.g. tetracycline to minimise the impact of antimicrobials on the
environmental resistome (Wellington et al. 2013). The ultimate effect of environmental ARGs
on human disease is an ongoing important research question (Bürgmann et al. 2018).

The use of metagenomics is a key strength of this study, allowing quantification of resistance
genes to a wide range of antibiotics and retrospective investigation if new resistance genes
emerge. The 24-hour composite samplers provide a representative sample of the hospital (Chau
et al. 2020), although hospital staff, outpatients and visitors will have also contributed to the
effluent. In addition, some patients will have moved around the hospital during the sampling
period. Although this study is limited to one hospital site at one time point the variation in
antimicrobial use and inpatient characteristics in each department has allowed us to treat them
as discrete treatment centres and draw conclusions about factors affecting ARG abundance.

There is little doubt that hospital resistant pathogens can be abundant in wastewater systems
(Gouliouris et al. 2019; Ludden et al. 2017; Maheshwari et al. 2016). However, using
metagenomic sequencing we show that resistance in hospital wastewater may quantitatively
reflect clinical isolate resistance for some bacterial species (*Enterococcaceae* and
Staphylococcaceae), although not all. As a surveillance tool this novel technique can represent
the burden of AMR carriage in hospital inpatients and hospital pipes for specific resistance
genes relating to important parenteral antimicrobials such as carbapenems and vancomycin. It
may also aid in identification of emerging patterns of ARG abundance and novel ARGs, and
how they may relate to changing patterns of transmission, infection control policies and
antimicrobial usage. Further longitudinal work evaluating the wastewater from multiple
hospital sites is needed to establish AMU/ARG relationships, optimal collection points and
sampling methods to be able to develop this as a surveillance technique.

In conclusion, we show in a multi-departmental study that the relationships between ARG
abundance in hospital wastewater and hospital AMU or clinical resistance levels may vary by
antimicrobial type and bacterial species. Our study emphasises in a novel way the ARG burden
from the high antimicrobial consuming and high resistance carriage environment of the hospital
and that promoting active antimicrobial stewardship, particularly of key parenteral
antimicrobials such as carbapenems and vancomycin, would impact the burden of
environmental AMR. Hospital wastewater is an important source of AMR into the
environment; this should be considered in environmental policy to reduce the flow of AMR
between different environmental reservoirs.

Acknowledgements

The authors would like to thank NHS Lothian estates and Scottish Water for their help with the

sampling and Mick Watson for his laboratory support.

Author contributions

MRP conceived the project and developed it with input from BvB, HL, FA and MW. MRP

facilitated sampling and DNA extraction with AW. PK, CP and BM provided clinical and

pharmaceutical databases and input. ARH performed antibiotic residue analysis. Bio-

informatics and analyses by HL, BvB, LMcN, BW, PM and MRP with input from FA and MW.

Manuscript drafted by MRP and HL with input from BVB and review and comments from all

authors.

Funding

This work was supported was funded by Academy of Medical Sciences (SGL016_1086 to

MRP), an Institutional Strategic Support Fund from University of Edinburgh (J22738 to

MRP) and by The Novo Nordisk Foundation (NNF16OC0021856: Global Surveillance of

Antimicrobial Resistance to HL, BvB, LMcN, MW, PM and FA).

STATEMENT OF CONTRIBUTION TO THE FIELD

Sewage is an attractive medium for surveillance of antimicrobial resistance (AMR). In this

study we interrogate the contribution of hospitals, as focal points of antimicrobial usage and

bacterial infections, to resistance in urban waste water. Previous studies have compared

resistance in hospital patients and their sewage effluent, but focus on only a single (or a few)

bacterial species and antimicrobials, therefore insufficiently addressing the diverse

microbiome(s) and resistome(s) in hospitals and wastewater.

In this study, we apply metagenomics to hospital wastewater and investigate the relationship

between the abundance of antimicrobial resistance genes (ARGs) in the sewage and clinical

activity. Clinical activity included antimicrobial usage and resistance in disease-causing

bacteria cultured from inpatients. This innovative analysis of sewage allows quantification of

both the full range of ARGs to all antimicrobials and specific ARGs of clinical interest.

We demonstrate variation between hospital departments in the abundance in ARGs in the

sewage which reflected differences in inpatients' resistant bacteria and antimicrobial usage.

Furthermore we show that the relationship between clinical activity and ARGs in wastewater

vary by resistance type and bacterial species. We suggest that detection of these relationships

is driven by ARGs to antibiotics only used in the hospital setting.

References

Aarestrup, Frank M., and Mark E. J. Woolhouse. 2020. 'Using Sewage for Surveillance of Antimicrobial Resistance A Global System Would Exploit Metagenomic Sequencing'. *Science* 367 (6478): 630–32.

Berendsen, Bjorn J. A., Robin S. Wegh, Joost Memelink, Tina Zuidema, and Linda A. M. Stolker. 2015. 'The Analysis of Animal Faeces as a Tool to Monitor Antibiotic Usage'. *Talanta* 132 (January): 258–68. https://doi.org/10.1016/j.talanta.2014.09.022.

Booth, Amy, Diana S. Aga, and Astrid L. Wester. 2020. 'Retrospective Analysis of the Global Antibiotic Residues That Exceed the Predicted No Effect Concentration for Antimicrobial Resistance in Various Environmental Matrices'. *Environment International* 141 (August): 105796. https://doi.org/10.1016/j.envint.2020.105796.

Buelow, Elena, Jumamurat R. Bayjanov, Eline Majoor, Rob JL Willems, Marc JM Bonten, Heike Schmitt, and Willem van Schaik. 2018. 'Limited Influence of Hospital Wastewater on the Microbiome and Resistome of Wastewater in a Community Sewerage System'. *FEMS Microbiology Ecology* 94 (7). https://doi.org/10.1093/femsec/fiy087.

Bürgmann, Helmut, Dominic Frigon, William H Gaze, Célia M Manaia, Amy Pruden, Andrew C. Singer, Barth F Smets, and Tong Zhang. 2018. 'Water and Sanitation: An Essential Battlefront in the War on Antimicrobial Resistance'. *FEMS Microbiology Ecology* 94 (9). https://doi.org/10.1093/femsec/fiy101.

Cerqueira, Gustavo C., Ashlee M. Earl, Christoph M. Ernst, Yonatan H. Grad, John P. Dekker, Michael Feldgarden, Sinéad B. Chapman, et al. 2017. 'Multi-Institute Analysis of Carbapenem Resistance Reveals Remarkable Diversity, Unexplained Mechanisms, and Limited Clonal Outbreaks'. *Proceedings of the National Academy of Sciences of the United States of America* 114 (5): 1135–40. https://doi.org/10.1073/pnas.1616248114.

Chau, K. K., L Barker, N Sims, E. B. Budgell, E Harriss, and D. W. Crook. 2020. 'Wastewater Surveillance of Antimicrobial Resistance in Human Populations : A Systematic Review'. *Preprints*, no. October. https://doi.org/10.20944/preprints202010.0267.v1.

Dabdoub, S. 2019. *Create BIOM-Format Tables (Http://Biom-Format.Org) from Kraken Output (Http://Ccb.Jhu.Edu/Software/Kraken/, Https://Github.Com/DerrickWood/Kraken).: Smdabdoub/Kraken-Biom*. https://github.com/smdabdoub/kraken-biom.

Department of Health and Social Care. 2019. *Tackling Antimicobiral Resistance 2019 - 2024: The UK's Five-Year National Action Plan*. ttps://www.gov.uk/government/publications/uk-5-year-action-plan-for-antimicrobial-resistance-2019-to-2024.

Drieux, Laurence, Sophie Haenn, Laurent Moulin, and Vincent Jarlier. 2016. 'Quantitative Evaluation of Extendedspectrum β-Lactamase-Producing Escherichia Coli Strains in the Wastewater of a French Teaching Hospital and Relation to Patient Strain'. *Antimicrobial Resistance and Infection Control* 5 (1): 1–5. https://doi.org/10.1186/s13756-016-0108-5.

European Centre for Disease Prevention and Control. 2020. *Antimicrobial Resistance in the EU/EEA (EARS-Net) - Annual Epidemiological Report 2019*. Stockholm.

Founou, Raspail Carrel, Luria Leslie Founou, and Sabiha Yusuf Essack. 2018. 'Extended Spectrum Beta-Lactamase Mediated Resistance in Carriage and Clinical Gram-Negative ESKAPE Bacteria: A Comparative Study between a District and Tertiary

Hospital in South Africa'. *Antimicrobial Resistance and Infection Control* 7: 134. https://doi.org/10.1186/s13756-018-0423-0.

Gouliouris, Theodore, Kathy E. Raven, Danesh Moradigaravand, Catherine Ludden, Francesc Coll, Beth Blane, Plamena Naydenova, et al. 2019. 'Detection of Vancomycin-Resistant Enterococcus Faecium Hospital-Adapted Lineages in Municipal Wastewater Treatment Plants Indicates Widespread Distribution and Release into the Environment'. *Genome Research* 29 (4): 626–34. https://doi.org/10.1101/gr.232629.117.

Gupta, Neil, Brandi M. Limbago, Jean B. Patel, and Alexander J. Kallen. 2011. 'Carbapenem-Resistant Enterobacteriaceae: Epidemiology and Prevention'. *Clinical Infectious Diseases* 53 (1): 60–67. https://doi.org/10.1093/cid/cir202.

Harrison, Xavier A. 2014. 'Using Observation-Level Random Effects to Model Overdispersion in Count Data in Ecology and Evolution'. *PeerJ* 2 (October). https://doi.org/10.7717/peerj.616.

Hendriksen, Rene S., Patrick Munk, Patrick Njage, Bram van Bunnik, Luke McNally, Oksana Lukjancenko, Timo Röder, et al. 2019. 'Global Monitoring of Antimicrobial Resistance Based on Metagenomics Analyses of Urban Sewage'. *Nature Communications* 10 (1): 1124. https://doi.org/10.1038/s41467-019-08853-3.

Human Microbiome Project. 2021. 'NIH Human Microbiome Project - Tools and Technology'. 2021. https://www.hmpdacc.org/hmp/resources/.

Hutinel, Marion, Patricia Maria Catharina Huijbers, Jerker Fick, Christina \AAhrén, Dan Göran Joakim Larsson, and Carl Fredrik Flach. 2019. 'Population-Level Surveillance of Antibiotic Resistance in Escherichia Coli through Sewage Analysis'. *Eurosurveillance* 24 (37): 1–11. https://doi.org/10.2807/1560-7917.ES.2019.24.37.1800497.

Ju, Feng, Karin Beck, Xiaole Yin, Andreas Maccagnan, Christa S. McArdell, Heinz P. Singer, David R. Johnson, Tong Zhang, and Helmut Bürgmann. 2019. 'Wastewater Treatment Plant Resistomes Are Shaped by Bacterial Composition, Genetic Exchange, and Upregulated Expression in the Effluent Microbiomes'. *The ISME Journal* 13 (2): 346–60. https://doi.org/10.1038/s41396-018-0277-8.

Klümper, Uli, Mario Recker, Lihong Zhang, Xiaole Yin, Tong Zhang, Angus Buckling, and William H. Gaze. 2019. 'Selection for Antimicrobial Resistance Is Reduced When Embedded in a Natural Microbial Community'. *The ISME Journal*, August, 1–11. https://doi.org/10.1038/s41396-019-0483-z.

Knudsen, Berith E., Lasse Bergmark, Patrick Munk, Oksana Lukjancenko, Anders Priemé, Frank M. Aarestrup, and Sünje J. Pamp. 2016. 'Impact of Sample Type and DNA Isolation Procedure on Genomic Inference of Microbiome Composition'. *MSystems* 1 (5). https://doi.org/10.1128/mSystems.00095-16.

Kotay, Shireen, Weidong Chai, William Guilford, Katie Barry, and Amy J. Mathers. 2017. 'Spread from the Sink to the Patient: In Situ Study Using Green Fluorescent Protein (GFP)-Expressing Escherichia Coli To Model Bacterial Dispersion from Hand-Washing Sink-Trap Reservoirs'. *Applied and Environmental Microbiology* 83 (8). https://doi.org/10.1128/AEM.03327-16.

Laxminarayan, Ramanan, Devi Sridhar, Martin Blaser, Minggui Wang, and Mark Woolhouse. 2016. 'Achieving Global Targets for Antimicrobial Resistance'. *Science* 353 (6302): 874–75. https://doi.org/10.1126/science.aaf9286.

Ludden, Catherine, Sandra Reuter, Kim Judge, Theodore Gouliouris, Beth Blane, Francesc Coll, Plamena Naydenova, et al. 2017. 'Sharing of Carbapenemase-Encoding Plasmids

between Enterobacteriaceae in UK Sewage Uncovered by MinION Sequencing'.
Microbial Genomics 3 (7). https://doi.org/10.1099/mgen.0.000114.

Maheshwari, Meenu, Nawar Hadi Yaser, Suraiya Naz, Mansha Fatima, and Iqbal Ahmad.
2016. 'Emergence of Ciprofloxacin-Resistant Extended-Spectrum β-Lactamase-
Producing Enteric Bacteria in Hospital Wastewater and Clinical Sources'. *Journal of
Global Antimicrobial Resistance* 5 (June): 22–25.
https://doi.org/10.1016/j.jgar.2016.01.008.

Mladenovic-Antic, S., B. Kocic, R. Velickovic-Radovanovic, M. Dinic, J. Petrovic, G.
Randjelovic, and R. Mitic. 2016. 'Correlation between Antimicrobial Consumption
and Antimicrobial Resistance of Pseudomonas Aeruginosa in a Hospital Setting: A 10-
Year Study'. *Journal of Clinical Pharmacy and Therapeutics* 41 (5): 532–37.
https://doi.org/10.1111/jcpt.12432.

Mouton, Johan W., and John N. van den Anker. 1995. 'Meropenem Clinical
Pharmacokinetics'. *Clinical Pharmacokinetics* 28 (4): 275–86.
https://doi.org/10.2165/00003088-199528040-00002.

Munk, Patrick, Berith Elkær Knudsen, Oksana Lukjancenko, Ana Sofia Ribeiro Duarte, Liese
Van Gompel, Roosmarijn E. C. Luiken, Lidwien A. M. Smit, et al. 2018. 'Abundance
and Diversity of the Faecal Resistome in Slaughter Pigs and Broilers in Nine European
Countries'. *Nature Microbiology* 3 (8): 898. https://doi.org/10.1038/s41564-018-
0192-9.

National Microbiology Laboratory. 2021. 'STARAMR'. 28 January 2021.
https://github.com/phac-nml/staramr.

Ory, Jerome, Genevieve Bricheux, Anne Togola, Jean Louis Bonnet, Florence Donnadieu-
Bernard, Laurence Nakusi, Christiane Forestier, and Ousmane Traore. 2016.
'Ciprofloxacin Residue and Antibiotic-Resistant Biofilm Bacteria in Hospital Effluent.'
Environmental Pollution (Barking, Essex : 1987) 214 (July): 635–45.
https://doi.org/10.1016/j.envpol.2016.04.033.

Paulshus, Erik, Inger Kühn, Roland Möllby, Patricia Colque, Kristin O'Sullivan, Tore Midtvedt,
Egil Lingaas, Rune Holmstad, and Henning Sørum. 2019. 'Diversity and Antibiotic
Resistance among Escherichia Coli Populations in Hospital and Community
Wastewater Compared to Wastewater at the Receiving Urban Treatment Plant'.
Water Research 161: 232–41. https://doi.org/10.1016/j.watres.2019.05.102.

Pehrsson, Erica C., Pablo Tsukayama, Sanket Patel, Melissa Mejía-Bautista, Giordano Sosa-
Soto, Karla M. Navarrete, Maritza Calderon, et al. 2016. 'Interconnected
Microbiomes and Resistomes in Low-Income Human Habitats'. *Nature* 533 (7602):
212–16. https://doi.org/10.1038/nature17672.

Rogues, A. M., C. Dumartin, B. Amadéo, A. G. Venier, N. Marty, P. Parneix, and J. P. Gachie.
2007. 'Relationship Between Rates of Antimicrobial Consumption and the Incidence
of Antimicrobial Resistance in Staphylococcus Aureus and Pseudomonas Aeruginosa
Isolates From 47 French Hospitals'. *Infection Control & Hospital Epidemiology* 28
(12): 1389–95. https://doi.org/10.1086/523280.

Safdar, Nasia, and Dennis G. Maki. 2002. 'The Commonality of Risk Factors for Nosocomial
Colonization and Infection with Antimicrobial-Resistant Staphylococcus Aureus,
Enterococcus, Gram-Negative Bacilli, Clostridium Difficile, and Candida'. *Annals of
Internal Medicine* 136 (11): 834–44.

Sandegren, Linus. 2014. 'Selection of Antibiotic Resistance at Very Low Antibiotic
    Concentrations'. *Upsala Journal of Medical Sciences* 119 (2): 103–7.
    https://doi.org/10.3109/03009734.2014.904457.

Santoro, Deborah O, Celia M C A Romao, and Maysa M Clementino. 2012. 'Decreased
    Aztreonam Susceptibility among Pseudomonas Aeruginosa Isolates from Hospital
    Effluent Treatment System and Clinical Samples.' *International Journal of
    Environmental Health Research* 22 (6): 560–70.
    https://doi.org/10.1080/09603123.2012.678000.

Sherry, Stephen. 2011. *Human Sequence Removal*. National Center for Biotechnology
    Information.
    https://www.hmpdacc.org/hmp/doc/HumanSequenceRemoval_SOP.pdf.

Sommer, Morten O.A., Gautam Dantas, and George M. Church. 2009. 'Functional
    Characterization of the Antibiotic Resistance Reservoir in the Human Microflora'.
    Science (New York, N.Y.) 325 (5944): 1128–31.
    https://doi.org/10.1126/science.1176950.

Talebi, M., M. R. Pourshafie, M. Katouli, and R. Möllby. 2008. 'Molecular Structure and
    Transferability of Tn1546-Like Elements in Enterococcus Faecium Isolates from
    Clinical, Sewage, and Surface Water Samples in Iran'. *Applied and Environmental
    Microbiology* 74 (5): 1350–56. https://doi.org/10.1128/AEM.02254-07.

Tornimbene, Barbara, Sergey Eremin, Martina Escher, Jolanta Griskeviciene, Sapna
    Manglani, and Carmem Lucia Pessoa-Silva. 2018. 'WHO Global Antimicrobial
    Resistance Surveillance System Early Implementation 2016-17.' *The Lancet.
    Infectious Diseases* 18 (3): 241–42. https://doi.org/10.1016/S1473-3099(18)30060-4.

Tuméo, E., H. Gbaguidi-Haore, I. Patry, X. Bertrand, M. Thouverez, and D. Talon. 2008. 'Are
    Antibiotic-Resistant Pseudomonas Aeruginosa Isolated from Hospitalised Patients
    Recovered in the Hospital Effluents?' *International Journal of Hygiene and
    Environmental Health* 211 (1–2): 200–204.
    https://doi.org/10.1016/j.ijheh.2007.02.010.

Vasilakopoulou, Alexandra, Polyxeni Karakosta, Sophia Vourli, Aikaterini Tarpatzi, Paraskevi
    Varda, Maria Kostoula, Anastasia Antoniadou, and Spyros Pournaras. 2020.
    'Gastrointestinal Carriage of Vancomycin-Resistant Enterococci and Carbapenem-
    Resistant Gram-Negative Bacteria in an Endemic Setting: Prevalence, Risk Factors,
    and Outcomes'. *Frontiers in Public Health* 8 (March): 55.
    https://doi.org/10.3389/fpubh.2020.00055.

Versporten, Ann, Peter Zarb, Isabelle Caniaux, Marie-Françoise Gros, Nico Drapier, Mark
    Miller, Vincent Jarlier, et al. 2018. 'Antimicrobial Consumption and Resistance in
    Adult Hospital Inpatients in 53 Countries: Results of an Internet-Based Global Point
    Prevalence Survey'. *The Lancet Global Health* 6 (6): e619–29.
    https://doi.org/10.1016/S2214-109X(18)30186-4.

Wellington, Elizabeth MH, Alistair BA Boxall, Paul Cross, Edward J Feil, William H Gaze, Peter
    M Hawkey, Ashley S Johnson-Rollings, et al. 2013. 'The Role of the Natural
    Environment in the Emergence of Antibiotic Resistance in Gram-Negative Bacteria'.
    The Lancet Infectious Diseases 13 (2): 155–65. https://doi.org/10.1016/S1473-
    3099(12)70317-1.

Wood, Derrick E., and Steven L. Salzberg. 2014. 'Kraken: Ultrafast Metagenomic Sequence
    Classification Using Exact Alignments'. *Genome Biology* 15 (3): R46.
    https://doi.org/10.1186/gb-2014-15-3-r46.

Woolhouse, Mark E.J., and Melissa J. Ward. 2013. 'Sources of Antimicrobial Resistance'.
Science 341 (6153): 1460–61. https://doi.org/10.1126/science.1243444.
Woolhouse, Mark E.J., Melissa Ward, Bram Van Bunnik, and Jeremy Farrar. 2015.
'Antimicrobial Resistance in Humans, Livestock and the Wider Environment'.
Philosophical Transactions of the Royal Society B: Biological Sciences 370 (1670).
https://doi.org/10.1098/rstb.2014.0083.
Yang, C M, M F Lin, P C Liao, H W Yeh, B V Chang, T K Tang, C Cheng, C H Sung, and M L Liou.
2009. 'Comparison of Antimicrobial Resistance Patterns between Clinical and Sewage
Isolates in a Regional Hospital in Taiwan.' *Letters in Applied Microbiology* 48 (5): 560–
65. https://doi.org/10.1111/j.1472-765X.2009.02572.x.
Zankari, E., H. Hasman, S. Cosentino, M. Vestergaard, S. Rasmussen, O. Lund, F. M.
Aarestrup, and M. V. Larsen. 2012. 'Identification of Acquired Antimicrobial
Resistance Genes'. *Journal of Antimicrobial Chemotherapy* 67 (11): 2640–44.
https://doi.org/10.1093/jac/dks261.
Zarfel, G, H Galler, G Feierl, D Haas, C Kittinger, E Leitner, A J Grisold, et al. 2013.
'Comparison of Extended-Spectrum-Beta-Lactamase (ESBL) Carrying Escherichia Coli
from Sewage Sludge and Human Urinary Tract Infection.' *Environmental Pollution
(Barking, Essex : 1987)* 173 (February): 192–99.
https://doi.org/10.1016/j.envpol.2012.09.019.
Zhao, Shengyuan, Meghan R. Perry, Sharon Kennedy, Julie Wilson, Margo E. Chase-Topping,
Eleanor Anderson, Michael C. Lockhart, and Mark E.J. Woolhouse. 2020. 'Risk Factors
for Carbapenemase-Producing Organisms among Inpatients in Scotland: A National
Matched Case–Control Study'. *Infection Control & Hospital Epidemiology*, December,
1–10. https://doi.org/10.1017/ice.2020.1351.

 Tables

Collection Point	Specialties	No. of wards	No. of pts	Average length of stay in days (s.d.)	Average age in years (s.d.)	DDD per 100 OBDs	No. of clinical isolates
CP1	Cardiology, Urology	3	46	4.9 (0.8)	62.6 (2.2)	123.7	19
CP3	Oncology, Haematology	7	67	3.7 (2.7)	62.1 (1.0)	200.5	27
CP4	Acute receiving unit	5	35	0.9 (0.7)	70.5 (2.3)	325.8	45
CP5	Neuroscience	3	59	3.3 (1.1)	53.5 (2.1)	73.5	8
CP6	Intensive care, Surgery, Medicine	3	70	7.6 (2.5)	66.6 (1.7)	223.8	17
CP7	Infectious Diseases, Surgery, Medicine	6	105	6.1 (3.2)	63.5 (0.8)	148.1	20
CP8	Respiratory, Medicine for the Elderly, Urology, Surgical High Dependency	6	133	12.8 (9.0)	69.0 (1.0)	116.4	25

Table 1. Demographics of hospital collection points. Standard deviation only represents

standard deviation of the average age and length of stay per week. Antimicrobial usage from

previous three months does not include antibiotics issued for outpatient prescriptions or in

theatres. Clinical isolates are from inpatients in the week surrounding wastewater collection.

Abbreviations: pts=patients, DDD=defined daily dose, OBDs=occupied bed days,

s.d.=standard deviation.

Figure legends

Figure 1. Hospital wastewater and community sewage resistome and microbiome
abundance composition. A) Principal coordinate analyses of resistome based on Bray-Curtis
dissimilarity. The percentage of variation explained is noted on the axis labels. B) Principal
coordinate analyses for the microbiome. C) Relative abundance of ARGs by antimicrobial
class. D) Relative abundance of the 19 most abundant bacterial genera in the wastewater and
sewage microbiome. Abbreviations: CP=collection point within hospital,
Seafield=community sewage works, TB=tuberculosis.

Figure 2. Heat map of 50 most abundant antimicrobial resistance genes (ARGs).
Relative abundance of ARGs (FPKM) were log transformed and both ARGs and collection
points were clustered using complete-linkage clustering. For ARGs clustering was based on
Pearson correlation coefficients, for collection points clustering was based on the BC-
dissimilarity matrix (Figure 1) which uses all genes.

Figure 3. Generalised linear mixed effects models for the relationship between
antimicrobial resistance gene abundance, hospital department antibiotic consumption
rates, and hospital department rates of resistance in clinical isolates.
A) Effect of antimicrobial usage (AMU) measured in defined daily dose per 100 occupied
bed days (DDD/100 OBDs) on antimicrobial resistance gene (ARG) abundance A.1.) The
main model, with a single coefficient for all resistance phenotypes. A.2.) Separate models
with coefficients for each antimicrobial. B) Association between antimicrobial resistance
gene abundance in the sewage and clinical resistance rates. B.1.) Main model, with a single
coefficient for all clinical isolate taxonomic family, stratified by sample type - urine or faecal
samples (All: Urine), and for resistance genes and any other sample source (All: Other). B.2.)
Separate models with coefficients for each isolate taxonomic family.

Fig 1.

Fig 2.

Sample Origin

cmlB1_1_AM296₄
ant(2_)-Ia_5_AY
aph(3_)-Ia_2_EU₂
aac(3)-Ib_1_L061
blaOXA-233_1_K
blaVIM-42_1_KP(
blaOXA-210_1_JI
erm(F)_1_M1473(
cfxA4_1_AY7699₂
tet(Q)_2_X58717
cfxA6_1_GQ3429
floR_2_AF118107
ant(6)-Ia_3_KF86
tet(W)_3_AJ4274
tet(O)_3_Y07780
tet(40)_2_AM419
tet(M)_9_X56353
oqxB_1_EU37091
aadA4_1_Z50802
blaTEM-4_1_LK3
tet(G)_2_AF1331₄
blaOXA-47_1_AY
aac(3)-IIIb_2_LLL
aac(3)-IIIc_1_L06
mef(A)_1_AJ9710
erm(B)_9_AF299₂
tet(C)_2_AY0462
aadA11_1_AY144
ere(A)_5_FN3968
tet(A)_3_AY1966∞
aac(3)-IIa_4_L22₆
aph(6)-Id_2_AF0₂
aph(3_)-Ib_1_M₂
mef(C)_1_AB571₂
aac(3)-Ia_1_X158
mph(G)_1_AB571
aac(3)-I_1_AJ877
tet(39)_1_KT3463
ere(D)_1_KP2657
blaOXA-334_1_K
msr(E)_1_FR7515
mph(E)_1_DQ839
aac(6_)-Ib-cr_1_D
tet(X)_1_GU0145
aac(6_)-Ib-Suzho₁
blaIMP-6_1_AB75
blaOXA-58_1_AY
sul2_12_AF49797
sul1_38_BX24835
ant(3_)-Ia_1_X0

Seafield CP1 CP4 CP6 CP7 CP8 CP3 CP5

Fig 3.
A.1

A.2