Secrets of the hospital underbelly: abundance of antimicrobial resistance genes in hospital

wastewater reflects hospital antimicrobial use and inpatient length of stay

Meghan R. Perry[1,2,3*], Bram van Bunnik[4*], Luke McNally[1,5,6], Bryan Wee[4], Patrick Munk[7],

Amanda Warr[8], Barbara Moore[3], Pota Kalima[3], Carol Philip[3], Ana Maria de Roda Husman[9],

Frank Aarestrup[7], Mark Woolhouse[1,4]

Meghan R Perry and Bram van Bunnik contributed equally to this work

Affiliations

1. Centre for Immunity, Infection and Evolution, University of Edinburgh, Edinburgh,

UK

2. Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

3. Regional Infectious Diseases Unit, Western General Hospital, Edinburgh, UK

4. Usher Institute, University of Edinburgh, Edinburgh, UK

5. Centre for Synthetic and Systems Biology, School of Biological Sciences, University

of Edinburgh, Edinburgh, UK

6. Institute of Evolutionary Biology, School of Biological Sciences, University of

Edinburgh, Edinburgh, UK

7. National Food Institute, Technical University of Denmark, Kgs. Lyngby 2800,

Denmark

8. Roslin Institute, University of Edinburgh, Edinburgh, UK

9. National Institute for Public Health and the Environment (RIVM), Bilthoven 3721, The

Netherlands

ABSTRACT

Introduction

Hospital wastewater is a potential major source of antimicrobial resistance (AMR). This study

uses metagenomics to ask how abundances of AMR genes in hospital wastewater are related

to clinical activity.

Methods

Sewage was collected over a 24-hour period from multiple wastewater collection points

representing different specialties within a tertiary hospital site and simultaneously from

community sewage works. High throughput shotgun sequencing was performed using Illumina

HiSeq4000. AMR gene abundances were correlated to hospital antimicrobial usage (AMU),

data on clinical activity and resistance prevalence in clinical isolates.

Findings

Microbiota and AMR gene composition varied between each collection point and overall AMR

gene abundance was higher in hospital wastewater than in community influent. The

composition of AMR genes correlated with microbiota composition (Procrustes analysis,

p=0.002). Increased antimicrobial consumption at a class level was associated with higher

AMR gene abundance within that class in wastewater (incidence rate ratio 2.80, C.I. 1.2-6.5,

p=0.016). Prolonged average patient length of stay was associated with higher total AMR gene

abundance in wastewater (incidence rate ratio 2.05, C.I. 1.39-3.01, p=0.0003). AMR gene

abundance at a class level within hospital wastewater did not reflect resistance patterns in the

181 clinical isolates grown from hospital inpatients over the time of wastewater sampling.

Conclusions

Hospital antimicrobial consumption and patient length of stay are important drivers of AMR

gene outflow into the environment. Using metagenomics to identify the full range of AMR

genes in hospital wastewater could represent a useful surveillance tool to monitor hospital

AMR gene outflow and guide environmental policy on AMR.

INTRODUCTION

In response to the antimicrobial resistance (AMR) crisis, a challenge for the research and medical communities is understanding the flow of AMR between different environmental niches[1] and deciding where to focus surveillance and interventions to inform effective policies and action[2]. There is an increasing interest in the contribution of hospital wastewater to AMR in the environment. Sewage treatment does not completely eradicate AMR genes and thus AMR genes can enter the food chain through water and the use of sewage sludge in agriculture[3,4]. As a complex matrix representing human bodily waste the potential of community sewage as a surveillance tool to monitor the global epidemiology of AMR has recently been explored[5].

Hospitals are epidemiologically important nodal points for concentrated antimicrobial consumption and are sources of resistant pathogens[6]. Secondary care surveillance, guided by national and international policies[7,8], is based on passive reporting of phenotypic laboratory results for specific pathogens or from screening samples on specific high risk patients[9,10]. These methods will not represent the full impact of antimicrobial use and inpatient activity on AMR carriage within a hospital, and thus risk of transmission, or capture all pertinent antimicrobial resistance genes (AMR genes). As hospital wastewater contains inpatient bodily waste we hypothesised that it could be used as a representation of inpatient community carriage of AMR and as such may be a useful surveillance tool.

Many previous studies have identified key pathogens and resistant genes in hospital wastewater[11-16] and some attempts have been made to correlate resistance of specific organisms in hospital clinical isolates with resistance in hospital wastewater with conflicting results[17-19]. There is currently a knowledge gap on how resistance in hospital wastewater quantitatively reflects clinical activity within hospitals. By applying the technique of metagenomics[5] to obtain

a universal view of AMR gene composition in hospital wastewater in this study we were able

to interrogate this relationship in a multi-departmental study.

METHODS

Sewage collection and antibiotic residue analysis

Sampling was performed in June 2017 on eight wastewater collection points (CP), representing

different clinical departments, identified to capture the effluent from the majority of the

Western General Hospital, Edinburgh (Supplementary Figure 1). Using composite sampling

machines, 100 mL of wastewater was sampled every 15 minutes over a 24-hour period thus

aiming to collect a representative sample of the hospital inpatient population. Simultaneously,

a 24-hour time proportional sample was collected at the inflow site to Seafield community

sewage works, which serves a population equivalent of 760,000 from Edinburgh and the

Lothians. Samples were transported from the site on dry ice and stored at -80^{o}C. Antibiotic

residue analysis was performed on 1L of composite hospital wastewaters and 1L of domestic

sewage using LC-MS/MS as previously described[5,20].

DNA extraction and analysis

DNA was extracted from sewage using the QIAamp Fast DNA Stool mini kit with an optimized

protocol as previously described[21] and sequenced on the HiSeq4000 platform (Illumina) using

2x 150bp paired-end sequencing. Paired reads were then assigned using Kraken2[22] and a

database of known vector sequences, UniVec_Core (downloaded 9[th] April 2019). Taxonomic

assignments were summarized at the genus level using kraken-biom[23]. One sample, CP2, was

heavily contaminated and removed from further analysis. Using the ResFinder database within

the MGMapper web tool, functional metagenomic antibiotic resistance determinants were

identified[24].

Data collection

Data was collected on clinical isolates from the week surrounding the hospital wastewater

sampling to represent pathogens in hospital inpatients. All types of clinical isolate were

included but duplicate samples from the same patient within a 48-hour period were omitted

from the final analysis. Antimicrobial usage was collated in the form of weekly pharmacy

issues to each ward over the 3 months prior to sampling and presented as defined daily dose

per 100 occupied bed days (DDD/100OBDs). Pharmacy issues for prescriptions for outpatient

use and for theatres were excluded.

Data analysis

All statistical analysis and plots were produced using R version 3.6.0[25].

Using the output from Kraken and Resfinder the relative abundance of AMR genes and

bacterial genera were calculated as FPKM (Fragments Per Kilobase of transcript per Million

mapped bacterial reads) to account for both sample-wise sequencing depth differences and

size-dependent probability of observing a reference[26]. Relative abundances and dissimilarity

matrices of bacterial genera and AMR genes were determined and the top 50 AMR genes were

visualised using a heatmap with gene-wise and collection point dendograms as previously

described[5]. Procrustes anaylsis was used to test the association between the resistome and

bacteriome dissimilarities.

Correlation between inpatient activity and AMR gene abundance

The source of variance in the abundance of AMR genes between the collection points was

investigated using a multilevel Poisson model with the dependent variable as counts of AMR

gene reads at each collection point aggregated at the class level, offset by the log of the average

gene-length per antibiotic class multiplied by the total bacterial reads per collection point.

Random effects of collection point, antimicrobial class and observation were included in the

model, the latter to model the over dispersion inherent to count data[27].

Each of the potential fixed effects were evaluated for correlation. If two variables were highly

correlated (>0.7) only one variable was chosen (based on lowest Akaike information criterion,

AIC). Using AMU in DDD per 100 OBD at the collection point level we accounted for co- and

cross-resistance by fitting both a measure of direct selection for resistance (effect of usage of

one antimicrobial class on AMR genes that confer resistance to that antimicrobial class) and

indirect selection (effect of total AMU on AMR gene abundance).

To assess the relationship between AMR in clinical isolates and AMR gene abundance in

hospital wastewater a binomial generalised linear mixed effects model was used including the

same three random effects with an addition of a random effect of species to control for inter-

species heterogeneity. Isolate type was included in the model as urinary/faecal or non-urinary

due to the different dynamics of inpatient bodily waste being represented in the wastewater

system. A similar model was additionally used to evaluate the relationship between AMR in

clinical isolates and AMU of the inpatient community.

Ethics

This study was conducted following approval from NHS Lothian Research and Development

committee under the sponsorship of University of Edinburgh. There was no direct patient

contact and therefore the study did not require ethical board approval.

RESULTS

The hospital departments served by the wastewater collection points differed by pattern of

antimicrobial use (Table 1, Supplementary Figure 2) and resistance in the 181 clinical isolates

identified in the week surrounding wastewater sampling (Supplementary Figure 3).

Metagenomics of wastewater

An average read pair count of 38.5 million (range 35.8-39.2 million) was obtained with an

average of 76% (range 67-81%) of reads allocated to bacteria from the seven hospital

wastewater samples and one community sewage sample (sequencing reads can be found

under Bioproject accession number PRJEB34410). An average of 0.08% of reads mapped to

AMR genes in the seven hospital wastewater samples versus 0.03% from Seafield

community sewage works (Supplementary Table 1).

One thousand and forty-seven unique bacterial genera were detected across all samples (range

705-965 genera per sample, Supplementary Table 2). The top nineteen genera accounted for

>60% of bacterial abundance in all samples (Figure 1, panel D). The most predominant genera

were *Pseudomonas* and *Acinetobacter*, mainly environmental species such as *Pseudomonas*

fluorsecens, Acinetobacter johnsonii, likely representing bacteria from biofilms within the

hospital pipes. When compared with the Seafield community sewage sample, there was a

difference in diversity with a higher predominance of gut associated bacteria including

Faecalibacterium, Bacteroides, Bifidobacterium and *Escherichia* in the hospital samples.

(Figure 1, Panel B & D).

AMR gene abundance and composition varied across different hospital collection points and

Seafield (Figures 1 panel A & C, Figure 2, Supplementary Figures 4 & 6). Apart from the

wastewater collected at CP4 which houses patients directly admitted from the community,

AMR gene abundance from hospital effluent was higher than AMR gene abundance in Seafield

community sewage works (Figure 2, Supplementary Fig 4). AMR gene composition was

strongly correlated with bacterial genus level composition (Procrustes, p=0.002)

(Supplementary Figure 5).

Over 60% of the resistome of hospital wastewater and community sewage samples is composed

of the 15 most abundant genes (Supplementary Figure 6), mainly belonging to the

aminoglycoside and macrolide antimicrobial classes (Figure 1, panel C). In total, we detected

168 different resistance genes belonging to ten different antimicrobial classes (Supplementary

Table 3). The genes detected included key AMR genes of interest to infection control including

blaOXA, blaIMP and genes of the VanA cassette.

Inpatient activity and AMR gene abundance

With increasing usage of an antimicrobial class there was a significant increase in the

abundance of AMR genes belonging to that class (incidence rate ratio (IRR) 2.80, C.I. 1.2-6.5,

p=0.016, Figure 3). No relationship between total AMU per collection point and total AMR

gene abundance was identified implying that cross- resistance is not extensively contributing

to AMR gene abundance. The abundance of AMR genes increased with increasing patient

length of stay (IRR 2.05, C.I. 1.39-3.01, p=0.0003, Figure 3). This result is not because of

increasing AMU with increasing LOS as these two variables are not correlated at collection

point level but instead likely reflects transmission amongst hospital inpatients. A minor but

significant interaction was identified between length of stay and DDDs per 100 OBDs which

implies that following a certain degree of antimicrobial exposure and hospital exposure through

length of stay there is no longer a cumulative effect on AMR gene abundance. The effect of

patient age was not significant.

Stratifying for species heterogeneity, AMR gene abundance at a class level within hospital

wastewater did not reflect resistance patterns in the clinical isolates from hospital inpatients

around the time of wastewater sampling (Supplementary Table 4). There was no difference

between the relationship of the directly excreted urinary and faecal isolates with AMR gene

abundance and other isolate types e.g. skin which would only reach the wastewater system

indirectly via run off from showers. At an antimicrobial class level the resistance observed in

clinical isolates did not reflect the antimicrobial usage of that class in the preceding 3 months.

Analysis of antibiotic residues in hospital effluent and community influent reflected the high

AMU within the hospital with an average 12-fold increased concentration in hospital effluent

(ranging between 4 and 13 μL^{-1}) for the five classes measured (Supplementary Figure 7). Our

residue data only represents the residue levels from the whole hospital and not individual

collection points and thus cannot be specifically correlated with AMR gene abundance.

DISCUSSION

This study identified that hospital AMU and patient length of stay impacts AMR gene

abundances in hospital effluent and has implications upstream for infection control in the

hospital and downstream for AMR in the environment. Overall, the distribution of bacterial

genera and AMR genes in our hospital wastewater samples and domestic sewage sample is

similar to previous described sewage composition in European regions[5,28].

There are different plausible explanations for the positive relationship between AMU and AMR

gene abundance. These include inpatient faecal and urinary carriage of AMR genes both from

direct antimicrobial exposure and from transmission between patients. However, exposure of

environmental bacteria in the hospital waste pipes, which may reside in biofilms for months,

to high levels of unmetabolised excreted antimicrobials from patients is also likely to play a

significant role[29]. Indeed, our data shows that the hospital antimicrobial residues are within

the minimum selection concentration range for some pathogen antimicrobial combinations e.g.

Escherichia coli and ciprofloxacin resistance[30]. A relationship has been previously observed at

a country level both between both AMU and antimicrobial residue levels and AMU and AMR

gene abundance in sewage[5].

The implications of the relationship between AMU and AMR gene abundance in wastewater

can be illustrated by the strong correlation between abundance of carbapenem resistance genes

and carbapenem consumption (R^2=0.87). Where a particular ward or department consumes

high levels of carbapenem then this work demonstrates that there could be high levels of

undetected faecal or urinary carriage of carbapenem resistance genes which could warrant more

stringent isolation of these patients, in fitting with concerns about "unsampled transmission

chains" in carbapenem-resistant *Enterobacteriaceae*[31]. Alternatively, or in addition to, if the

70% renal excretion of unchanged meropenem is selecting out resistant organisms in the pipes

for release into the environment then procedures for the bodily waste of patients on meropenem

may need to be reconsidered.

The positive effect of increasing length of stay on AMR gene abundance is in keeping with

prolonged duration of inpatient stay being a risk factor for carriage and infection with resistant

microorganisms[32–34]. That length of stay remains as a signal even despite controlling for

antimicrobial consumption supports the theory of transmission of antimicrobial resistant

organisms amongst patients and their local environment during their inpatient stay. This has

further implications for hospital infection control management of long stay patients as they

may be more likely to harbour key AMR genes[32,35] and be a source for transmission.

As the majority of AMR genes are carried by non-pathogenic commensal bacteria[36] the

advantage of using metagenomics is that we are able to capture AMR genes carried by the wide

variety of bacterial genera identified. Although shotgun sequencing cannot identify which

bacteria are carrying which AMR genes our results indicate that the AMR genes present are

highly correlated with the bacteria identified at that collection point (Supplementary Fig 7).

This can explain why levels of AMR genes for aminoglycosides, tetracyclines and macrolides

may be higher than expected in relation to phenotypic resistance in clinical isolates; the

composition of bacterial genera selected out may have intrinsic or high levels of resistance to

these antimicrobial classes. The potential for transfer of AMR genes within the sewage network

onto human pathogens has been demonstrated indicating the importance of obtaining a

universal view of AMR genes[16].

No quantitative relationship was observed between clinical isolates stratified by organism and

isolate type and AMR gene abundance in hospital wastewater. There are multiple possible

explanations for this. The clinical isolates grown were only 7% of all microbiological samples

taken and only represent a selected group of hospital patients whereas the hospital wastewater

represents the carriage of the whole inpatient population. Faecal or urinary resistance carriage

is often present on non-pathogenic organisms and does not always progress to clinical disease

represented by clinical isolate growth. In addition, other biological niches of human resistance

carriage that will not well be represented in our hospital wastewater samples. The lack of

relationship could also be due to the inherent issues of correlating phenotypic resistance and

genotypic resistance and the loss of resolution in allocating resistances into antimicrobial

classes rather than individual drugs. In addition, it may be because different relationships exist

at an organism and AMR gene level in sewage which are lost in trying to compare the two

microbial communities. A recent study using EARS-net clinical isolate data and urban sewage

samples found correlations in only a small proportion of pathogen-antimicrobial relationships

studied[37]. Due to conflicting relationships in the literature[11,12,14,37], further detailed work on

specific organisms, isolate types and resistance in sewage is required to interrogate these

relationships further.

No relationship was identified between AMU in the previous 3 months and resistance in

clinical isolates stratified by species and isolate type. This can also be explained by many of

the reasons discussed above. Previous literature examining this has only found a relationship

between specific organisms and antibiotics over a much longer time period[38,39].

There was a higher abundance of AMR genes in the hospital wastewater samples when

compared to Seafield community sewage works. This is likely due to dilution, a general loss

of AMR-gene carrying human commensal bacteria in the environment of sewerage system[40]

and exposure to lower levels of antibiotic residues. When comparing sewage influent in paired

communities with and without a hospital Buelow *et al* concluded that there was minimal effect

of a hospital on community influent[28]. However, in other work, hospital wastewater impacted

resistome composition at an individual gene level[41]. It is worth noting that the AMR gene

abundance in our urban sewage influent (Seafield community sewage works), which receives

the hospital wastewater from our studied hospital and another large tertiary hospital, is higher

than the hospital wastewater collected in CP4, the admissions unit which is our closest

representation of community AMR carriage. Concern has been raised relating to the impact of

hospital wastewater on urban influent and effluent and specific water treatments for hospital

wastewater have been called for. This work highlights the concept that physicians should

consider, where possible, prescribing of environmentally degradable antimicrobials e.g. beta-

lactams over those antimicrobials which have persistent residues which are mobile between

environmental niches e.g. tetracycline to minimise the impact of antimicrobials on the

environmental resistome[42]. The ultimate effect of environmental AMR genes on human health

is an ongoing important research question[29].

One of the strengths of this study is the use of metagenomics to quantify resistance genes to a

wide range of antibiotics. This technique would also allow for retrospective investigation if

new resistance genes emerge. However, when, as in this analysis, antibiotics and AMR genes

are grouped together at a class level detail can be lost and there are incidences when looking

for specific pathogens and resistance genes is warranted. The use of 24-hour composite

samplers is a strong technique to give a representative sample of the hospital but it must be

considered that hospital staff, outpatients and visitors will have also contributed to the effluent.

In addition, a proportion of patients will have moved around the hospital during the period

studied. Although this study is limited to one hospital site at one time point the variation in

antimicrobial use and inpatient characteristics in each department has allowed us to treat them

as discrete treatment centres and draw conclusions about factors affecting AMR gene

abundance.

There is little doubt that hospital resistant pathogens can be abundant in wastewater

systems[15,16,18]. The results from this study imply that the place for measuring AMR genes using

this novel tool of using metagenomics to analyse hospital wastewater is not to comment on

resistance in clinical isolates within the hospital. Rather its utility could fall on representing

carriage of AMR in hospital inpatients and hospital pipes and identification of emerging key

AMR genes in relation to changing patterns of transmission, infection control policies and

antimicrobial usage. Further work evaluating using spot samples and optimal collection points

of hospital wastewater would be required.

In conclusion, we show in a multi-departmental study that AMR gene abundance in hospital

wastewater is influences by hospital AMU and length of inpatient stay. Our study further

emphasises the AMR gene transmission risks of exposure to high antimicrobial consuming and

high resistance carriage environments. Hospital wastewater is an important source of AMR

into the environment and should be considered in environmental policy to reduce the flow of

AMR between different environmental reservoirs.

Acknowledgements

The authors would like to thank NHS Lothian estates and Scottish Water for their help with the

sampling and Mick Watson for his laboratory support.

Author contributions

MRP conceived the project and developed it with input from BvB, FA and MW. MRP

facilitated sampling and DNA extraction with AW. PK, CP and BM provided clinical and

pharmaceutical databases and input. ARH facilitated antibiotic residue analysis. Bio-

informatics and analyses by BvB, LMcN, BW, PM and MRP with input from FA and MW.

Manuscript drafted by MRP with input from BVB and review and comments from all authors.

Financial support

MRP was funded by Academy of Medical Sciences (SGL016_1086) and an Institutional

Strategic Support Fund from University of Edinburgh. BvB, LMcN, MW, PM and FA were

funded by The Novo Nordisk Foundation (NNF16OC0021856: Global Surveillance of

Antimicrobial Resistance).

References

1.   Woolhouse, M., Ward, M., van Bunnik, B. & Farrar, J. Antimicrobial resistance in

humans, livestock and the wider environment. *Philos. Trans. R. Soc. B Biol. Sci.* **370**,

(2015).

2.   Laxminarayan, R., Sridhar, D., Blaser, M., Wang, M. & Woolhouse, M. Achieving global

targets for antimicrobial resistance. *Science* **353**, 874–875 (2016).

3.   Woolhouse, M. E. J. & Ward, M. J. Sources of Antimicrobial Resistance. *Science* **341**,

1460–1461 (2013).

4.   Bréchet, C. *et al.* Wastewater Treatment Plants Release Large Amounts of Extended-

Spectrum β-Lactamase–Producing Escherichia coli Into the Environment. *Clin. Infect.*

Dis. **58**, 1658–1665 (2014).

5.   Hendriksen, R. S. *et al.* Global monitoring of antimicrobial resistance based on

metagenomics analyses of urban sewage. *Nat. Commun.* **10**, 1124 (2019).

6.   Versporten, A. *et al.* Antimicrobial consumption and resistance in adult hospital

inpatients in 53 countries: results of an internet-based global point prevalence survey.

Lancet Glob. Health **6**, e619–e629 (2018).

7.   Hsia, Y. *et al.* Use of the WHO Access, Watch, and Reserve classification to define

patterns of hospital antibiotic use (AWaRe): an analysis of paediatric survey data from 56

countries. *Lancet Glob. Health* **7**, e861–e871 (2019).

8.   Overview | Antimicrobial stewardship: systems and processes for effective antimicrobial

medicine use | Guidance | NICE. Available at: https://www.nice.org.uk/guidance/ng15.

(Accessed: 25th July 2019)

9.   Tornimbene, B. *et al.* WHO Global Antimicrobial Resistance Surveillance System early

implementation 2016–17. *Lancet Infect. Dis.* **18**, 241–242 (2018).

10. UK 5-year action plan for antimicrobial resistance 2019 to 2024. *GOV.UK* Available at:

https://www.gov.uk/government/publications/uk-5-year-action-plan-for-antimicrobial-

resistance-2019-to-2024. (Accessed: 28th May 2019)

11. Tuméo, E. *et al.* Are antibiotic-resistant Pseudomonas aeruginosa isolated from

hospitalised patients recovered in the hospital effluents? *Int. J. Hyg. Environ. Health* **211**,

200–204 (2008).

12. Lozniewski *et al.* Comparison of sludge and clinical isolates of Listeria monocytogenes.

Lett. Appl. Microbiol. **32**, 336–339 (2001).

13. Drieux, L., Haenn, S., Moulin, L. & Jarlier, V. Quantitative evaluation of extended-

spectrum β-lactamase-producing Escherichia coli strains in the wastewater of a French

teaching hospital and relation to patient strain. *Antimicrob. Resist. Infect. Control* **5**, 9

(2016).

14. Talebi, M., Pourshafie, M. R., Katouli, M. & Möllby, R. Molecular Structure and

Transferability of Tn1546-Like Elements in Enterococcus faecium Isolates from Clinical,

Sewage, and Surface Water Samples in Iran. *Appl. Environ. Microbiol.* **74**, 1350–1356

(2008).

15. Gouliouris, T. *et al.* Detection of vancomycin-resistant Enterococcus faecium hospital-

adapted lineages in municipal wastewater treatment plants indicates widespread

distribution and release into the environment. *Genome Res.* **29**, 626–634 (2019).

16. Ludden, C. *et al.* Sharing of carbapenemase-encoding plasmids between

Enterobacteriaceae in UK sewage uncovered by MinION sequencing. *Microb. Genomics*

3, (2017).

17. Yang, C. m. *et al.* Comparison of antimicrobial resistance patterns between clinical and

sewage isolates in a regional hospital in Taiwan. *Lett. Appl. Microbiol.* **48**, 560–565

(2009).

18. Maheshwari, M., Yaser, N. H., Naz, S., Fatima, M. & Ahmad, I. Emergence of

ciprofloxacin-resistant extended-spectrum β-lactamase-producing enteric bacteria in

hospital wastewater and clinical sources. *J. Glob. Antimicrob. Resist.* **5**, 22–25 (2016).

19. Santoro, D. O., Romão, C. M. C. A. & Clementino, M. M. Decreased aztreonam

susceptibility among Pseudomonas aeruginosa isolates from hospital effluent treatment

system and clinical samples. *Int. J. Environ. Health Res.* **22**, 560–570 (2012).

20. Berendsen, B. J. A., Wegh, R. S., Memelink, J., Zuidema, T. & Stolker, L. A. M. The

analysis of animal faeces as a tool to monitor antibiotic usage. *Talanta* **132**, 258–268

(2015).

21. Knudsen, B. E. *et al.* Impact of Sample Type and DNA Isolation Procedure on Genomic

Inference of Microbiome Composition. *mSystems* **1**, (2016).

22. Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification

using exact alignments. *Genome Biol.* **15**, R46 (2014).

23. Dabdoub, S. *Create BIOM-format tables (http://biom-format.org) from Kraken output*

(http://ccb.jhu.edu/software/kraken/, https://github.com/DerrickWood/kraken).:

smdabdoub/kraken-biom. (2019).

24. Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob.*

Chemother. **67**, 2640–2644 (2012).

25. R: The R Project for Statistical Computing. Available at: https://www.r-project.org/.

(Accessed: 20th June 2019)

26. Munk, P. *et al.* Abundance and diversity of the faecal resistome in slaughter pigs and

broilers in nine European countries. *Nat. Microbiol.* **3**, 898 (2018).

27. Harrison, X. A. Using observation-level random effects to model overdispersion in count

data in ecology and evolution. *PeerJ* **2**, (2014).

28. Buelow, E. *et al.* Limited influence of hospital wastewater on the microbiome and

resistome of wastewater in a community sewerage system. *FEMS Microbiol. Ecol.* **94**,

(2018).

29. Bürgmann, H. *et al.* Water and sanitation: an essential battlefront in the war on

antimicrobial resistance. *FEMS Microbiol. Ecol.* **94**, (2018).

30. Sandegren, L. Selection of antibiotic resistance at very low antibiotic concentrations.

Ups. J. Med. Sci. **119**, 103–107 (2014).

31. Cerqueira, G. C. *et al.* Multi-institute analysis of carbapenem resistance reveals

remarkable diversity, unexplained mechanisms, and limited clonal outbreaks. *Proc. Natl.*

Acad. Sci. U. S. A. **114**, 1135–1140 (2017).

32. Safdar, N. & Maki, D. G. The commonality of risk factors for nosocomial colonization

and infection with antimicrobial-resistant Staphylococcus aureus, enterococcus, gram-

negative bacilli, Clostridium difficile, and Candida. *Ann. Intern. Med.* **136**, 834–844

(2002).

33. Gupta, N., Limbago, B. M., Patel, J. B. & Kallen, A. J. Carbapenem-Resistant

Enterobacteriaceae: Epidemiology and Prevention. *Clin. Infect. Dis.* **53**, 60–67 (2011).

34. Founou, R. C., Founou, L. L. & Essack, S. Y. Extended spectrum beta-lactamase

mediated resistance in carriage and clinical gram-negative ESKAPE bacteria: a

comparative study between a district and tertiary hospital in South Africa. *Antimicrob.*

Resist. Infect. Control **7**, 134 (2018).

35. Ny, P., Nieberg, P. & Wong-Beringer, A. Impact of carbapenem resistance on

epidemiology and outcomes of nonbacteremic Klebsiella pneumoniae infections. *Am. J.*

Infect. Control **43**, 1076–1080 (2015).

36. Sommer, M. O. A., Dantas, G. & Church, G. M. Functional characterization of the

antibiotic resistance reservoir in the human microflora. *Science* **325**, 1128–1131 (2009).

37. Pärnänen, K. M. M. *et al.* Antibiotic resistance in European wastewater treatment plants

mirrors the pattern of clinical antibiotic resistance prevalence. *Sci. Adv.* **5**, (2019).

38. Mladenovic-Antic, S. *et al.* Correlation between antimicrobial consumption and

antimicrobial resistance of Pseudomonas aeruginosa in a hospital setting: a 10-year study.

J. Clin. Pharm. Ther. **41**, 532–537 (2016).

39. Rogues, A. M. *et al.* Relationship Between Rates of Antimicrobial Consumption and the

Incidence of Antimicrobial Resistance in Staphylococcus aureus and Pseudomonas

aeruginosa Isolates From 47 French Hospitals. *Infect. Control Hosp. Epidemiol.* **28**,

1389–1395 (2007).

40. Pehrsson, E. C. *et al.* Interconnected microbiomes and resistomes in low-income human

habitats. *Nature* **533**, 212–216 (2016).

41. Lamba, M. *et al.* Carbapenem resistance exposures via wastewaters across New Delhi.

Environ. Int. **119**, 302–308 (2018).

42. Wellington, E. M. *et al.* The role of the natural environment in the emergence of

antibiotic resistance in Gram-negative bacteria. *Lancet Infect. Dis.* **13**, 155–165 (2013).

Collection Point (CP)	Specialties	No. of wards	No. of pts	Average length of stay in days (s.d.)	Average age in years (s.d.)	DDD per 100 OBD	No. of clinical isolates
CP1	Cardiology, Urology	3	46	4.9 (0.8)	62.6 (2.2)	123.7	19
CP3	Oncology, Haematology	7	67	3.7 (2.7)	62.1 (1.0)	200.5	27
CP4	Acute receiving unit	5	35	0.9 (0.7)	70.5 (2.3)	325.8	45
CP5	Neuroscience	3	59	3.3 (1.1)	53.5 (2.1)	73.5	8
CP6	Intensive care, Surgery, Medicine	3	70	7.6 (2.5)	66.6 (1.7)	113.8	17
CP7	Infectious Diseases, Surgery, Medicine	6	105	6.1 (3.2)	63.5 (0.8)	148.1	20
CP8	Respiratory, Medicine for the Elderly, Urology, Surgical High Dependency	6	133	12.8 (9.0)	69.0 (1.0)	116.4	25

453   **Table 1 Demographics of hospital collection points**

454   Standard deviation only represents standard deviation of the average age and length of stay per

455   week. Antimicrobial usage does not include antibiotics issued for outpatient prescriptions or in

456   theatres.

457   DDD defined daily dose, OBD occupied bed day, s.d. standard deviation

Figure legends

Figure 1. **Hospital wastewater and community sewage resistome and microbiome**

abundance composition.

A) Principal coordinate analyses of resistome based on Bray-Curtis dissimilarity. The

percentage of variation explained is noted on the axis labels. B) Principal coordinate analyses

for the microbiome. C) Relative abundance of AMR genes by antimicrobial class. D) Relative

abundance of the 19 most abundant bacterial genera in the wastewater and sewage microbiome.

Abbreviations: CP=collection point within hospital, Seafield=community sewage works,

TB=tuberculosis.

Figure 2 **Heat map of 50 most abundant antimicrobial resistance genes.**

Relative abundance of AMR genes (FPKM) were log transformed and both AMR genes and

collection points were clustered using complete-linkage clustering. For AMR genes clustering

was based on Pearson correlation coefficients, for collection points clustering was based on the

BC-dissimilarity matrix (Figure 1) which uses all genes.

Figure 3. Generalised linear mixed effects model for the relationship between hospital

department characteristics and antimicrobial resistance gene abundance.

Effect of antimicrobial usage (AMU) measured in defined daily dose per 100 occupied bed

days (DDD/100 OBDs) on AMR gene abundance, controlling for cross –resistance by

including total AMU per collection point, average age and average length of stay (LOS),

including an interaction effect between AMU and LOS.

*= $p < 0.05$, LOS = length of stay, AB = antibiotics, CP = collection point.

489   Figure 1

493   Figure 2

498 **Figure 3**