1 **The day-to-day experiences of caring for children with Osteogenesis Imperfecta: A**
2 **qualitative descriptive study**
3

Authors

Aimee R. Castro[a,b] MSc
Jessica Marinello[a,b], BSc
Khadidja Chougui[b], BA
Marilyn Morand[b], BSc
Claudette Bilodeau[b], BSc RN
Frank Rauch[b], MD
Argerie Tsimicalis[a,b], PhD RN

12
13 [a]*Ingram School of Nursing, McGill University, Montréal, Canada*
14 [b]*Shriners Hospitals for Children-Canada, Montréal, Canada*
15

Acknowledgements

Funding: This work was generously supported by the Tunis Shriners, the Newton
Foundation, Scotiabank®, and the McGill University Ingram School of Nursing
Summer Bursary Programs of 2018 and 2019. Dr. Tsimicalis is supported by a
Chercheur-Boursier Junior 1 from the Fonds de Recherche Québec-Santé (Quebec
Medical Research Council). Ms. Castro is supported by a Fonds de Recherche Québec-
Santé Formation de Doctorat (doctoral research award).
23
Colleagues: Several Shriners clinicians and caregiving stakeholders also provided
invaluable feedback on the interview guides (Johanne Brunelle, RN; Sylvie-Anne
Plourde, RN; Catherine Dubé, BSW; Trudy Wong, MSW; Mary Curtin, BA; Maria
Caruso, BA, DipEd, CertSpEd; Marie Donato, BA, DipEd, CertEd; and Angela
Gugliotti, BA). Finally, this study could not have been conducted without the caregiver
participants who took time to share their knowledge and expertise.

Declaration of Interest

The authors report no conflicts of interest.
32

Abstract

Purpose: Osteogenesis Imperfecta (OI) is a rare genetic condition known to cause

bone fragility. Family caregivers, such as parents, of children with OI play an

important role in helping these children live well at home. This study aimed to

explore the day-to-day experiences of caregivers who are caring for children with

OI.

Materials and Methods: A qualitative descriptive study was conducted. Eighteen

adult caregivers of children with OI were recruited at a children's hospital in

Montréal, Canada to participate in individual interviews. The interviews were

transcribed and inductively thematically analysed.

Results: The following caregiving themes were identified in these interviews:

regular day-to-day caregiving activities, including morning routines, evening

routines, and the facilitation of their child's mobilization; periods that made the

caregiving routine more challenging, such as fractures, surgeries, and pain; and

the long-term strategies caregivers employed to support day-to-day care, such as

managing the environment, accessing medical and school resources, and

coordinating care and respite.

Conclusions: The results showcase what being a caregiver for a child with OI

involves on a day-to-day basis. The recommendations include suggestions for

future clinical, policy, and research endeavours to better support family caregivers

of children with OI.

Keywords: caregiving; osteogenesis imperfecta; rare diseases; chronic care;

brittle bone disease; parenting; paediatric care

<u>Introduction</u>

Osteogenesis Imperfecta (OI) is a rare genetic condition affecting 1 in 10 000

births that is predominantly caused by an alteration in the genes responsible for collagen

production (Marini et al., 2017; Trejo & Rauch, 2016). It is characterized by increased

bone fragility, but the condition can also cause teeth and soft tissue abnormalities,

discoloured sclera, hearing loss, shortened stature, and skeletal deformities (Marini et

al., 2017; Trejo & Rauch, 2016). OI Types I – IV are the original classifications of OI,

although other types have been added since (Trejo & Rauch, 2016). OI Type I is the

most common form of the condition and has the mildest disease severity, with fewer

fractures expected over the life course. OI Type II is lethal in the neonatal period. OI

Type III is the most severe form of OI that is compatible with a longer lifespan. OI

Type IV is of moderate disease severity. OI Types V, VI, and VII are newer

classifications that present with different tissue and bone mineralization phenotypes,

and have mild to moderate severity (Marini et al., 2017; Trejo & Rauch, 2016). At

present, there is no cure for OI. Bisphosphonate therapy is used to improve bone mass

density (Trejo & Rauch, 2016). Surgery, physical therapy, and occupational therapy are

other common treatment modalities (Dogba et al., 2013; Marini et al., 2017).

When a child has OI, a parent usually adopts the role of primary caregiver to

assist with their child's everyday needs (Vanz, Felix, da Rocha, & Schwartz, 2015).

Challenges unique to OI caregiving include the constant risk of fractures, over-

protectionism, social isolation, home adaptations, and suspicions of child abuse

(Arabaci et al., 2015; Bernehall & Brodin, 2002; De Carmoy, 2004; Dogba, Rauch, Tre,

Glorieux, & Bedos, 2014). OI parents have reported lower scores in the environmental

quality of life domain (i.e., the domain which includes factors such as environmental

safety and access to social supports) compared to the general population of parents

(Szczepaniak-Kubat, Kurnatowska, Jakubowska-Pietkiewicz, & Chlebna-Sokol, 2012;

Vanz et al., 2015). The OI caregiver experience may generate positive experiences for

families, such as the development of an optimistic mindset and being inspired by their

children's resilience (Dogba et al., 2013). However, for many families, a child's OI

diagnosis appears to be a life-altering event requiring various and ongoing parental

adjustments (Arabaci et al., 2015; Dogba et al., 2013; Murphy, Christian, Caplin, &

Young, 2007).

It is important to understand the day-to-day experiences of caregivers of children

with unique needs like OI in order to develop services that are adapted to the specialized

daily care needs of these families (McCann, Bull, & Winzenberg, 2012; Pearlin,

Mullan, Semple, & Skaff, 1990). While several studies on the OI family experience

have included some analysis of the day-to-day lives of OI caregivers (Bernehall &

Brodin, 2002; De Carmoy, 2004; Deatrick, Knafl, & Walsh, 1988; Dogba et al., 2013;

Dogba et al., 2014; Hill, Baird, & Walters, 2014; Santos, Pires, Soares, & Barros,

2017), none of these studies focused primarily on the minutiae and routines of daily

care.

Therefore, the purpose of this study was to explore the day-to-day experiences

of caregivers who are caring for children with OI. We interviewed 18 adult caregivers

of children with OI about their day-to-day OI caregiving experiences. They described

their routine activities of day-to-day caregiving, their experiences during challenging

periods of caregiving, and the long-term strategies they used to support their child's

day-to-day care needs. These findings have important implications for clinical practice,

policy, and research.

<u>Materials and methods</u>

Design

This qualitative study was part of a larger one which sought to explore the views

of OI caregivers on using Internet-based technologies to support their caregiving needs

(manuscript available as a preprint) (Castro, Chougui, Bilodeau, & Tsimicalis, 2019).

The study methodology was qualitative description (Sandelowski, 2000). Ethical

approval was received from McGill University's Institutional Review Board (Study

Number A05-B39-17B) and the study site prior to recruitment.

Participant recruitment

A purposive sampling strategy was used to recruit family caregivers of children

who were being treated for different types of OI at the study site. 19 caregivers were

approached in-person, and all verbally agreed to participate, although one caregiver was

lost to follow-up. Inclusion criteria included all family caregivers of patients being

treated for OI at the hospital site, who considered themselves to be the primary

caregivers of the child, and who spoke either English or French. All participants

provided written consent to participate in the study, with the understanding that they

would not be identifiable in any presentations of the research. All participant data has

been anonymized.

Data collection and analysis

A demographic data survey was filled out by each participant. Individual semi-

structured interviews were subsequently conducted by a member of the research team,

in either English or French depending on the caregiver's preference. The interviews

were conducted by either author AC or KC, at the study site or by telephone or

videoconferencing. Key questions relevant to day-to-day care from the larger study's

interview guide are listed in Table 1. The interview guide was reviewed by key

informant clinicians prior to the participant interviews. Interviews were audiotaped,

field notes were recorded during and after the interviews, and a reflective journal was

kept by author AC. The surveys were analyzed using descriptive statistics. The

transcribed data and field notes were uploaded into Excel, where they were first

inductively open-coded by three of the authors (AC, JM, and KC), and then further

analyzed primarily by AC using (Braun & Clarke, 2006)'s six steps for thematic

analysis. Interviews conducted in French were first open-coded into English by authors

KC and JM who are fluent in both English and French. In frequent meetings with the

principal investigator (AT), the research team discussed the ongoing development

of the codebook by AC and established the final themes. These codebooks and

meeting notes were recorded in a research journal, as were the ongoing research

reflections of AC.

Table 1. Key interview questions and prompts related to day-to-day care.

Main questions:
1) Can you describe what a regular day looks like when caring for your child with OI?
2) And then can you describe a more challenging day recently?
Prompts:
-What are some of the duties or responsibilities of caring for a child with OI?
-How has your caring for your child changed over time?
-What areas of your family's life do you feel are going well?
-What are some areas that you would like to see improve?

-What happens on a day with a bone break? What happens in the days that follow?
-Do you feel you know about community resources for families of children with chronic conditions?
-How has caregiving affected your lifestyle and schedule? Your job? Your relationships? Your personal care?
-How do you feel about your social support?
-Do you have access to extra help and/or care around the house?

Results

Demographic Data

Eighteen caregivers from 14 families participated. Thirteen caregivers were

women and five were men. Three families resided outside of Canada. Four of the 14

families shared that they had family incomes of less than $50 000 per year, and all 18

caregivers had received at least some post-secondary education. These interviews

resulted in 13.05 hours of audio recordings and over 8000 spreadsheet lines of coded

transcript. Their children's OI types included I, III, IV, and VI; types III and IV were

the most common. Some children with OI required no mobilizing assistance, while

others used a wheelchair full-time. Further demographic data are listed in Table 2.

Table 2. Demographic data.

Demographic Trait	**n (individuals, unless otherwise indicated)**
Individuals interviewed	18
Number of families represented	14
Language of interview (as caregiver preferred)	
English	12
French	6

Median caregiver age in years (range)	37.5 (24-57)
Caregiver's sex (/18 caregivers)	
Female	13
Male	5
Marital Status (/18 caregivers)	
Married or common-law	14
Single (never married)	2
Separated or divorced	2
Residential region (/14 families)	
Local (< 2 hours' drive from Montreal)	7
Within Québec, not local	1
Other Canadian region	3
International	3
Highest level of education (/18 caregivers)	
Some post-secondary (university or college)	5
Received university or college degree/diploma	11
Postgraduate	2
Estimated family income (/14 families)	
Less than $25 000	3
$25 000 - $50 000	1
$50 000 - $80 000	1
More than $80 000	7
Do not know	1
Prefers not to answer	1
Relationship to child with OI (/18 caregivers)	
Mother	11
Father	5
Other: legal guardian	2
Child's OI type (/14 families)	
I	3
III	4
IV	6
VI	1
Number of families with more than 1 child with OI in the household	3
Ages of children with OI (17 children with OI)	
Baby (0-12 months)	2
Toddler (13 months - 3 years old)	1
Pre-school (4-5 years old)	1

School-aged (6-12 years old)	11
Teenager (13-18 years old)	2
Range in number of fractures / surgeries per lifetime in the children with OI Note: Several families did not mention the number of fractures or surgeries their children had experienced. This question was not explicitly included in our interview guide or questionnaire.	1 - 200 fractures / 0 - 40 surgeries
How would you describe the health of your child with OI right now? (/18 caregivers)	
Excellent	5
Very good	4
Good	8
Fair	1
Poor	0
Family history of OI (/14 families)	
Yes	3
No	11

Note: This table has been adapted for this report from the demographic data table

portrayed in (Castro et al., 2019). The two tables are similar but not identical.

Caregiving Themes

We inductively derived the following themes relating to the day-to-day

experiences of OI caregiving: regular day-to-day caregiving activities, challenging

periods, and the long-term strategies caregivers used to support day-to-day care.

Regular day-to-day caregiving activities

OI caregivers were asked to share what a regular OI caregiving day looked like.

Key components of these day-to-day activities included following morning and evening

routines to assist with their child's activities of daily living (ADLs), and facilitating

mobilization.

Morning and evening routines. Mornings were particularly challenging for

caregivers, when many ADLs tended to occur all at once, and all tasks needed to be

completed before school and paid work began. Mornings consisted of waking everyone

up, making lunches, helping children get dressed, and using the bathroom. Some

children needed assistance in dressing with their leg braces or corsets. Children needing

the most assistance with ADLs were younger children and those with more severe types

of OI. Chronic pain management was also a part of the morning routine for at least two

families. Most families chose to drive their children to school, although one child used

an adapted bus. One mother said her son would walk to school if it was nice outside, but

if his backpack was heavy or the sidewalk was icy, she would drive him to school,

instead. Another family explained that wheelchair ramps were often too icy or snowy

during winter to be useful. Other caregivers shared that they did not use the adapted

school bus because it arrived too early to be useful for them in their busy mornings. So,

sometimes, services meant to be accessible were not truly accessible for these families.

Evenings were described as being more relaxed than mornings. In the evenings,

caregivers made dinner, had their children complete chores as they were able to, played

games with their children, talked about school, did homework, and listened to

audiobooks as a family. Caregivers took their children to specialized and frequent

physical therapy and occupational therapy appointments. Some children engaged in

adapted sports, various artistic classes, and other low-impact activities. Older children

with fewer accessibility challenges would go outside to play with friends on their own.

Facilitating mobilization. Mobilizing or transferring their children during these

activities was conducted carefully. Some school-aged children would crawl or slide

around on their buttocks, reserving walkers and wheelchairs for more fatiguing times.

One child with a milder form of OI needed her family's regular encouragement to move

on her own, particularly up stairs where she had more difficulties. Transfers presented

with some of the greatest difficulties for families. Many of the children needed at least

some assistance in transferring to and from the bed, car, wheelchair, or toilet. One legal

guardian shared that "just moving him from point a to point b is most challenging" (Guardian [Mother], Family 8, Type VI). While they typically lifted their school-aged child into the truck, it could be a challenge to lift him as he was getting bigger and heavier, especially during icy and slippery winters. For families that had children with more severe types of OI, all of these activities facilitating mobilization needed to be accomplished without causing further pain or fractures. In contrast, families with Type I OI generally reported not noticing any significant differences in the day-to-day requirements of caring for their child living with OI, compared to caring for a child without OI.

Challenging Periods

OI caregivers were constantly aware that the regular daily routines they had developed could be disrupted by fractures, surgeries, and pain. These challenging periods increased the care management needed.

Fractures. The initial stress of bone breaks came from their unexpectedness. The first fractures in their child's life caused some caregivers to panic, although caregivers said they learned to adapt. One mother shared that a challenging day for her was when she was called from her office due to a fracture at school:

> The challenging time is when I'm called from the office that there's a fracture. Sometimes she fractures in school, sometimes she fractures when she comes back. And I will have to quickly rush home to go see the severity of the fracture, whether it's just a crack, or it's a real fracture where we have to go to the hospital. So basically, that's when it becomes challenging. (Mother, Family 6, Type III)

She explained that she did not like going to the hospital because she said the hospitals in her country did not understand OI; she felt stigmatized in hospitals.

With more severe forms of OI, a fracture could happen practically spontaneously. One caregiver explained how easily a femur fracture occurred:

He was jumping on my bed. And, you know, like a normal four-year-old would
do. We were doing laundry, he was jumping on the bed, and with that, we heard
a *snap*. And he is *screaming*, leg bevelled out, and we're like "well, that's
strange." And then we went to the hospital, and they're like, "okay, it's a femur
fracture." And that happened about two other times after that. Then they're like,
"okay, we're going to test him [for OI]" (Guardian [Older Sister], Family 8,
Type VI)

Another caregiver shared that whenever her child fractured, she immediately assessed

the fracture and administered appropriate medications. Then, her child relaxed for the

rest of day. Other parents shared that in the case of severe fractures, they would go to

the hospital and wait for treatment and recovery.

Challenges of surgery. Surgeries presented unique challenges for families.

Hospital boredom was often a challenge for parents, especially if they had to fly to

Montréal for surgery and wait several days or weeks before returning home. A few

caregivers discussed how challenging planning their lives around surgeries could be –

especially when, after arranging their work and family schedules around a specific date,

the surgery date was sometimes changed by the hospital.

Care coordination for treatment created additional challenges for the three

international families interviewed. Caregivers shared that accessing OI treatment in

their own countries could be difficult, so a few of them received pro-bono treatment

supplies through the international OI community. For a mother whose daughter needed

a few surgeries over several months, finding a Montréal sponsor to live with (as may

sometimes be required) was a significant care coordination challenge for her. She said

she had to somehow find a sponsor to live with for several months, in a city far from her

own, while also arranging care with her husband and their other children who would

stay behind in her home country. She explained:

So I leave [the other children] with my husband when I came to the [Montreal
OI treatment centre] in August . . . I think from the way it is, [the treatments]

1 will end by December . . . You understand, I have my kids calling me, 'Mommy,
2 when are you coming back?' . . . it's a very long time. (Mother, Family 13, Type
3 III)

4 ***Caregiving after a fracture or surgery.*** Caregiving became more difficult after a

5 fracture or a surgery, with the child's limb or limbs immobilized and unable to bear any

6 weight. Immobility required more assistance with ADLs and made transfers more

7 challenging. Leg fractures were particularly problematic because the child was then

8 non-weight-bearing on that limb, making transfers to and from places - such as the

9 bathtub, toilet, or car - heavy and complex for the caregiver. When children fractured,

10 time for self-care and outside excursions could become more difficult:

11 Um, especially when they're fractured, it's not as easy to get out. You don't
12 want to be transferring them when they're in pain, getting them into their car-
13 seat. And when you're the only person at home 'cause your husband's working,
14 if you need to go to the grocery store, then you have to take them with you. So
15 sometimes you're not necessarily getting yourself ready in the morning because
16 you're just trying to get them into a position where they can get out of the house
17 at some point or for an appointment. (Mother, Family 13, Type III)
18

19 After surgery, the family's routine would change. One mother explained that

20 after a fracture, her child would sleep in the same bedroom as her, receive more pain

21 medication, and stay home for a few days post-surgery. Fractures and surgeries created

22 stress for caregivers during ADLs, as they were worried about any jostling during care

23 work that could harm the child. As a result, caregivers tended to minimize activity and

24 stay at home during these periods.

25 ***Pain.*** For some families of children with OI, life could revolve around pain

26 management. One mother explained that when her school-aged son was in pain, he

27 would call for "Mommy!" a lot more frequently (Mother, Family 5, Type III). Another

28 mother shared that her child was often in physical pain, particularly experiencing

29 regular headaches and back pain. As a caregiver, this mother had to evaluate the pain

level two to three times per day and administer pain medications as needed. Two

parents also shared that their children experienced regular back pain in spite of their

fractures being healed. Periods of increased pain, then, created frequent disruptions to

OI caregivers' daily routines.

Long-Term Strategies for Supporting Day-to-Day Care

A comment by one mother embodied the management of day-to-day care:

Dealing with OI is most likely all the time, how are we going to do things? You
know? How are we going to surpass this limitation? (Mother, Family 5, Type III)
9
Caregivers learned strategies which would make future day-to-day caregiving work

easier. They learned to manage the environment, access medical and school resources,

and coordinate care and respite.

Managing the environment. Generally, caregivers tried to balance their desire

to protect their child from harm, while also minimizing the limits placed on their child

day-to-day. Managing care for caregivers often meant "managing the environment more

than the kid":

We make sure that there is no chair or anything that could actually fall on her.
the remote for the TV are never actually accessible if she's standing from the
couch, because . . . it's heavy. It's an old remote. We do play stuff differently,
without saying "no you can't" or "don't do that" or "don't touch that". It's really
just managing the environment more than the kid . . . [The] environment is our
responsibility. (Mother, Family 1, Type IV)
23
Managing the environment included adapting their housing to be safer (for instance, by

installing extra-cushioned carpeting), coordinating with schools to manage the care of

their child safely, and searching for low-impact activities that their children could

participate in. With these environmental management strategies, future day-to-day care

work was made easier, because their homes and schools were more prepared for a child

with brittle bones to thrive in, and their children had low-impact activities and skills that

they could develop.

Accessing medical and school supports. Caregivers described spending a lot of their caregiving time researching information and resources about OI, particularly for accessing medical and school supports. Once specialized medical and school supports were established, caregivers felt more at ease in their day-to-day care work.

Accessing medical support. For several caregivers, just because they had a diagnosis of OI did not mean they had adequate medical support and information for ongoing care. Caregivers shared that it was only once they found the Montréal OI treatment centre that they felt supported in their child's medical care. Healthcare providers outside of the treatment centre – such as emergency staff and general pediatricians at other hospitals – did not always know how to treat OI. Some parents shared that when other health professionals tried to treat their children, the clinicians fractured their child's other bones in the process. One Québecoise mother believed not enough information was initially shared with her about her rights and available resources. She explained that when her daughter was a pre-schooler, the two of them spent an entire year avoiding all outings because they did not know how to mobilize her growing daughter beyond using a baby stroller. They were socially isolated and home-bound until she was told that her daughter had the right to access a properly fitted wheelchair.

Access to the greater OI community was an important resource for finding long-term medical supports. Several caregivers spent a lot of time seeking resources through their OI network. For instance, a few caregivers managed to receive regular OI treatment materials in their own country via the international OI community when they could not access those materials at home. Once caregivers managed to access long-term OI medical supports, either with an established OI social network or – ideally – with a specialized OI treatment centre nearby, caregivers felt more supported in their day-to-

day caregiving activities, because they had access to people who were knowledgeable

about OI; they were no longer caregiving alone.

Accessing school supports. For some families, finding daycare centres and

schools that were comfortable with a child with OI was a challenging task. Once they

found an accommodating school, several families described having to go into the school

every year to share with the new educators how to care for a child with OI. These new

school caregivers had to be taught:

How do we immobilize a fracture, how do we give the medication, and how do
we put him on the toilet seat . . . how do we take care of him? (Mother, Family
5, Type III)
11
Most children were attending public schools. However, they still required access to a

school aide person who could help them in the washroom, carry books, push the

wheelchair, or write for them, as needed.

A few caregivers were using more specialized school services. One child moved

to Montréal with her mother to attend a school for children with special needs, while the

child's father and siblings remained in their hometown several hours away. Another

family was choosing to homeschool their child instead of sending him to a public

school, because they wanted to be sure he was mobilizing enough during the day. They

feared that a public school would be too scared of him breaking a bone that they would

not allow him to mobilize enough.

Other routine medical and school resource challenges included the availability

of services, language barriers, and Internet access. Even among financially secure

families, some families could not find appropriate daycare centres for their child with

more severe OI. One mother explained that there was a long waiting list for daycare for

children with special needs. Language barriers could also affect resource accessibility.

Some families noted that the lack of OI resources available in languages outside of

English (such as French) were limited. Access to the greater OI community through the

Internet could also determine whether families managed to find a specialized OI

treatment centre and other OI resources.

Coordinating care and accessing respite. Caregivers described many care

coordination tasks related to their child's OI. These included the following tasks:

arranging and attending numerous medical appointments and treatments, remembering

to bring the wheelchair or walker whenever they went out, managing respite care, being

flexible with schedules in case of surgery or fractures, keeping track of medical records,

seeking out community and health resources, and filling out medical paperwork and

reimbursement forms. All of these activities required significant time and coordination

skills on the part of caregivers, which frustrated some of the caregivers. However, once

these coordination activities were accomplished, they made caregivers' future day-to-

day care work easier. For instance, by keeping track of their child's medical records and

being flexible with their schedules, caregivers were able to quickly adapt if medical

emergencies arose.

Most caregivers expressed chronic challenges with accessing respite care,

explaining that they did not trust others to care for their child with OI. One mother said

that learning to trust others is "sort of an issue" (Mother, Family 1, Type IV). Several

caregivers shared that they were not comfortable hiring a babysitter for their child with

OI, or even leaving their child with friends or family. Still, some families had

grandparents and friends who were trying to learn how to care for their child in order to

provide these caregivers with parenting breaks. Two mothers suggested that more

trusted respite care opportunities, such as kids' events or summer camps organized by

OI treatment centres, would be very helpful. If caregivers had easy access to people

they trusted to care for their child, day-to-day care was made easier because caregivers

had a reliable back-up caregiver. However, most caregivers did not have anyone they

fully trusted to provide this care.

<u>Discussion</u>

4   This study offers a comprehensive overview of the day-to-day experiences of

caring for a child with brittle bones, the challenging periods that can disrupt day-to-day

routines, and the long-term strategies they used to make day-to-day OI caregiving

easier. Understanding the unique daily experiences of caregivers of children with rare

and chronic conditions like OI is important knowledge for helping clinicians and policy

makers to build better supports for these caregivers. Such supports could improve

caregivers' daily experiences of caregiving, and therefore improve their child's care

(Luijkx, Putten, & Vlaskamp, 2017; McCann et al., 2012).

12   The limited existing OI literature suggests that OI parents often have more tasks

to accomplish in one day than most parents do. Even on typical days without fractures

or surgery, a key difference is the child's independence in completing ADLs. For more

severe types of OI, in particular, the multiple physical transfers needed for toileting,

clothing, and transport increased the energy and amount of time it took to accomplish

these ADLs (Deatrick et al., 1988; Dogba et al., 2014). Facilitating mobilization and

transportation for those living with rare and physical conditions is a common but

challenging caregiving task (Andrews, 2013). However, in addition to managing the

task of mobilization, OI caregivers of children with more severe types of OI also

experience the unique and constant fear of fracturing their child while completing

transfers and other day-to-day tasks (Brodin, 1993). Our study corroborates these

reports of the experiences of OI day-to-day caregiving. In our study, caregivers shared

that mornings were the most intense period of the day due to the number of ADLs

needing attendance by a strict deadline (i.e., when school and caregivers' paid work

began). Furthermore, and in accordance with the research literature, facilitating their

child's mobilization and transportation was one of the most significant components of

caregivers' day-to-day routines.

For OI caregivers in our study, periods which presented particular challenges to

OI caregivers' day-to-day routines were periods with fractures, surgeries, and pain.

These time periods were also described as some of the most challenging aspects of OI

care in other studies of OI families (De Carmoy, 2004; Deatrick et al., 1988; Dogba et

al., 2013; Santos et al., 2017). However, while the caregivers in our study

acknowledged facing challenges and at times feeling overwhelmed, most appeared to be

successfully managing – if not always thriving – as parents and caregivers; this was true

even for the parents of children with more severe OI. Caregivers explained that once

they found the study site (a specialized OI treatment centre), they felt more supported as

caregivers. These results correspond with the results of another study which was also

conducted at our study site (Dogba et al., 2013). However, these results contrast with

other studies on OI caregiving which described caregivers feeling burned out, hopeless,

and overwhelmed by the unique challenges of caring for children with brittle bones

(Arabaci et al., 2015; Bozkurt et al., 2014; De Carmoy, 2004). These latter studies do

not appear to have been conducted in settings with access to specialized OI services.

This contrast may speak to the psychosocial benefits that access to specialized OI

medical resources can have on OI families.

The literature suggests that managing the environment, accessing medical and

school supports, coordinating care, and finding trusted respite services are all common

care management strategies of chronic caregivers (Pelentsov, Laws, & Esterman, 2015;

Samia, O'Sullivan, Fallon, Aboueissa, & Hepburn, 2018). Other researchers have noted

that OI caregivers spend significant amounts of time trying to gain reliable access to OI

medical and school supports, social supports, and respite care (Arabaci et al., 2015;

Bernehall & Brodin, 2002). These strategies for facilitating day-to-day OI care were

described by caregivers in our study, as well.

Implications for clinicians and policymakers

This study has implications for pediatric clinicians. Finding information about

OI, accessing treatment (especially for families who lived far away from the treatment

center), finding knowledgeable local healthcare providers, and coordinating available

community and school supports, were all challenges faced by the caregivers in our

study. Clinicians and clinical researchers should work to develop clinical home-care

tools to provide long-term supports for OI families. For instance, the study site has

developed a splint kit that helps caregivers splint their children's fractures at home; such

a kit is particularly valuable when families do not have knowledgeable healthcare

professionals nearby to assist them (Leclair, 2018). Similar resources could further

support caregivers in helping their children to thrive with OI.

OI medical services should be more accessible. For instance, OI specialists

should be easier to find. OI caregivers shared that their day-to-day caregiving

experiences were improved once they had access to the study site and specialized OI

care. However, a few caregivers also shared that the time taken to travel to

appointments, to complete surgeries, and to fill out reimbursement forms was too time-

consuming and complicated. Institutions and policymakers should work to ensure that

all medical services are accessible, efficient, and easy to navigate, with as little family

separation as possible.

Policymakers should focus on building supports for trusted respite care and

other high-quality services. Policymakers should work to build more comprehensive

and qualified respite support services for caregivers of care recipients with complex

needs. A few caregivers in this study suggested that child care services provided by

specialized treatment centres, such as special drop off days or summer camps, would be

appreciated. Another option might be to develop OI caregiving teaching materials to

help caregivers teach their friends and extended family on how to care for their child

day-to-day. A few of the caregivers mentioned that even when they had community

supports like adapted buses and wheelchair ramps, those services often did not meet

their unique caregiving needs, so they did not use them. Policymakers should work to

ensure that such services are adapted and truly accessible to families with complex care

needs, so that these families can learn to trust the services developed for them.

Strengths and Limitations

Some of our study's methodological strengths are that it provides rich

descriptions of the study setting, participant demographics, and methods for

transferability purposes. Given the time and resources available for the study, we

collected and analysed a large amount of interview data. Other strengths of our study

are its contributions towards addressing the paucity of literature available on OI

caregiving, as well as on the supportive care needs of parents of children with a rare

condition (Pelentsov et al., 2015). While corroborating much of the current OI

caregiving literature, our study also adds to our knowledge of OI caregiving by

providing more detailed data on what day-to-day OI caregiving looks like. This study

appears to be the first to focus on the day-to-day tasks and routines of OI caregiving.

Understanding the daily life of family caregivers is important because the differences

required in care for caregivers of children with unique health needs can cause

significant caregiver distress (Pelentsov et al., 2015). Furthermore, understanding their

unique needs can support the development of evidence-based health and social supports

for these families (Collins et al., 2016; Luijkx et al., 2017).

This study has several limitations. With the time and resources available for the

study, only 18 caregivers from 14 families were interviewed. The sample could have

benefited from more socioeconomic caregiver diversity. All caregivers had some post-

secondary education, and a majority of the families (n=7) reported incomes of more

than $80 000 per year. Only one caregiver was under the age of thirty, and all but two

parents said they had another adult helping them at home, such as a spouse, adult child,

or grandparent. The needs of younger and single-parent OI caregivers should be

researched further. Finally, all of these families had access to the study site, which

specializes in OI treatment. Researchers should explore the experiences of families who

have more limited educational and financial resources, and who do not have access to a

specialized centre for OI. Since the majority of the caregivers in this study had school-

aged children (n=11 of the 17 children), more research should be conducted on the

caregiving experiences of those caring for younger children, as well as those caring for

teenagers. Further qualitative and quantitative research should be conducted to build on

this study's work to better understand the diverse range of OI caregivers' day-to-day

caregiving experiences.

Conclusion

In conclusion, the rich data collected from these interviews helped elucidate the

day-to-day activities and challenges faced by OI caregivers. The results complement

previous research in the caregiving and OI research fields, while also adding a deeper

understanding of the day-to-day experiences of OI caregivers. This knowledge may help

clinicians and healthcare systems offer improved services to these families. It also

provides further direction for future research initiatives to help promote the health and

well-being of OI caregivers.

References

Andrews, M. (2013). Measuring exertion during caregiving of children and young adults with cerebral palsy who require assistance for mobility and self-care. *Phys Occup Ther Pediatr, 33*(3), 300.

Arabaci, L. B., Bozkurt, S., Vara, S., Ozen, S., Darcan, S., & Simsek, D. G. (2015). Difficulties experienced by caregivers of patients diagnosed with osteogenesis imperfecta (OI): Example of a hospital. *J Pak Med Assoc, 65*(7), 764-770.

Bernehall, C. I., & Brodin, J. (2002). What families with children with brittle bones want to tell. *Child: Care, Health and Development, 28*(4), 309-315. Retrieved from http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed8&NEWS=N&AN=34846908 https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2214.2002.00282.x

Bozkurt, S., Baysan Arabaci, L., Vara, S., Ozen, S., Goksen, D., & Darcan, S. (2014). The impact of psycho-educational training on the psychosocial adjustment of caregivers of osteogenesis imperfecta patients. *J Clin Res Pediatr Endocrinol, 6*(2), 84-92. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/24932601

Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qual Res Psychol, 3*(2), 77-101. doi:10.1191/1478088706qp063oa

Brodin, J. (1993). Children and adolescents with brittle bones — psycho-social aspects. *Child: Care Health Dev, 19*(5), 341-347. doi:10.1111/j.1365-2214.1993.tb00738.x

Castro, A., Chougui, K., Bilodeau, C., & Tsimicalis, A. (2019). Exploring the Views of Osteogenesis Imperfecta Caregivers on Internet-Based Technologies: A Qualitative Descriptive Study. *JMIR Preprints*. doi:10.2196/preprints.15924

Collins, A., Hennessy-Anderson, N., Hosking, S., Hynson, J., Remedios, C., & Thomas, K. (2016). Lived experiences of parents caring for a child with a life-limiting condition in Australia: A qualitative study. *Palliat Med, 30*(10), 950-959.

De Carmoy, R. J. (2004). Children and adolescents with brittle bones: How can they overcome their disability? *Neuropsychiatrie de l'Enfance et de l'Adolescence, 52*(6), 422-429. doi:http://dx.doi.org/10.1016/j.neurenf.2004.07.004

Deatrick, J. A., Knafl, K. A., & Walsh, M. (1988). The process of parenting a child with a disability: normalization through accommodations. *J Adv Nurs, 13*(1), 15-21.

Dogba, M. J., Bedos, C., Durigova, M., Montpetit, K., Wong, T., Glorieux, F. H., & Rauch, F. (2013). The impact of severe osteogenesis imperfecta on the lives of young patients and their parents - a qualitative analysis. *BMC Pediatr, 13*(1), 153. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/24074180

Dogba, M. J., Rauch, F., Tre, G., Glorieux, F. H., & Bedos, C. (2014). Shaping and managing the course of a child's disease: parental experiences with osteogenesis imperfecta. *Disability & Health Journal, 7*(3), 343-349. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/24947576

Hill, C. L., Baird, W. O., & Walters, S. J. (2014). Quality of life in children and adolescents with Osteogenesis Imperfecta: a qualitative interview based study. *Health & Quality of Life Outcomes, 12*(1), 54. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/24742068

Leclair, A. (Producer). (2018, 2018/08/25). EXCLUSIVE: Saskatchewan boy creates splint kit for fellow patients at Shriners Hospital for Children. Retrieved from https://globalnews.ca/news/4068922/exclusive-saskatchewan-boy-creates-splint-kit-for-fellow-patients-at-montreal-shiners-hospital/

Luijkx, J., Putten, A. A. J., & Vlaskamp, C. (2017). Time use of parents raising children with severe or profound intellectual and multiple disabilities. *Child: Care Health Dev, 43*(4), 518-526. doi:10.1111/cch.12446

Marini, J. C., Forlino, A., Bachinger, H. P., Bishop, N. J., Byers, P. H., Paepe, A., . . . Semler, O. (2017). Osteogenesis imperfecta. *Nat Rev Dis Primers, 3*, 17052. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/28820180

McCann, D., Bull, R., & Winzenberg, T. (2012). The daily patterns of time use for parents of children with complex needs: A systematic review. *J Child Health Care, 16*(1), 26-52.

Murphy, N. A., Christian, B., Caplin, D. A., & Young, P. C. (2007). The health of caregivers for children with disabilities: caregiver perspectives. *Child: Care Health Dev, 33*(2), 180-187. doi:doi:10.1111/j.1365-2214.2006.00644.x

Pearlin, L. I., Mullan, J. T., Semple, S. J., & Skaff, M. M. (1990). Caregiving and the Stress Process: An Overview of Concepts and Their Measures. *The Gerontologist, 30*(5), 583-594. doi:10.1093/geront/30.5.583

Pelentsov, L. J., Laws, T. A., & Esterman, A. J. (2015). The supportive care needs of parents caring for a child with a rare disease: A scoping review. *Disability & Health Journal, 8*(4), 475-491. doi:https://doi.org/10.1016/j.dhjo.2015.03.009

Samia, L. W., O'Sullivan, A., Fallon, K. C., Aboueissa, A.-M., & Hepburn, K. W. (2018). Building on self-efficacy for experienced family caregivers: The Savvy Advanced Program. *The Gerontologist*. doi:10.1093/geront/gny016

Sandelowski, M. (2000). Whatever happened to qualitative description? *Res Nurs Health, 23*(4), 334-340. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/10940958

Santos, M. C. D., Pires, A. F., Soares, K., & Barros, L. (2017). Family experience with osteogenesis imperfecta type 1: the most distressing situations. *Disabil Rehabil*, 1-7. doi:https://dx.doi.org/10.1080/09638288.2017.1334236

Szczepaniak-Kubat, A., Kurnatowska, O., Jakubowska-Pietkiewicz, E., & Chlebna-Sokol, D. (2012). Assessment of quality of life of parents of children with osteogenesis imperfecta. *Adv Clin Exp Med, 21*(1), 99-104. Retrieved from http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed14&NEWS=N&AN=364542406

Trejo, P., & Rauch, F. (2016). Osteogenesis imperfecta in children and adolescents-new developments in diagnosis and treatment. *Osteoporos Int, 27*(12), 3427-3437. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/27492436

Vanz, A. P., Felix, T. M., da Rocha, N. S., & Schwartz, I. V. (2015). Quality of life in caregivers of children and adolescents with Osteogenesis Imperfecta. *Health & Quality of Life Outcomes, 13*, 41. doi:http://dx.doi.org/10.1186/s12955-015-0226-4