Travel time to health facilities as a marker of geographical accessibility across heterogeneous land coverage in Peru

Gabriel Carrasco-Escobar[1,2,*], Edgar Manrique[1], Kelly Tello-Lizarraga[3], J. Jaime

Miranda[4,5]

* Corresponding Author: Gabriel Carrasco Escobar, MSc, PhD(c): gabriel.carrasco@upch.pe

[1] Health Innovation Lab, Institute of Tropical Medicine "Alexander von Humboldt", Universidad
Peruana Cayetano Heredia, Lima, Peru

[2] Division of Infectious Diseases, Department of Medicine, University of California San Diego,
La Jolla, CA, USA

[3] Facultad de Salud Publica y Administración, Universidad Peruana Cayetano Heredia, Lima, Peru

[4] CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano
Heredia, Lima, Peru.

[5] School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru

Keywords: Travel time, geographic accessibility, health care accesibility

v4.0

ABSTRACT

The geographical accessibility to health facilities is conditioned by the topography and environmental conditions overlapped with different transport facilities between rural and urban areas. To better estimate the travel time to the most proximate health facility infrastructure and determine the differences across heterogeneous land coverage types, this study explored the use of a novel cloud-based geospatial modeling approach and use as a case study the unique geographical and ecological diversity in the Peruvian territory. Geospatial data of 145,134 cities and villages and 8,067 health facilities in Peru were gathered with land coverage types, roads infrastructure, navigable river networks, and digital elevation data to produce high-resolution (30 m) estimates of travel time to the most proximate health facility across the country. This study estimated important variations in travel time between urban and rural settings across the 16 major land coverage types in Peru, that in turn, overlaps with socio-economic profiles of the villages. The median travel time to primary, secondary, and tertiary healthcare facilities was 1.9, 2.3, and 2.2 folds higher in rural than urban settings, respectively. Also, higher travel time values were observed in areas with a high proportion of the population with unsatisfied basic needs. In so doing, this study provides a new methodology to estimate travel time to health facilities as a tool to enhance the understanding and characterization of the profiles of accessibility to health facilities in low- and middle-income countries (LMIC), calling for a service delivery redesign to maximize high quality of care.

v4.0

1. INTRODUCTION

Despite growing consensus to combat inequalities in the accessibility to healthcare around the world, large disparities in healthcare accessibility remain as a problem in countries with an ongoing rural-to-urban transition. According to the 'Tracking Universal Health Coverage: 2017 Global Monitoring Report', half of the worldwide population lacks essential health services (World Health Organization & World Bank, 2017). To overcome the disadvantage of marginalized populations, the international community through the United Nations (UN) have stated 17 Sustainable Development Goals (SDG) targeted by 2030 (UN General Assembly, 2015). From these goals, the interface between goal 3, — 'Good health and well-being',; and goal 10, — 'Reduced inequalities', play an important role to foster and couple endeavors towards ensured access to healthcare services.

The broad term 'accessibility', when referring to healthcare, focuses on multiple domains such as the provision of healthcare facilities, supply chain, quality and effective services, human resources, and on the demand side, health-seeking behaviors (Agbenyo, Marshall Nunbogu, & Dongzagla, 2017; Peters et al., 2008). All these characteristics pointed to the ability of a population to receive appropriate, affordable and quality medical care when needed (Kanuganti, Sarkar, Singh, & Arkatkar, 2015). Importantly, in rural and high poverty areas the most common reasons that prevents the access to healthcare are the geographical accessibility, availability of the right type of care, financial accessibility, and acceptability of service (Al-Taiar, Clark, Longenecker, & Whitty, 2010; Peters et al., 2008). This study focuses on the travel time to health facilities as an important component of the geographical (or physical) accessibility to healthcare.

Several studies in developing countries report that geographical accessibility is the main factor that prevents the use of primary healthcare (Al-Taiar et al., 2010; Kanuganti et al., 2015; Müller, Smith, Mellor, Rare, & Genton, 1998; Noor, Zurovac, Hay, Ochola, & Snow, 2003; Perry & Gesler, 2000; Stock, 1983), and not only conditions the ability of the population for health seeking, but also the capacity of the health system to implement prevention and control strategies with adequate coverage. However, fewer studies have explored the heterogeneity in geographical accessibility across areas with contrasting land coverage (Bashshur, Shannon, & Metzner, 1971; Comber, Brunsdon, & Radburn, 2011), i.e. the marked variation in the topography and

v4.0

environment conditions overlapped with different transport facilities between rural and urban

areas that may influence the geographical accessibility across these areas. The geographical

accessibility to health services has a direct impact on health outcomes since determine the timely

response to patients that seek care, community-based campaigns (i.e. vaccination, iron

supplements to combat anemia, etc.), or deliver first response to accidents or natural disasters.

Previous studies highlighted the importance of geographical or physical accessibility using a

variety of methods (Comber et al., 2011; Delamater, Messina, Shortridge, & Grady, 2012;

Huerta Munoz & Källestål, 2012; Ouko, Gachari, Sichangi, & Alegana, s. f.). The emergence of

'Precision Public Health' driven by estimates of a wide range of health indicators at a high

spatial resolution is defined as the use of the best available data to target more effectively and

efficiently interventions of all kinds to those most in need (Chowkwanyun, Bayer, & Galea,

2018; Dowell, Blazes, & Desmond-Hellmann, 2016; Horton, 2018; Tatem, 2018). This approach

may be favorable since traditionally government's reports aggregates data at administrative units,

in a way that obscure the prioritization of resources. A recent study used a precision public

health approach to estimate the geographical accessibility to major cities (Weiss et al., 2018),

however, this approach has not yet been used for estimating the geographical accessibility to

health facilities in developing countries.

This study sought to estimate the travel time to the most proximate health facility in rural and

urban areas across heterogeneous land coverage types in Peru as a means to help resources

prioritization, disease surveillance, as well as prevention and control strategies. Multiple sources

of geospatial data were fitted with a novel cloud-based geospatial modeling approach (Weiss

et al., 2018) to produce high-resolution (30 m) estimates of travel time to the most proximate

health facility across the country. These estimates were then compared between urban and rural

settings and across 16 major land coverage types in Peru.

2.   MATERIALS AND METHODS

2.1. Study design

Ecological study using the Peruvian registry of villages and health facilities to model the travel

time required for individuals in each village to reach the most proximate health facility (shortest

v4.0

travel time) in a two-step process. First, a friction surface was computed. Several geospatial

datasets (land coverage types, boundaries of restricted areas, roads infrastructure, navigable river

networks, and topography) were used to construct a surface (i.e. raster or grid) of a given spatial

resolution (i.e. 30m per pixel) where the value of each pixel (or cell) contains the time required

to travel one meter in that given area. Secondly, this friction surface and the geolocation of the

health facilities were used to infer the travel time to the the most proximate (lowest time) health

facility using a cumulative cost function. As a result, the travel time estimate for the most

proximate health facility was computed for the entire country. The computed values were then

summarized in a 500m-radius from the geolocation of cities and villages; per district, province or

department; by urban/rural areas; and across 16 major land coverage types defined by the

Ministry of Environment (MEnv).

2.2. Study area

This study was conducted using nationwide data from Peru, located on the Pacific coast of South

America. Peru encompasses an area of 1,285,216 Km_2 and 32,162,184 inhabitants divided in 25

departments and 1,722 districts. Major ecological areas in the country are divided into the coast,

highlands, and jungle (**Figure 1A**), however this study explore a higher granularity of ecological

areas with more than 60 unique land coverage areas (**Supplementary Information 1**) that were

officially classified in Peru. This classification was based on ecological, topographic, and climate

characteristics, that in turn are important for the calculation of travel time since each category

requires a different displacement effort.

2.3. Data Sources

The datasets were divided according to its use in the construction of the friction surface and the

travel time map.

a) Friction surface construction

The land coverage types were derived from satellite images from MODIS MCD12Q1 product

(Friedl et al., 2010). The MODIS collection includes seventeen land coverage types including

urban and rural areas inferred by the spectral signature of the satellite images. The boundaries of

v4.0

the national protected natural areas were included using data provided by the MEnv. The road

infrastructure in all districts was provided by the Peruvian Ministry of Transportation (MTrans),

and the navigable river network was derived from the HydroSHEDS Flow Accumulation dataset

(Lehner, Verdin, & Jarvis, 2006). The estimates of the friction surface (minutes required to travel

one meter) were adjusted by the slope of the terrain. This means that, the travel time required to

cross an area will be proportionally dependent to the slope of the terrain. The slope for each area

was calculated using the SRTM Digital Elevation Data (Jarvis, Reuter, Nelson, & Guevara,

2008) produced by NASA.

b) Travel time estimation

The target locations used for the cumulative cost function were the health facilities (clinics) of

the Ministry of Health (MH). This data was obtained from the geo-localization registry of health

facilities (RENAES in spanish) (**Figure 1B**). The MH organize the health facilities in three

categories according to the complexity of services they provide (from primary healthcare to

specialized hospitals). The primary level includes basic health facilities with no laboratory, the

secondary level includes health facilities with laboratory, and tertiary level includes hospitals and

higher complexity services. Finally, travel time estimates were extracted for each city and village

(**Figure 1C**). The most updated geo-localization of villages was provided by the Ministry of

Education (MEd) in a recent census of cities and villages, and education facilities.

2.4. Data Analysis

a) Friction surface construction

The estimation of travel time were conducted in Google Earth Engine (GEE) (Gorelick et al.,

2017). A surface grid was constructed using the information about land coverage, road

infrastructure, and river network. All datasets were converted into aligned grids with a 30-meter

resolution. Each dataset contained the information of the speed of movement in each feature. All

the layers were then combined with the fastest mode of movement taking precedence (Km h-1).

The speed assigned for each category of land cover were obtained from elsewhere (Weiss et al.,

2018). A data transformation was conducted, so each pixel within the 2D grid contained the cost

(time) to moving through the area encompassed in the pixel, herein referred to as 'friction

v4.0

surface'. Slope adjustment was carried out using the Tobler's Hiking Function (Tobler, 1993)

and the speed was penalized (reduced) in urban and national protected areas to account for

vehicular traffic and restricted displacement, respectively.

b) Travel time estimation

To calculate the travel time from the villages to the most proximate health facility, the

cumulative cost function was used in GEE to generate the accessibility map. The cumulative cost

function is a least-cost-path algorithm, briefly, all possible paths were analyzed iteratively and

the weighted cost (in this case, weighted by time) was then minimized. The minimum travel time

to the most proximate health facility was computed for each pixel in the grid, then the median

travel time was summarized in a 500m-radius from the geolocation of each city or village

(**Supplementary Information 2**). Values between the 5% and 95% percentile range were

considered to avoid extreme values. Since a health facility could be located in the 30m2

corresponding to the pixel spatial resolution of the estimates, a baseline 10-minutes travel time

was considered. The analysis was carried out for each health facility category. After GEE

processing, all data outputs were imported and analyzed using R v.3.6.0 (R Development Core

Team, R Foundation for Statistical Computing, Vienna, Australia).

The computed travel time was then summarized per district, province or department; by

urban/rural areas; and across 16 major land coverage types defined by the MEnv. Urban/rural

status was defined based on the MODIS land coverage satellite images (described previously in

2.3 Data Sources). To better detail the large diversity of land coverage types in Peru, a shortlist

of 16 eco-regions provided by the MEnv (**Supplementary Information 1**) was used to

summarize the travel time in these areas. In addition, the distribution of travel time relative to the

proportion of population with unsatisfied basic needs (UBN) ─ a multidimensional poverty

measurement developed by the United Nation's Economic Commission for Latin America and

the Caribbean (ECLAC) ─ per department was computed with data provided by the Ministry of

Economy (MEco).

3. RESULTS

Travel time to health facilities

v4.0

For this study, we gathered geo-referenced data on 145,134 villages (**Figure 1B**) and 8,067

health facilities (**Figure 1C**) across the 1,722 districts in the Peruvian territory. The health

facility density (number of health facilities divided by the total population) in Peru was 2.58 per

10,000 inhabitants with variations between major ecological areas, from 1.35 in the coast, 4.56 in

the highlands, to 5.21 in the jungle.

Friction and travel time maps were reconstructed in Google Earth Engine using the described

local datasets at a spatial resolution of 30 meters per pixel (**Supplementary Information 2**).

Country-wide median travel time from each village to the most proximate health facility varies

according to category: primary healthcare = 39 minutes (IQR=20 – 93), secondary healthcare =

152 minutes (IQR=75 – 251), and tertiary healthcare = 448 minutes (IQR=302 – 631).

Importantly, maximum travel time reached 7,819, 12,429, and 35,753 minutes for primary,

secondary, and tertiary levels, respectively (**Figure 2**).

Urban/rural and ecological settings

High heterogeneity was observed in contrasting land coverage areas. The median travel time was

5.3 fold higher in rural (85 minutes; IQR=11–7,819) than in urban settings (16 minutes; IQR =

11–835) to a primary healthcare facility; 3.2 fold higher in rural (226 minutes; IQR = 11–12,429)

than in urban settings (70 minutes; IQR = 11–3,386) to a secondary healthcare facility; and 2.4

fold higher in rural (568 minutes; IQR = 11–35,753) than in urban settings (235 minutes; IQR =

11–10,048) to a tertiary healthcare facility. A larger variation in travel time to primary healthcare

was observed in rural compared to urban areas, and conversely, a larger variation in travel time

to tertiary healthcare was observed in urban compared to rural areas (**Figure 3**). The district-

level stratified averages in **Figure 2** show that there were also strong heterogeneities within

major ecological regions. The north-east part of the Amazon Region, which borders with

Colombia and Brazil, presented the largest country-wide travel times to the most proximate

health facilities. The largest travel times to the most proximate health facilities within the

Highlands Region was observed in the southern areas of the Andes, and in the coast was

observed in the southern coast. Contrasting distributions of travel time to the most proximate

health facility was observed between the 16 eco-regions defined by the MEnv (**Figure 3**).

v4.0

Travel time to health facilities relative to UBN

When the travel time to most proximate health facilities was distributed relative to the proportion

of the population with unsatisfied basic needs at department level (administrative level 1), a

positive trend was observed (**Figure 4**). The slope of this relation was increased in geographical

accessibility to tertiary health facilities in comparison to primary or secondary health facilities.

4. DISCUSSION

The present study explored the use of novel cloud-based geospatial modeling approach fitted

with detailed local geospatial data to accurately estimate the travel time to the most proximate

health facility across a highly diverse geographical and ecological settings as observed in Peru.

This study showed the first quantification of heterogeneities in travel time to the most proximate

health facility as a surrogate of geographical accessibility in the Latin American region. Most of

the differences in travel time arose from heterogeneous land coverage profiles and the contrast

between urban and rural areas. This is particularly important due to the fact that in Peru and in

most LMIC, the most detailed data is available at a coarse administrative level that deter the

resource planning and healthcare provision in these countries. Another direct implication of the

utility of this approach is providing yet another angle of disadvantages amongst the most

underserved, now in terms of access to healthcare as measured by distance and time, one of

multiple aspects of high-quality healthcare.

In settings with a scattered distribution of villages, timely access to health facilities is a

cornerstone to improve the health status of impoverished populations and a first step to provide

high quality care (Kruk et al., 2018; Kruk, Pate, & Mullan, 2017). Although the use of big data

and high-detail datasets paves the way for a comprehensive quantification of geographical

accessibility in terms of distance and travel time, these technologies were not previously applied

to estimate geographical accessibility to health facilities until recently (Tatem, 2018). Using this

analytical approach, this study demonstrated that the population in the Jungle area have less

accessibility since healthcare services are reachable at longer trajectories and travel time,

understood as less geographical accessibility. The dramatic heterogeneity in travel time to the

most proximate health facility observed in this study corresponds to the contrasting landscape

v4.0

composition in the coast, highlands, and jungle regions. A dense road network was observed in

the Coast, facilitating access to multiple services including healthcare as reported in other studies

in India and Africa (Kanuganti et al., 2015; Strano, Viana, Sorichetta, & Tatem, 2018).

Conversely, sparse road coverage was observed in the Highlands and only the two major cities in

the Jungle region had roads.

Consistently with previous studies (Bashshur et al., 1971; Comber et al., 2011), this study

determined the heterogeneity in travel time to the most proximate health facility across areas

with contrasting land coverage types. Despite that this fact is widely accepted, few attempts have

been made to quantify these heterogeneities. In addition, asymmetries were identified when the

travel time to the most proximate health facility was compared along socio-economic profiles

based on the unsatisfied basic needs index proposed by the United Nations Development

Programme (UNDP). Uneven trends of greater travel time to health facilities (lower geographical

accessibility) were observed among villages with higher rates of unmet basic needs. These

results are consistent with previous reports of negative trends in geographical access to

healthcare facilities in low-income populations (Kiwanuka et al., 2008; Meyer, Luong,

Mamerow, & Ward, 2013; Peters et al., 2008; Tanser, Gijsbertsen, & Herbst, 2006).

It is important to highlight that the analysis conducted in this study did not take into account

variability due to climatic factors that may prevent displacement to health facilities (i.e. floods or

landslides). However, Highlands and Jungle areas are more prone to this kind of natural disaster,

leading to a conservative estimation of travel time in these areas. Traffic, which may greatly

influence the estimates in the large cities, was not considered in the analysis and potentially

cause an underestimation of the travel time to health facilities. In addition, seasonal variability

may greatly affect some displacement routes such as rivers; however, only navigable rivers were

considered in this approach and the availability to displace through this rivers are less affected by

seasonality. Another important consideration about the least-cost-path algorithm used in this

analysis is that we infer the lowest travel time boundary to reach a health facility. This

consideration relies on the assumption that the villagers opt for this route despite the cost and

danger of the route in addition to its availability, as explained above.

v4.0

In addition, the data reported here was generated at a meso-scale, with a spatial resolution of 30

meters. At this scale and resolution, some important details could be lost and affect the travel

time estimations. For instance, in some settings the travel time might be increased due to

meandering rivers or roads that follow the morphology of the terrain. The model assumes that

transit flows in a direct manner, meaning that zigzagging routes may cause our approach

underestimate the real travel time to reach a health facility. Despite these possible shortcomings,

the proposed approach provided conservative yet useful estimates of travel times to health

facilities that are important for planning of prevention and control strategies for multiple health-

related events. This approach demonstrates that curation and alignment of geospatial data from

multiple governmental institutions are important for national decision-making. In addition, the

use of mapping and modeling techniques, and 'big data' were recognized as critical for better

planning (Buckee et al., 2018; Hay, George, Moyes, & Brownstein, 2013; Tatem, 2018);

however, a remaining challenge is the implementation of these approaches into routine disease

prevention and control programs (Buckee et al., 2018; Hay et al., 2013).

This study acknowledges the relevance of other components of health access that may play an

important role in the underlying phenomena. The sole presence of clinic infrastructure does not

assure a proper healthcare delivery. Supply chain, human resources, financial accessibility,

acceptability of services, and availability of treatment are some remaining barriers once

geographical accessibility is overcome (Agbenyo et al., 2017; Al-Taiar et al., 2010; Johar,

Soewondo, Pujisubekti, Satrio, & Adji, 2018). Further studies were suggested to get a

comprehensive understanding of the accessibility to healthcare in Peru and other LMIC.

5. CONCLUSION

This study used a new methodology to estimate the travel time to most proximate health facilities

as a first step to understanding and characterizing the geographical accessibility profiles in Peru.

Contrasting patterns were observed across heterogeneous land coverage areas and urban and

rural settings. These findings are important as first steps for tailoring strategies to deliver

appropriate, affordable and quality healthcare to impoverished populations.

v4.0

REFERENCES

Agbenyo, F., Marshall Nunbogu, A., & Dongzagla, A. (2017). Accessibility mapping of health facilities in rural Ghana. *Journal of Transport & Health*, *6*, 73-83. https://doi.org/10.1016/j.jth.2017.04.010

Al-Taiar, A., Clark, A., Longenecker, J. C., & Whitty, C. J. (2010). Physical accessibility and utilization of health services in Yemen. *International Journal of Health Geographics*, *9*(1), 38. https://doi.org/10.1186/1476-072X-9-38

Bashshur, R. L., Shannon, G. W., & Metzner, C. A. (1971). Some Ecological Differentials in the Use of Medical Services. *Health Services Research*, *6*(1), 61-75.

Buckee, C. O., Cardenas, M. I. E., Corpuz, J., Ghosh, A., Haque, F., Karim, J., … Zaw, W. (2018). Productive disruption: Opportunities and challenges for innovation in infectious disease surveillance. *BMJ Global Health*, *3*(1), e000538. https://doi.org/10.1136/bmjgh-2017-000538

Chowkwanyun, M., Bayer, R., & Galea, S. (2018). «Precision» Public Health—Between Novelty and Hype. *The New England Journal of Medicine*, *379*(15), 1398-1400. https://doi.org/10.1056/NEJMp1806634

Comber, A. J., Brunsdon, C., & Radburn, R. (2011). A spatial analysis of variations in health access: Linking geography, socio-economic status and access perceptions. *International Journal of Health Geographics*, *10*, 44. https://doi.org/10.1186/1476-072X-10-44

v4.0

Delamater, P. L., Messina, J. P., Shortridge, A. M., & Grady, S. C. (2012). Measuring geographic

access to health care: Raster and network-based methods. *International Journal of Health*

Geographics, *11*(1), 15. https://doi.org/10.1186/1476-072X-11-15

Dowell, S. F., Blazes, D., & Desmond-Hellmann, S. (2016). Four steps to precision public health.

Nature News, *540*(7632), 189. https://doi.org/10.1038/540189a

Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., & Huang,

X. (2010). MODIS Collection 5 global land cover: Algorithm refinements and

characterization of new datasets. *Remote Sensing of Environment*, *114*(1), 168-182.

https://doi.org/10.1016/j.rse.2009.08.016

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google

Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of*

Environment, *202*, 18-27. https://doi.org/10.1016/j.rse.2017.06.031

Hay, S. I., George, D. B., Moyes, C. L., & Brownstein, J. S. (2013). Big Data Opportunities for

Global Infectious Disease Surveillance. *PLOS Medicine*, *10*(4), e1001413.

https://doi.org/10.1371/journal.pmed.1001413

Horton, R. (2018). Offline: In defence of precision public health. *The Lancet*, *392*(10157), 1504.

https://doi.org/10.1016/S0140-6736(18)32741-7

Huerta Munoz, U., & Källestål, C. (2012). Geographical accessibility and spatial coverage

modeling of the primary health care network in the Western Province of Rwanda.

v4.0

International Journal of Health Geographics, *11*(1), 40. https://doi.org/10.1186/1476-072X-11-40

Jarvis, A., Reuter, H. I., Nelson, A., & Guevara, E. (2008). *Hole-filled SRTM for the globe Version 4.*

Johar, M., Soewondo, P., Pujisubekti, R., Satrio, H. K., & Adji, A. (2018). Inequality in access to health care, health insurance and the role of supply factors. *Social Science & Medicine*, *213*, 134-145. https://doi.org/10.1016/j.socscimed.2018.07.044

Kanuganti, S., Sarkar, A. K., Singh, A. P., & Arkatkar, S. S. (2015). Quantification of accessibility to health facilities in rural areas. *Case Studies on Transport Policy*, *3*(3), 311-320. https://doi.org/10.1016/j.cstp.2014.08.004

Kiwanuka, S. N., Ekirapa, E. K., Peterson, S., Okui, O., Rahman, M. H., Peters, D., & Pariyo, G. W. (2008). Access to and utilisation of health services for the poor in Uganda: A systematic review of available evidence. *Transactions of The Royal Society of Tropical Medicine and Hygiene*, *102*(11), 1067-1074. https://doi.org/10.1016/j.trstmh.2008.04.023

Kruk, M. E., Gage, A. D., Arsenault, C., Jordan, K., Leslie, H. H., Roder-DeWan, S., … Pate, M. (2018). High-quality health systems in the Sustainable Development Goals era: Time for a revolution. *The Lancet Global Health*, *6*(11), e1196-e1252. https://doi.org/10.1016/S2214-109X(18)30386-3

v4.0

Kruk, M. E., Pate, M., & Mullan, Z. (2017). Introducing The Lancet Global Health Commission

on High-Quality Health Systems in the SDG Era. *The Lancet Global Health*, *5*(5), e480-

e481. https://doi.org/10.1016/S2214-109X(17)30101-8

Lehner, B., Verdin, K., & Jarvis, A. (2006). HydroSHEDS technical documentation, version 1.0.

World Wildlife Fund US, Washington, DC, 1-27.

Meyer, S. B., Luong, T. C., Mamerow, L., & Ward, P. R. (2013). Inequities in access to healthcare:

Analysis of national survey data across six Asia-Pacific countries. *BMC Health Services*

Research, *13*, 238. https://doi.org/10.1186/1472-6963-13-238

Müller, I., Smith, T., Mellor, S., Rare, L., & Genton, B. (1998). The effect of distance from home

on attendance at a small rural health centre in Papua New Guinea. *International Journal of*

Epidemiology, *27*(5), 878-884.

Noor, A. M., Zurovac, D., Hay, S. I., Ochola, S. A., & Snow, R. W. (2003). Defining equity in

physical access to clinical services using geographical information systems as part of

malaria planning and monitoring in Kenya. *Tropical Medicine & International Health*,

8(10), 917-926. https://doi.org/10.1046/j.1365-3156.2003.01112.x

Ouko, J. J. O., Gachari, M. K., Sichangi, A. W., & Alegana, V. (s. f.). Geographic information

system-based evaluation of spatial accessibility to maternal health facilities in Siaya

County, Kenya. *Geographical Research*, *0*(0). https://doi.org/10.1111/1745-5871.12339

Perry, B., & Gesler, W. (2000). Physical access to primary health care in Andean Bolivia. *Social*

Science & Medicine, *50*(9), 1177-1188. https://doi.org/10.1016/S0277-9536(99)00364-0

v4.0

Peters, D. H., Garg, A., Bloom, G., Walker, D. G., Brieger, W. R., & Rahman, M. H. (2008).

Poverty and access to health care in developing countries. *Annals of the New York Academy*

of Sciences, *1136*, 161-171. https://doi.org/10.1196/annals.1425.011

Stock, R. (1983). Distance and the utilization of health facilities in rural Nigeria. *Social Science &*

Medicine (1982), *17*(9), 563-570.

Strano, E., Viana, M. P., Sorichetta, A., & Tatem, A. J. (2018). Mapping road network

communities for guiding disease surveillance and control strategies. *Scientific Reports*,

8(1), 4744. https://doi.org/10.1038/s41598-018-22969-4

Tanser, F., Gijsbertsen, B., & Herbst, K. (2006). Modelling and understanding primary health care

accessibility and utilization in rural South Africa: An exploration using a geographical

information   system.   *Social   Science   &   Medicine*,   *63*(3),   691-705.

https://doi.org/10.1016/j.socscimed.2006.01.015

Tatem, A. J. (2018). Innovation to impact in spatial epidemiology. *BMC Medicine*, *16*(1), 209.

https://doi.org/10.1186/s12916-018-1205-5

Tobler, W. (1993). *Three Presentations on Geographical Analysis and Modeling: Non- Isotropic*

Geographic Modeling; Speculations on the Geometry of Geography; and Global Spatial

Analysis (93-1). Recuperado de https://escholarship.org/uc/item/05r820mz

UN General Assembly. (2015, octubre 21). Transforming our world: The 2030 Agenda for

Sustainable   Development.   Recuperado   9   de   octubre   de   2018,   de

http://www.refworld.org/docid/57b6e3e44.html

v4.0

Weiss, D. J., Nelson, A., Gibson, H. S., Temperley, W., Peedell, S., Lieber, A., … Gething, P. W.

(2018). A global map of travel time to cities to assess inequalities in accessibility in 2015.

Nature, *553*(7688), 333-336. https://doi.org/10.1038/nature25181

World Health Organization, & World Bank. (2017). *Tracking universal health coverage: 2017*

Global   Monitoring   Report.   Recuperado   de

http://www.who.int/healthinfo/universal_health_coverage/report/2017/en/

Acknowledgements.

We thank the various ministries of Peru for making such useful data freely-available to researchers.

Gabriel Carrasco-Escobar was supported by NIH/Fogarty International Center Global Infectious

Diseases Training Program (D43 TW007120). J. Jaime Miranda acknowledges having received

support from the Alliance for Health Policy and Systems Research (HQHSR1206660), the Bernard

Lown Scholars in Cardiovascular Health Program at Harvard T.H. Chan School of Public Health

(BLSCHP-1902), Bloomberg Philanthropies, FONDECYT via CIENCIACTIVA/CONCYTEC,

British Council, British Embassy and the Newton-Paulet Fund (223-2018, 224-2018),

DFID/MRC/Wellcome Global Health Trials (MR/M007405/1), Fogarty International Center

(R21TW009982, D71TW010877), Grand Challenges Canada (0335-04), International

Development Research Center Canada (IDRC 106887, 108167), Inter-American Institute for

Global Change Research (IAI CRN3036), Medical Research Council (MR/P008984/1,

MR/P024408/1, MR/P02386X/1), National Cancer Institute (1P20CA217231), National Heart,

Lung and Blood Institute (HHSN268200900033C, 5U01HL114180, 1UM1HL134590), National

Institute of Mental Health (1U19MH098780), Swiss National Science Foundation (40P740-

v4.0

409   Author's contributions

410   Conceived and designed the study: G.C.E. and J.J.M. Data collection: G.C.E., K.T.L., and E.M.

411   Analyzed the data: G.C.E., K.T.L., and E.M. Wrote the manuscript: G.C.E. and J.J.M. All authors

412   reviewed and approved the final manuscript.

414   Competing interests

415   The authors declare no competing interests.

417   Data Availability

418   Raw datasets and codes are available at google earth engine repository, details in the

419   Supplementary information section.

v4.0

422 **Figure 1. Study area.** A) Major ecological areas (coast, andes, and jungle) in Peru. Solid lines

423 represent the 25 Departments (administrative level 1). B) Spatial location of primary, secondary,

424 and tertiary health facilities. C) Spatial location of villages. Maps were produced using QGIS,

425 and the base map was obtained OpenTopoMap (http://www.opentopomap.org), under CC BY-

426 SA 3.0.

v4.0

428 **Figure 2. Country-wide map of travel time to health facilities for 2018.** District-level average

429 travel time to each category of healthcare facilities. A) Primary healthcare. B) Secondary

430 healthcare. C) Tertiary healthcare. Color scale in logarithmic scale

432
433

v4.0

 Figure 3. Distribution of travel time to most proximate health facility. Estimates across the

16 eco-regions defined by the Peruvian Ministry of environment and rural/urban settings for

primary, secondary and tertiary healthcare. Y-axis in logarithmic scale.

v4.0

441 **Figure 4. Median travel time to each health facility category relative to the proportion of**

442 **population with unsatisfied basic needs per department.** Y-axis in logarithmic scale

v4.0