1 **Title Page**
2
3 **Article Type:** Research Article
4
5 **Title:** Potential impact of outpatient stewardship interventions on antibiotic exposures of
6 bacterial pathogens
7

8 **Authors:** Christine Tedijanto,[1] Yonatan H Grad,[2,3] Marc Lipsitch[1,2]
9
10 **Author Affiliations:** [1] Center for Communicable Disease Dynamics, Department of
11 Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA;
12 [2] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public
13 Health, Boston, Massachusetts, USA; [3] Division of Infectious Diseases, Brigham and Women's
14 Hospital, Harvard Medical School, Boston, Massachusetts, USA
15
16 **Corresponding Author:** Christine Tedijanto, ctedijanto@g.harvard.edu
17

18 **Manuscript Word Count:** 4,632
19

Abstract

The relationship between antibiotic stewardship and population levels of antibiotic resistance remains unclear. In order to better understand shifts in selective pressure due to stewardship, we use publicly available data to estimate the effect of changes in prescribing on exposures to frequently used antibiotics experienced by potentially pathogenic bacteria that are asymptomatically colonizing the microbiome. We quantify this impact under four hypothetical stewardship strategies. In one scenario, we estimate that elimination of all unnecessary outpatient antibiotic use could avert 6 to 48% (IQR: 17 to 31%) of exposures across pairwise combinations of sixteen common antibiotics and nine bacterial pathogens. All scenarios demonstrate that stewardship interventions, facilitated by changes in clinician behavior and improved diagnostics, have the opportunity to broadly reduce antibiotic exposures across a range of potential pathogens. Concurrent approaches, such as vaccines aiming to reduce infection incidence, are needed to further decrease exposures occurring in "necessary" contexts.

Introduction

Antibiotic consumption is a known driver of antibiotic resistance. In developed nations, over 80% of antibiotic consumption for human health occurs in the outpatient setting (European Centre for Disease Prevention and Control, 2018; Public Health Agency of Canada, 2018; Public Health England, 2017; Swedres-Svarm, 2017), and US-based studies conducted across different subpopulations have estimated that 23-40% of outpatient prescriptions may be inappropriate (Chua, Fischer, & Linder, 2019; Fleming-Dutra et al., 2016; Olesen, Barnett, MacFadden, Lipsitch, & Grad, 2018). Inappropriate antibiotic use leads to increased risk of adverse events (Linder, 2008; Shehab, Patel, Srinivasan, & Budnitz, 2008), disruption of colonization resistance and other benefits of the microbial flora (Buffie & Pamer, 2013; Khosravi & Mazmanian, 2013), and bystander selection for antibiotic resistance, with little to no health gains for the patient. Recent work on bystander selection estimates that, for 8 of 9 potential pathogens of interest, over 80% of their exposures to commonly used antibiotic classes in the outpatient setting occur when the organisms are asymptomatically colonizing the microbiome, not causing disease (Tedijanto, Olesen, Grad, & Lipsitch, 2018). A corollary of extensive bystander selection is that reductions in use will prevent antibiotic exposures for species throughout the microbiome.

We use publicly available data to quantify potentially avertable exposures of bacterial species to commonly used antibiotics under hypothetical changes in prescribing practice. The set of scenarios included here are intended as thought experiments to explore the upper bounds of avertable antibiotic exposures. Reductions in antibiotic consumption have historically had variable impacts on resistance levels, likely dependent on setting, baseline consumption and resistance patterns, and fitness costs (Andersson & Hughes, 2010; Lipsitch, 2001). In addition,

treatment strategies are primarily guided by randomized controlled trials assessing immediate

clinical outcomes, with less careful consideration given to the microbiome-wide effect of such

decisions. We apply analytical methods to parse out species-level effects of changes in

prescribing practice in order to better understand the potential impact of antibiotic stewardship.

Methods

Scenarios of interest

This analysis includes sixteen antibiotics that are frequently prescribed in the outpatient setting

and nine potentially pathogenic bacterial species that are commonly carried in the normal human

microbiome. For each antibiotic-species pair, we estimate the proportion of antibiotic exposures

experienced by that species that could be averted under four hypothetical scenarios. The

scenarios range from broad elimination of unnecessary prescribing to focused modifications of

antibiotic use for specific indications as follows:

1) Eliminate unnecessary antibiotic use across all outpatient conditions.

2) Eliminate all antibiotic use for outpatient respiratory conditions for which antibiotics are

not indicated.

3) Eliminate all antibiotic use for acute sinusitis.

4) Prescribe nitrofurantoin for all cases of cystitis in women.

We use the results reported by Fleming-Dutra and colleagues as estimates of unnecessary

antibiotic use in ambulatory care settings (Fleming-Dutra et al., 2016). In this study, the authors

convened a group of experts to estimate the proportion of appropriate antibiotic use by condition

and age group based on clinical guidelines. When guidelines could not be used for this task, the

rate of appropriate antibiotic prescribing was estimated by benchmarking against the lowest-

prescribing US region. Based on the same paper, we consider viral upper respiratory tract

infection, influenza, non-suppurative otitis media, viral pneumonia, bronchitis, and allergy and

asthma to be outpatient respiratory conditions for which antibiotics are not indicated.

Antibiotic treatment of acute bacterial sinusitis is currently guideline-recommended (Rosenfeld

et al., 2015). However, bacteria are an infrequent cause of acute sinusitis, and due to the self-

limiting nature of the syndrome, evidence to support antibiotic treatment is weak (Burgstaller,

Steurer, Holzmann, Geiges, & Soyka, 2016; Fokkens, Lund, & Mullol, 2007). The cause of

sinusitis (whether bacterial, viral, or noninfectious) can be very difficult to distinguish in

practice, and as a result of this challenge and others, antibiotics continue to be prescribed at over

80% of US outpatient visits with a primary diagnosis of acute sinusitis (Smith et al., 2013). In

contrast, antibiotics are always recommended for urinary tract infections (UTIs). Despite being

recommended as a second-line therapy for cystitis due to concerns about resistance,

fluoroquinolones are the most common treatment, prescribed at 40% of outpatient visits for

uncomplicated UTI (Kabbani et al., 2018). We explore the hypothetical scenario of treating all

cases of cystitis in women with nitrofurantoin, a recommended first-line therapy with good

potency against common uropathogens, low levels of resistance, and decreased risk of collateral

damage to the intestinal microbiome due to its propensity to concentrate in the bladder (Gupta et

al., 2011; Gupta, Scholes, & Stamm, 1999; Stewardson et al., 2015). Nitrofurantoin is considered

clinically appropriate unless the patient has chronic kidney disease, is showing signs of early

pyelonephritis, or has had a prior urinary isolate resistant to nitrofurantoin (Hooton & Gupta,

2019). In our analysis, we exclude patients with a simultaneous diagnosis of pyelonephritis. We

assume that other contraindications are fairly rare in the general population.

Data sources and methodology

Extending our recent work on bystander selection (Tedijanto et al., 2018), we used the 2015

National Ambulatory Medical Care Survey and National Hospital Ambulatory Medical Care

Survey (NAMCS/NHAMCS) (National Center for Health Statistics, 2015a, 2015b), the Human

Microbiome Project (HMP) (Huttenhower et al., 2012; The Human Microbiome Project

Consortium, 2012), and assorted carriage and etiology studies (Bäckhed et al., 2015; Bluestone,

Stephenson, & Martin, 1992; Bogaert et al., 2011; Brook, Thompson, & Frazier, 1994; Celin,

Bluestone, Stephenson, Yilmaz, & Collins, 1991; Chira & Miller, 2010; Edlin, Shapiro, Hersh, &

Copp, 2013; Ginsburg et al., 1985; Gunnarsson, Holm, & Söderström, 1997; Gupta et al., 1999;

Jack M. Gwaltney, Scheld, Sande, & Sydnor, 1992; Hammitt et al., 2006; Holgerson, Öhman,

Rönnlund, & Johansson, 2015; Susan S. Huang et al., 2009; Jain et al., 2015; Arch G. Mainous,

Hueston, Everett, & Diaz, 2006; Pettigrew et al., 2012; Regev-Yochay et al., 2004; Verhaegh et

al., 2010; Loretta Wubbel et al., 1999; Yassour et al., 2016) (details in Figure 1 – Source Data

File 1) to estimate national outpatient antibiotic exposures by drug, species, and condition.

NAMCS/NHAMCS are annual cross-sectional surveys designed to sample outpatient visits in

the United States, and up to five diagnoses and thirty medications may be associated with each

visit. We used methodology developed by Fleming-Dutra and colleagues (Fleming-Dutra et al.,

2016), and applied in other studies (Olesen, Barnett, MacFadden, Lipsitch, et al., 2018), to group

diagnosis codes into conditions and link antibiotic prescriptions with the most likely indication.

For this analysis, visits with a diagnosis of acute cystitis (ICD-9-CM: 595.0), unspecified cystitis

(595.9) or unspecified UTI (599.0), without a concurrent diagnosis of pyelonephritis (590.1,

590.8), renal abscess (590.2), or kidney infection (590.9), were considered to be associated with

cystitis. In this analysis, we maintain the assumption that one antibiotic prescription is equivalent

to one exposure; antibiotic exposures experienced by a given species and associated with a given

condition are roughly estimated as the product of antibiotic prescriptions for that condition and

species carriage prevalence, which is dependent on disease etiology (target exposures) and

asymptomatic carriage prevalence (bystander exposures). For diagnoses where etiology was not

readily available, we assumed that none of our species of interest were causative agents. These

assumptions have been enumerated in detail in prior work (Tedijanto et al., 2018). We applied

proportions of unnecessary antibiotic prescribing by condition and age group estimated by

Fleming-Dutra and colleagues based on expert opinion, clinical guidelines, and regional

variability in use (Fleming-Dutra et al., 2016). For relevant scenarios (1 and 2), we applied the

proportions of unnecessary use evenly across all antibiotic prescriptions. Antibiotics and

antibiotic classes are identified by the Lexicon Plus® classification scheme (*Lexicon Plus*,

2008).

The proportion of avertable antibiotic exposures for each species is defined by Equation 1. The

equation adopts previously described notation (Tedijanto et al., 2018) with modifications. A

listing of all variables and descriptions can be found in Equation 1 – Figure Supplement 1. Let *a*

represent antibiotic, *s* represent species, *i* represent ICD-9-CM diagnosis code, and *g* represent

age group. Throughout the analysis we have weighted outpatient visits to be nationally

representative using the sampling and nonresponse weights provided in NAMCS/NHAMCS. Let

X_{as} be the number of avertable exposures by antibiotic and species, T_{as} be the total number of

exposures by antibiotic and species, d_{aig} be the number of prescriptions of antibiotic *a* associated

with diagnosis code *i* in age group *g*, p_{sig} be the carriage prevalence of species *s* among those

with diagnosis code *i* in age group *g*, and q_{aig} be the proportion of avertable exposures by

diagnosis code and age group in the given scenario. For example, in the scenario assessing

elimination of all unnecessary antibiotic use, q_{aig} is the proportion of avertable antibiotic use by

diagnosis and age group (Fleming-Dutra et al., 2016). Alternatively, in the scenario assessing

elimination of non-nitrofurantoin treatment for cystitis, q_{aig} is 1 when a is not nitrofurantoin, i is

a diagnosis code associated with cystitis, and the patient is female, and 0 elsewhere. Carriage

prevalences (p_{sig}) are assumed to be constant within three age groups (under 1 year, 1-5 years,

over 5 years old) (Tedijanto et al., 2018), while proportions of avertable antibiotic use (q_{aig}) were

reported for three different age groups (0-19 years, 20-64 years, 65 years old and over) (Fleming-

Dutra et al., 2016). G is the smallest set of age groups that accounts for this granularity (under 1

year, 1-5 years, 6-19 years, 20-64 years, 65 years old and over). For antibiotic prescriptions that

occurred at visits without any ICD-9-CM diagnosis codes (i=0), we applied the carriage

prevalence among healthy individuals.

Equation 1. Proportion of avertable exposures by species and antibiotic. Equation 1 – Figure
Supplement 1 includes further details on all variables.
162

$$\frac{X_{as}}{T_{as}} = \frac{\sum_{g=1}^{G} \sum_{i=0}^{I} d_{aig} \times p_{sig} \times q_{aig}}{\sum_{g=1}^{G} \sum_{i=0}^{I} d_{aig} \times p_{sig}}$$

In scenarios where unnecessary use is eliminated (1 and 2), we assume that only bystander

exposures are affected. Equivalently, this presumes that perfect discrimination between bacterial

and non-bacterial etiologies is possible. This results in a slight modification to the numerator of

Equation 1 -- p_{sig} is changed to p_{s0g} as all eliminated exposures would have occurred during

asymptomatic carriage. For cases where e_{sig} is 1, we assume zero avertable exposures. In

addition, we make slight modifications to q_{aig} when we estimate the proportion of bacterial cases

to be larger than the proportion of necessary prescriptions (Figure 1 – Source Data File 2)

(Bluestone et al., 1992; Brook, 2016; Celin et al., 1991). As a sensitivity analysis, we include the

proportion of avertable exposures for each bacteria-antibiotic pair under Scenario 1 if antibiotic

use was eliminated equally across both target and bystander exposures (Figure 1 – Figure

Supplement 1). All analysis was conducted in R version 3.6.1.

Results

177 Results for all four scenarios are depicted in Figure 1.

Figure 1. Heatmaps showing the estimated percentage of species exposures to each antibiotic or
antibiotic class that could be averted if *A:* unnecessary antibiotic prescriptions across all
outpatient conditions, *B:* all antibiotic use for outpatient respiratory conditions for which
antibiotics are not indicated, *C:* all antibiotic use for acute sinusitis, or *D:* non-nitrofurantoin
treatment of cystitis in women was eliminated. Drug class results include prescriptions of all
antibiotics in that class, as identified by the Lexicon Plus® classification system. Sensitivity and
other additional analyses are shown in Figure 1 – Figure Supplements 1-4.
Abbreviations: *Antibiotics (y-axis):* AMX-CLAV=amoxicillin-clavulanate,, MACR/LINC=macrolides/
lincosamides, TMP-SMX=sulfamethoxazole-trimethoprim; *Organisms (x-axis):* EC=*E. coli*, HI=*H. influenzae*,
KP=*K. pneumoniae*, MC=*M. catarrhalis*, PA=*P. aeruginosa*, SA=*S. aureus*, SAg=*S. agalactiae*, SP=*S. pneumoniae*,
SPy=*S. pyogenes*; PY=person-years

Scenario 1

We estimate that elimination of unnecessary antibiotic prescriptions across all outpatient

conditions would prevent 6 to 48% (IQR: 17 to 31%) of antibiotic-species exposures (Figure

1A). The smallest reduction is associated with *S. pyogenes* exposures to cefdinir and the largest

with *H. influenzae*, *E. coli,* and *P. aeruginosa* exposures to azithromycin. If all unnecessary

antibiotic use could be prevented, over 30% of exposures to amoxicillin-clavulanate, penicillin,

azithromycin, clarithromycin, levofloxacin, and doxycycline across most potential pathogens of

interest could be averted. Of particular interest, elimination of all unnecessary prescribing in the

outpatient setting could reduce exposures of *S. pneumoniae* to penicillins and macrolides by 27%

and 37%, respectively, and of *S. aureus* to penicillins and quinolones by 27% and 21%. For *E.*

coli and *K. pneumoniae*, approximately one-quarter of exposures to cephalosporins and one-fifth

of exposures to quinolones could be averted.

Scenario 2

Scenario 2, elimination of all antibiotic use for outpatient respiratory conditions where

antibiotics are not indicated, is a subset of Scenario 1 and accounts for a substantial portion of

avertable exposures included in the first scenario. As in Scenario 1, the results are primarily

driven by drug and depend on the amount of use of that drug for the affected conditions (in this

case, conditions for which antibiotics are not indicated). Pathogen characteristics affecting the

proportion of avertable exposures can be better understood by looking across all species for a

single drug. For example, we focus on the row for azithromycin. Based on NAMCS/NHAMCS

2015, 32% of all azithromycin use is associated with the affected conditions. Since these

conditions are never caused by our organisms of interest, the maximum proportion of avertable

exposures for any species is 32%. However, the proportion of avertable exposures is modulated

by the number of azithromycin exposures that occur for treatment of other conditions – for

example, a lower proportion of avertable exposures for *S. pyogenes*, *M. catarrhalis*, and *S.*

agalactiae indicates that they have relatively frequent exposure to azithromycin during treatment

of other conditions that are unaffected in this scenario, often both as a bystander and causative

pathogen.

Scenario 3

In Scenario 3, we explore elimination of all antibiotic use for acute sinusitis. Overall, acute

sinusitis accounts for just 3% of prescriptions of our included antibiotic classes, resulting in the

low proportions of avertable exposures across most bacteria-antibiotic combinations. However,

proportions of avertable exposures are relatively high for amoxicillin-clavulanate and

clarithromycin, antibiotics for which a large proportion of their use is due to acute sinusitis (15%

and 9%, respectively). In this scenario, we make the assumption that all use for acute sinusitis is

unnecessary. In reality, a small number of cases are truly bacterial and antibiotics are indicated

for patients with persistent, severe or worsening symptoms (Rosenfeld et al., 2015). We explore

varying levels of unnecessary use based on etiological data in Figure 1 – Figure Supplement 2

(Aitken & Taylor, 1998; Benninger, Holzer, & Lau, 2000; Fleming-Dutra et al., 2016; Sande &

Gwaltney, 2004; Snow, Mottur-Pilson, & Hickner, 2001; Wald, Guerra, & Byers, 1991;

Williams, Simel, Roberts, & Samsa, 1992).

Scenario 4

In this scenario, we consider a program to prescribe only nitrofurantoin for all cases of acute

cystitis in women, replacing all other antibiotics that are currently in use. As a result of this

change, we estimate that exposures of *E. coli* and *K. pneumoniae* to cephalosporins could be

reduced by 16% and 17%, respectively, and exposures to quinolones could decrease by 26% and

28%. Uniquely in this scenario among the ones we considered, several of our species of interest

cause a large proportion of UTI cases (*E. coli*, *K. pneumoniae*, and *P. aeruginosa*). To explore

the effects of being a causative pathogen on the proportion of avertable exposures, we focus on

the results for ciprofloxacin. Overall, 27% of ciprofloxacin use is associated with acute cystitis in

women. Organisms that are not causative pathogens of cystitis have proportions of avertable

exposures close to or less than 27%, but a substantially higher proportion of exposures could be

averted among causative pathogens - 33%, 36%, and 44% of ciprofloxacin exposures to *E. coli*,

K. pneumoniae, and *P. aeruginosa*, respectively. Although *E. coli* is the most common cause of

UTIs, its estimated proportion of avertable exposures in this scenario is lower than that of *P.*

aeruginosa, likely due to the larger number of antibiotic exposures incurred by *E. coli* during

asymptomatic carriage.

This scenario also allows us to observe the impact of differences in disease incidence, antibiotic

use patterns, and carriage prevalence across age groups on the proportion of avertable exposures.

For example, *M. catarrhalis* is asymptomatically carried in children much more frequently than

in adults. Since cystitis occurs primarily in adults, the proportion of avertable exposures is often

lower for *M. catarrhalis* compared to other species due to bystander exposures in children that

are largely unaffected by modified prescribing practices for cystitis. This age group effect can be

substantial for antibiotics that are commonly given to children, especially when the antibiotic is

also used to treat a condition caused by *M. catarrhalis* (e.g. cefdinir and otitis media), but is less

influential for antibiotics that are infrequently prescribed to children, such as ciprofloxacin and

levofloxacin.

Overall, the proportion of avertable exposures is driven by pathogen, syndrome, and prescribing

characteristics. Because our organisms of interest experience the vast majority of their antibiotic

exposures as bystanders, the metric is largely dependent on the proportion of antibiotic use that is

associated with the conditions affected in the given scenario. In a scenario where all antibiotic

use for affected conditions is eliminated, this proportion would be equal to the proportion of

avertable exposures for organisms exposed to antibiotics only during carriage (never pathogens).

Being a causative pathogen for any of the affected conditions increases the proportion of

avertable exposures. In contrast, being a causative pathogen for unaffected conditions increases

unaffected antibiotic exposures and thus decreases the proportion of avertable exposures. For

pathogens responsible for multiple syndromes, both factors may be at play. Finally,

asymptomatic carriage prevalence is positively associated with both affected and unaffected

antibiotic exposures, and, as a result, its overall effect on the proportion of avertable exposures is

dependent on the bacteria-antibiotic combination of interest.

Discussion

We quantify the species-level impact of changes in antibiotic consumption as the proportion of

antibiotic exposures experienced by common bacterial pathogens that could be averted under

four hypothetical scenarios. In the scenario where unnecessary antibiotic use for all outpatient

conditions is eliminated, we find that up to 48% of exposures (of *H. influenzae*, *E. coli,* and *P.*

aeruginosa to azithromycin) could be avoided. In addition, impact of the intervention across

antibiotics and species is widespread, with half of antibiotic-species pairs expected to experience

a reduction in exposures of over one-fifth (22%). For conditions which require antibiotic

treatment, such as UTIs, switching to antibiotics with decreased collateral damage to the

microbiome, such as nitrofurantoin, may be an effective strategy for reducing antibiotic

exposures across species.

In three out of four situations we assess, antibiotics are considered entirely unnecessary for some

or all cases. In the future, this method may also be extended to measure net changes in exposures

resulting from more nuanced scenarios where one antibiotic is substituted for another. We did

not assess such changes for nitrofurantoin, as its effects on the microbiome outside of the bladder

are thought to be minimal (Stewardson et al., 2015). The reductions in use presented here may be

practically infeasible due to challenges including similar clinical presentation of viral and

bacterial infections, laboratory processing times that prevent identification of causative

pathogens during outpatient visits, individual considerations such as allergies (Sakoulas, Geriak,

& Nizet, 2018) or heightened risk of adverse events, misdiagnosis (Filice et al., 2015; Tomas,

Getman, Donskey, & Hecker, 2015), and patient-driven demand (Vanden Eng et al., 2003). Even

if elimination of unnecessary use were fully realized, our results imply that the majority of

species' antibiotic exposures occur in the context of "necessary" antibiotic use. These findings

underscore the importance of considering bystander effects and the need for a multi-pronged

approach to programs aimed at controlling antibiotic resistance. Rapid diagnostics, diagnostics

that accurately discriminate between bacterial and non-bacterial causes, patient education, and

improved decision-making tools and other interventions to motivate changes in clinician

behavior can enhance responsible antibiotic consumption and reduce unnecessary antibiotic use;

these should be implemented simultaneously with prevention measures such as infection control,

access to clean water and sanitation, safe sex interventions, and vaccination, which aim to reduce

infection incidence and thus any antibiotic use.

At the time of this analysis, NHAMCS data from hospital outpatient departments was not

available for 2015. In the 2010-2011 NAMCS/NHAMCS data, visits to hospital outpatient

departments accounted for 36% of included sampled visits, but just 9% of all antibiotic mentions.

Since the proportions of avertable exposures including and excluding these visits for 2010-2011

were similar for Scenario 1 (Figure 1 – Figure Supplement 3), we assumed that the existing 2015

data without hospital outpatient department visits was representative of the NAMCS/NHAMCS

population for the purposes of our analysis. Certain outpatient encounters are also outside the

scope of NAMCS/NHAMCS, including federal facility visits, telephone contacts, house calls,

long-term care stays, urgent care, retail clinics, and hospital discharge prescriptions.

Additional limitations of the included datasets and method for estimating antibiotic exposures

have been previously enumerated (Tedijanto et al., 2018). Notably, the NAMCS/NHAMCS does

not provide information to link medications with diagnoses, so we adopted a published tiered

system to pair antibiotic use at each visit with the most likely indication (Fleming-Dutra et al.,

2016). For visits with multiple diagnoses, all antibiotic use was attributed to the single most-

likely indication. This method tends to overestimate antibiotic use for conditions for which

antibiotics are almost always indicated, which may lead to clinically unusual diagnosis-treatment

linkages. In the same way, antibiotic prescriptions are underestimated for diagnoses for which

antibiotic use is not indicated, potentially leading to downward bias in avertable exposures for

these conditions. Additionally, NAMCS/NHAMCS antibiotic use data does not include

prescription details such as duration or dose. Future studies with more granular data may extend

the methods presented here to account for such factors.

At the time of this analysis, Fleming-Dutra et al. remained the most up-to-date source of

unnecessary antibiotic prescribing by condition. Recent work with slightly different methods

found a lower proportion of inappropriate antibiotic prescriptions overall (23.2% compared to

30%) but did not report their estimates by condition (Chua et al., 2019). Although comparable

estimates of unnecessary prescribing over time are unavailable, multiple studies have shown that

declines in antibiotic use from approximately 2011 to 2016 in the United States have been

primarily due to pediatric prescribing, implying that, at least for adults, levels of unnecessary

prescribing likely remained similar (Durkin et al., 2018; King, Bartoces, Fleming-Dutra, Roberts,

& Hicks, 2019; Klevens et al., 2019; Olesen, Barnett, MacFadden, Lipsitch, et al., 2018). We

also assume that the proportion of unnecessary use is constant across all antibiotic use for the

same condition as it is difficult to identify specific antibiotic prescriptions that were unwarranted

without detailed chart review. A study among outpatients in the Veterans Affairs medical system

found that among prescriptions for community-acquired pneumonia, sinusitis, and acute

exacerbations of chronic bronchitis, the highest proportion of macrolide use was inappropriate

(27%), followed by penicillins (22%) and quinolones (12%) (Tobia, Aspinall, Good, Fine, &

Hanlon, 2008). Similar studies are needed to understand which antibiotics are frequently used

inappropriately for other indications and settings.

Finally, it is important to note that the reduction in antibiotic exposures estimated here does not

translate to the same reduction in the prevalence of resistance or in the morbidity and mortality

attributable to resistance. Although population-level antibiotic consumption has been positively

correlated with levels of antibiotic resistance, the impact of changing consumption is not well-

understood and is likely to vary widely by bacteria-antibiotic combination (or even strain-

antibiotic combination) based on resistance mechanisms, fitness costs, and co-selection, among

other factors (Andersson & Hughes, 2010; Pouwels et al., 2017). Antibiotic consumption is also

highly heterogeneous and the impact of stewardship on resistance may depend on the affected

patient populations (Olesen, Barnett, MacFadden, Brownstein, et al., 2018). Additionally, in this

analysis, we give each exposure equal weight. However, the selective pressure imposed by a

single exposure depends on a number of variables, including pharmacokinetics,

pharmacodynamics, distribution of bacteria across body sites, and bacterial population size. For

example, we might expect that the probability of resistance scales with population size, and thus

that an exposure received by an individual with higher bacterial load will have a larger impact on

resistance. Further research, integrating knowledge from clinical, ecological, and evolutionary

spheres, is needed to elucidate the relationship between antibiotic use and selective pressures and

ultimately between use and resistance at the population level (MacLean & San Millan, 2019).

Reductions in antibiotic consumption are necessary to preserve the potency of these drugs.

Quantifying changes in species-level exposures due to stewardship programs is one more step

towards understanding how changes in antibiotic use correspond to antibiotic resistance. The

methods presented here may be easily extended to incorporate other data sources, such as claims,

or to assess more specific stewardship programs. We show that while improved prescribing

practices have the potential to prevent antibiotic exposures experienced by bacterial species

throughout the microbiome, complementary efforts to facilitate appropriate antibiotic

consumption and decrease overall infection incidence are required to substantially avert

exposures.

Acknowledgements
We thank Dr. Lauri Hicks for her helpful comments on this manuscript.
381
382   **Competing Interests**
383   ML has received consulting income from Affinivax, Antigen Discovery, Merck, and Pfizer and
384   research grants through Harvard School of Public Health from Pfizer and PATH. YHG has
385   received consulting income from Merck and GlaxoSmithKline. The authors have no other
386   relationships or activities that could appear to have influenced the submitted work.
387
Funding Sources
This work was supported by Grant U54GM088558 (Models of Infectious Disease Agent Study,
Center for Communicable Disease Dynamics) from the National Institute of General Medical
Sciences, Grant R01AI132606 from the National Institute of Allergy and Infectious Diseases (to
YHG), Grant CK000538-01 from the Centers for Disease Control and Prevention, and the Doris
Duke Charitable Foundation. CT is supported by Grant T32AI007535 from the National Institute
of Allergy and Infectious Diseases. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institute of General Medical
Sciences, National Institutes of Health, National Institute of Allergy and Infectious Diseases,
Centers for Disease Control and Prevention, Department of Health and Human Services, or the
Doris Duke Charitable Foundation.
399

400   **Figure Supplements**
401
402   *Equation 1 – Figure Supplement 1.* Notation, descriptions, and sources for variables in Equation
403   1.
404
405   *Figure 1 – Figure Supplement 1.* Sensitivity analysis of proportions of avertable exposures
406   across all outpatient conditions when proportion of unnecessary use is applied equally across
407   target and bystander exposures (Equation 1 without modification).
408
409   *Figure 1 – Figure Supplement 2.* Sensitivity analyses for Scenario 3 (elimination of all antibiotic
410   use for acute sinusitis).
411
412   *Figure 1 – Figure Supplement 3.* Proportions of avertable exposures under Scenario 1
413   (elimination of all unnecessary antibiotic use across outpatient conditions) using 2010-2011
414   NAMCS/NHAMCS data with and without NHAMCS outpatient department data.
415
416

Source Data Files

Figure 1 – Source Data File 1. Summary of etiology studies.

Figure 1 – Source Data File 2. Modifications to proportions of unnecessary antibiotic prescriptions based on published etiology studies and antibiotic use in NAMCS/NHAMCS. Modifications were made if the reported proportion of bacterial cases for a given condition exceeded the estimated proportion of appropriate prescriptions reported in Fleming-Dutra et al.

Figure 1 – Source Data File 3. Estimated antibiotic class exposures by exposed and causative species.

References

References

Aitken, M., & Taylor, J. A. (1998). Prevalence of Clinical Sinusitis in Young Children Followed Up by Primary

Care Pediatricians. *Archives of Pediatrics & Adolescent Medicine*, *152*(3), 244–248.

https://doi.org/10.1001/archpedi.152.3.244

Andersson, D. I., & Hughes, D. (2010). Antibiotic resistance and its cost: Is it possible to reverse resistance? *Nature*

Reviews Microbiology, *8*(4), 260–271. https://doi.org/10.1038/nrmicro2319

Bäckhed, F., Roswall, J., Peng, Y., Feng, Q., Jia, H., Kovatcheva-Datchary, P., … Wang, J. (2015). Dynamics and

Stabilization of the Human Gut Microbiome during the First Year of Life. *Cell Host & Microbe*, *17*(5),

690–703. https://doi.org/10.1016/j.chom.2015.04.004

Benninger, M. S., Holzer, S. E. S., & Lau, J. (2000). Diagnosis and treatment of uncomplicated acute bacterial

rhinosinusitis: Summary of the Agency for Health Care Policy and Research evidence-based report.

Otolaryngology–Head and Neck Surgery, *122*(1), 1–7. https://doi.org/10.1016/S0194-5998(00)70135-5

Bluestone, C. D., Stephenson, J. S., & Martin, L. M. (1992). Ten-year review of otitis media pathogens. *The*

Pediatric Infectious Disease Journal, *11*(8), S7–S11. https://doi.org/10.1097/00006454-199208001-00002

Bogaert, D., Keijser, B., Huse, S., Rossen, J., Veenhoven, R., Gils, E. V., … Sanders, E. (2011). Variability and

Diversity of Nasopharyngeal Microbiota in Children: A Metagenomic Analysis. *PLoS ONE*, *6*(2), e17035.

https://doi.org/10.1371/journal.pone.0017035

Brook, I. (2016). Microbiology of chronic rhinosinusitis. *European Journal of Clinical Microbiology & Infectious*

Diseases, *35*(7), 1059–1068. https://doi.org/10.1007/s10096-016-2640-x

Brook, I., Thompson, D. H., & Frazier, E. H. (1994). Microbiology and Management of Chronic Maxillary Sinusitis.

Archives of Otolaryngology–Head & Neck Surgery, *120*(12), 1317–1320.

Buffie, C. G., & Pamer, E. G. (2013). Microbiota-mediated colonization resistance against intestinal pathogens.

Nature Reviews Immunology, *13*(11), 790–801. https://doi.org/10.1038/nri3535

Burgstaller, J. M., Steurer, J., Holzmann, D., Geiges, G., & Soyka, M. B. (2016). Antibiotic efficacy in patients with

a moderate probability of acute rhinosinusitis: A systematic review. *European Archives of Oto-Rhino-*

Laryngology, *273*(5), 1067–1077. https://doi.org/10.1007/s00405-015-3506-z

Celin, S. E., Bluestone, C. D., Stephenson, J., Yilmaz, H. M., & Collins, J. J. (1991). Bacteriology of Acute Otitis

Media in Adults. *JAMA*, *266*(16), 2249–2252. https://doi.org/10.1001/jama.266.16.2249

Chira, S., & Miller, L. G. (2010). *Staphylococcus aureus* is the most common identified cause of cellulitis: A

systematic review. *Epidemiology and Infection*, *138*(3), 313–317.

https://doi.org/10.1017/S0950268809990483

Chua, K.-P., Fischer, M. A., & Linder, J. A. (2019). Appropriateness of outpatient antibiotic prescribing among

privately insured US patients: ICD-10-CM based cross sectional study. *BMJ*, *364*(k5092).

https://doi.org/10.1136/bmj.k5092

Durkin, M. J., Jafarzadeh, S. R., Hsueh, K., Sallah, Y. H., Munshi, K. D., Henderson, R. R., & Fraser, V. J. (2018).

Outpatient Antibiotic Prescription Trends in the United States: A National Cohort Study. *Infection Control*

& Hospital Epidemiology, *39*(5), 584–589. https://doi.org/10.1017/ice.2018.26

Edlin, R. S., Shapiro, D. J., Hersh, A. L., & Copp, H. L. (2013). Antibiotic resistance patterns of outpatient pediatric

urinary tract infections. *The Journal of Urology*, *190*(1), 222–227.

https://doi.org/10.1016/j.juro.2013.01.069

European Centre for Disease Prevention and Control. (2018). *Antimicrobial consumption: Annual epidemiological*

report for 2017. Stockholm.

Filice, G. A., Drekonja, D. M., Thurn, J. R., Hamann, G. M., Masoud, B. T., & Johnson, J. R. (2015). Diagnostic

Errors that Lead to Inappropriate Antimicrobial Use. *Infection Control & Hospital Epidemiology*, *36*(8),

949–956. https://doi.org/10.1017/ice.2015.113

Fleming-Dutra, K. E., Hersh, A. L., Shapiro, D. J., Bartoces, M., Enns, E. A., File, T. M., … Hicks, L. A. (2016).

Prevalence of Inappropriate Antibiotic Prescriptions Among US Ambulatory Care Visits, 2010-2011.

JAMA, *315*(17), 1864–1873. https://doi.org/10.1001/jama.2016.4151

Fokkens, W., Lund, V., & Mullol, J. (2007). European position paper on rhinosinusitis and nasal polyps 2007. A

summary for otorhinolaryngologists. *Rhinology*, *45*(97).

Ginsburg, C. M., McCracken, G. H., Crow, S. D., Dildy, B. R., Morchower, G., Steinberg, J. B., & Lancaster, K.

(1985). Seroepidemiology of the Group-A Streptococcal Carriage State in a Private Pediatric Practice.

American Journal of Diseases of Children, *139*(6), 614–617.

https://doi.org/10.1001/archpedi.1985.02140080084039

Gunnarsson, R. K., Holm, S. E., & Söderström, M. (1997). The prevalence of beta-haemolytic streptococci in throat

specimens from healthy children and adults: Implications for the clinical value of throat cultures.

Scandinavian Journal of Primary Health Care, *15*(3), 149–155.

https://doi.org/10.3109/02813439709018506

Gupta, K., Hooton, T. M., Naber, K. G., Wullt, B., Colgan, R., Miller, L. G., … Soper, D. E. (2011). International

Clinical Practice Guidelines for the Treatment of Acute Uncomplicated Cystitis and Pyelonephritis in

Women: A 2010 Update by the Infectious Diseases Society of America and the European Society for

Microbiology and Infectious Diseases. *Clinical Infectious Diseases*, *52*(5), e103–e120.

https://doi.org/10.1093/cid/ciq257

Gupta, K., Scholes, D., & Stamm, W. E. (1999). Increasing prevalence of antimicrobial resistance among

uropathogens causing acute uncomplicated cystitis in women. *JAMA*, *281*(8), 736–738.

https://doi.org/10.1001/jama.281.8.736

Gwaltney, Jack M., Scheld, W. M., Sande, M. A., & Sydnor, A. (1992). The Microbial Etiology and Antimicrobial

Therapy of Adults with Acute Community-Acquired Sinusitis—A 15-Year Experience at the University of

Virginia and Review of Other Selected Studies. *Journal of Allergy and Clinical Immunology*, *90*(3), 457–

462. https://doi.org/10.1016/0091-6749(92)90169-3

Hammitt, L. L., Bruden, D. L., Butler, J. C., Baggett, H. C., Hurlburt, D. A., Reasonover, A., & Hennessy, T. W.

(2006). Indirect Effect of Conjugate Vaccine on Adult Carriage of *Streptococcus pneumoniae*: An

Explanation of Trends in Invasive Pneumococcal Disease. *The Journal of Infectious Diseases*, *193*(11),

1487–1494. https://doi.org/10.1086/503805

Holgerson, P. L., Öhman, C., Rönnlund, A., & Johansson, I. (2015). Maturation of oral microbiota in children with

or without dental caries. *PLoS ONE*, *10*(5), 1–20. https://doi.org/10.1371/journal.pone.0128534

Hooton, T. M., & Gupta, K. (2019). Acute simple cystitis in women. In *UpToDate, Post, TW (Ed)*. Retrieved from

https://www.uptodate.com/contents/acute-simple-cystitis-in-women

Huang, Susan S., Hinrichsen, V. L., Stevenson, A. E., Rifas-Shiman, S. L., Kleinman, K., Pelton, S. I., …

Finkelstein, J. A. (2009). Continued Impact of Pneumococcal Conjugate Vaccine on Carriage in Young

Children. *Pediatrics*, *124*(1), e1–e11. https://doi.org/10.1542/peds.2008-3099

Huttenhower, C., Gevers, D., Knight, R., Abubucker, S., Badger, J. H., Chinwalla, A. T., … White, O. (2012).

Structure, function and diversity of the healthy human microbiome. *Nature*, *486*(7402), 207–214.

https://doi.org/10.1038/nature11234

515 Jain, S., Self, W. H., Wunderink, R. G., Fakhran, S., Balk, R., Bramley, A. M., … Finelli, L. (2015). Community-

516   Acquired Pneumonia Requiring Hospitalization among U.S. Adults. *New England Journal of Medicine*,

517   *373*(5), 415–427. https://doi.org/10.1056/NEJMoa1500245

518 Kabbani, S., Hersh, A. L., Shapiro, D. J., Fleming-Dutra, K. E., Pavia, A. T., & Hicks, L. A. (2018). Opportunities

519   to Improve Fluoroquinolone Prescribing in the United States for Adult Ambulatory Care Visits. *Clinical*

520   *Infectious Diseases*, *67*(1), 134–136. https://doi.org/10.1093/cid/ciy035

521 Khosravi, A., & Mazmanian, S. K. (2013). Disruption of the gut microbiome as a risk factor for microbial

522   infections. *Current Opinion in Microbiology*, *16*(2), 221–227. https://doi.org/10.1016/j.mib.2013.03.009

523 King, L. M., Bartoces, M., Fleming-Dutra, K. E., Roberts, R. M., & Hicks, L. A. (2019). Changes in US Outpatient

524   Antibiotic Prescriptions from 2011-2016. *Clinical Infectious Diseases*. https://doi.org/10.1093/cid/ciz225

525 Klevens, R. M., Caten, E., Olesen, S. W., DeMaria, A., Troppy, S., & Grad, Y. H. (2019). Outpatient antibiotic

526   prescribing in Massachusetts, 2011-2015. *Open Forum Infectious Diseases*, *6*(5), ofz169.

527   https://doi.org/10.1093/ofid/ofz169

528 *Lexicon Plus*. (2008). Retrieved from https://www.cerner.com/solutions/drug-database

529 Linder, J. A. (2008). *Editorial Commentary:* Antibiotics for Treatment of Acute Respiratory Tract Infections:

530   Decreasing Benefit, Increasing Risk, and the Irrelevance of Antimicrobial Resistance. *Clinical Infectious*

531   *Diseases*, *47*(6), 744–746. https://doi.org/10.1086/591149

532 Lipsitch, M. (2001). The rise and fall of antimicrobial resistance. *Trends in Microbiology*, *9*(9), 438–444.

533   https://doi.org/10.1016/S0966-842X(01)02130-8

534 MacLean, R. C., & San Millan, A. (2019). The evolution of antibiotic resistance. *Science*, *365*(6458), 1082–1083.

535   https://doi.org/10.1126/science.aax3879

536 Mainous, Arch G., Hueston, W. J., Everett, C. J., & Diaz, V. A. (2006). Nasal Carriage of *Staphylococcus aureus*

537   and Methicillin-Resistant *S aureus* in the United States, 2001-2002. *The Annals of Family Medicine*, *4*(2),

538   132–137. https://doi.org/10.1370/afm.526

539 National Center for Health Statistics. (2015a). *National Ambulatory Medical Care Survey—Public-use data file and*

540   *documentation*. Retrieved from ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/NAMCS

National Center for Health Statistics. (2015b). *National Hospital Ambulatory Medical Care Survey—Public-use*

data file and documentation. Retrieved from

ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/NHAMCS

Olesen, S. W., Barnett, M. L., MacFadden, D. R., Brownstein, J. S., Hernández-Díaz, S., Lipsitch, M., & Grad, Y.

H. (2018). The distribution of antibiotic use and its association with antibiotic resistance. *ELife*, *7*(e39435).

https://doi.org/10.7554/eLife.39435.001

Olesen, S. W., Barnett, M. L., MacFadden, D. R., Lipsitch, M., & Grad, Y. H. (2018). Trends in outpatient antibiotic

prescribing practice among US older adults, 2011-2015: An observational study. *BMJ*, *362*, k3155.

https://doi.org/10.1136/bmj.k3155

Pettigrew, M. M., Laufer, A. S., Gent, J. F., Kong, Y., Fennie, K. P., & Metlay, J. P. (2012). Upper respiratory tract

microbial communities, acute otitis media pathogens, and antibiotic use in healthy and sick children.

Applied and Environmental Microbiology, *78*(17), 6262–6270. https://doi.org/10.1128/AEM.01051-12

Pouwels, K. B., Batra, R., Patel, A., Edgeworth, J. D., Robotham, J. V., & Smieszek, T. (2017). Will co-trimoxazole

resistance rates ever go down? Resistance rates remain high despite decades of reduced co-trimoxazole

consumption. *Journal of Global Antimicrobial Resistance*, *11*, 71–74.

https://doi.org/10.1016/j.jgar.2017.07.013

Public Health Agency of Canada. (2018). *Canadian Antimicrobial Resistance Surveillance System 2017 Report*.

Public Health England. (2017). *English Surveillance Programme for Antimicrobial Utilisation and Resistance*

(ESPAUR) 2017.

Regev-Yochay, G., Dagan, R., Raz, M., Carmeli, Y., Shainberg, B., Derazne, E., … Rubinstein, E. (2004).

Association Between Carriage of *Streptococcus pneumoniae* and *Staphylococcus aureus* in Children.

JAMA, *292*(6), 716–720. https://doi.org/10.1001/jama.292.6.716

Rosenfeld, R. M., Piccirillo, J. F., Chandrasekhar, S. S., Brook, I., Kumar, K. A., Kramper, M., … Corrigan, M. D.

(2015). Clinical Practice Guideline (Update): Adult Sinusitis. *Otolaryngology–Head and Neck Surgery*,

152(2S), S1–S39. https://doi.org/10.1177/0194599815572097

Sakoulas, G., Geriak, M., & Nizet, V. (2018). Is a Reported Penicillin Allergy Sufficient Grounds to Forgo the

Multidimensional Antimicrobial Benefits of β-Lactam Antibiotics? *Clinical Infectious Diseases*, *68*(1),

157–164. https://doi.org/10.1093/cid/ciy557

Sande, M. A., & Gwaltney, J. M. (2004). Acute Community-Acquired Bacterial Sinusitis: Continuing Challenges

and Current Management. *Clinical Infectious Diseases*, *39*, S151–S158. https://doi.org/10.1086/421353

Shehab, N., Patel, P. R., Srinivasan, A., & Budnitz, D. S. (2008). Emergency Department Visits for Antibiotic-

Associated Adverse Events. *Clinical Infectious Diseases*, *47*(6), 735–743. https://doi.org/10.1086/591126

Smith, S. S., Evans, C. T., Tan, B. K., Chandra, R. K., Smith, S. B., & Kern, R. C. (2013). National burden of

antibiotic use for adult rhinosinusitis. *Journal of Allergy and Clinical Immunology*, *132*(5), 1230–1232.

https://doi.org/10.1016/j.jaci.2013.07.009

Snow, V., Mottur-Pilson, C., & Hickner, J. M. (2001). Principles of Appropriate Antibiotic Use for Acute Sinusitis

in Adults. *Annals of Internal Medicine*, *134*(6), 495. https://doi.org/10.7326/0003-4819-134-6-200103200-

00016

Stewardson, A. J., Gaïa, N., François, P., Malhotra-Kumar, S., Delémont, C., Martinez de Tejada, B., … Lazarevic,

V. (2015). Collateral damage from oral ciprofloxacin versus nitrofurantoin in outpatients with urinary tract

infections: A culture-free analysis of gut microbiota. *Clinical Microbiology and Infection*, *21*(4), 344.e1-

344.e11. https://doi.org/10.1016/j.cmi.2014.11.016

Swedres-Svarm. (2017). *Consumption of antibiotics and occurrence of antibiotic resistance in Sweden*. Public

Health Agency of Sweden and National Veterinary Institute.

Tedijanto, C., Olesen, S. W., Grad, Y. H., & Lipsitch, M. (2018). Estimating the proportion of bystander selection

for antibiotic resistance among potentially pathogenic bacterial flora. *Proceedings of the National Academy

of Sciences*, *115*(51), E11988–E11995. https://doi.org/10.1073/pnas.1810840115

The Human Microbiome Project Consortium. (2012). A framework for human microbiome research. *Nature*,

486(7402), 215–221. https://doi.org/10.1038/nature11209

Tobia, C. C., Aspinall, S. L., Good, C. B., Fine, M. J., & Hanlon, J. T. (2008). Appropriateness of antibiotic

prescribing in veterans with community-acquired pneumonia, sinusitis, or acute exacerbations of chronic

bronchitis: A cross-sectional study. *Clinical Therapeutics*, *30*(6), 1135–1144.

https://doi.org/10.1016/j.clinthera.2008.06.009

Tomas, M. E., Getman, D., Donskey, C. J., & Hecker, M. T. (2015). Overdiagnosis of Urinary Tract Infection and

Underdiagnosis of Sexually Transmitted Infection in Adult Women Presenting to an Emergency

Department. *Journal of Clinical Microbiology*, *53*(8), 2686–2692. https://doi.org/10.1128/JCM.00670-15

Vanden Eng, J., Marcus, R., Hadler, J. L., Imhoff, B., Vugia, D. J., Cieslak, P. R., … Besser, R. E. (2003).

Consumer Attitudes and Use of Antibiotics. *Emerging Infectious Diseases*, *9*(9), 1128–1135.

https://doi.org/10.3201/eid0909.020591

Verhaegh, S. J. C., Lebon, A., Saarloos, J. A., Verbrugh, H. A., Jaddoe, V. W. V., Hofman, A., … van Belkum, A.

(2010). Determinants of *Moraxella catarrhalis* colonization in healthy Dutch children during the first 14

months of life. *Clinical Microbiology and Infection*, *16*(7), 992–997. https://doi.org/10.1111/j.1469-

0691.2009.03008.x

Wald, E. R., Guerra, N., & Byers, C. (1991). Upper Respiratory Tract Infections in Young Children: Duration of and

Frequency of Complications. *Pediatrics*, *87*(2), 129–133.

Williams, J. W., Simel, D. L., Roberts, L., & Samsa, G. P. (1992). Clinical Evaluation for Sinusitis. *Annals of*

Internal Medicine, *117*(9), 705–710. https://doi.org/10.7326/0003-4819-117-9-705

Wubbel, Loretta, Muniz, L., Ahmed, A., Trujillo, M., Carubelli, C., McCoig, C., … McCracken, G. H. (1999).

Etiology and treatment of community-acquired pneumonia in ambulatory children. *The Pediatric Infectious*

Disease Journal, *18*(2), 98–104.

Yassour, M., Vatanen, T., Siljander, H., Hämäläinen, A.-M., Härkönen, T., Ryhänen, S. J., … Xavier, R. J. (2016).

Natural history of the infant gut microbiome and impact of antibiotic treatments on strain-level diversity

and stability. *Science Translational Medicine*, *8*(343), 129–139.

https://doi.org/10.1126/scitranslmed.aad0917

615
616

625 **Equation 1 – Figure Supplement 1.** Notation, descriptions, and sources for variables
626 in Equation 1.
627

Variable	Definition	Source
d_{aig}	Number of prescriptions (using nationally representative weights) of antibiotic a associated with ICD9-CM diagnosis code i in age group g	NAMCS/NHAMCS 2015
e_{sig}	Proportion of cases of condition defined by ICD9-CM diagnosis code i in age group g caused by species s	Published etiology studies (see Figure 1 – Source Data File 1)
p_{s0g}	Asymptomatic carriage prevalence of species s in age group g	Human Microbiome Project (HMP) and published carriage studies
p_{sig}	Carriage prevalence of species s among individuals diagnosed with ICD9-CM code i in age group g	$e_{sig} + (1 - e_{sig})p_{s0g}$
q_{aig}	Proportion of prescriptions of antibiotic a associated with ICD9-CM diagnosis code i in age group g that are avertable under the given scenario	Based on article by Fleming-Dutra et al. (Fleming-Dutra et al., 2016) with adjustments as described in Figure 1 – Source Data File 2; note that in the nitrofurantoin-specific scenario, this variable is 1 for all antibiotic use for cystitis (ICD codes defined in main text) in women of all age groups, and 0 otherwise
X_{as}	Number of exposures of antibiotic a experienced by species s that are avertable under the given scenario	$\sum_{g=1}^{G} \sum_{i=0}^{I} d_{aig} \times p_{sig} \times q_{aig}$
T_{as}	Total number of exposures of antibiotic a experienced by species s	$\sum_{g=1}^{G} \sum_{i=0}^{I} d_{aig} \times p_{sig}$

Figure 1 – Figure Supplement 1. Sensitivity analysis of proportions of avertable exposures across all outpatient conditions when proportion of unnecessary use is applied equally across target and bystander exposures (Equation 1 without modification).

	EC	HI	KP	MC	PA	SA	SAg	SP	SPy	Weighted prescription rate (per 10,000 PY)
AMOXICILLIN	27	27	25	23	26	26	26	25	36	810.5
AMX-CLAV	37	37	35	30	37	36	36	35	51	415.3
PENICILLIN	29	29	29	32	29	29	30	30	44	83.6
PENICILLINS	29	29	27	25	29	29	29	27	40	**1381.7**
AZITHROMYCIN	49	49	48	44	48	49	48	48	51	869.9
CLINDAMYCIN	18	18	18	20	18	19	19	18	20	243.8
CLARITHROMYCIN	27	27	26	25	27	27	26	27	27	51.0
MACR/LINC	39	39	38	39	38	38	39	38	41	**1332.0**
CIPROFLOXACIN	13	15	13	20	10	15	15	15	15	547.2
LEVOFLOXACIN	30	32	30	33	26	32	33	32	33	285.0
MOXIFLOXACIN	17	17	17	20	17	17	17	17	16	71.4
QUINOLONES	19	20	18	24	15	21	21	21	20	**1026.8**
CEPHALEXIN	20	20	19	21	18	20	21	20	26	462.6
CEFDINIR	24	25	22	17	21	24	22	21	58	123.7
CEPHALOSPORINS	25	27	24	20	23	26	25	26	48	**1116.7**
DOXYCYCLINE	31	31	31	32	30	31	31	31	31	259.7
TETRACYCLINE	7	7	7	7	7	7	7	7	7	12.2
MINOCYCLINE	23	23	23	23	23	23	23	23	23	110.8
TETRACYCLINES	28	28	28	30	27	28	28	28	28	**383.1**
TMP-SMX	16	18	17	24	13	19	20	19	19	398.5
NITROFURANTOIN	12	13	12	14	10	14	14	14	14	191.8

Legend: 0 >0 ≥5 ≥10 ≥15 ≥20 ≥25 ≥30 ≥35 ≥40 ≥45 ≥50 NA

Penicillins
Macrolides/lincosamides
Quinolones
Cephalosporins
Tetracyclines
Sulfamethoxazole-trimethoprim
Nitrofurantoin

Abbreviations: *Antibiotics (y-axis):* AMX-CLAV=amoxicillin-clavulanate, MACR/LINC=macrolides/lincosamides, TMP-SMX=sulfamethoxazole-trimethoprim; *Organisms (x-axis):* EC=*E. coli*, HI=*H. influenzae*, KP=*K. pneumoniae*, MC=*M. catarrhalis*, PA=*P. aeruginosa*, SA=*S. aureus*, SAg=*S. agalactiae*, SP=*S. pneumoniae*, SPy=*S. pyogenes*; PY=person-years

A. Use estimates of unnecessary prescribing for acute sinusitis from Fleming-Dutra et al. Based on
Fleming-Dutra et al. (Fleming-Dutra et al., 2016), we estimate that the proportion of unnecessary
antibiotic prescriptions for acute sinusitis is 18%, 100%, and 34% for individuals who are 0-19, 20-64, and
over 65 years old, respectively (see Figure 1 – Source Data File 2).

B. Use estimates of bacterial etiology to approximate necessary antibiotic use for acute sinusitis.
The following upper bound estimates for the proportion of acute sinusitis cases with bacterial etiology
were used: 2% for adults and 13% for children (applied to 0-19 year old age group) (Sande & Gwaltney,
2004; Snow et al., 2001; Wald et al., 1991).

C. Use estimates of bacterial etiology among those presenting to primary care clinics to
approximate necessary antibiotic use. Patients visiting the clinic are likely to have more persistent,
severe, or worsening symptoms, which may be indications of bacterial etiology. Studies have estimated
that approximately 38% of adults and 17% of children (upper bounds) visiting primary care for symptoms
of acute upper respiratory tract infection have bacterial rhinosinusitis (Aitken & Taylor, 1998; Benninger et
al., 2000; Williams et al., 1992).

662
663
664 **Abbreviations:** *Antibiotics (y-axis):* AMX-CLAV=amoxicillin-clavulanate, MACR/LINC=macrolides/
665 lincosamides, TMP-SMX=sulfamethoxazole-trimethoprim; *Organisms (x-axis):* EC=*E. coli*, HI=*H.*
666 *influenzae*, KP=*K. pneumoniae*, MC=*M. catarrhalis*, PA=*P. aeruginosa*, SA=*S. aureus*, SAg=*S.*
667 *agalactiae*, SP=*S. pneumoniae*, SPy=*S. pyogenes*; PY=person-years
668
669

Figure 1 – Figure Supplement 3. Proportions of avertable exposures under Scenario 1
(elimination of all unnecessary antibiotic use across outpatient conditions) using 2010-
2011 NAMCS/NHAMCS data with *(left)* and without *(right)* NHAMCS outpatient
department data.

Abbreviations: *Antibiotics (y-axis):* AMX-CLAV=amoxicillin-clavulanate, MACR/LINC=macrolides/
lincosamides, TMP-SMX=sulfamethoxazole-trimethoprim; *Organisms (x-axis):* EC=*E. coli*, HI=*H.*
influenzae, KP=*K. pneumoniae*, MC=*M. catarrhalis*, PA=*P. aeruginosa*, SA=*S. aureus*, SAg=*S.*
agalactiae, SP=*S. pneumoniae*, SPy=*S. pyogenes*; PY=person-years
679

Figure 1 – Source Data File 1. Summary of etiology studies.

Publication	Study description	Etiology results
Bacteriology of Acute Otitis Media in Adults (Celin et al., 1991)	34 outpatients aged 18 years or older presenting at the emergency department and diagnosed with acute otitis media; middle-ear aspirates were collected for culture	*H. influenzae*: 26% *M. catarrhalis*: 3% *S. aureus*: 3% *S. pneumoniae*: 21% *S. pyogenes*: 3%
Ten-year review of otitis media pathogens (Bluestone et al., 1992)	2807 effusions from children with acute otitis media were collected	*H. influenzae*: 23% *M. catarrhalis*: 14% *S. aureus*: 1% *P. aeruginosa*: 1% *S. pneumoniae*: 35% *S. pyogenes*: 3%
The microbial etiology and antimicrobial therapy of adults with acute community-acquired sinusitis: A fifteen-year experience at the University of Virginia and review of other selected studies (Gwaltney et al., 1992)	339 adult patients with acute community-acquired sinusitis; sinus aspirates were collected prior to treatment	To estimate the proportion of acute sinusitis cases caused by each organism, we multiplied the proportion of bacterial cultures in which each organism was identified (shown below) by the proportion of all acute sinusitis cases estimated to be bacterial in nature (2%) (Fokkens et al., 2007): *H. influenzae*: 35% *M. catarrhalis*: 4% *S. aureus*: 3% *S. pneumoniae*: 41%
Microbiology and Management of Chronic Maxillary Sinusitis (Brook et al., 1994)	68 patients aged 18 to 67 years who underwent the Caldwell-Luc procedure for chronic sinusitis and whose cultures showed bacterial growth; specimens were taken from the infected sinus during surgery	*E. coli*: 2.9% *H. influenzae*: 4.4% *K. pneumoniae*: 2.9% *M. catarrhalis*: 11.8% *S. aureus*: 11.8% *S. agalactiae*: 5.9% *S. pneumoniae*: 5.9% *S. pyogenes*: 7.4%
Increasing Prevalence of Antimicrobial Resistance Among Uropathogens Causing Acute Uncomplicated Cystitis in Women (Gupta et al., 1999)	4802 women aged 18 to 50 years with an outpatient diagnosis of acute cystitis; urine samples were collected	<u>Female</u> *E. coli*: 86% *Klebsiella* spp.: 3%
Etiology and treatment of community-acquired pneumonia in ambulatory children (Wubbel et al., 1999)	174 ambulatory patients age 6 months to 16 years with radiographic and clinical evidence of pneumonia; blood, nasopharyngeal, and pharyngeal samples were collected	*S. pneumoniae*: 27%

Staphylococcus aureus is the most common identified cause of cellulitis: a systematic review (Chira & Miller, 2010)	808 patients with cellulitis who were included in 16 studies published between 1966 and 2006; studies were eligible for inclusion if they used needle aspiration and/or punch biopsy and excluded if they pertained to ocular, odontogenic, pelvic, or surgical site-associated cellulitis	*E. coli*: 0.4% *H. influenzae*: 0.1% *P. aeruginosa*: 0.5% *S. aureus*: 8% *S. agalactiae*: 0.5% *S. pyogenes*: 4.3%
Antibiotic Resistance Patterns of Outpatient Pediatric Urinary Tract Infections (Edlin et al., 2013)	25,418 urinary isolates from patients younger than 18 years in the outpatient setting from 195 United States hospitals in The Surveillance Network	<u>Female</u> *E. coli*: 83% *Klebsiella* spp.: 4% *P. aeruginosa*: 2% <u>Male</u> *E. coli*: 50% *Klebsiella* spp.: 10% *P. aeruginosa*: 7%
Community-Acquired Pneumonia Requiring Hospitalization among U.S. Adults (Jain et al., 2015)	2259 patients who were hospitalized for community-acquired pneumonia (based on radiographic evidence) and had specimens for both bacterial and viral testing; specimens may have been taken from blood, acute-phase serum, urine, nasopharynx, oropharynx, sputum, pleural fluid, endotracheal aspirate, or broncho-alveolar-lavage samples	*H. influenzae*: 0.6% *Pseudomonas* spp.: 0.4% *S. aureus*: 1.6% *S. pneumoniae*: 5.1% *S. pyogenes*: 0.3%

Figure 1 – Source Data File 2. Modifications to proportions of unnecessary antibiotic
prescriptions based on published etiology studies and antibiotic use in
NAMCS/NHAMCS. Modifications were made if the estimated proportion of bacterial
cases for a given condition exceeded the estimated proportion of appropriate
prescriptions reported in Fleming-Dutra et al. (Fleming-Dutra et al., 2016).

Age group	Fleming-Dutra estimate	New estimate	Explanation
Pharyngitis			
0-19 years old	34%	Pharyngitis: 47%; Strep throat: 0%	Assuming that all strep-associated use of our antibiotics of interest (based on NAMCS/NHAMCS) was necessary, we estimated the proportion of non-strep pharyngitis antibiotic use that would have to be eliminated to maintain the overall estimate reported by Fleming-Dutra et al. (Fleming-Dutra et al., 2016).
20-64 years old	75%	Updated to 51% overall; Pharyngitis: 100%; Strep throat: 0%	
>64 years old	NA (assume 0%)	NA (assume 0%)	No change.
Sinusitis			
0-19 years old	9%	Acute: 18%; Chronic: 0%	While acute sinusitis is rarely bacterial, bacteria are believed to play a much more prominent role in the etiology of chronic sinusitis (Brook, 2016). Thus, we assume that antibiotic use would only be eliminated from acute sinusitis and maintain the overall estimate of unnecessary use reported by Fleming-Dutra et al.(Fleming-Dutra et al., 2016) when possible.
20-64 years old	51%	Updated to 40% overall; Acute: 100%; Chronic: 0%	
>64 years old	16%	Acute: 34%; Chronic: 0%	
Suppurative otitis media			
0-19 years old	10%	10%	No change. Approximately 84% of pediatric cases were found to have bacterial growth (Bluestone et al., 1992), so 10% is a reasonable estimate for avertable use.
20-64 years old	33%	26%	Based on an etiology study in adults, only 26% of cases were found to have no bacterial growth (Celin et al., 1991).
>64 years old	NA (assume 0%)	NA (assume 0%)	No change.

689 **Figure 1 – Source Data File 3.** Estimated antibiotic class exposures by exposed and
690 causative species.
691
692 The following tables contain the estimated number of annual antibiotic exposures (in 100,000s) in the
693 outpatient setting in the United States by the species being exposed (y-axis) and the species that is the
694 cause of antibiotic use (x-axis). The values in each table, excluding the "Other" and "Overall" columns
695 were calculated using the following expression:
696

$$\sum_{g=1}^{G}\sum_{i=0}^{I} d_{aig} \times e_{s_{c}ig} \times p_{s_{e}0g}$$

698
Let *a* represent antibiotic class, s_c represent causative species, s_e represent exposed species, *i* represent
ICD-9-CM diagnosis code, and *g* represent age group (under 1 year, 1-5 years, 6-19 years, 20-64 years,
65 years old and over). Then, d_{aig} is the number of prescriptions of antibiotic class *a* associated with
diagnosis code *i* in age group *g*, $e_{s_{c}ig}$ is the proportion of cases with diagnosis code *i* that are estimated to
be caused by causative species s_c in age group *g*, and $p_{s_{e}0g}$ is the healthy carriage prevalence of
exposed species s_e among individuals in age group *g*. When s_c and s_e are the same organism, $p_{s_{e}0g}$ is set
to 1. For each variable, we use the same data sources and estimation methods described in the main text
(NAMCS/NHAMCS 2015, Human Microbiome Project, and published literature).
707
The diagonals, shaded in light gray, are "target" antibiotic exposures for the pathogen of interest; in other
words, these are antibiotic exposures that the organism experienced because it was causing disease.
The "Overall" column is all exposures to the antibiotic class of interest experienced by each species and
is equivalent to T_{as} in Equation 1 of the main text. The "Other" column is the difference between the
"Overall" column and the sum of the remaining columns. Note that this column represents exposures due
to non-bacterial causes as well as bacterial organisms other than our nine potential pathogens of interest.
714
The "Any antibiotic" table includes all prescriptions for drugs classified as antibiotics based on the Lexicon
Plus® Therapeutic Classification Scheme (*Lexicon Plus*, 2008). Any drug with a Level 1 description of
"anti-infectives", but with a Level 2 description that was not "amebicides", "antihelmintics", "antifungals",
"antimalarial agents", or "antiviral agents" was included as an antibiotic. The "Any included antibiotic
class" table includes all prescriptions for drugs in the classes shown in the following tables. Antibiotic
classes were defined using the Level 2 categories in Lexicon Plus®.
721
These tables are designed to facilitate a better understanding of the "nuts and bolts" of this work and to
enable calculation of other metrics of interest. For example, the proportion of bystander exposures as
calculated by the authors in an earlier paper (Tedijanto et al., 2018) for any given antibiotic and species
pair can be calculated as one minus the target exposures (diagonal value) over the "Overall" value. These
tables can also be used to understand how interventions focused on a single species (e.g. vaccines) may
reduce antibiotic exposures experienced by another species.
728
729
730

Any antibiotic

Exposed species	Causative species										Overall
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	
E. coli	150.2	25.1	7.7	15.9	4.2	12.5	3.6	34.5	43.5	1838.4	2135.6
H. influenzae	103.9	31	7.9	16	4.4	12.9	3.7	34.4	44.3	1892.7	2151.2
K. pneumoniae	11.5	4.2	11.4	2.7	0.5	1.5	0.4	6.2	5.4	220.8	264.7
M. catarrhalis	4.9	7.3	0.4	19.6	0.5	1.1	0.3	12.2	6	139	191.3
P. aeruginosa	2.8	0.5	0.2	0.3	6.1	0.3	0.1	0.6	1.1	50.3	62.4
S. aureus	18.8	5.3	1.4	3.4	0.8	18.2	0.7	7.5	8.2	349.4	413.9
S. agalactiae	4.2	1.6	0.3	1	0.2	0.6	5.2	2.4	2.1	80.6	98.2
S. pneumoniae	39.2	13.3	3	8.5	1.8	5.1	1.4	40.3	19.1	743.6	875.4
S. pyogenes	7.1	1.3	0.5	0.8	0.3	0.8	0.2	1.6	60.8	125.3	198.8

731
732

Any included antibiotic class

Exposed species	Causative species										Overall
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	
E. coli	143.3	23.8	7.3	15	4	11.6	3.3	32.8	42.9	1042.9	1327
H. influenzae	99.2	29.5	7.6	15.1	4.1	11.9	3.4	32.8	43.7	1072.1	1319.4
K. pneumoniae	11	4	10.9	2.5	0.5	1.4	0.4	6	5.3	126.1	168.1
M. catarrhalis	4.7	6.9	0.4	18.5	0.4	1	0.3	11.6	5.9	88.8	138.5
P. aeruginosa	2.7	0.5	0.2	0.3	5.8	0.3	0.1	0.6	1.1	28.2	39.8
S. aureus	18	5.1	1.4	3.2	0.8	16.9	0.6	7.2	8.1	198.3	259.5
S. agalactiae	4	1.5	0.3	1	0.2	0.5	4.9	2.3	2.1	46.6	63.3
S. pneumoniae	37.4	12.6	2.9	8	1.7	4.8	1.4	38.4	18.8	427.4	553.4
S. pyogenes	6.7	1.2	0.5	0.8	0.3	0.8	0.2	1.5	59.9	70.5	142.5

733
734

Penicillins

Exposed species	Causative species										Overall
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	
E. coli	2.7	13.7	0.8	8.5	0.4	3.9	1.4	18.9	23.6	247.9	321.7
H. influenzae	2	16.3	0.8	8.5	0.4	3.9	1.4	18.8	23.8	253.1	329.1
K. pneumoniae	0.3	2.5	1.1	1.5	0.1	0.5	0.2	3.7	3.1	32	44.9
M. catarrhalis	0.3	4.6	0.1	9.9	0.2	0.6	0.2	7.4	4.6	34.3	62.2
P. aeruginosa	<0.1	0.3	<0.1	0.2	0.5	0.1	<0.1	0.3	0.6	6.3	8.3
S. aureus	0.4	3	0.1	1.9	0.1	5.3	0.3	4.3	4.5	47.9	67.8
S. agalactiae	0.1	0.9	<0.1	0.6	<0.1	0.2	2	1.4	1.3	12.3	18.8
S. pneumoniae	0.9	7.7	0.3	4.8	0.3	1.7	0.6	21.3	11.1	109.1	157.9
S. pyogenes	0.1	0.7	<0.1	0.4	<0.1	0.2	0.1	0.8	31.5	15.9	49.8

Macrolides/lincosamides

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	3.3	4.1	0.6	3	0.2	3	1	5.7	3.9	264.9	289.7
H. influenzae	2.3	5.6	0.6	3	0.2	3.1	1.1	5.8	4	272.8	298.4
K. pneumoniae	0.2	0.6	0.9	0.4	<0.1	0.4	0.1	1	0.5	31.8	36
M. catarrhalis	0.1	0.7	<0.1	4.1	<0.1	0.2	<0.1	1.4	0.4	19.6	26.6
P. aeruginosa	0.1	0.1	<0.1	0.1	0.2	0.1	<0.1	0.1	0.1	7.3	8.1
S. aureus	0.4	0.9	0.1	0.6	<0.1	4.5	0.2	1.2	0.7	50.4	59.1
S. agalactiae	0.1	0.2	<0.1	0.2	<0.1	0.1	1.5	0.3	0.2	11.6	14.3
S. pneumoniae	0.8	1.9	0.2	1.3	0.1	1.2	0.4	7.3	1.6	106.9	121.8
S. pyogenes	0.2	0.2	<0.1	0.2	<0.1	0.2	0.1	0.3	5.6	18.1	24.9

736
737

Quinolones

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	61.3	0.7	2.6	0.5	1.4	0.5	0.2	1	0.3	168.7	237.2
H. influenzae	42.1	1	2.7	0.5	1.5	0.5	0.2	1	0.3	174.3	224.1
K. pneumoniae	4.5	0.1	3.9	0.1	0.2	0.1	<0.1	0.1	<0.1	19.1	28.1
M. catarrhalis	1.4	0.1	0.1	0.7	0.1	<0.1	<0.1	0.2	<0.1	7.2	9.8
P. aeruginosa	1.2	<0.1	0.1	<0.1	2.1	<0.1	<0.1	<0.1	<0.1	4.8	8.2
S. aureus	7.6	0.1	0.5	0.1	0.3	0.8	<0.1	0.2	0.1	31.5	41.1
S. agalactiae	1.6	<0.1	0.1	<0.1	0.1	<0.1	0.3	<0.1	<0.1	6.9	9.1
S. pneumoniae	15.5	0.3	1	0.2	0.5	0.2	0.1	1.3	0.1	65	84.3
S. pyogenes	2.9	<0.1	0.2	<0.1	0.1	<0.1	<0.1	0.1	0.5	11.9	15.8

738
739

Cephalosporins

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	38.3	4.6	1.7	2.4	1.1	2.4	0.4	6.6	14	185.7	257.3
H. influenzae	26.8	5.6	1.8	2.4	1.1	2.5	0.4	6.5	14.5	190.8	252.3
K. pneumoniae	3	0.7	2.5	0.4	0.1	0.3	<0.1	1.1	1.6	21.9	31.9
M. catarrhalis	1.8	1.4	0.1	2.8	0.1	0.2	<0.1	2.5	0.9	15.7	25.6
P. aeruginosa	0.7	0.1	<0.1	<0.1	1.6	0.1	<0.1	0.1	0.4	5	8.1
S. aureus	4.9	0.9	0.3	0.5	0.2	3.6	0.1	1.4	2.6	35	49.5
S. agalactiae	1.1	0.3	0.1	0.2	0.1	0.1	0.6	0.5	0.6	8.3	11.7
S. pneumoniae	10.4	2.5	0.7	1.4	0.5	1	0.1	7.5	5.6	76	105.5
S. pyogenes	1.8	0.2	0.1	0.1	0.1	0.2	<0.1	0.3	20.7	12.6	36.2

Tetracyclines

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	0.7	0.3	0.1	0.1	<0.1	0.3	0.1	0.3	0.2	78.6	80.6
H. influenzae	0.5	0.4	0.1	0.1	<0.1	0.3	0.1	0.3	0.2	81.3	83.3
K. pneumoniae	0.1	<0.1	0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	8.8	9.1
M. catarrhalis	<0.1	<0.1	<0.1	0.2	<0.1	<0.1	<0.1	<0.1	<0.1	2.8	3
P. aeruginosa	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	2.2	2.3
S. aureus	0.1	0.1	<0.1	<0.1	<0.1	0.5	<0.1	<0.1	<0.1	14.6	15.4
S. agalactiae	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.1	<0.1	<0.1	3.2	3.3
S. pneumoniae	0.2	0.1	<0.1	<0.1	<0.1	0.1	<0.1	0.4	0.1	29.9	30.9
S. pyogenes	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.3	5.6	6

741
742

Sulfamethoxazole-trimethoprim

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	23.1	0.3	1.1	0.5	0.6	1.5	0.3	0.4	0.8	67	95.5
H. influenzae	16.1	0.5	1.1	0.5	0.6	1.5	0.3	0.4	0.9	68.8	90.5
K. pneumoniae	1.8	<0.1	1.6	0.1	0.1	0.2	<0.1	<0.1	0.1	9	12.9
M. catarrhalis	0.9	<0.1	0.1	0.7	<0.1	0.1	<0.1	<0.1	<0.1	8.1	10
P. aeruginosa	0.4	<0.1	<0.1	<0.1	0.9	<0.1	<0.1	<0.1	<0.1	1.7	3.2
S. aureus	2.9	0.1	0.2	0.1	0.1	2.2	0.1	0.1	0.2	13.3	19.1
S. agalactiae	0.7	<0.1	<0.1	<0.1	<0.1	0.1	0.4	<0.1	<0.1	3.2	4.5
S. pneumoniae	6.1	0.1	0.4	0.2	0.2	0.6	0.1	0.6	0.3	29	37.6
S. pyogenes	1.1	<0.1	0.1	<0.1	<0.1	0.1	<0.1	<0.1	1.2	4.3	6.9

743
744

Nitrofurantoin

Exposed species	Causative species										
	E. coli	H. influenzae	K. pneumoniae	M. catarrhalis	P. aeruginosa	S. aureus	S. agalactiae	S. pneumoniae	S. pyogenes	Other	Overall
E. coli	13.8	<0.1	0.6	<0.1	0.3	<0.1	<0.1	<0.1	<0.1	30.1	44.9
H. influenzae	9.5	<0.1	0.6	<0.1	0.3	<0.1	<0.1	<0.1	<0.1	31.2	41.6
K. pneumoniae	1	<0.1	0.9	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	3.4	5.3
M. catarrhalis	0.3	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	1.1	1.4
P. aeruginosa	0.3	<0.1	<0.1	<0.1	0.5	<0.1	<0.1	<0.1	<0.1	0.9	1.6
S. aureus	1.7	<0.1	0.1	<0.1	0.1	<0.1	<0.1	<0.1	<0.1	5.6	7.5
S. agalactiae	0.4	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	1.2	1.6
S. pneumoniae	3.5	<0.1	0.2	<0.1	0.1	<0.1	<0.1	<0.1	<0.1	11.5	15.4
S. pyogenes	0.7	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	2.1	2.9

745
746