Living alone, loneliness and lack of emotional support as predictors of suicide and self-harm: a nine-year follow up of the UK Biobank cohort

Authors: Richard J. Shaw[1], Breda Cullen[1], Nicholas Graham[1], Donald M. Lyall[1], Daniel Mackay[1], Chukwudi Okolie[2], Robert Pearsall[1,3], Joey Ward[1], Ann John[2], Daniel J. Smith[1]

1. Institute of Health and Wellbeing, University of Glasgow, UK.

2. Institute of Life Sciences, Medical School, Swansea University, Swansea, UK.

3. Department of Psychiatry, Monklands Hospital, Airdrie, UK.

Declarations of interest: none.

Correspondence to:

Dr. Richard Shaw

Address: Institute of Health and Wellbeing, University of Glasgow, Public Health, 1 Lilybank Gardens, Glasgow, G12 8RZ, UK.

Telephone: +44 (0) 141 330 2334.

Email: Richard.Shaw@glasgow.ac.uk.

Highlights

- First cohort study to investigate loneliness's relationship with deaths by suicide
- Loneliness is associated with a modest increased risk of death by suicide for men
- For men, living with a partner reduces the risk of death by suicide
- Loneliness increases the risk of hospitalization for self-harm for men and women

Abstract

Background: The association between loneliness and suicide is poorly understood. We investigated how living alone, loneliness and emotional support were related to suicide and self-harm in a longitudinal design.

Methods: Between 2006 and 2010 UK Biobank recruited and assessed in detail over 0.5 million people in middle age. Data were linked to prospective hospital admission and mortality records. Adjusted Cox regression models were used to investigate relationships between living arrangements, loneliness and emotional support, and both suicide and self-harm as outcomes.

Results: For men, both living alone (Hazard Ratio (HR) 2.16, 95%CI 1.51-3.09) and living with non-partners (HR 1.80, 95%CI 1.08-3.00) were associated with death by suicide, independently of loneliness, which had a modest relationship with suicide (HR 1.43, 95%CI 0.1.01-2.03). For women, there was no evidence that living arrangements, loneliness or emotional support were associated with death by suicide. Associations between living alone and self-harm were explained by health for women, and by health, loneliness and emotional support for men. In fully adjusted models, loneliness was associated with hospital admissions for self-harm in both women (HR 1.89, 95%CI 1.57-2.28) and men (HR 1.74, 95%CI 1.40-2.16).

Limitations: Loneliness and emotional support were operationalized using single item measures.

Conclusions: For men - but not for women - living alone or living with a non-partner increased the risk of suicide, a finding not explained by subjective loneliness. Overall, loneliness may be more important as a risk factor for self-harm than for suicide. Loneliness also appears to lessen the protective effects of cohabitation.

Keywords

Loneliness, Suicide, Self-harm, Living arrangements, Emotional support.

INTRODUCTION

Loneliness, defined as *the subjective perception of a lack of contact with other people* (HM Government, 2018; Perlman and Peplau, 1981), is associated with premature mortality (Elovainio et al., 2017; Rico-Uribe et al., 2018), physical and mental ill-health, worse cognitive function (Cacioppo et al., 2014; Hakulinen et al., 2018; Hawkley and Cacioppo, 2010; Solmi et al., 2020) and increased use of health services (Dreyer et al., 2018). Loneliness was already a major concern due to the consequences of population ageing and greater numbers of people living alone, before the UK government recently prioritised loneliness by making it a ministerial responsibility (HM Government, 2018). Fears around the consequences of physical distancing in relation to Covid-19 have only increased those concerns (Gunnell et al 2020.; Holmes et al., 2020). However, it is currently not clear whether subjective loneliness *per se* is the primary reason why people living alone may be at increased risk of suicidal behaviour.

Living alone and loneliness indicate relationships and social connections, nonetheless they are separate constructs with overlapping features (Smith and Victor, 2019). Living alone is distinct from cohabiting relationships, as well as residency with non-partners such as parents, children or friends, who might be expected to be sources of emotional, financial and practical support (Amato, 2014; van Hedel et al., 2018). However, clearly people living alone may engage with others outside the household and potentially receive emotional support from other sources. In contrast, loneliness is the *subjective perception* of a lack of contact with other people (Hawkley and Cacioppo, 2010).

Living alone has been consistently linked with self-harm and suicide (Conejero et al., 2016; Frisch and Simonsen, 2013). Theories such as the Interpersonal Theory of Suicide hypothesise that loneliness and emotional support are central to the development of suicidal behaviour (Stanley et al., 2016; Van Orden et al., 2010). However, there is little empirical research in this area (Van Orden et al., 2010). Identifying robust risk factors for suicidal behaviour is methodologically challenging for a number of reasons, not least because suicide is a rare event (Klonsky et al., 2016; Stickley and Koyanagi, 2016). Most studies of loneliness and suicidal behaviour have used self-reported measures of suicidality (Bennardi et al., 2019), which may be prone to reporting biases (Beutel et al., 2017; Stickley and Koyanagi, 2016), and only a few studies (mostly case-control studies) have investigated loneliness as a potential cause of deaths by suicide (Courtin and Knapp, 2017; Holt-Lunstad et al., 2015).

The extensive data within the UK Biobank cohort represents a unique opportunity to overcome these methodological challenges. The cohort consists of over half a million people, and the baseline questionnaire included detailed questions on living arrangements, loneliness and emotional support, in addition to key sociodemographic and health data. These data have also been linked prospectively to hospital episode statistics and mortality records (Sudlow et al., 2015).

Our primary hypothesis was that living alone may represent an independent risk factor for self-harm and suicide. We also set out to assess whether any observed association between living alone and suicidal behaviour might be explained by subjective loneliness or by perceived lack of emotional support.

METHODS
Data
All adults aged between 40-70 years who were registered with the UK National Health Service (NHS) and living within 25 miles of 22 assessment centres across England, Scotland and Wales were invited

to participate in UK Biobank at baseline. The achieved sample of 502,536 people had a response rate of 5.5% (Sudlow et al., 2015) and an age range of between 37 and 73. Baseline assessments took place between 2006 and 2010 (Sudlow et al., 2015) and consisted of a single visit lasting approximately two to three hours, including a computerised touch screen questionnaire, nurse interviews, and physical measurements. For all participants, the date and cause of death were sought from death certificates held within the National Health Service Information Centre (England and Wales), and National Health Service Central Register Scotland. At the time of analysis, mortality data were available up to the middle of February 2018 for England and Wales and until June 2017 for Scotland. All participants who attended a Biobank assessment centre in England (n = 448,811) were linked to Hospital Episode statistics (HES) to obtain hospital admission records for self-harm up to March 2015. This study is covered by the generic ethical approval for UK Biobank studies from the National Health Service National Research Ethics Service (June 17, 2011; Ref 11/NW/0382). Participants provided electronic informed consent for the baseline assessments and the register linkage.

Outcomes

Death by suicide was defined as the act of intentionally ending one's own life (Nock et al., 2008) and was ascertained from death records using ICD 10 codes X60-X84 (intentional self-harm), Y10-34 (undetermined cause), as used by the UK Office for National Statistics (2016). Participants dying from other causes of death were censored at time of death.

Hospital admissions for self-harm were defined as any act of intentional self-poisoning or self-injury carried out by an individual, irrespective of the motivation or suicidal intent (National Collaborating Centre for Mental Health, 2012). This was assessed using the first admission for self-harm following attendance at the UK Biobank baseline assessment centre. Hospitalization for self-harm was assessed using ICD 10 codes X60 to-84 and Z91.5, for diagnosis and causes of admissions.

Main exposures of interest

Living arrangements (alone; husband, wife or partner; other) were assessed using data from the baseline touch screen questionnaire. Participants were asked how many people lived in their household. If there was more than one person, the participant was then asked how people were related to them. If any member of the household was a spouse or partner, participants were classified as living with a husband, wife or partner. The other category included both relatives and unrelated people.

Loneliness was assessed using a single question taken from the baseline touchscreen questionnaire: "Do you often feel lonely?" (Responses: yes; no; do not know; prefer not to answer). This item was taken from a longer scale and has previously been shown to be associated with health outcomes (McCormack et al., 2014).

Emotional Support was assessed with the question "2-4 times a week; about once a week; about once a month; once every few months; never or almost never; do not know; prefer not answer.)

Covariates

We included baseline sociodemographic and health variables which might confound relationships between the key variables of interest and death by suicide and hospital admissions for self-harm.

The socio-demographic variables included were: sex, age (continuous), and self-reported measures of ethnicity (derived here into White British, other), current employment status, (employed, retired, other), education (degree; professional; NVQ, HND or HNC; A level; O level; CSE; none) ever having a

same sex partner (none, at least one), and area deprivation indicated using the Townsend Index (continuous) for the participant's postcode at recruitment.

The health variables included were: number of physical morbidities participants reported to the interviewing nurse (zero, one, two, three or more); Body Mass Index (BMI) based on measurements made at the assessment centre (normal or underweight, overweight, obese); self-reported measures of ever seeing a GP for depression (yes, no); taking psychotropic medication at baseline (yes/no); alcohol consumption (daily/almost daily, 3-4 days a week, 1-2 times a week /once a month, special occasions/never, former); and smoking status (never, previous, current).

Statistical analysis

Prior research (Kyung-Sook et al., 2018), and based on statistical interactions we found in preliminary analyses, indicated that gender modifies the relationship between the main exposures of interest and suicidality. Consequently, we carried out the analyses stratified by gender. Cox proportional hazards regression was used to investigate deaths by suicide. The proportional hazards assumption was tested for using Schoenfeld residuals. For hospital admissions due to self-harm, however, the proportional hazards assumption was not met for loneliness, so data were reanalysed using a Royston Parmar model (Royston and Lambert, 2011), with Akaike information criterion from preliminary analyses indicating that two knots should be used to model the baseline hazard, and single time varying parameter, for loneliness.

Six different models are presented for both deaths by suicide and hospital admissions for self-harm. The first three models are presented for each of living arrangements, loneliness, and emotional support separately. Models 1 are univariable regression models only including each of the main independent variables. Models 2 adjust for all sociodemographic variables, and Models 3 additionally adjust for the health variables. Model 4 adds loneliness and Model 5 adds emotional support to Model 3. In Model 6, all variables were included.

We accounted for missing data using multiple imputation by chained equations, generating twenty imputed data sets. Imputation models were stratified by gender and included age, living arrangements, loneliness, emotional support, all variables used in the models, the Nelson-Aalen estimate of cumulative hazard, survival status (Cleves et al., 2016), and additional variables to improve model fit, including household income, participation in social groups, contact with friends and family, parental depression, limiting longstanding illness and self-rated health. These variables were not included in the main models because they either had comparatively high rates of missing data which limited their utility in preliminary complete case analysis, or, in the case of health variables, might mediate the relationship between our exposures of interest and outcomes. Our models were fitted to each imputed data set and combined in accordance with Rubin's rules. All analyses were carried out using Stata 16.0.

RESULTS

Sociodemographic characteristics of study participants by gender are shown in table 1. With respect to the main independent variables of interest, men were more likely to cohabit, whereas women were more likely to live alone or with non-partner(s). Women were somewhat more likely to report often feeling lonely, and men were much more likely to report that they never had any emotional support. Men were much more likely to have died by suicide (n=181) than women (n=85). Among potential confounders, men were more likely than woman to be in both the most and least advantaged categories of the socioeconomic measures, and women generally had poorer health.

Death by suicide

The results of Cox proportional hazard models for deaths by suicide are shown in table 2. For men there were initially strong relationships, relative to cohabitation, between living alone or with a non-partner and death by suicide (model 1). These associations were attenuated after adjusting for sociodemographic factors (sex, age, ethnicity, employment status, area deprivation education, and ever had a same sex relationship (model 2) and health measures (physical morbidities, BMI, ever seen GP for depression, psychotropic medication, alcohol consumption and smoking status) (model 3). Adjusting for loneliness (model 4) or emotional support (model 5) only led to a slight attenuation of associations, and in the final fully adjusted model (model 6) both living alone (Hazard Ratio (HR) 2.02, 95% CI 1.40 to 293) and living with a person who was not a partner (HR 1.72, 95% CI 1.03 to 2.88) were associated with death by suicide.

The relationship between loneliness and low levels of emotional support and death by suicide, although relatively strong in unadjusted models (model 1), fell after adjustment for sociodemographic factors and health. Once living arrangements were accounted for, loneliness (model 4), and lower levels of emotional support (model 5) were only modestly associated with death by suicide. In contrast, for women, there was little evidence of any association between death by suicide and living arrangements, loneliness or emotional support.

Finally, in fully adjusted models, we conducted interaction tests to assess whether the relationship between living arrangements and death by suicide was modified by loneliness or emotional support. For men, a Wald test indicated a significant interaction (p=0.002) between living arrangements and loneliness, presented in Figure 1a. Men who often experienced loneliness or those who were not lonely and living alone, or with a non-partner only, had three times the risk of dying by suicide compared to those who cohabit and are not lonely.

Hospital admissions for self-harm

The results for the Royston Parmar models for associations between hospital admissions for self-harm and living arrangements, loneliness and perceived emotional support are shown in table 3.

For men, all of living arrangements, loneliness and perceived emotional support were associated with hospital admissions for self-harm in unadjusted analyses. The strength of these associations was reduced after adjusting for sociodemographic characteristics (model 2) and health measures (model 3). Both loneliness (model 4) and lower levels of emotional support (model 5) explained part of the relationship between living alone and self-harm. In the final model, there was no evidence of an association between living alone and self-harm. In contrast, both lower levels of emotional support and loneliness were associated with increased risk of hospitalization for self-harm.

Women differed from men in that the living arrangements categories had weaker relationships with self-harm within the unadjusted model (model 1). Furthermore, the associations for women were explained by health (model 3). For women, loneliness (model 4) and lower levels of perceived emotional support (model 5) were associated with increased risk of self-harm, independent of other factors (with the caveat that the relationship between lack of emotional supports and self-harm could be explained by loneliness).

For men, adjusting for all confounders, we found a significant (p=0.023) interaction indicating that loneliness modified the relationship between living arrangements and hospitalization for self-harm. Overall, loneliness removed any protective effects of cohabitation over living alone, such that men who were often lonely had similarly increased HRs of hospitalization of around 2, irrespective of their living arrangements. In contrast, among men who did not report loneliness, living alone was associated with a modest increase in risk of hospitalization for self-harm (HR 1.32, 95% CI 0.95 to

1.82) and a greater increase was found for those living only with non-partners (HR 1.88, 95% CI 1.24 to 2.83) (see Figure 1b).

DISCUSSION

Our goal was to investigate the association between living arrangements, loneliness, perceived emotional support and subsequent risk of suicidal behaviours, within a large general population cohort in middle-age.

For men, given that both living alone and living with a non-partner were both associated with an increased risk of death by suicide, it is possible that having a partner is protective against death by suicide. Subjective loneliness and perceived emotional support had modest relationships with death by suicide and these variables explained little of the relationship between living alone and death by suicide. For women, none of living arrangements, loneliness, nor emotional support were associated with death by suicide.

For both genders, loneliness and emotional support were associated with increased risk of hospitalization for self-harm. However, associations between living arrangements and self-harm were more limited, being explained by health for women, and health, loneliness and emotional support for men.

Our study has some important novel findings. While loneliness has been linked to suicidal ideation and attempts (Stickley and Koyanagi, 2016), including within case-control studies (Sinclair et al., 2005), a recent systematic review (Solmi et al., 2020) suggests that our study is the first longitudinal study of its kind to investigate the relationship between loneliness and deaths by suicide. The absence of prior research is not a surprise given the methodological challenges of studying such rare outcomes as suicide (Van Orden et al., 2010).

Our finding that deaths by suicide and hospital admissions for self-harm for men were associated with living arrangements, is consistent with the literature for marital status that finds that married or cohabiting people have lower risks of suicide compared to single people (Conejero et al., 2016; Frisch and Simonsen, 2013). Our study adds to this literature by showing that men appear to actively benefit from living with a partner (not just from not living alone). Demographic factors, such as the older age of the UK Biobank sample, could explain why we did not find any associations between living arrangements and suicide and self-harm for women. Kyung-Sook et al. (2018) found associations between marital status and suicide only among younger women.

An original contribution of our study with respect to living arrangements is that most studies of suicidal behaviour focus on the concept that it is *living alone* that is harmful (Turecki and Brent, 2016). However, our results indicate that both men who lived alone *and with non-partners* were at increased risk of dying by suicide and self-harm. Apart from Frisch and Simonsen's (2013) study, which indicated that people living in households with more than nine people had increased risk of suicide, the focus has generally been on living alone as a risk factor, rather than other relationships. However, this is to some extent consistent with a wider literature which has found that those living alone and those living with people other than a partner are more likely to have anxiety or depressive disorders (Joutsenniemi et al., 2006).

Another original contribution with respect to living arrangements is that our results are the first to indicate that the associations between living alone and self-harm might be explained by loneliness and emotional support. However, this does not appear to be the case for deaths by suicide. Such

findings, particularly if replicated elsewhere, may have implications for theories that try to explain the psychological and social antecedents of suicidal behaviour.

Implications

As noted above, the rarity of death by suicide as an outcome has made it very difficult to investigate some theorised risks for death by suicide, as well as other relatively infrequent outcomes such as hospitalization for self-harm (Klonsky et al., 2016; Van Orden et al., 2010). Many of the theories proposed to explain the development of suicidal behaviour are based on studies investigating risk factors for other forms of suicidal behaviour, such as self-reported suicidal ideation or attempts. Clearly, the assumption that the risk factors for all types of suicidal thoughts and behaviours will be the same is problematic (DeJong et al., 2010; Klonsky et al., 2016). From a public health perspective, it is important to identify potentially causal and modifiable factors that may differ between death by suicide and other suicidal behaviours.

This study clearly shows that, for men, living arrangements, loneliness and emotional support are important risk factors for both death by suicide and hospitalization for self-harm. For women, loneliness and lack of emotional support are important risk factors for hospitalization for self-harm. Our results are less certain with respect to the role that living arrangements, loneliness and emotional support might play in deaths by suicide for women. Given the large differences in rates of death by suicide between men and women (Scourfield and Evans, 2015) and systematic review findings that the relationship between marital status and suicide is moderated by gender (Kyung-Sook et al., 2018), *a priori* we decided to analyse the results separately by gender. The large gender difference in death by suicide rates indicates that either the effects of risk factors or their prevalence differ by gender. From our stratified analyses, we find much more modest associations (around half the strength of equivalent associations for men) between living arrangements and loneliness and death by suicide in unadjusted models for women, and these associations are largely explained by socioeconomic factors, health and mental health at baseline. The confidence intervals for the sample do not completely rule out the possibility of there being associations between death by suicide and living arrangements, loneliness and emotional support for women within the general population. To carry out improved analyses for women would require much larger samples, such as national-level UK census linked to hospital and mortality records. Even then, only some aspects of our analyses could be replicated as these administrative data lack many important variables, particularly loneliness.

Our findings on the relationship between death by suicide and living arrangements, loneliness and emotional support for men provide evidence that these are not simply distal risk factors: the associations persist after adjusting for measures of physical and mental health. With respect to death by suicide, the protective effects of living with a partner are particularly important. Cohabiting with a partner appears to be protective against death by suicide even after adjusting for socioeconomic and demographic factors, physical health, mental health, loneliness and emotional support. This could be the case for a number of reasons but operationalizing loneliness or emotional support using single item measures is unlikely to be the explanation. The loneliness and emotional support measures we used were strongly associated with self-harm. An alternative methodological explanation is that men who die by suicide may be less willing to seek treatment for poor mental health and that the risk of suicide is greater among those who never see a GP, compared to those who see a GP once a year (Windfuhr et al., 2016), hence there may be residual confounding due to poor health. However, that would not in itself explain why men who are not living with a partner have increased risk of poor mental health, that in turn leads to death by suicide.

There are also theoretical reasons that could explain why having a partner is protective against death by suicide. One possibility is that the benefits of having a partner may be linked to men's sense of masculinity and self-image (Scourfield et al., 2012), rather than emotional support or companionship. Given the absence of any protective effects of non-partners, and the removal of the protective effects of having a partner among people who are lonely, some of the risk might be due to the concept of "perceived burdensomeness" from the Interpersonal Theory of Suicide (Van Orden et al., 2010). Components of perceived burdensomeness, which include self-hatred and feeling so flawed that one becomes a liability to others, could be theorized as being active drivers for death by suicide. Perceived burdensomeness could arise in situations in which living arrangements suggested a dysfunctional relationship (such as being lonely while cohabiting) or living in situations in which men are unable to fulfil traditional male roles that require having a partner (Scourfield and Evans, 2015). It is possible that perceived burdensomeness may not just be a driver of suicidal behaviour in general, but also a driver towards more lethal self-harm behaviours.

For hospitalization for self-harm, in contrast to death by suicide, loneliness and emotional support appear to be the more important than living arrangements. Associations exist after adjusting for sociodemographic and mental and physical health factors, and for men loneliness and emotional wellbeing explain most of the relationship between living alone and hospitalization for self-harm. This is consistent with the idea that loneliness and lack of emotional support are mechanisms through which living alone, or at least without a partner, could increase risk of suicide. This is also consistent with the concept of 'thwarted belongingness' from the Interpersonal Theory of Suicide, which suggests that loneliness alongside the absence of reciprocal caring relationships can lead to self-destructive behaviours (Stanley et al., 2016; Van Orden et al., 2010). However, given that loneliness appears to remove any protective effects of cohabitation for men, it also raise questions about the extent to which a lonely person can also feel that they do or do not have a reciprocal caring relationship.

Our findings are consistent with some differences in the risks for different types of suicidal behaviour (DeJong et al., 2010; Klonsky et al., 2016). One possibility is that while thwarted belongingness and perceived burdensomeness both drive suicidal behaviour, the latter more strongly drives individuals towards more lethal methods. However, our results may also be consistent with other theories for suicidality. An alternative is that loneliness could be considered a form of emotional dysregulation. Emotional dysregulation theory proposes that while emotional dysregulation is an important factor in self-harm, some aspects of emotional dysregulation may be protective against something that is as daunting and as fearful as lethal self-harm (Stanley et al., 2016).

Strengths and limitations
The key strength of this study is that UK Biobank had a baseline sample of more than 500,000 people. This very large sample provided an opportunity to study death by suicide and hospitalization for self-harm, which are both rare outcomes. However, UK Biobank data do have some potential limitations. The recruitment of such a large sample is only justifiable if it collects data on a broad range of topics, many of which are necessarily operationalised using single item questions. The use of single item measures for loneliness and perceived emotional support might be considered a weakness of our study, but single-item measures are considered valid (Stickley and Koyanagi, 2016) and have been recommended for the study of loneliness (HM Government, 2018). Another limitation is that UK Biobank had an invitation response rate of only 5.5%, and, compared to the general population UK Biobank is less economically deprived with some evidence of a healthy volunteer selection bias (Fry et al., 2017).

Our study uses observational data with the inherent limitations for inferring causality. We have adjusted for a broad range of potential confounding variables at baseline and we had follow-up data on deaths by suicide and hospitalisation for self-harm. However, our baseline measures were only recorded at a single time point and the causal direction is impossible to determine conclusively. It is likely that the relationship between the social connection measures and baseline health is bidirectional. However, given that the peak age of onset for major depressive disorder is early adulthood (Myrna M. Weissman et al., 2016), we decided to focus on a somewhat conservative approach, prioritising relationships presented in models that have adjusted for both mental health and sociodemographic measures. We were also limited in our analyses by only having self-harm hospital admission data for England (these data were not available for Wales or Scotland). It is also the case that many who self-harm do not seek help from services or, when they do, are not admitted but are reviewed as out-patients (Gunnell et al., 2005).

Finally, the target sample of this population was those aged 40 to 70 living in the United Kingdom, and the sample contained only a handful of people outside this age range. These results may not be generalizable to other age groups or to other cultures where attitudes to suicide, or other societal level risk factors for suicide, such as the availability of fire arms, may be very different.

CONCLUSIONS

This study raises several questions for future exploration. Our results suggest that addressing loneliness in the general population may reduce the risk of self-harm but, for death by suicide, there is a much more complex (and likely sex-specific) relationship between loneliness, living arrangements and perceived emotional support. Overall, this work demonstrates that for men (but not for women) living alone or with a non-partner increases the risk of suicide, a finding not explained by perceived loneliness. It appears likely that loneliness may be more important as a risk factor for self-harm than for suicide. These findings may reflect differences in the theoretical pathways for death by suicide and self-harm.

AUTHOR STATEMENTS

Contributors

RJS, DJS, BC and AJ developed the research question. NG, JW, BC, RJS, AJ, DML, RP assisted with deriving and coding the variables. RJS carried out the analysis. BC, DML and DM provided advice on statistical methodology. AJ and CO carried out the literature review. RJS, DJS, AJ and CO drafted the manuscript. All authors edited and agreed the final manuscript.

Funding sources

This project was supported by two MRC Mental Health Data Pathfinder Awards [grant numbers MC_PC_17217), MC_PC_17211].

Acknowledgements

We thank all participants in the UK Biobank study. The project was completed using UK Biobank project code 6553 (PI: Smith). UK Biobank was established by the Wellcome Trust, Medical Research Council, Department of Health, Scottish Government and Northwest Regional Development Agency. UK Biobank has also had funding from the Welsh Assembly Government and the British Heart Foundation. Data collection was funded by UK Biobank.

REFERENCES

Amato, P.R., 2014. Marriage, cohabitation and mental health. Fam. Matters 96, 5-13.

Bennardi, M., Caballero, F.F., Miret, M., Ayuso-Mateos, J.L., Haro, J.M., Lara, E., Arensman, E., Cabello, M., 2019. Longitudinal Relationships Between Positive Affect, Loneliness, and Suicide Ideation: Age-Specific Factors in a General Population. Suicide Life. Threat. Behav. 49, 90-103.

Beutel, M.E., Klein, E.M., Brähler, E., Reiner, I., Jünger, C., Michal, M., Wiltink, J., Wild, P.S., Münzel, T., Lackner, K.J., Tibubos, A.N., 2017. Loneliness in the general population: prevalence, determinants and relations to mental health. BMC Psychiatry 17, 97.

Cacioppo, S., Capitanio, J.P., Cacioppo, J.T., 2014. Toward a neurology of loneliness. Psychol. Bull. 140, 1464-1504.

Cleves, M., Gould, W.W., Marchenko, Y.V., 2016. An Introduction to Survival Analysis Using Stata: Revised Third Edition. Stata Press, College Station, Texas.

Conejero, I., Lopez-Castroman, J., Giner, L., Baca-Garcia, E., 2016. Sociodemographic Antecedent Validators of Suicidal Behavior: A Review of Recent Literature. Curr. Psychiat. Rep. 18, 94.

Courtin, E., Knapp, M., 2017. Social isolation, loneliness and health in old age: a scoping review. Health Soc. Care Comm. 25, 799-812.

DeJong, T.M., Overholser, J.C., Stockmeier, C.A., 2010. Apples to oranges?: A direct comparison between suicide attempters and suicide completers. J. Affect. Disord. 124, 90-97.

Dreyer, K., Steventon, A., Fisher, R., Deeny, S.R., 2018. The association between living alone and health care utilisation in older adults: a retrospective cohort study of electronic health records from a London general practice. BMC Geriatr. 18, 269.

Elovainio, M., Hakulinen, C., Pulkki-Råback, L., Virtanen, M., Josefsson, K., Jokela, M., Vahtera, J., Kivimäki, M., 2017. Contribution of risk factors to excess mortality in isolated and lonely individuals: an analysis of data from the UK Biobank cohort study. Lancet Public Health 2, e260-e266.

Frisch, M., Simonsen, J., 2013. Marriage, cohabitation and mortality in Denmark: national cohort study of 6.5 million persons followed for up to three decades (1982–2011). Int. J. Epidemiol. 42, 559-578.

Fry, A., Littlejohns, T.J., Sudlow, C., Doherty, N., Adamska, L., Sprosen, T., Collins, R., Allen, N.E., 2017. Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. Am. J. Epidemiol. 186, 1026-1034.

Gunnell, D., Appleby, L., Arensman, E., Hawton, K., John, A., Kapur, N., Khan, M., O'Connor, R.C., Pirkis, J., COVID-19 Suicide Prevention Research Collaboration, Suicide risk and prevention during the COVID-19 pandemic, 2020. Lancet Psychiatry. published online April 21. https://doi.org/10.1016/S2215-0366(20)30171-1.

Gunnell, D., Bennewith, O., Peters, T.J., House, A., Hawton, K., 2005. The epidemiology and management of self-harm amongst adults in England. J. Public Health 27, 67-73.

Hakulinen, C., Pulkki-Råback, L., Virtanen, M., Jokela, M., Kivimäki, M., Elovainio, M., 2018. Social isolation and loneliness as risk factors for myocardial infarction, stroke and mortality: UK Biobank cohort study of 479 054 men and women. Heart 104, 1536-1542.

Hawkley, L.C., Cacioppo, J.T., 2010. Loneliness Matters: A Theoretical and Empirical Review of Consequences and Mechanisms. Ann. Behav. Med. 40, 218-227.

HM Government, 2018. A connected society A strategy for tackling loneliness - laying the foundations for change. Department for Digital, Culture, Media and Sport, London.

Holmes, E.A., O'Connor, R.C., Perry, V.H., Tracey, I., Wessely, S., Arseneault, L., Ballard, C., Christensen, H., Cohen Silver, R., Everall, I., Ford, T., John, A., Kabir, T., King, K., Madan, I., Michie, S., Przybylski, A.K., Shafran, R., Sweeney, A., Worthman, C.M., Yardley, L., Cowan, K., Cope, C., Hotopf, M., Bullmore, E., 2020. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. Lancet Psychiatry. published online April 15. https://doi.org/10.1016/S2215-0366(20)30168-1.

Holt-Lunstad, J., Smith, T.B., Baker, M., Harris, T., Stephenson, D., 2015. Loneliness and Social Isolation as Risk Factors for Mortality:A Meta-Analytic Review. Perspect Psychol Sci 10, 227-237.

Joutsenniemi, K., Martelin, T., Martikainen, P., Pirkola, S., Koskinen, S., 2006. Living arrangements and mental health in Finland. J. Epidemiol. Community Health 60, 468-475.

Klonsky, E.D., May, A.M., Saffer, B.Y., 2016. Suicide, Suicide Attempts, and Suicidal Ideation. Annu. Rev. Clin. Psychol. 12, 307-330.

Kyung-Sook, W., SangSoo, S., Sangjin, S., Young-Jeon, S., 2018. Marital status integration and suicide: A meta-analysis and meta-regression. Soc. Sci. Med. 197, 116-126.

McCormack, A., Edmondson-Jones, M., Fortnum, H., Dawes, P., Middleton, H., Munro, K.J., Moore, D.R., 2014. The prevalence of tinnitus and the relationship with neuroticism in a middle-aged UK population. J. Psychosom. Res. 76, 56-60.

Myrna M. Weissman, Priya Wickramaratne, Marc J. Gameroff, Virginia Warner, Daniel Pilowsky, Rajni Gathibandhe Kohad, Helena Verdeli, Jamie Skipper, Ardesheer Talati, 2016. Offspring of Depressed Parents: 30 Years Later. Am. J. Psychiatry 173, 1024-1032.

National Collaborating Centre for Mental Health, 2011. Self-harm: longer-term management. NICE clinical guidelines 133, London: National Institute for Clinical Excellence, 2011.

Nock, M.K., Borges, G., Bromet, E.J., Cha, C.B., Kessler, R.C., Lee, S., 2008. Suicide and Suicidal Behavior. Epidemiol. Rev. 30, 133-154.

Office for National Statistics (ONS), 2016. Suicides in the UK: 2014 registrations. Office for National Statistics Newport.

Perlman, D., Peplau, L.A., 1981. Toward a social psychology of loneliness. Pers Relationships 3, 31-56.

Rico-Uribe, L.A., Caballero, F.F., Martín-María, N., Cabello, M., Ayuso-Mateos, J.L., Miret, M., 2018. Association of loneliness with all-cause mortality: A meta-analysis. PLOS ONE 13, e0190033.

Royston, P., Lambert, P.C., 2011. Flexible Paramatric Survial Analysis Using Stata: Beyond the Cox Model Stata Press, College Station, Texas.

Scourfield, J., Evans, R., 2015. Why Might Men Be More at Risk of Suicide After a Relationship Breakdown? Sociological Insights. Am. J. Mens Health 9, 380-384.

Scourfield, J., Fincham, B., Langer, S., Shiner, M., 2012. Sociological autopsy: An integrated approach to the study of suicide in men. Soc. Sci. Med. 74, 466-473.

Sinclair, J.M.A., Harriss, L., Baldwin, D.S., King, E.A., 2005. Suicide in depressive disorders: A retrospective case-control study of 127 suicides. J. Affect. Disord. 87, 107-113.

Smith, K.J., Victor, C., 2019. Typologies of loneliness, living alone and social isolation and their associations with physical and mental health Ageing Soc. 39, 1709-1730.

Solmi, M., Veronese, N., Galvano, D., Favaro, A., Ostinelli, E.G., Noventa, V., Favaretto, E., Tudor, F., Finessi, M., Shin, J.I., Smith, L., Koyanagi, A., Cester, A., Bolzetta, F., Cotroneo, A., Maggi, S., Demurtas, J., De Leo, D., Trabucchi, M., 2020. Factors Associated With Loneliness: An Umbrella Review Of Observational Studies. J. Affect. Disord. 271, 131-138.

Stanley, I.H., Hom, M.A., Rogers, M.L., Hagan, C.R., Joiner, T.E., 2016. Understanding suicide among older adults: a review of psychological and sociological theories of suicide. Aging Ment. Health 20, 113-122.

Stickley, A., Koyanagi, A., 2016. Loneliness, common mental disorders and suicidal behavior: Findings from a general population survey. J. Affect. Disord. 197, 81-87.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman, T., Collins, R., 2015. UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. PLOS Med. 12, e1001779.

Turecki, G., Brent, D.A., 2016. Suicide and suicidal behaviour. Lancet 387, 1227-1239.

van Hedel, K., Martikainen, P., Moustgaard, H., Myrskylä, M., 2018. Cohabitation and mental health: Is psychotropic medication use more common in cohabitation than marriage? SSM Popul. Health 4, 244-253.

Van Orden, K.A., Witte, T.K., Cukrowicz, K.C., Braithwaite, S.R., Selby, E.A., Joiner Jr, T.E., 2010. The interpersonal theory of suicide. Psychol. Rev. 117, 575-600.

Windfuhr, K., While, D., Kapur, N., Ashcroft, D.M., Kontopantelis, E., Carr, M.J., Shaw, J., Appleby, L., Webb, R.T., 2016. Suicide risk linked with clinical consultation frequency, psychiatric diagnoses and psychotropic medication prescribing in a national study of primary-care patients. Psychol. Med. 46, 3407-3417.

Table 1: Characteristics for Men and Women in UK Biobank

	Women n	Women %	Men n	Men %
Social Connection				
Living arrange'				
Alone	53,747	19.7	39,188	17.1
Cohabitation	188,588	69.0	174,503	76.2
Non Partner	28,787	10.5	13,330	5.8
Missing	2,290	0.8	2,113	0.9
loneliness				
No	210,917	77.2	190,330	83.1
Yes	57,045	20.9	34,371	15.0
Missing	5,440	2.0	4,433	1.9
Emotional Support				
Almost Daily	140,692	51.5	117,881	51.5
2-4 times a week	30,321	11.1	16,462	7.2
Once a Week	33,579	12.3	19,751	8.6
Monthly	30,276	11.1	22,965	10.0
Never	29,284	10.7	42,449	18.5
Missing	9,250	3.4	9,626	4.2
Socio Demographic				
Age				
35/44	27,835	10.2	23,968	10.5
45/54	80,110	29.3	62,305	27.2
55/64	116,860	42.7	95,446	41.7
65/74	48,597	17.8	47,415	20.7
Ethnicity				
White British	240,265	87.9	202,345	88.3
Other	31,871	11.7	25,278	11.0
Missing	1,266	0.5	1,511	0.7
Employment status				
Employed	149,212	54.6	137,958	60.2
Retired	95,565	35.0	71,420	31.2
Other	27,105	9.9	18,323	8.0
Missing	1,520	0.6	1,433	0.6
Deprivation				
Least deprived	54,571	20.0	46,093	20.1
4th Quintile	54,568	20.0	45,537	19.9
3rd Quintile	55,333	20.2	45,057	19.7
2nd Quintile	55,355	20.3	45,020	19.7
Most deprived	53,248	19.5	47,131	20.6
Missing	327	0.1	296	0.1
Education				
Degree	84,541	30.9	76,635	33.4
Professional	40,982	15.0	31,452	13.7
NVQ, HND or HNC	28,304	10.4	34,682	15.1
A level	15,758	5.8	11,477	5.0
O level	41,639	15.2	23,045	10.1
CSE	10,863	4.0	7,745	3.4
None	45,968	16.8	39,307	17.2
Missing	5,347	2.0	4,791	2.1
Ever same sex Relationship				
None	235,296	86.1	197,726	86.3
At least one	6,923	2.5	8,894	3.9
Missing	31,184	11.4	22,514	9.8

	Women n	Women %	Men n	Men %
Centre Country				
England	243,756	89.2	205,057	89.5
Wales	9,701	3.6	8,177	3.6
Scotland	19,945	7.3	15,900	6.9
Health				
Multi morbidity[1]				
Zero	99,438	36.4	83,795	36.6
One	90,116	33.0	76,252	33.3
Two	49,995	18.3	42,382	18.5
Three or more	33,853	12.4	26,705	11.7
missing	na[1]		Na	
Ever Depressed				
No	159,662	58.4	168,388	73.5
Yes	110,878	40.6	58,590	25.6
Missing	2,862	1.1	2,156	0.9
Psychotropic Meds				
No	235,483	86.1	209,628	91.5
Yes	28,474	10.4	14,217	6.2
Missing	9,445	3.5	5,289	2.3
Alcohol usage				
Daily / almost daily	43,867	16.0	57,907	25.3
3 to 4 times week	55,899	20.5	59,546	26.0
1 -2 times a week	105,686	38.7	79,469	34.7
Special event or never	57,137	20.9	23,263	10.2
Former	9,983	3.7	8,125	3.6
Missing	830	0.3	824	0.4
Smoking				
Never	162,064	59.3	111,473	48.7
Previous	85,458	31.3	87,612	38.2
Current	24,367	8.9	28,612	12.5
Missing	1,513	0.6	1,437	0.6
Obesity				
Normal & Under	107,822	39.4	57,340	25.0
Overweight	99,859	36.5	112,249	49.0
Obese	64,262	23.5	57,899	25.3
Missing	1,459	0.5	1,646	0.7
Suicidality				
Hospital admission for Self-harm[2]				
Events	589		471	
Person yrs follow up	1,458,045		1,220,987	
Died by suicide				
Events	85		181	
Person yrs follow up	2,444,031		2,027,824	

1. Data are only available for those reporting conditions,

2 Analyses only include those who attended a Biobank centre in England at baseline.

Table 2: Death by suicide hazard ratios for loneliness, living arrangements and perceived social support, in unadjusted and adjusted models for men and women in the UK Biobank study

	Model 1 HR 95% (CI)	Model 2 HR 95% (CI)	Model 3 HR 95% (CI)	Model 4 HR 95% (CI)	Model 5 HR 95% (CI)	Model 6 HR 95% (CI)
Men						
Living arrangements (Cohabitating)						
Alone	3.57 (2.61 to 4.87)***	2.72 (1.93 to 3.82)***	2.35 (1.66 to 3.32)***	2.16 (1.51 to 3.09)***	2.14 (1.50 to 3.09)***	2.02 (1.40 to 2.93)***
Non partner	2.91 (1.80 to 4.73)***	2.11 (1.28 to 3.49)***	1.94 (1.17 to 3.20)**	1.80 (1.08 to 3.00)*	1.81 (1.09 to 3.01)*	1.72 (1.03 to 2.88)*
Loneliness (No)						
Yes	3.20 (2.35 to 4.36)***	2.45 (1.77 to 3.40)***	1.71 (1.22 to 2.41)**	1.43 (1.01 to 2.03)*		1.39 (0.97 to 1.99)+
Emotional support (daily)						
2-4 times a week	1.89 (1.11 to 3.21)*	1.69 (0.99 to 2.88)+	1.51 (0.88 to 2.57)		1.29 (0.75 to 2.21)	1.25 (0.73 to 2.15)
Once a week	2.50 (1.59 to 3.92)***	2.21 (1.40 to 3.49)**	2.00 (1.26 to 3.15)**		1.62 (1.02 to 2.60)*	1.55 (0.97 to 2.49)+
Monthly	1.60 (0.98 to 2.64)+	1.45 (0.88 to 2.40)	1.30 (0.78 to 2.14)		1.08 (0.65 to 1.80)	1.02 (0.61 to 1.71)
Never	1.86 (1.26 to 2.73)***	1.60 (1.08 to 2.36)*	1.51 (1.02 to 2.22)*		1.25 (0.83 to 1.87)	1.17 (0.77 to 1.76)
Women						
Living arrangements (Cohabitating)						
Alone	1.58 (0.96 to 2.61)+	1.51 (0.89 to 2.54)	1.21 (0.71 to 2.06)	1.24 (0.72 to 2.12)	1.23 (0.72 to 2.11)	1.25 (0.72 to 2.15)
Non partner	1.45 (0.76 to 2.79)	1.24 (0.63 to 2.43)	1.01 (0.51 to 1.99)	1.03 (0.52 to 2.04)	1.02 (0.52 to 2.02)	1.03 (0.52 to 2.05)
Loneliness (No)						
Yes	1.49 (0.91 to 2.43)	1.42 (0.86 to 2.34)	0.92 (0.55 to 1.54)	0.89 (0.53 to 1.50)		0.92 (0.54 to 1.57)
Emotional support (daily)						
2-4 times a week	1.33 (0.70 to 2..54)	1.25 (0.66 to 2.38)	1.08 (0.57 to 2.05)		1.04 (0.54 to 2.00)	1.05 (0.55 to 2.02)
Once a week	1.21 (0.64 to 2.29)	1.18 (0.62 to 2.24)	0.97 (0.51 to 1.84)		0.93 (0.49 to 1.79)	0.95 (0.49 to 1.82)
Monthly	0.78 (0.35 to 1.74)	0.78 (0.35 to 1.75)	0.66 (0.30 to 1.48)		0.65 (0.29 to 1.45)	0.66 (0.29 to 1.49)
Never	1.13 (0.57 to 2.25)	1.17 (0.58 to 2.34)	1.00 (0.50 to 2.02)		0.98 (0.49 to 1.98)	1.00 (0.49 to 2.06)

*** p ≤ 0.001, ** p ≤0.01, * p ≤0.05, + p ≤0.1.
Model 1 univariable cox regression, model 2 adjusts for socio-demographic factors (sex, age, ethnicity, employment status, deprivation and highest qualification, ever had same sex relationship), model 3 is model 2 plus adjusting for health measures (number physical morbidities, ever depressed, on psychotropic medication, BMI, alcohol consumption, smoking). Model 4 is model 3 plus mutual adjustment for loneliness and living arrangements. Model 5 is model 3 plus mutual adjustment for living arrangements and perceived social support. Model 6 is model 3 plus mutual adjustment for living arrangements, loneliness and perceived social support.

Table 3: Admissions to hospital for self-harm hazard ratios for loneliness, living arrangements and perceived social support, in unadjusted and adjusted models for men and women in the UK Biobank study

	Model 1 HR 95% (CI)	Model 2 HR 95% (CI)	Model 3 HR 95% (CI)	Model 4 HR 95% (CI)	Model 5 HR 95% (CI)	Model 6 HR 95% (CI)
Men						
Living arrangements (Cohabitating)						
Alone	3.14 (2.57 to 3.84)***	1.67 (1.34 to 2.09)***	1.37 (1.09 to 1.72)**	1.17 (0.93 to 1.48)	1.17 (0.93 to 1.48)	1.06 (0.83 to 1.34)
Non partner	3.56 (2.70 to 4.70)***	1.77 (1.32 to 2.38)***	1.64 (1.23 to 2.20)***	1.44 (1.07 to 1.94)*	1.44 (1.10 to 1.95)*	1.33 (0.98 to 1.79)+
Loneliness (No)						
Yes	5.32 (4.43 to 6.40)***	3.13 (2.57 to 3.82)***	2.01 (1.63 to 2.47)***	1.91 (1.55 to 2.37)***		1.74 (1.40 to 2.16)***
Per log day X Loneliness	0.97 (0.93 to 1.01)+	0.97 (0.93 to 1.01)+	0.97 (0.93 to 1.01)+	0.97 (0.93 to 1.01)+		0.97 (0.93 to 1.01)+
Emotional support (daily)						
2-4 times a week	1.95 (1.37 to 2.77)***	1.68 (1.18 to 2.40)**	1.47 (1.03 to 2.10)*		1.41 (0.99 to 2.02)+	1.34 (0.94 to 1.92)+
Once a week	2.46 (1.82 to 3.32)***	2.01 (1.48 to 2.72)***	1.79 (1.32 to 2.42)***		1.69 (1.24 to 2.32)***	1.56 (1.14 to 2.14)**
Monthly	2.13 (1.58 to 2.88)***	1.84 (1.34 to 2.49)***	1.60 (1.18 to 2.17)**		1.53 (1.13 to 2.09)**	1.39 (1.02 to 1.90)*
Never	2.98 (2.38 to 3.74)***	2.16 (1.72 to 2.72)***	1.95 (1.54 to 2.46)***		1.86 (1.46 to 2.36)***	1.65 (1.29 to 2.10)***
Women						
Living arrangements (Cohabitating)						
Alone	1.75 (1.44 to 2.11)***	1.67 (1.37 to 2.04)***	1.18 (0.96 to 1.44)	1.04 (0.85 to 1.28)	1.15 (0.93 to 1.41)	1.04 (0.85 to 1.28)
Non partner	2.00(1.60 to 2.50)***	1.21 (0.96 to 1.53)	0.94 (0.74 to 1.19)	0.85 (0.67 to 1.08)	0.92 (0.73 to 1.17)	0.85 (0.67 to 1.08)
Loneliness (No)						
Yes	4.35 (3.60 to 5.13)***	3.25 (2.74 to 3.85)***	1.92 (1.61 to 2.29)***	1.93 (1.61 to 2.31)***		1.89 (1.57 to 2.28)***
Per log day X Loneliness	0.95 (0.93 to 0.98)***	0.95 (0.93 to 0.98)***	0.95 (0.93 to 0.98)***	0.95 (0.93 to 0.98)***		0.95 (0.93 to 0.98)***
Emotional support (daily)						
2-4 times a week	1.21 (0.92 to 1.60)	1.18 (0.90 to 1.56)	0.99 (0.75 to 1.31)		0.98 (0.74 to 1.30)	0.91 (0.69 to 1.21)
Once a week	1.52 (1.19 to 1.94)***	1.50 (1.17 to 1.92)***	1.20 (0.94 to 1.54)		1.18 (0.92 to 1.52)	1.05 (0.82 to 1.35)
Monthly	1.26 (0.98 to 1.66)+	1.27 (0.97 to 1.67)+	1.03 (0.78 to 1.35)		1.02 (0.78 to 1.34)	0.88 (0.67 to 1.16)
Never	2.20 (1.76 to 2.76)***	1.94 (1.54 to 2.43)***	1.50 (1.19 to 1.89)***		1.49 (1.18 to 1.88)***	1.24 (0.98 to 1.58)+

*** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$, + $p \leq 0.1$.
Model 1 Univariable Royston Parmar parmetric Hazard models, model 2 adjusts for socio-demographic factors (sex, age, ethnicity, employment status, deprivation and highest qualification, ever had same sex relationship). Model 3 is model 2 plus adjusting for health measures (number physical morbidities, ever depressed, on psychotropic medication, BMI, alcohol consumption, smoking). Model 4 is model 3 plus mutual adjustment for loneliness and living arrangements. Model 5 is model 3 plus mutual adjustment for living arrangements and perceived social support. Model 6 is model 3 plus mutual adjustment for living arrangements, loneliness and perceived social support.

Figure 1: The interactions between living arrangements and loneliness in the prediction of deaths by suicide and hospital admissions for self-harm for men in the UK Biobank Study.