DNA methylation biomarker selected by an ensemble machine learning approach

predicts mortality risk in an HIV-positive veteran population

Chang Shu[1,2], Amy C. Justice[2,3], Xinyu Zhang[1,2], Vincent C. Marconi[4], Dana B.

Hancock[5], Eric O. Johnson[5,6], Ke Xu[1,2]

1. Department of Psychiatry, Yale School of Medicine, New Haven, CT, 06516

2. Connecticut Veteran Healthcare System, West Haven, CT, 06515

3. Department of Internal Medicine, Yale School of Medicine, New Haven, CT, 06516

4. Division of Infectious Disease, Emory University School of Medicine, Atlanta GA

5. GenOmics, Bioinformatics, and Translational Research Center, Biostatistics and

Epidemiology Division, RTI International, Research Triangle Park, NC, USA

6. Fellow Program, RTI International, Research Triangle Park, NC, USA.

All corresponding address to

Ke Xu, M.D., Ph.D.

Associate Professor of Psychiatry

Yale School of Medicine

Email: ke.xu@yale.edu

Tel: 203-932-5711x7430

Word count: 3645

Abstract: 249

Figures: 6

Tables: 3

Abstract

Background: With the improved life expectancy of people living with HIV (PLWH), identifying vulnerable subpopulations at high risk of mortality is important for clinical care. Evidences showed that DNA methylation (DNAm) is associated with aging and mortality in non-HIV populations. Here, we aimed to establish a panel of DNAm biomarkers that can predict mortality risk among PLWH.

Methods: 1,081 HIV-positive participants from the Veterans Aging Cohort Study (VACS) were divided into training (N=460), validation (N=114), and testing (N=507) sets. VACS index was used as a measure of mortality risk among PLWH. Model training and fine-tuning were conducted using the ensemble method in the training and validation sets and prediction performance was assessed in the testing set. The survival analysis comparing the predicted high and low mortality risk groups was conducted. The Gene Ontology enrichment analysis of the predictive CpG sites was performed.

Results: We selected a panel of 393 CpGs for the ensemble prediction model. The prediction model showed excellent performance in predicting high mortality risk with an auROC of 0.809 (95%CI: 0.767-0.851) and a balanced accuracy of 0.653 (95%CI: 0.611, 0.693) in the testing set. The predicted high mortality risk group was significantly associated with 10-year mortality (hazard ratio=1.79, p=4E-05) compared with low mortality risk group. These 393 CpGs were located in 280 genes enriched in immune and inflammation responses pathways.

Conclusions: We identified a panel of DNAm features associated with mortality risk in PLWH. These DNAm features may serve as predictive biomarkers for mortality risk among PLWH.

48   **Keywords:** DNA methylation, HIV, mortality risk, machine learning prediction

Introduction

Combination antiretroviral therapy has significantly improved the life expectancy of people living with HIV (PLWH), but there is still a gap in life expectancy between PLWH and the general population(1-4). It is important to identify vulnerable groups with a high risk of mortality among PLWH and to deliver early interventions and clinical care for those patients. Previous studies have demonstrated that the Veterans Aging Cohort Study (VACS) index is significantly associated with mortality and is considered a measure of mortality risk among PLWH (5-7). The VACS index is a composite score summing HIV progression measures and general organ injury indicators of the kidneys and liver, which may be able to capture the early stage of elevated risk for mortality.

A large body of evidence has demonstrated that epigenetic modification is influenced by internal and external environmental changes and is associated with the early stages of pathophysiological processes (8-11). DNA methylation (DNAm), one of the most widely studied epigenetic marks, is strongly correlated with aging (12-14), substance use (e.g., cigarette smoking and alcohol consumption) (15-21), and a variety of diseases (8-11, 22, 23). Since DNAm is relatively stable and easy to detect in body fluids obtained through noninvasive procedures, DNAm marks have emerged as robust biomarkers for disease diagnosis (24), disease subtype classification (25, 26) and treatment response monitoring (27, 28).

Since DNAm biomarkers are objectively measured and can reflect pathological processes of disease progression, DNAm can be used to identify individual vulnerability and mortality risk among PLWH. In some cases, DNAm alterations can occur before clinical diagnosis. For example, a longitudinal study of DNAm showed that most DNA

methylome changes occurred 80-90 days before clinically detectable glucose elevation

(29). As another example, mitochondrial epigenetic changes can indicate early-stage

prediabetes (30). Although the clinical diagnosis of myocardial infarction has been well

established, DNA methylation in the blood has utility in the diagnosis and monitoring of

cardiac pathologies and in the study of normal human cardiac physiology and

development (31). These studies support the utility of DNAm features as biomarkers of

risk for future onset of disease. Here, we apply this approach to identify individuals with

a high risk of mortality in an HIV-positive population.

DNAm plays an important role in HIV infection and disease progression. We

previously reported the association of two CpG sites in the promoter region of *NLRC5*

with HIV infection (32). *NLRC5* is a major transcriptional activator of the MHC class I

gene. DNAm has also been linked to HIV comorbid diseases, such as diabetes and

kidney function (33, 34). Furthermore, aging is significantly associated with thousands

of CpGs in the epigenome, and the epigenetic clock and DNAm age are becoming

widely recognized (12-14). DNAm marks are predictive of mortality in non-HIV

populations (35-40). Therefore, we hypothesized that DNAm is associated with mortality

risk among PLWH and that DNAm signatures in the blood can serve as biomarkers to

predict mortality among HIV-positive individuals.

Machine learning methods have been widely applied to select DNAm features

that are informative for the clinical diagnosis and classification of complex diseases (20,

41, 42). Ensemble machine learning methods can aggregate multiple machine learning

models (base models) and usually provide better prediction outcomes than single base

models (43, 44). An ensemble approach has been shown to perform well in

personalized medicine and disease outcomes, such as in cancer and diabetes (45, 46).

In this study, by applying an ensemble machine learning approach, we aimed to

identify DNAm features that can serve as biomarkers of mortality risk among HIV-

positive individuals. Here, the VACS index was used as a measure of mortality risk in an

HIV-positive population (5-7, 47). These predictive DNAm biomarkers can potentially be

used for informing future clinical care and providing new insights into the epigenetic

mechanism of mortality risk among HIV-positive patients.

Methods

An overview of our analytical approach is shown in the flowchart in **Figure 1**. Our

prediction model was built with training and validation sets profiled on the 450K array

and then independently evaluated with the testing set profiled on the EPIC array. Briefly,

we first applied an ensemble learning approach to build a machine learning model to

predict high or low mortality risk, and we then examined the association of the selected

CpG features with mortality by a survival analysis. Then, we conducted a Gene

Ontology enrichment analysis for the selected DNAm features. Last, we conducted a

meta-analysis of the epigenome-wide association (EWA) on the VACS index of the

entire sample.

Study population

All participants in sample sets 1 and 2 (**Figure 1**) were from the VACS that is a

prospective cohort study of veterans focusing on the clinical outcomes of HIV infection

(5). DNA samples were extracted from the peripheral blood of 1,081 HIV-positive men

from the VACS. Participants in sample set 1 were randomly partitioned into a training

set (80%, N=460) and a validation set (20%, N=114), and sample set 2 was used as the

independent testing set (N=507). Table 1 shows the demographic and clinical

information on patient age, sex, race, smoking status, CD4 count, viral load, HIV

medication adherence, VACS index, and mortality in the training, validation and testing

sets. The training and validation sets included slightly older individuals and more African

Americans than the testing set. The VACS index was slightly lower in the testing set

than in the training and validation sets. There were no significant differences in sex,

smoking, HIV medication adherence, CD4 count, $\log_{10}$ HIV-1 viral load or 10-year

mortality across the three sample sets.

Mortality risk

The VACS index is a well-established score for mortality risk among PLWH (5-7).

The VACS index is scored by summing preassigned points for age, CD4 count, HIV-1

RNA, hemoglobin, platelets, aspartate and alanine transaminase (AST and ALT),

creatine, estimated glomerular filtration rate (eGFR), and viral hepatitis C infection (7).

High mortality risk among PLWH was defined as a VACS index score > 40, which was

based on a prior observation that the predicted 3-year mortality of 10% for this group

was significantly higher than that for the group with VACS index scores ≤ 40(48).

Prediction models were developed by machine learning methods to predict high

mortality risk (VACS index > 40) and low mortality risk (VACS index ≤ 40) groups

among PLWH.

Genome-wide DNAm profiling and quality control

DNA samples in sample set 1 were profiled by Infinium Human Methylation 450K

BeadChip (HM450K, Illumina Inc., CA, USA), and DNA samples in sample set 2 were

profiled by the Infinium Human Methylation EPIC BeadChip (Illumina Inc., CA, USA)

(**Figure 1**). DNAm for the training and validation sets were evaluated using the same

quality control (QC) protocol (49) in the R package *minfi* (50). In detail, CpG sites on sex

chromosomes and within 10 base pairs of a single nucleotide polymorphism were

removed. The detection p-value threshold was set at 10^{-12} for both sample sets 1 and 2.

After QC, 408,583 CpG sites common between the HM450K and EPIC arrays were

used for analysis to ensure the same coverage between the two sets. DNA methylation

among the common CpG sites was highly correlated between the HM450K and EPIC

arrays (r=0.986). Proportions of 6 cell types (CD4+ T cells, CD8+ T cells, natural killer T

cells, B cells, monocytes and granulocytes) were estimated for all participants in sample

sets 1 and 2 using an established method (51).

Feature selection of CpG sites in the training set

We first preselected a panel of CpG sites associated with high mortality risk

among PLWH based on the EWA of the VACS index score in the training set. CpG sites

with p<0.001 were preselected to build the prediction models. A liberal cutoff of p<0.001

was arbitrarily set to ensure a sufficient number of predictive DNAm features to build the

prediction models. The variable importance (a score between 0-100) of each

preselected CpG site was ranked by elastic-net regularized generalized linear models

(GLMNET) by the R package *caret* (52) based on 100 bootstraps, where each bootstrap

included 70% of all samples. CpG sites with zero variable importance for 80% of the

bootstraps were considered to be nonpredictive features and were removed from further

model development. The remaining CpG sites were ranked based on the median

importance ranking among 100 bootstraps and were divided into 20 groups. Each CpG

group was used to build machine learning models.

Developing machine learning prediction models for mortality risk among PLWH

1) Model development in the training set: We developed an ensemble method

that aggregated the prediction results from four base machine learning models: random

forest (RF), GLMNET, support vector machines (SVM) and k-nearest neighbors (k-NN)

(53-56) using the model choice of "rf", "glmnet", "svmLinear" and "knn" in the R package

caret (52). These four base models have been commonly used in predicting binary

outcomes and have expanded the diversity of algorithms (53-56). Ten-fold cross-

validation was used in the model training process to minimize overfitting. These four

base models were independently trained to predict mortality risk among PLWH in the

training set and then aggregated by the ensemble method using the R package

caretEnsemble (ver. 2.0.1) (57). The prediction performance of each ensemble model

was evaluated by using area under the receiver operating characteristic curve (auROC)

and the area under the precision-recall curve (auPRC).

2) Final CpG group selection in the validation set: The CpG group with the

highest auROC in the validation set was selected as the final feature group for the

ensemble model.

3) Independent evaluation in the testing set: Using the ensemble model and the

final feature group, we predicted the high mortality risk group and evaluated prediction

performance in the testing set by using auROC and balanced accuracy. Balanced

accuracy was defined as the average accuracy obtained on each class, as shown in the

following formula (58). Balanced accuracy was used in this study to avoid biased

accuracy due to imbalanced samples (58). The 95% confidence interval of balanced

accuracy was estimated by 1,000 stratified bootstraps of the testing set.

$$Balanced\ accuracy$$

$$= \frac{1}{2}\left(\frac{True\ positive}{True\ positive + False\ negative} + \frac{True\ negative}{True\ negative + False\ positive}\right)$$

Survival analysis

By using the final ensemble model, we classified each individual in the entire

sample as having as high or low mortality risk. Kaplan-Meier survival curves presented

10-year survival probability by high or low mortality risk group. Survival analysis was

conducted using a Cox proportional hazards model on 10-year mortality comparing the

high and low mortality risk groups. We used age as time scale *t*, and our model adjusted

for sex, race, smoking, self-reported HIV medication adherence, $\log_{10}$ of HIV viral load

and CD4 count.

$$h(t) = h_0(t)\exp\left(\beta_1 predicted\ mortality\ risk\ +\ \beta_2\ sex\ +\ \beta_3\ race\ +\ \beta_4\ smoking\right.$$

$$\left. +\ \beta_5\ HIV\ medication\ adherence\ +\ \beta_6\ \log_{10}(viral\ load) + \beta_7\ CD4\ count\right)$$

Biological interpretation of the predictive panel of CpG sites on mortality risk
among PLWH

204   We performed Gene Ontology (GO) enrichment analysis using *missMethyl* to

205   adjust for bias by different numbers of CpG sites per gene (59, 60). Genes that harbor

206   or are located near at least one predictive CpG site were used for GO analysis.

Epigenome-wide association analysis on mortality risk among PLWH in all

samples

Since DNA methylation of two sample sets was measured by two different

platforms, we performed EWA on high and low mortality risk groups separately in

sample sets 1 and 2, and we then conducted a meta-analysis to detect epigenome-wide

signals in the entire sample. In each EWA, we used a two-step linear model approach

as previously described (49). The EWA model adjusted for confounding factors,

including age, sex, race, smoking, cell type proportions and control principle

components. EWA meta-analysis of sample sets 1 and 2 was conducted using METAL

(61). The weights of effect size were the inverse of the corresponding standard errors

for the meta-analysis (61). CpG sites with Bonferroni corrected p-value < 0.05 were

considered statistically significant.

Results

Feature selection and ensemble model training for mortality risk among PLWH

High mortality risk among PLWH was defined as VACS index > 40 based on

previous literature showing a predicted 3-year mortality of 10% for this group (48).

Prediction models were developed by machine learning methods to predict high

mortality risk (VACS index > 40) and low mortality risk groups (VACS index ≤ 40)

among PLWH.

A panel of 856 CpGs with p<0.001 was preselected based on EWA in the training

set. We ranked these candidate predictors by median GLMNET importance ranking

among 100 bootstraps using 70% of the training sample. We excluded 178 CpG sites

that had a variable importance score of zero among 80% of the bootstraps. A final panel

of 678 CpG sites were selected and formed into 20 groups based on importance

ranking to determine the best performing CpG group for the ensemble model (**Figure 2**).

In the training set, we used 4 common machine learning classification models

(RF, GLMNET, SVM, and k-NN) as our base models for the ensemble method (53-56)

and trained them independently for the 20 groups of CpGs. The performance metrics of

GLMNET, RF and SVM were mostly comparable in terms of auROC and auPRC, and

they plateaued to 1 with an increasing number of CpGs in the training set (**Figure 3**,

Figure 4). The performance of k-NN was poorer than the other 3 methods, but its

auROC and auPRC remained above 0.9 in the training set. An ensemble model

combining the prediction results of all 4 base models was used (**Figure 3**, **Figure 4**).

In the validation set, the performances of 4 base models varied. Three models,

GLMNET, RF, and SVM, showed good performance with both auROC and auPRC > 0.8,

but the performance of k-NN was poor with auROC < 0.8 (**Figure 3**, **Figure 4**). The

ensemble model showed the best performance at 393 CpG sites with an excellent

performance of auROC (0.829). The accuracy of this prediction model was 0.807, and

the balanced accuracy that accounted for class imbalances between participants at high

and low risk of mortality was 0.782. Thus, the ensemble model with 393 DNAm features

was used as the final prediction model (**Table S1**).

In the testing set, the ensemble model with 393 DNAm features showed excellent

performance with auROC of 0.809 (95%CI: 0.767-0.851), prediction accuracy of 0.761

and balanced accuracy of 0.653 (95%CI: 0.611, 0.693) (**Figure 5**), suggesting that our

ensemble model with a panel of 393 features was able to differentiate between high and

low mortality risk in an HIV-positive population.

Survival analysis

By using our ensemble model that predicts individual mortality risk, we found that

participants predicted to have a high risk of mortality showed significantly lower 10-year

survival probability than participants predicted to have a low risk of mortality (**Figure 6**).

Using a Cox proportional-hazards model adjusting for baseline age, sex, race, viral load,

CD4 count and antiviral medication adherence, participants who were predicted to have

a high risk of mortality remained to have an elevated risk of mortality compared with

those predicted to have a low risk of mortality, with a hazard ratio (HR) of 1.79 (95%CI:

1.35-2.37, p=4E-05).

Biological interpretations of predictive CpGs by Gene Ontology enrichment

analysis

The 393 predictive CpGs were located in or near 280 genes. The top 8 enriched

pathways based on these 280 genes included response to virus (p= 4.26E-05), defense

response (p=1.29E-04), cytokine receptor binding (p=1.48E-04) and regulation of

response to interferon-gamma (p= 4.15E-04) (**Table 2**). Our findings suggested that the

selected 393 CpG sites are biologically relevant to HIV pathogenesis and progression.

Epigenome-wide association on mortality risk among PLWH

A meta-analysis of EWA of sample sets 1 and 2 identified 208 epigenome-wide

significant CpGs after Bonferroni correction (**Figure S2**). These significant CpG sites

were located in 112 genes and included genes previously reported as being associated

with HIV pathogenesis. For example, cg07839457 in *NLRC5* was previously reported to

be associated with HIV infection (32). Interestingly, 30 of 208 CpG sites were also in the

panel of machine learning predictive CpGs (**Table 3**). Twenty out of the 30 overlapping

CpG sites were negatively associated with mortality risk, while 10 CpG sites were

positively associated with mortality risk. Some of the overlapping CpGs were located in

viral response genes, such as *IFITM1* and *PARP9*.

Discussion

In this study, we presented evidence that DNAm marks in blood were predictive

of mortality risk in an HIV-positive population. We identified a panel of 393 CpG sites

that were highly predictive for high vs. low risk of mortality among PWLH. We also found

that our predicted mortality risk based on the ensemble model was strongly associated

with mortality in HIV-positive individuals in the VACS cohort. In addition, the selected

393 DNAm features were located in genes enriched in HIV pathogenesis and

progression. Thus, we identified a panel of 393 DNAm biomarkers that may enhance

the understanding of the epigenetic mechanisms of mortality risk among PLWH.

We demonstrated that a machine learning approach can predict mortality risk

among HIV-positive individuals across different methylation data sets. One of the

challenges of a machine learning method is overfitting. We attempted to limit overfitting

in the development of the ensemble prediction models in several ways: 1) model

development and final model and feature selection were conducted separately in the

training and validation sets, 2) 10-fold cross-validation was performed during the

training, 3) the ensemble model was used to aggregate prediction results from multiple

base models instead of arbitrarily choosing a specific machine learning prediction model,

and 4) the final model was evaluated in an independent test set. We further observed

that the performance of different models generated by the individual machine learning

method varied. The ensemble-based modeling method outperformed some base

models and could aggregate prediction results from all base models. Furthermore, our

prediction model was built on training and validation sets profiled on the 450K array,

and it was independently evaluated in the testing set that was profiled on the EPIC array.

Of note, removing batch effects is an important step in EWA analysis. Methods have

been applied to address the bias related to batches (62). Here, our goal was to select a

set of generalizable methylation markers that are relatively stable regardless of batch,

cohort, and other confounding factors. Without removing batch effects within the set of

CpG sites common to both arrays, our model still showed good predictive performance,

indicating that our model is generalizable, regardless of methylation platform. Our

results suggested that the ensemble prediction model is relatively stable and robust.

Our results showed that the CpG-based ensemble prediction model is strongly

associated with mortality in an HIV-positive population. This finding is consistent with

previous literature showing that DNAm marks in blood can predict mortality in non-HIV

populations (40). We found that 30 out of 393 CpGs reached epigenome-wide

significance. The majority of the 393 CpG sites are located within or near genes that are

involved in known HIV pathology and progression. For example, cg22930808 on

PARP9 and cg07107453 on *IFI44* were selected by the machine learning prediction

model and reached epigenome-wide significance; both genes are involved in HIV

pathogenesis. Both cg22930808 and cg07107453 are less methylated in the high

mortality risk group than in the low mortality risk group. *IFI44* is an interferon-alfa

inducible protein and is associated with infection by several viruses. A previous study

showed that higher expression of *IFI44* facilitated HIV-1 latency (63), which may

increase mortality risk. In addition, cg12359279 is located in the *MX1* gene (Interferon-

Induced GTP-Binding Protein Mx1). *MX1* encodes a GTP-metabolizing protein that is

induced by interferon I and II and is involved in interferon gamma signaling and the Toll-

like signaling pathway.

Studies have shown that comorbidity and aging are associated with HIV-related

excess mortality (64-66), and some immune biomarkers can partially explain HIV-

related excess mortality risk (67). Since the VACS index included measures of immune

function and indicators for general organ injury, it is plausible that predictive CpG sites

(based on the VACS index) are located near genes involved in immune system

development or other functions, such as liver or kidney functions. Among the 393

predictive CpG sites, no CpG site located on the genes overlapped with previously

reported liver or kidney diseases. However, two of the 393 CpG sites, cg16249932 and

cg00463367, are located near genes related to immune system development (*MAEA*

and *GATA3*).

The biological relevance of these 393 CpG sites was further supported by the

Gene Ontology enrichment analysis. The top enriched pathways, such as response to

virus and cytokine receptor binding, may indicate important biological pathways that

lead to increased risk of mortality among PLWH.

We acknowledge that there are several limitations in this study. We defined high

mortality risk by a cutoff of 40 for the VACS index based on previous literature, with a

predicted 3-year mortality of 10% for individuals with a VACS index score >40 (48); the

cutoff for defining high mortality risk may vary in other populations. Additionally, the

generalizability of our prediction model may be limited because our samples were

predominantly middle-aged men. All samples in our study were HIV-positive, and we

cannot infer whether the identified CpGs are relevant to predicting mortality in an HIV-

negative population. Last, we are unable to validate the array-based DNA methylation of

the selected CpG sites by machine learning or to identify significant CpG sites by EWAS

through a different platform. However, in a different sample, we found a high correlation

of PBMC methylome between EPIC and the methyl-capture sequencing platform

(r=0.986) on the same 4 peripheral blood monocyte cells. Future studies in other

populations are warranted to validate our ensemble prediction model and the selected

methylation features.

Conclusions

We identified a panel of 393 predictive DNAm features in blood that was predictive of

mortality risk among PLWH. These DNAm features may serve as biomarkers to identify

individuals at high risk for mortality and may help to prioritize genes to better understand

the mechanisms of mortality risk in an HIV-positive population. These DNAm features

have the potential to be used for monitoring HIV progression in future clinical care.

Figures

Figure 1: Flowchart of analytical procedures for selecting CpG sites in the peripheral

blood methylome, machine learning prediction models to predict high and low mortality

risk groups, survival analysis, Gene Ontology enrichment analysis, and epigenome-wide

association analysis.

Figure 2: Variable importance ranking of predictive machine learning CpG sites.

Variable importance is a score between 0 and 100, as calculated by elastic-net

regularized generalized linear models (GLMNET). We obtained variable importance

scores from 100 bootstraps. The top 20 ranked CpG sites and 20 bootstraps are shown.

Figure 3: Area under the receiver operating characteristic (auROC) curve in the training and validation sets.

Figure 4: Area under the precision-recall curve (auPRC) in the training and validation sets**.**

Figure 5: Receiver operating characteristic (auROC) curve in the testing set.

384
385

386 **Figure 6**: Kaplan-Meier curves of predicted high and low mortality risk groups among

387 people living with HIV.

388
389

 Tables

Table 1: Study sample characteristics

	Training set (N=460)	Validation set (N=114)	Testing set (N=507)	p value[*]
Age (year, mean ± sd)	52.56 (7.54)	51.21 (8.09)	50.86 (7.67)	0.002
Female (%)	6 (1.3)	1 (0.9)	11 (2.2)	0.452
Race (%)				
Caucasian	45 (9.8)	9 (7.9)	46 (9.1)	
African Americans	392 (85.2)	103 (90.4)	409 (80.7)	0.003
Other	23 (5.0)	2 (1.8)	52 (10.3)	
Smokers (%)	360 (59.4)	309 (58.4)	294 (58.0)	0.719
HIV treatment adherence (%)	362 (78.7)	85 (74.6)	382 (75.3)	0.399
CD4 count	432.68 (291.43)	411.82 (281.55)	450.61 (280.05)	0.287
log 10 HIV-1 viral load	2.76 (1.22)	2.68 (1.17)	2.83 (1.22)	0.399
VACS index (mean ± sd)	30.62 (20.35)	35.50 (21.83)	35.46 (22.35)	0.001
10-year mortality (%)	123 (26.7)	32 (28.1)	121 (23.9)	0.477

[*]ANOVA test is used for continuous variables, chi-square test is used for categorical variables

Table 2: Gene ontology term enrichment analysis of the selected 393 CpG sites that predict mortality risk among HIV-positive population

Term	Total genes	Predictive Genes	P value
tumor necrosis factor receptor superfamily binding	46	7	8.33E-06
response to virus	303	15	4.26E-05
defense response	1505	39	1.29E-04
mitochondrial DNA metabolic process	18	4	1.43E-04
cytokine receptor binding	267	12	1.48E-04
regulation of response to interferon-gamma	25	4	4.15E-04
regulation of interferon-gamma-mediated signaling pathway	25	4	4.15E-04
cell-cell adhesion mediator activity	50	6	4.43E-04
intrinsic component of the cytoplasmic side of the plasma membrane	7	3	4.76E-04
immune response	1896	45	4.97E-04

391
392

Table 3: Overlapping CpG sites between machine learning selected CpG sites and epigenome-wide significant CpG sites on mortality risk among people living with HIV

probe	Chr	Position	Nearest gene	Variable Importance	Meta Effect (SE)	Meta P	Refgene group	Relation to CpG island
cg01971407	11	313624	*IFITM1*	9.9	-0.0399 (0.0039)	8.05E-25	TSS1500	N_Shelf
cg22930808	3	122281881	*PARP9;DTX3L*	13.7	-0.1057 (0.0105)	7.24E-24	5UTR;TSS1500	N_Shore
cg23570810	11	315102	*IFITM1*	13.0	-0.0638 (0.0065)	1.15E-22	Body	N_Shore
cg14864167	8	66751182	*PDE7A*	12.0	-0.0832 (0.0085)	1.54E-22	Body	N_Shelf
cg01190666	20	62204908	*PRIC285*	19.4	-0.0343 (0.0035)	1.79E-22	5UTR	N_Shore
cg11702942	8	144102584	*LY6E;LY6E*	7.8	-0.0382 (0.004)	7.52E-22	Body;Body	S_Shore
cg03607951	1	79085586	*IFI44L*	23.0	-0.0683 (0.0072)	3.58E-21	TSS1500	
cg03848588	9	32525008	*DDX58*	14.4	-0.0274 (0.0029)	6.02E-21	Body	N_Shore
cg04582010	11	313120	*IFITM1*	22.1	-0.0455 (0.0052)	2.19E-18	TSS1500	S_Shore
cg18394552	5	159428643		24.4	0.0342 (0.0043)	1.57E-15		
cg03753191	13	43566902	*EPSTI1;EPSTI1*	4.1	-0.018 (0.0023)	2.43E-15	TSS1500	S_Shore
cg17267239	1	173640200	*ANKRD45*	21.8	-0.0186 (0.0025)	1.22E-13	TSS1500	S_Shore
cg12461141	11	5710654	*TRIM22*	37.6	-0.0274 (0.0037)	1.37E-13	TSS1500	
cg09251764	17	6659070	*XAF1;XAF1*	13.8	-0.0142 (0.002)	3.93E-13	TSS200	
cg05626226	4	106515450	*FLJ20184*	11.5	0.0279 (0.004)	2.45E-12	Body	
cg22959742	10	13913931	*FRMD4A*	16.4	0.0289 (0.0041)	3.08E-12	Body	
cg16936953	17	57915665	*TMEM49*	11.3	-0.0379 (0.0056)	1.95E-11	Body	
cg18181703	17	76354621	*SOCS3*	32.6	-0.0215 (0.0033)	1.15E-10	Body	N_Shore
cg07107453	1	79114976	*IFI44*	4.8	-0.0292 (0.0046)	2.26E-10	TSS1500	
cg25114611	6	35696870	*FKBP5;LOC285847*	12.8	-0.0125 (0.002)	4.78E-10	TSS1500;Body	S_Shore
cg01059398	3	172235808	*TNFSF10*	26.7	-0.0289 (0.0047)	7.85E-10	Body	
cg19459791	15	65363022		15.1	0.0159 (0.0026)	1.20E-09		S_Shelf
cg26282236	12	1025755	*RAD52*	30.2	0.0243 (0.0041)	3.63E-09	Body	

cg06357748	12	1025529	*RAD52*	10.7	0.0277 (0.0047)	4.12E-09	Body	
cg04442417	20	62191507	*PRIC285;PRIC285*	26.6	0.0201 (0.0035)	1.22E-08	Body	Island
cg14602222	12	1025663	*RAD52*	13.2	0.0241 (0.0042)	1.42E-08	Body	
cg26724018	11	5716255	*TRIM22*	11.9	-0.0171 (0.003)	1.49E-08	5UTR	
cg03084350	3	38065265	*PLCD1;PLCD1;PLCD1*	18.0	0.0114 (0.002)	2.05E-08	Body	N_Shore
cg00569896	4	204382		0.2	0.0254 (0.0047)	5.78E-08		N_Shore
cg12126344	1	12207564		8.4	-0.011 (0.002)	7.36E-08		

Legends

Figure 1: Flowchart of analytical procedures for selecting CpG sites in the peripheral blood methylome, machine learning prediction models to predict high and low mortality risk groups, survival analysis, Gene Ontology enrichment analysis, and epigenome-wide association analysis.

Figure 2: Variable importance ranking of predictive machine learning CpG sites. Variable importance is a score between 0 and 100, as calculated by elastic-net regularized generalized linear models (GLMNET). We obtained variable importance scores from 100 bootstraps. The top 20 ranked CpG sites and 20 bootstraps are shown.

Figure 3: Area under the receiver operating characteristic (auROC) curve in the training and validation sets.

Figure 4: Area under the precision-recall curve (auPRC) in the training and validation sets**.**

Figure 5: Receiver operating characteristic (auROC) curve in the testing set.

Figure 6: Kaplan-Meier curves of predicted high and low mortality risk groups among people living with HIV.

Figure S1: Distribution of Veteran Aging Cohort Study (VACS) index between sample sets 1 and 2.

Figure S2: Epigenome-wide association (EWA) on mortality risk among PLWH: quantile-quantile plot, Manhattan plot and overlapping CpG sites between EWA significant CpGs and predictive CpGs.

Table 1: Study sample characteristics

Table 2: Gene ontology term enrichment analysis of the selected 393 CpG sites that

predict mortality risk among people living with HIV

Table 3: Overlapping CpG sites between machine learning selected CpG sites and

epigenome-wide significant CpG sites on mortality risk among people living with HIV

Table S1: Predictive CpG sites by variance importance

Table S2: Epigenome-wide significant CpG sites in meta-analysis of EWAS on mortality

risk among people living with HIV

List of abbreviations

AUC: Area Under Curve

CI: Confidence interval

DMR: differentially methylated region

DNA: Deoxyribonucleic acid

DNAm: DNA methylation

DAVID: Database for Annotation, Visualization and Integrated Discovery

EWA: epigenome-wide association

FDR: False discovery rate

FWER: Family-wise error rate

GLMNET: elastic-net regularized generalized linear models

GO: Gene ontology

HIV: Human immunodeficiency virus

HM450K: Human Methylation 450K BeadChip

k-NN: k-nearest neighbors

NK: Natural killer

PC: Principal component

PLWH: people living with HIV

QC: Quality control

SVM: Support Vector Machines

VACS: Veterans Aging Cohort Study

XGBoost: Extreme Gradient Boosting Tree

Declarations

Ethics approval and consent to participate

The study was approved by the Committee of the Human Research Subject Protection

at Yale University and the Institutional Research Board Committee of the Connecticut

Veteran Healthcare System. All subjects provided written consent.

Availability of data and materials

Demographic and clinical variables and DNAm data for the VACS samples were

submitted to GEO dataset (GSE117861) and are available to the public. All codes for

analysis are also available upon request to the corresponding author.

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by the National Institute on Drug Abuse (R03DA039745,

R01DA038632, R01DA047063, R01DA047820).

Authors' contributions

CS was responsible for data analysis and manuscript preparation. ACJ provided DNA

samples and clinical data and contributed to manuscript preparation. XZ was

responsible for the bioinformatics data processing. VM was involved in clinical data

collection and manuscript preparation. DH and EJ contributed to the analytical approach

and to manuscript preparation. KX was responsible for the study design, study protocol,

sample preparation, data analysis, interpretation of findings, and manuscript preparation.

Acknowledgments

References

1.   Wandeler G, Johnson LF, Egger M. Trends in life expectancy of HIV-positive adults on
antiretroviral therapy across the globe: comparisons with general population. Curr Opin HIV
AIDS. 2016;11(5):492-500.
2.   Teeraananchai S, Kerr S, Amin J, Ruxrungtham K, Law M. Life expectancy of HIV-positive
people after starting combination antiretroviral therapy: a meta-analysis. HIV Medicine.
2017;18(4):256-66.
3.   Marcus JL, Chao CR, Leyden WA, Xu L, Quesenberry CP, Jr., Klein DB, et al. Narrowing the
Gap in Life Expectancy Between HIV-Infected and HIV-Uninfected Individuals With Access to
Care. Journal of acquired immune deficiency syndromes (1999). 2016;73(1):39-46.
4.   Gueler A, Moser A, Calmy A, Günthard HF, Bernasconi E, Furrer H, et al. Life expectancy
in HIV-positive persons in Switzerland: matched comparison with general population. AIDS
(London, England). 2017;31(3):427.
5.   Justice AC, Dombrowski E, Conigliaro J, Fultz SL, Gibson D, Madenwald T, et al. Veterans
aging cohort study (VACS): overview and description. 2006;44(8 Suppl 2):S13.
6.   Justice AC, Modur S, Tate JP, Althoff KN, Jacobson LP, Gebo K, et al. Predictive accuracy
of the Veterans Aging Cohort Study (VACS) index for mortality with HIV infection: a north
American cross cohort analysis. Journal of acquired immune deficiency syndromes (1999).
2013;62(2):149.
7.   Tate JP, Justice AC, Hughes MD, Bonnet F, Reiss P, Mocroft A, et al. An internationally
generalizable risk index for mortality after one year of antiretroviral therapy. AIDS (London,
England). 2013;27(4):563-72.
8.   Lam K, Pan K, Linnekamp JF, Medema JP, Kandimalla R. DNA methylation based
biomarkers in colorectal cancer: a systematic review. Biochimica et Biophysica Acta (BBA)-
Reviews on Cancer. 2016;1866(1):106-20.
9.   Teroganova N, Girshkin L, Suter CM, Green MJ. DNA methylation in peripheral tissue of
schizophrenia and bipolar disorder: a systematic review. BMC genetics. 2016;17(1):27.
10.   Bakusic J, Schaufeli W, Claes S, Godderis L. Stress, burnout and depression: A systematic
review on DNA methylation mechanisms. Journal of Psychosomatic Research. 2017;92:34-44.
11.   Li M, D'Arcy C, Li X, Zhang T, Joober R, Meng X. What do DNA methylation studies tell us
about depression? A systematic review. Translational psychiatry. 2019;9(1):68.
12.   Horvath S. DNA methylation age of human tissues and cell types. Genome biology.
2013;14(10):3156.
13.   Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of
ageing. Nature Reviews Genetics. 2018:1.
14.   Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, et al. Genome-wide
methylation profiles reveal quantitative views of human aging rates. Molecular cell.
2013;49(2):359-67.
15.   Liu C, Marioni RE, Hedman ÅK, Pfeiffer L, Tsai P-C, Reynolds LM, et al. A DNA
methylation biomarker of alcohol consumption. Molecular psychiatry. 2018;23(2):422.
16.   Breitling LP, Yang R, Korn B, Burwinkel B, Brenner H. Tobacco-smoking-related
differential DNA methylation: 27K discovery and replication. The American Journal of Human
Genetics. 2011;88(4):450-7.

17.   Gao X, Zhang Y, Breitling LP, Brenner H. Relationship of tobacco smoking and smoking-
related DNA methylation with epigenetic age acceleration. Oncotarget. 2016;7(30):46878-89.
18.   Joubert BR, Håberg SE, Nilsen RM, Wang X, Vollset SE, Murphy SK, et al. 450K
epigenome-wide scan identifies differential DNA methylation in newborns related to maternal
smoking during pregnancy. Environmental health perspectives. 2012;120(10):1425-31.
19.   Lee KW, Pausova Z. Cigarette smoking and DNA methylation. Frontiers in genetics.
2013;4:132.
20.   Zhang X, Hu Y, Aouizerat BE, Peng G, Marconi VC, Corley MJ, et al. Machine learning
selected smoking-associated DNA methylation signatures that predict HIV prognosis and
mortality. 2018;10(1):155.
21.   Zhang R, Miao Q, Wang C, Zhao R, Li W, Haile CN, et al. Genome-wide DNA methylation
analysis in alcohol dependence. Addiction biology. 2013;18(2):392-403.
22.   Kraiczy J, Nayak KM, Howell KJ, Ross A, Forbester J, Salvestrini C, et al. DNA methylation
defines regional identity of human intestinal epithelial organoids and undergoes dynamic
changes during development. Gut. 2019;68(1):49-61.
23.   Nano J, Ghanbari M, Wang W, de Vries PS, Dhana K, Muka T, et al. Epigenome-Wide
Association Study Identifies Methylation Sites Associated With Liver Enzymes and Hepatic
Steatosis. Gastroenterology. 2017;153(4):1096-106.e2.
24.   Delpu Y, Cordelier P, Cho W, Torrisani J. DNA methylation and cancer diagnosis.
International journal of molecular sciences. 2013;14(7):15029-58.
25.   Figueroa ME, Lugthart S, Li Y, Erpelinck-Verschueren C, Deng X, Christos PJ, et al. DNA
methylation signatures identify biologically distinct subtypes in acute myeloid leukemia. Cancer
cell. 2010;17(1):13-27.
26.   Holm K, Hegardt C, Staaf J, Vallon-Christersson J, Jönsson G, Olsson H, et al. Molecular
subtypes of breast cancer are associated with characteristic DNA methylation patterns. Breast
cancer research. 2010;12(3):R36.
27.   Berdasco M, Esteller M. Clinical epigenetics: seizing opportunities for translation. Nature
Reviews Genetics. 2019;20(2):109-27.
28.   Mohammad HP, Barbash O, Creasy CL. Targeting epigenetic modifications in cancer
therapy: erasing the roadmap to cancer. Nat Med. 2019;25(3):403-18.
29.   Chen R, Xia L, Tu K, Duan M, Kukurba K, Li-Pook-Than J, et al. Longitudinal personal DNA
methylome dynamics in a human with a chronic condition. Nat Med. 2018;24(12):1930-9.
30.   Zheng LD, Linarelli LE, Brooke J, Smith C, Wall SS, Greenawald MH, et al. Mitochondrial
Epigenetic Changes Link to Increased Diabetes Risk and Early-Stage Prediabetes Indicator. Oxid
Med Cell Longev. 2016;2016:5290638.
31.   Zemmour H, Planer D, Magenheim J, Moss J, Neiman D, Gilon D, et al. Non-invasive
detection of human cardiomyocyte death using methylation patterns of circulating DNA. Nat
Commun. 2018;9(1):1443.
32.   Zhang X, Justice AC, Hu Y, Wang Z, Zhao H, Wang G, et al. Epigenome-wide differential
DNA methylation between HIV-infected and uninfected individuals. Epigenetics.
2016;11(10):750-60.
33.   Chen J, Huang Y, Hui Q, Mathur R, Gwinn M, So-Armah K, et al. Epigenetic Associations
with Estimated Glomerular Filtration Rate (eGFR) among Men with HIV Infection. Clin Infect Dis.
2019.

34. Mathur R, Hui Q, Huang Y, Gwinn M, So-Armah K, Freiberg MS, et al. DNA Methylation
Markers of Type 2 Diabetes Mellitus Among Male Veterans With or Without Human
Immunodeficiency Virus Infection. J Infect Dis. 2019;219(12):1959-62.
35. Marioni RE, Shah S, McRae AF, Chen BH, Colicino E, Harris SE, et al. DNA methylation age
of blood predicts all-cause mortality in later life. Genome biology. 2015;16(1):25.
36. Marioni RE, Harris SE, Shah S, McRae AF, von Zglinicki T, Martin-Ruiz C, et al. The
epigenetic clock and telomere length are independently associated with chronological age and
mortality. International journal of epidemiology. 2016;45(2):424-32.
37. Perna L, Zhang Y, Mons U, Holleczek B, Saum K-U, Brenner H. Epigenetic age
acceleration predicts cancer, cardiovascular, and all-cause mortality in a German case cohort.
Clinical epigenetics. 2016;8(1):64.
38. Christiansen L, Lenart A, Tan Q, Vaupel JW, Aviv A, McGue M, et al. DNA methylation
age is associated with mortality in a longitudinal Danish twin study. Aging Cell. 2016;15(1):149-
54.
39. Fransquet PD, Wrigglesworth J, Woods RL, Ernst ME, Ryan J. The epigenetic clock as a
predictor of disease and mortality risk: a systematic review and meta-analysis. Clin Epigenetics.
2019;11(1):62.
40. Zhang Y, Wilson R, Heiss J, Breitling LP, Saum K-U, Schöttker B, et al. DNA methylation
signatures in peripheral blood strongly predict all-cause mortality. Nature communications.
2017;8:14617.
41. Holder LB, Haque MM, Skinner MK. Machine learning for epigenetics and future medical
applications. Epigenetics. 2017;12(7):505-14.
42. Crowgey EL, Marsh AG, Robinson KG, Yeager SK, Akins RE. Epigenetic machine learning:
utilizing DNA methylation patterns to predict spastic cerebral palsy. BMC Bioinformatics.
2018;19(1):225.
43. Dietterich TG, editor Ensemble methods in machine learning. International workshop on
multiple classifier systems; 2000: Springer.
44. Dietterich TG. Ensemble learning. The handbook of brain theory and neural networks.
2002;2:110-25.
45. Moon H, Ahn H, Kodell RL, Baek S, Lin C-J, Chen JJ. Ensemble methods for classification
of patients for personalized medicine with high-dimensional data. Artificial Intelligence in
Medicine. 2007;41(3):197-207.
46. Alghamdi M, Al-Mallah M, Keteyian S, Brawner C, Ehrman J, Sakr S. Predicting diabetes
mellitus using SMOTE and ensemble machine learning approach: The Henry Ford Exercise
Testing (FIT) project. PloS one. 2017;12(7).
47. Bebu I, Tate J, Rimland D, Mesner O, Macalino GE, Ganesan A, et al. The VACS Index
Predicts Mortality in a Young, Healthy HIV Population Starting Highly Active Antiretroviral
Therapy. J Acquir Immune Defic Syndr. 2014;65(2):226-30.
48. Justice AC, Modur SP, Tate JP, Althoff KN, Jacobson LP, Gebo KA, et al. Predictive
accuracy of the Veterans Aging Cohort Study index for mortality with HIV infection: a North
American cross cohort analysis. Journal of acquired immune deficiency syndromes (1999).
2013;62(2):149-63.

49. Lehne B, Drong AW, Loh M, Zhang W, Scott WR, Tan S-T, et al. A coherent approach for
analysis of the Illumina HumanMethylation450 BeadChip improves data quality and
performance in epigenome-wide association studies. 2015;16(1):37.
50. Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al. Minfi:
a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA
methylation microarrays. 2014;30(10):1363-9.
51. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al.
DNA methylation arrays as surrogate measures of cell mixture distribution. BMC Bioinformatics.
2012;13(1):86.
52. Kuhn MJJoss. Building predictive models in R using the caret package. 2008;28(5):1-26.
53. Hastie T, Tibshirani R, Friedman J, Franklin J. The elements of statistical learning: data
mining, inference and prediction. The Mathematical Intelligencer. 2005;27(2):83-5.
54. Kotsiantis SB, Zaharakis I, Pintelas P. Supervised machine learning: A review of
classification techniques. Emerging artificial intelligence applications in computer engineering.
2007;160:3-24.
55. Chen T, Guestrin C, editors. Xgboost: A scalable tree boosting system. Proceedings of the
22nd acm sigkdd international conference on knowledge discovery and data mining; 2016: ACM.
56. Ogutu JO, Piepho H-P, Schulz-Streeck T. A comparison of random forests, boosting and
support vector machines for genomic selection. BMC Proc. 2011;5 Suppl 3(Suppl 3):S11-S.
57. Deane-Mayer ZA, Knowles J. caretEnsemble: Ensembles of caret models. R package
version. 2016;2(0).
58. Brodersen KH, Ong CS, Stephan KE, Buhmann JM, editors. The balanced accuracy and its
posterior distribution. 2010 20th International Conference on Pattern Recognition; 2010: IEEE.
59. Phipson B, Maksimovic J, Oshlack A. missMethyl: an R package for analyzing data from
Illumina's HumanMethylation450 platform. Bioinformatics. 2015;32(2):286-8.
60. Geeleher P, Hartnett L, Egan LJ, Golden A, Raja Ali RA, Seoighe C. Gene-set analysis is
severely biased when applied to genome-wide methylation data. Bioinformatics.
2013;29(15):1851-7.
61. Willer CJ, Li Y, Abecasis GRJB. METAL: fast and efficient meta-analysis of genomewide
association scans. 2010;26(17):2190-1.
62. Chen D-P, Lin Y-C, Fann CSJ. Methods for identifying differentially methylated regions for
sequence- and array-based data. Briefings in Functional Genomics. 2016;15(6):485-90.
63. Power D, Santoso N, Dieringer M, Yu J, Huang H, Simpson S, et al. IFI44 suppresses HIV-1
LTR promoter activity and facilitates its latency. Virology. 2015;481:142-50.
64. Hentzien M, Dramé M, Allavena C, Jacomet C, Valantin M-A, Cabié A, et al. Impact of
age-related comorbidities on five-year overall mortality among elderly HIV-infected patients in
the late HAART era—role of chronic renal disease. The journal of nutrition, health & aging.
2016;20(4):408-14.
65. Langebeek N, Kooij KW, Wit FW, Stolte IG, Sprangers MAG, Reiss P, et al. Impact of
comorbidity and ageing on health-related quality of life in HIV-positive and HIV-negative
individuals. AIDS (London, England). 2017;31(10):1471-81.
66. Hentzien M, Dramé M, Delpierre C, Allavena C, Cabié A, Cuzin L, et al. HIV-related excess
mortality and age-related comorbidities in patients with HIV aged≥ 60: a relative survival
analysis in the French Dat'AIDS cohort. BMJ open. 2019;9(1):e024841.

67.   So-Armah KA, Tate JP, Chang C-CH, Butt AA, Gerschenson M, Gibert CL, et al. Do
Biomarkers of Inflammation, Monocyte Activation, and Altered Coagulation Explain Excess
Mortality Between HIV Infected and Uninfected People? Journal of acquired immune deficiency
syndromes (1999). 2016;72(2):206-13.
654