

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

1 **Genomic Epidemiology of Methicillin-Resistant *Staphylococcus aureus* in Two Cohorts of**  
2 **High-Risk Military Trainees**

3  
4 Robyn S. Lee PhD<sup>1,2\*</sup>, Eugene V. Millar PhD<sup>3,4</sup>, Alanna Callendrello BA<sup>2</sup>, Caroline E. English  
5 BA<sup>3,4</sup>, Emad M. Elassal MS<sup>3,4</sup>, Michael W. Ellis MD<sup>5</sup>, Jason W. Bennett MD MSPH<sup>6,7</sup>,  
6 William P. Hanage PhD<sup>2</sup>

- 7 1. Epidemiology Division, Dalla Lana School of Public Health, University of Toronto, Toronto, Canada  
8 2. The Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H. Chan  
9 School of Public Health, Boston, MA, USA  
10 3. Infectious Disease Clinical Research Program, Department of Preventive Medicine and Biostatistics,  
11 Uniformed Services University of the Health Sciences, Bethesda, MD, USA  
12 4. Henry M. Jackson Foundation for the Advancement of Military Medicine, Bethesda, MD, USA  
13 5. University of Toledo College of Medicine and Life Sciences, Toledo, OH, USA  
14 6. Walter Reed Army Institute of Research, Silver Spring, MD, USA  
15 7. Department of Medicine, Uniformed Services University of the Health Sciences, Bethesda, MD, USA  
16

17 \*Address correspondence to:  
18 Dr. Robyn S. Lee, PhD  
19 Epidemiology Division  
20 Dalla Lana School of Public Health, University of Toronto  
21 Health Sciences Building  
22 155 College Street, Room HS 586  
23 Toronto, Ontario, Canada  
24 M5T 3M7  
25 Email: robyn.s.c.lee@gmail.com

26 Alternative corresponding author:  
27 Dr. William P. Hanage, PhD  
28 Harvard T.H. Chan School of Public Health  
29 677 Huntington Avenue  
30 Kresge Building, 5<sup>th</sup> floor  
31 Boston, MA, USA  
32 02115  
33 whanage@hsph.harvard.edu  
34 617-432-7955

35  
36 Text word count: 3039

37 Figures: 5

38 Tables: 3

39 Key words: Methicillin-resistant *Staphylococcus aureus* ; genomic epidemiology ; military ; community-acquired  
40 MRSA ; transmission

41  
42 Summary (40/40 words): US Infantry trainees were followed through training for MRSA skin and soft tissue  
43 infection, swabbing for colonization at 2-4 week intervals. Sequencing suggests serial importations of diverse strains  
44 on base, followed by transmission mostly within platoons, involving multiple anatomical sites.

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 45 **Abstract – 246/250 words**

46 **Background.** MRSA skin and soft tissue infection (SSTI) is a significant cause of morbidity in  
47 military trainees. To guide interventions, it is critical we understand the epidemiology of MRSA  
48 in this population.

49 **Methods.** Two cohorts ('companies') of US Army Infantry trainees (N=343) at Fort Benning,  
50 GA, USA, were followed during their training cycles (Jun.-Dec. 2015). Trainees had nares,  
51 oropharynx, perianal and inguinal areas swabbed for MRSA colonization at five ~2-4 week  
52 intervals, and monitored for SSTI throughout training. Epidemiological data were collected.  
53 Isolates were sequenced using Illumina HiSeq and NovaSeq. Single-nucleotide polymorphisms  
54 and clusters were identified. Multi-locus sequence type (MLST) and antimicrobial resistance  
55 genes were predicted from *de novo* assemblies.

56 **Results.** 87 trainees were positive at least once for MRSA (12 had SSTI, 2 without any  
57 colonization). Excluding those positive at baseline, 43.7% were colonized within the first month  
58 of training. 244/254 samples were successfully sequenced (including all SSTI). ST8 (n=135,  
59 100% of SSTI), ST5 (n=81) and ST87 (n=21) were the most represented. Three main Clusters  
60 were identified, largely corresponding to these STs. Sub-analyses within Clusters showed  
61 multiple importations of MRSA, with transmission subsequently predominantly within, rather  
62 than between, platoons in each company. Over 50% of trainees were colonized only at other  
63 anatomical sites; restricting analyses to nares missed substantial transmission.

64 **Conclusions.** Serial importations of MRSA into this high-risk setting likely contribute to the  
65 ongoing burden of MRSA colonization and infection among military trainees. Sampling multiple  
66 anatomical sites is critical for comprehensive characterization of MRSA transmission.

67

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 68 **Introduction**

69 Community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) is a major  
70 cause of skin and soft tissue infections (SSTI) [1]. Military personnel have an elevated risk of  
71 MRSA SSTI compared to the general population with new recruits disproportionately affected  
72 [2, 3]. SSTI outbreaks compromise not only training cycles, but also operational readiness  
73 among deployed forces [4]. The annual cost of SSTIs to the US military is approximately 14-32  
74 million US dollars [5]. Past SSTI prevention strategies in military populations have included  
75 nasal decolonization [6] as well as chlorhexidine-based body wash [7, 8], based on the principle  
76 that MRSA colonization precedes and is necessary for infection [9]. However, neither strategy  
77 prevented SSTI in these high-risk populations [6, 8].

78  
79 To develop effective prevention strategies, it is critical we understand the epidemiology and  
80 transmission dynamics of MRSA in this population. Recently, whole genome sequencing (WGS)  
81 has been used to describe the epidemiology of MRSA colonization [4] and infection among US  
82 Army trainees [10]. A key limitation of this study was that WGS was restricted to trainees with  
83 pulse-field gel electrophoresis (PFGE) type USA300 MRSA, the most common cause of CA-  
84 MRSA SSTI. As such, the conclusions that could be drawn about overall MRSA transmission  
85 dynamics, and prevalence of different strains circulating in this population, were greatly limited.  
86 To address this concern, we conducted a prospective cohort study of MRSA in the same trainee  
87 population, following them throughout their training cycle for colonization and infection events.  
88 All MRSA-positive samples from multiple anatomical sites were included.

89

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

90 Our study found a high prevalence of MRSA colonization at baseline, comprising a genetically  
91 diverse population imported from across the US by recruits entering basic training. This was  
92 followed by substantial person-to-person transmission, particularly among soldiers belonging to  
93 the same ‘ platoons’ (i.e., those in the same training company who share the same living  
94 quarters). Moreover, the observed transmission events frequently involved different anatomical  
95 sites - irrespective of nasal colonization. As nasal swabs are routinely and almost exclusively  
96 used to detect MRSA colonization in clinical settings, this suggests MRSA transmission may be  
97 drastically under-estimated with current infection control methodology – a finding with  
98 significant implications for the control of MRSA.

99

### 100 **Methods**

101 **Overview.** We conducted a longitudinal cohort study of SSTI among US Army Infantry trainees  
102 at Fort Benning, Georgia, from June-December 2015. Two companies (designated A and B)  
103 were included, which commenced the 14-week training cycle at two different timepoints during  
104 peak SSTI seasons (June and September, respectively). Infantry training companies are  
105 comprised of ~200 soldiers, further segregated into four platoons comprised of ~50 soldiers. See  
106 **Supplementary Methods** for additional detail.

107

108 **MRSA screening.** MRSA colonization swabs were obtained at baseline and at ~2-4 week  
109 intervals over five study visits. [6] Anterior nasal, oropharyngeal, inguinal and perianal swabs  
110 were collected (see **Supplementary Methods**). Swabs were initially processed at the Benning  
111 Martin Army Community Hospital clinical microbiology laboratory as previously described [8].

112

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

113 ***Epidemiologic data collection.*** Trainees completed questionnaires at baseline for demographic  
114 and MRSA risk factor information, and at follow-up visits to collect data on personal hygiene  
115 practices.

116  
117 ***SSTI case definition.*** Trainees were considered to have probable SSTI if they presented with  
118 cellulitis, abscess, folliculitis, impetigo, paronychia, infected blister, or pilonidal cyst; SSTI cases  
119 were defined as those with *S. aureus* cultured from the clinical site. Case detection is described  
120 in the **Supplementary Methods**.

121  
122 ***Genome sequencing and bioinformatics.*** See **Supplementary Methods** for details. In brief,  
123 genomes were sequenced using the Illumina HiSeq and NovaSeq platforms. As most samples  
124 were known to be USA300, we initially aligned reads to the USA300 TCH1516 reference  
125 genome (National Center for Biotechnology Information [NCBI] Accession CP000730.1) using  
126 the Burrows-Wheeler Aligner MEM algorithm (v.0.7.15, [12]) and identified variants compared  
127 to this reference. Prophage sequences were identified using Phaster (<http://phaster.ca>, [13]) and  
128 SNPs in these regions were masked.

129  
130 Concatenated SNP alignments were used to generate maximum likelihood trees with IQ-TREE  
131 (v.1.6.8, [14]). To infer a more accurate phylogeny for the analysis including all STs, we masked  
132 predicted recombination sites using Gubbins (v.2.3.4, [15]). Hierarchical Bayesian Analysis of  
133 Population Structure (RhierBAPS v.1.1.0, [16]) was used to discriminate clusters of closely-  
134 related samples in R (v.3.5.2). Analyses were then repeated for each major Cluster identified  
135 using a closely-related reference genome.

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

136 *De novo* assemblies were also generated, using SPAdes (v.3.8.0, [17]) and Unicycler (v.0.4.7,  
137 Wick R, <https://github.com/rrwick/Unicycler>). Sequence Types (STs) according to the multi-  
138 locus sequence typing (MLST) scheme [18] were predicted using ‘mlst’ (v.2.10, Seemann T,  
139 <https://github.com/tseemann/mlst>) and Abricate (v.0.8, Seemann T,  
140 <https://github.com/tseemann/abricate>) was used with ResFinder [19] and the Virulence Factor  
141 DataBase (VFDB, [20]) to detect genes associated with antimicrobial resistance and virulence,  
142 respectively. *mupA* and *qacA* genes were predicted using gene-puller.py  
143 (<https://github.com/kwongj/gene-puller>).

144  
145 **Statistical analysis.** Categorical comparisons were done using the Chi-Square test where cell  
146 counts > 5, or Fisher’s Exact test if otherwise. Distributions were compared using the Wilcoxon-  
147 Mann-Whitney test. Conditional logistic regression was used to compare hygiene variables  
148 between trainees who became colonized with time-matched, MRSA-negative controls  
149 (**Supplementary Methods**). Analyses were done in Stata15 (StataCorp, College Station, TX,  
150 USA).

151  
152 **Data availability.** Sequence data is on NCBI’s Sequence Read Archive under BioProject  
153 PRJNA587530.

154  
155 **Ethics approval.** The study was approved by the Uniformed Services University Institutional  
156 Review Board (IDCRP-090) and Harvard T.H. Chan School of Public Health Institutional  
157 Review Board (IRB17-0802).

158

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 159 **Results**

160 We followed two Infantry training companies in the summer and fall of 2015. 343 trainees  
161 participated (132 and 211 in companies A and B, respectively), with 86.6% (297/343) completed  
162 all study visits (**Supplementary Results**). 87 (25.4%) of trainees were MRSA-positive during  
163 the study; 85 had at least one positive colonization swab and 12 were diagnosed with MRSA  
164 SSTI. Among the latter, only seven were colonized prior to or at the same time as their SSTI  
165 (five with the infecting strain); three were colonized only at later timepoints, and two were never  
166 colonized (**Supplementary Table 1**).

167  
168 Among those colonized, thirty-five had a single positive sample; the oropharynx and perianal  
169 sites were most frequently colonized, each representing 31.4% of swabs, followed by nasal  
170 (20.0%), and inguinal (17.1%) sites. Fifty trainees were colonized at multiple anatomical sites  
171 and/or timepoints. The number of repeat positive samples did not differ significantly by  
172 anatomical site of isolation ( $p = 0.16$ ), although inguinal swabs tended to be less frequently  
173 positive (**Figure 1**).

174  
175 *Risk factors for colonization or infection*

176 There were no significant differences in age, self-reported race, or MRSA risk factors at baseline  
177 (**Table 1**) comparing those with colonization/infection to those who were never MRSA-positive.  
178 Epidemiological characteristics of SSTI cases are described in the **Supplementary Results**.

179

180

181

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 182 *Timing of MRSA acquisition*

183 Twenty-seven trainees were colonized at baseline (see **Supplementary Results**). Excluding  
184 these, most trainees appeared to have acquired MRSA shortly after starting training (**Figure 2**);  
185 43.7% of all positive samples were collected during the first month of training. There were no  
186 significant differences in personal hygiene practices over the same timepoint between trainees  
187 with new colonization and those who were never positive (**Supplementary Results**,  
188 **Supplementary Table 2**).

189

### 190 *Genomic analyses*

191 High-quality sequencing data was available for 244/254 (96.1%) MRSA-positive samples from  
192 85 individuals; further analyses presented are restricted to this subset. ST8 (USA300) was the  
193 dominant strain identified (**Table 2**), and included all cases of SSTI. The virulence factor  
194 Panton-Valentine Leukocidin was present in all clinical isolates (as demonstrated by intact *lukS*  
195 and *lukF* genes) and nearly half of colonization isolates (**Table 2**). All samples were PCR-  
196 positive for *mecA*, while 98.8% were predicted to have *mecA in silico*; 2/3 isolates lacking *mecA*  
197 *in silico* were different STs than other isolates from those trainees, indicating that mixed  
198 genotypes may have been present at sampling (**Supplementary Table 3**). To assess the potential  
199 utility of mupirocin and chlorhexidine for decolonization, *mupA* and *qacA* genes were predicted.  
200 *mupA* was only detected in colonization isolates (all ST5) from four trainees (and one ST8  
201 clinical isolate from another trainee), while *qacA* was detected in colonization isolates from three  
202 trainees. Correlation between PFGE and ST are shown in **Supplementary Table 4**.

203

204



## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 205 *Strain replacement*

206 Strain replacement was rare (**Supplementary Figure 1**), occurring only four times during the  
207 study (excluding co-colonization). Two instances involved different STs; trainee 2013 was  
208 initially colonized with an ST5 strain but developed an ST8-SSTI, while 2031 was colonized  
209 with ST8 but later colonized with ST5. The remaining instances involved trainees 1007 and  
210 2061. These trainees had isolates from the same ST that were separated from their later isolates  
211 by > 97 core SNPs, suggestive of exogenous re-infection.

212

### 213 *Overall transmission analyses*

214 See **Supplementary Results** for quality metrics. The recombination-adjusted maximum  
215 likelihood tree is shown in **Figure 3**, excluding one highly divergent isolate (1134\_5N, see  
216 **Supplementary Results**). Five Clusters were identified, with three containing >1 isolate;  
217 Cluster-1 (n=139, 97.1% ST8); Cluster-2 (n=81, all ST5); and Cluster-4 (n=21, all ST87).  
218 Pairwise adjusted core SNP distances between isolates indicate that transmission was  
219 predominantly within, rather than between, platoons in the same Companies (**Figure 4**).

220

### 221 *Sub-analyses of transmission, by Cluster*

222 To provide the highest resolution, analyses were re-run for each Cluster using a closely-related  
223 reference (**Supplementary Table 5**). As ST8s represent the largest proportion of samples  
224 collected, the maximum likelihood tree for Cluster 1 is shown in **Figure 5** (see **Supplementary**  
225 **Figures 2 and 3** for Clusters 2 and 4, respectively). Within Cluster 1, there were nine sub-groups  
226 each representing isolates from 1 to 24 trainees. These sub-groups were separated from one  
227 another by a minimum of 34 SNPs (**Table 3**). In nearly all sub-groups, there were isolates from

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

228 different trainees that shared close genetic relationships, indicative of recent transmission (**Table**  
229 **3**). Overlaying company and platoon (**Figure 5**) suggests that such transmission was largely  
230 restricted to the same platoons, with a small amount across platoons in the same company.  
231 Isolates from different companies were only found in the largest sub-group (1-1), with as few as  
232 6 core SNPs separating them (including repeat isolates per trainee, the median core SNP distance  
233 was 71 across companies, IQR 35-100). Given these companies were separated in time and  
234 space, this suggests undetected transmission via an intermediate source.

235  
236 Nearly all transmission appears to be within rather than between platoons for Clusters 2 and 4 as  
237 well (**Supplementary Figures 2 and 3**). The exception to this was sub-group 1 in Cluster 4,  
238 where multiple platoons are implicated; however, all six trainees in this sub-group were  
239 colonized at baseline (**Supplementary Results and Supplementary Table 6**). There was no  
240 evidence of transmission across companies in either Cluster.

241

### 242 *Anatomical sites and their role in transmission*

243 The nares are commonly thought to play a key role in transmission, but > 50% of trainees  
244 (regardless of Cluster/ST) were positive only at other anatomical sites (**Table 3**). To examine the  
245 impact of this, we repeated our transmission analyses only including isolates from the nares and  
246 SSTI (**Supplementary Table 7**) and found that transmission was substantially under-estimated,  
247 with as many as 100% of trainees being missed in some Clusters/sub-groups.

248

249

250

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

251 *ST8s in context of previous analyses*

252 We compared our contemporaneous samples to historical clinical and colonization isolates from  
253 trainees with ST8-associated SSTI at Fort Benning [4, 10] (described further in the  
254 **Supplementary Results**), and found historical isolates largely clustered independently of more  
255 recent samples (**Supplementary Figure 4**). The median core pairwise distance between these  
256 isolates and our 2015 samples was 111 SNPs (IQR 80-137). 99% of pairs (14,187/14,385) were  
257 separated by > 15 SNPs, suggesting a single reservoir of MRSA is unlikely to be driving ongoing  
258 transmission of ST8-MRSA on base.

259

### 260 **Discussion**

261 MRSA SSTI continues to be a major source of morbidity in military trainees in the United States  
262 [2, 3] and other countries [21, 22]. In this study, we integrated genomics with comprehensive  
263 epidemiologic data to investigate MRSA transmission in US Army Infantry trainees at Fort  
264 Benning, GA, a group known to be at increased risk for MRSA colonization and infection.

265

266 We followed individuals prospectively through training for MRSA colonization and SSTI,  
267 including all positive colonization isolates from these trainees over time. We found there were  
268 multiple STs present at baseline, with subsequent transmission predominantly within, rather than  
269 between platoons - reinforcing the importance of close proximity for MRSA transmission.

270 Consistent with previous ST8 analyses [4, 10], there appeared to be little transmission across  
271 training classes (i.e., companies) over time, suggesting no single endemic strain in this context.

272 By contrast, the persistent burden of MRSA colonization and infection is likely due to the serial  
273 importation of CA-MRSA strains with each incoming group of new trainees. With > 30,000

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

274 trainees from multiple US states and territories cycling through Fort Benning annually, the  
275 diversity of MRSA strains being introduced into this single geographic site is likely considerable.

276

277 Surprisingly, a large proportion of trainees (27/343, [7.9%]) were positive at baseline or positive  
278 shortly after training onset. This suggests that particular phases of training may serve as critical  
279 ‘control points’ wherein interventions to reduce transmission and/or SSTI could be implemented.

280 As no specific risk factors for MRSA colonization were identified, we propose that risk of  
281 acquisition may be predominantly mediated by exposure to circulating strains and shared training  
282 activities, rather than individual hygiene practices - though it is possible the small numbers may  
283 preclude detection of significant differences between our comparison groups.

284

285 The nares are considered the primary reservoir for *S. aureus* [23]. In the current study, however,  
286 we found that more than half (55.3%) of those colonized were exclusively positive at other  
287 anatomical sites. Screening of these different sites is therefore critical to accurately detect  
288 colonization and resolve transmission. This finding may have important implications for the  
289 control of MRSA in military trainees, as MRSA prevalence, transmission, and success of  
290 decolonization efforts, have previously been largely estimated from nasal carriage alone [24]. A  
291 randomized controlled trial among Army combat medics in training [6] found that, while  
292 intranasal mupirocin reduced nasal colonization with MRSA, SSTI rates remained unaffected.  
293 Similar results were found in a large-scale, field-based evaluation of chlorhexidine-based body  
294 wash [7, 8]. One possibility is that inoculation via fomites may contribute to SSTI, however,  
295 another possibility is that trainees colonized exclusively at non-nasal sites (not included in these  
296 studies due to the design) served as a constant reservoir for re-colonization/infection. These

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

297 explanations need not be mutually exclusive. This suggests that, given the negligible levels of  
298 resistance to either treatment, mupirocin coupled with chlorhexidine gluconate washes may yet  
299 be viable tools for SSTI reduction in this population, potentially coupled with oral antimicrobials  
300 [25]. However, for such interventions to work, this would likely require *all* colonized individuals  
301 are identified and treated, regardless of anatomical site.

302

303 This study has a number of major strengths, including a high participation rate with > 85% of  
304 participants completing all visits. Because this was conducted in a military training setting, the  
305 schedule, location, type of activities, and risk of MRSA acquisition were relatively uniform for  
306 trainees across all platoons at all visits. Additionally, the prospective design, with screening for  
307 colonization and collection of corresponding epidemiological data at regular intervals throughout  
308 training allowed us to detect such events as they occurred and assess the potential association  
309 with specific behaviours that preceded them. This degree of follow-up is rarely feasible outside  
310 of a hospital setting. Another strength was the screening of multiple anatomical sites over time;  
311 this not only allowed us to more accurately detect colonization than previous studies in this  
312 population [6, 7, 24] but examine the role they play in transmission. Finally, previous studies  
313 focused exclusively on ST8, with sampling exclusively from cases of SSTI and select controls  
314 [4, 10]; by including all STs, we were able to provide the first fully comprehensive analysis of  
315 population structure and transmission dynamics of MRSA circulating in this setting.

316

317 There are several limitations of this work. Due to resource requirements, we focused recruitment  
318 on two (out of 37) training classes from Jun. - Dec. 2015. Though our analysis suggests limited  
319 transmission across companies (i.e., classes), it is possible some transmission events were not

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

320 detected. Permanent staff on base (e.g. drill sergeants, health care personnel, etc.) may also  
321 represent a reservoir of MRSA, though our analysis suggest this is less likely (as there were  
322 multiple circulating strains, with evidence of importation from outside the base). Staff may also  
323 be source of unrecognized transmission, potentially even during in-processing, but unfortunately,  
324 were not included in the current study. Similarly, sampling of high touch common surfaces (e.g.  
325 countertops, weight benches, wrestling mats, etc.) was not conducted as part of this investigation.  
326 Future studies should address these gaps, to better understand the role these may play in MRSA  
327 transmission dynamics.

328

329 We also note that the included cohorts were purposely chosen based on peak MRSA SSTI  
330 seasonality [8], with some studies reporting increased colonization in warmer months [26, 27].  
331 MRSA prevalence in our classes may therefore be higher than would be found in other cohorts.  
332 However, baseline prevalence of MRSA colonization among combat medic trainees [6] in Jan. –  
333 Dec. 2005 was only 3.9% (based on nasal carriage) and restricting our estimates to nasal carriage  
334 yielded a similar proportion (2.0%,  $p=0.08$ ), suggesting our findings are likely generalizable to  
335 other training courses and/or periods. Another potential limitation of this work is that, for privacy  
336 reasons, perianal and inguinal swabs were collected by the trainees themselves, rather than by  
337 study personnel. Though they were instructed beforehand, it is possible these swabs were not  
338 collected in a consistent fashion and may be more likely to yield false negative results. However,  
339 we did not find a significant difference in frequency of positivity across anatomical sites. Finally,  
340 personal hygiene practices were self-reported at routine study visits for the preceding month,  
341 potentially leading to misclassification due to poor recall. However, as colonization status was

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

342 not known at the time of collecting this information (and was not reported to trainees during the  
343 study), we would not expect this misclassification to be differential.

344

345 Through this in-depth study, we identified critical timepoints in military training associated with  
346 MRSA colonization and highlight the importance of different anatomical sites in mediating this  
347 transmission. Given our historical reliance on nasal screening for detection of MRSA  
348 colonization, and subsequent interventions to prevent infection, this work may have important  
349 implications not only for control of MRSA in the military, but in-patient settings as well. In order  
350 to halt MRSA transmission in the military, and other vulnerable populations, increased screening  
351 to include these other anatomical sites may be necessary; however, cost and feasibility of such  
352 interventions must also be investigated.

353

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

354 **Acknowledgements.**

355 We are thankful for the participation of the US Army Infantry Trainees in our study. We are  
356 indebted to the study team of clinical research coordinators, laboratory personnel, and data  
357 management staff for their dedication to the project. High-performance computing was done  
358 using the Odyssey cluster from the Faculty of Arts and Science, Harvard University. RSL, EVM,  
359 AC, CEE, EE, MWE, JWB and WPH have no conflict to declare.

360

361 **Funding.** This work (IDCRP-090) was conducted by the Infectious Disease Clinical Research  
362 Program (IDCRP), a Department of Defense (DoD) program executed by the Uniformed  
363 Services University of the Health Sciences (USU) through a cooperative agreement with The  
364 Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc. (HJF). This  
365 project has been supported with federal funds from the National Institute of Allergy and  
366 Infectious Diseases, National Institutes of Health (NIH), under Inter-Agency Agreement Y1-AI-  
367 5072 and from the Defense Health Program, U.S. Department of Defense, under award  
368 HU0001190002. This work was also supported by an award from the Military Infectious Disease  
369 Research Program [grant number 64763] to MWE. The genomic epidemiology sub-study was  
370 supported by an R01 grant from the National Institutes of Health [grant number R01AI128344]  
371 awarded to WPH and the Canadian Institutes of Health Research [grant number MFE 152448]  
372 awarded to RSL.

373

374 **Contributions.** EVM, MWE, and JB designed and ran the parent longitudinal cohort study.  
375 WPH and RSL conceived the current genomic study. RSL designed and was responsible for the  
376 bioinformatics and epidemiological analyses, designing/preparing the figures and tables, and



RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

377 writing the manuscript. CEE, EE, and AC were responsible for isolating the bacteria, and culture,  
378 and AC prepared the sequencing libraries. WPH reviewed the initial draft. All authors were  
379 involved in interpreting the data, and reviewed and provided feedback on the manuscript. RSL  
380 takes full responsibility for the content of the paper and decision to submit for publication. Please  
381 note, the views expressed are those of the authors and do not necessarily reflect the official  
382 policy or position of the Henry M. Jackson Foundation for the Advancement of Military  
383 Medicine, the Uniformed Services University, the Department of the Army, the Department of  
384 Defense, nor the U.S. Government. Mention of trade names, commercial products, or  
385 organizations does not imply endorsement by the U.S. Government.

386

387

388

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

### 389 **References**

- 390 1. David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*:  
391 epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev*  
392 **2010**; 23(3): 616-87.
- 393 2. Landrum ML, Neumann C, Cook C, et al. Epidemiology of *Staphylococcus aureus* blood  
394 and skin and soft tissue infections in the US military health system, 2005-2010. *JAMA*  
395 **2012**; 308(1): 50-9.
- 396 3. America DoDUSo. Cellulitis and abscess, active components, US Armed Forces, 2002-  
397 2005. , **2006**.
- 398 4. Millar EV, Rice GK, Ellassal EM, et al. Genomic Characterization of USA300 Methicillin-  
399 Resistant *Staphylococcus aureus* (MRSA) to Evaluate Intraclass Transmission and  
400 Recurrence of Skin and Soft Tissue Infection (SSTI) Among High-Risk Military Trainees.  
401 *Clin Infect Dis* **2017**; 65(3): 461-8.
- 402 5. Lee BY, Singh A, David MZ, et al. The economic burden of community-associated  
403 methicillin-resistant *Staphylococcus aureus* (CA-MRSA). *Clin Microbiol Infect* **2013**;  
404 19(6): 528-36.
- 405 6. Ellis MW, Griffith ME, Dooley DP, et al. Targeted intranasal mupirocin to prevent  
406 colonization and infection by community-associated methicillin-resistant *Staphylococcus*  
407 *aureus* strains in soldiers: a cluster randomized controlled trial. *Antimicrob Agents*  
408 *Chemother* **2007**; 51(10): 3591-8.
- 409 7. Millar EV, Chen WJ, Schlett CD, et al. Frequent use of chlorhexidine-based body wash  
410 associated with a reduction in methicillin-resistant *Staphylococcus aureus* nasal  
411 colonization among military trainees. *Antimicrob Agents Chemother* **2015**; 59(2): 943-9.
- 412 8. Ellis MW, Schlett CD, Millar EV, et al. Hygiene strategies to prevent methicillin-resistant  
413 *Staphylococcus aureus* skin and soft tissue infections: a cluster-randomized controlled  
414 trial among high-risk military trainees. *Clin Infect Dis* **2014**; 58(11): 1540-8.
- 415 9. Miller LG, Diep BA. Clinical practice: colonization, fomites, and virulence: rethinking the  
416 pathogenesis of community-associated methicillin-resistant *Staphylococcus aureus*  
417 infection. *Clin Infect Dis* **2008**; 46(5): 752-60.
- 418 10. Millar EV, Rice GK, Schlett CD, et al. Genomic epidemiology of MRSA infection and  
419 colonization isolates among military trainees with skin and soft tissue infection.  
420 *Infection* **2019**.
- 421 11. Fritz SA, Hogan PG, Camins BC, et al. Mupirocin and Chlorhexidine Resistance in  
422 *Staphylococcus aureus* in Patients with Community-Onset Skin and Soft Tissue  
423 Infections. *Antimicrobial Agents and Chemotherapy* **2013**; 57(1): 559-68.
- 424 12. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.  
425 *arXiv* **2013**: 1303.3997v2.
- 426 13. Arndt D, Grant JR, Marcu A, et al. PHASTER: a better, faster version of the PHAST phage  
427 search tool. *Nucleic Acids Res* **2016**; 44(W1): W16-21.
- 428 14. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective  
429 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*  
430 **2015**; 32(1): 268-74.

## RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

- 431 15. Croucher NJ, Page AJ, Connor TR, et al. Rapid phylogenetic analysis of large samples of  
432 recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **2015**;  
433 43(3): e15.
- 434 16. Tonkin-Hill G, Lees JA, Bentley SD, Frost SDW, Corander J. RhierBAPS: An R  
435 implementation of the population clustering algorithm hierBAPS. *Wellcome Open Res*  
436 **2018**; 3: 93.
- 437 17. Bankevich A, Nurk S, Antipov D, et al. SPAdes: a new genome assembly algorithm and its  
438 applications to single-cell sequencing. *J Comput Biol* **2012**; 19(5): 455-77.
- 439 18. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for  
440 characterization of methicillin-resistant and methicillin-susceptible clones of  
441 *Staphylococcus aureus*. *J Clin Microbiol* **2000**; 38(3): 1008-15.
- 442 19. Zankari E, Hasman H, Cosentino S, et al. Identification of acquired antimicrobial  
443 resistance genes. *J Antimicrob Chemother* **2012**; 67(11): 2640-4.
- 444 20. Chen L, Zheng D, Liu B, Yang J, Jin Q. VFDB 2016: hierarchical and refined dataset for big  
445 data analysis--10 years on. *Nucleic Acids Res* **2016**; 44(D1): D694-7.
- 446 21. Aamot HV, Eskonsipo PKJ, Jorgensen SB, Blomfeldt A. *Staphylococcus aureus*  
447 colonization during military service: a prospective cohort study. *Clin Microbiol Infect*  
448 **2018**; 24(7): 744-8.
- 449 22. Nivesvivat T, Janthayanont D, Mungthin M, et al. Methicillin-susceptible *Staphylococcus*  
450 *aureus* skin infections among military conscripts undergoing basic training in Bangkok,  
451 Thailand, in 2014. *BMC Res Notes* **2016**; 9: 179.
- 452 23. Mulcahy ME, McLoughlin RM. Host-Bacterial Crosstalk Determines *Staphylococcus*  
453 *aureus* Nasal Colonization. *Trends Microbiol* **2016**; 24(11): 872-86.
- 454 24. Ellis MW, Hospenthal DR, Dooley DP, Gray PJ, Murray CK. Natural history of community-  
455 acquired methicillin-resistant *Staphylococcus aureus* colonization and infection in  
456 soldiers. *Clin Infect Dis* **2004**; 39(7): 971-9.
- 457 25. Simor AE PE, McGeer A, Konvalinka A, LOeb M, Devlin HR, Kiss A. Randomized  
458 Controlled Trial of Chlorhexidine Gluconate for Washing, Intranasal Mupirocin,  
459 and Rifampin and Doxycycline Versus No Treatment for the Eradication of Methicillin-Resistant  
460 *Staphylococcus aureus* Colonization. *Clin Infect Dis* **2007**; 44(178-85).
- 461 26. Leekha S, Diekema DJ, Perencevich EN. Seasonality of staphylococcal infections. *Clin*  
462 *Microbiol Infect* **2012**; 18(10): 927-33.
- 463 27. Mork RL, Hogan PG, Muenks CE, et al. Comprehensive modeling reveals proximity,  
464 seasonality, and hygiene practices as key determinants of MRSA colonization in exposed  
465 households. *Pediatr Res* **2018**; 84(5): 668-76.
- 466 28. SNP-sites: rapid efficient extraction of SNPs from multi- FASTA alignments.
- 467 29. Cheng L, Connor TR, Siren J, Aanensen DM, Corander J. Hierarchical and spatially explicit  
468 clustering of DNA sequences with BAPS software. *Mol Biol Evol* **2013**; 30(5): 1224-8.
- 469 30. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and  
470 annotation of phylogenetic and other trees. *Nucleic Acids Res* **2016**; 44(W1): W242-5.
- 471 31. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y, McInerney G. ggtree: anrpackage for  
472 visualization and annotation of phylogenetic trees with their covariates and other  
473 associated data. *Methods in Ecology and Evolution* **2017**; 8(1): 28-36.
- 474

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

475

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

476  
477  
478  
479  
480

**Tables.**

**Table 1. Clinical and epidemiological characteristics of trainees at baseline (N=343)**

Characteristic	Colonized and/or SSTI (n=87)	Never colonized and no SSTI (n=256)	P value
Age, average (SD)	19.2 (1.7)	19.4 (2.5)	0.42 <sup>a</sup>
Hispanic ethnicity, n (%)	15 (17.2)	40 (15.6)	0.72 <sup>b</sup>
Race, n (%)			
White	64 (73.6)	182 (71.1)	
Black	8 (9.2)	23 (9.0)	
Asian	0 (0)	6 (2.3)	
American Indian	1 (1.1)	5 (2.0)	
Pacific Islander	0 (0)	3 (1.2)	
Other <sup>c</sup>	7 (8.0)	20 (7.8)	
> 1 race reported	7 (8.0)	17 (6.6)	0.88 <sup>d</sup>
Lived alone prior to training, n (%)	5 (5.7)	10 (3.9)	0.54 <sup>d</sup>
Lived with pets/livestock, n (%)	64 (73.6)	177 (69.1)	0.44 <sup>b</sup>
Antibiotics in previous 6 months, n (%)	6 (6.9) <sup>e</sup>	16 (6.3)	0.83 <sup>b</sup>
Admitted to hospital in past year, n (%)	1 (1.1)	6 (2.3)	0.68 <sup>d</sup>
Worked in a hospital in past year, n (%)	0 (0)	8 (3.1)	1.0 <sup>d</sup>
Lived with someone who was hospitalized/worked in hospital in past year, n (%)	1 (1.1)	69 (27.0)	0.47 <sup>d</sup>
Known/suspected SSTI/MRSA infection in past year, n (%)	2 (2.3)	2 (0.8)	0.27 <sup>b</sup>
Prior contact with MRSA SSTI in 3 months prior to start of training, n (%)	1 (1.1)	4 (1.6)	1.00 <sup>b</sup>

481 Note, all trainees on base were male during the study year. SSTI = skin and soft tissue infection ; MRSA =  
482 methicillin-resistant *S. aureus*. <sup>a</sup> T-test with unequal standard deviations. <sup>b</sup> Chi-square test with 1 df. <sup>c</sup> Includes 28  
483 individuals who self-reported as “Hispanic” and selected race “Other”. <sup>d</sup> Fisher’s Exact test. <sup>e</sup> One individual with  
484 prior antibiotic exposure had an SSTI; none of those with SSTI had hospitalization, prior SSTI themselves or contact  
485 with someone with SSTI in the three months prior to arrival to Fort Benning.

486  
487  
488

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

489 **Table 2. Genomic characteristics of included isolates (N=244)**

490

<i>In silico</i> prediction	Colonization, n=232	Clinical, n=12
MLST, n (%)		
ST8	123 (53.0)	12 (100)
ST5	81 <sup>a</sup> (34.9)	0 (0)
ST87	21 (9.1)	0 (0)
Other STs <sup>b</sup>	4 (1.7)	0 (0)
SLV of ST8	1 (0.4)	0 (0)
Novel STs <sup>c</sup>	2 (0.9)	0 (0)
lukS/lukF-positive, n (%)	108 (46.6)	12 (100)

491 ‘MLST’ refers to multi-locus sequence type, while ‘SLV’ refers to a single locus variant, wherein one allele differs  
492 from the listed ST. Three colonization samples were PCR-positive for *mecA* but negative *in silico*: 1133\_2P,  
493 1134\_5N, 2069\_2I. <sup>a</sup>This includes isolate 2014\_3P which has two alleles of the *tpi* gene (1,99). <sup>b</sup>This includes  
494 ST15, ST30, ST45, ST72. <sup>c</sup>Includes a triple locus variant of ST8.

495

496

497 **Table 3. Pairwise core single nucleotide polymorphism (SNP) distances within sub-groups of Clusters 1, 2 and 4**  
 498

Cluster	Sub-group	Number of isolates	Number of trainees	SNP distance between trainees, median (range)	SNP distance within trainees, median (range)	Number of trainees with positive at baseline, n (%)	Number of trainees with nasal samples, n (%)	Number of trainees with clinical samples, n (%)
1	1-1	69	24	7 (0-383)	0 (0-9)	4 (16.7)	12 (50.0)	6 (25.0)
	1-2	16	8	4 (0-9)	0 (0-4)	0 (0)	3 (37.5)	-
	1-3	7	2	2 (1-3)	0 (0-1)	1 (50.0)	1 (50.0)	-
	1-4	11	5	1 (0-3)	0 (0-2)	0 (0)	2 (40.0)	3 (60.0)
	1-5	4	2	159 (157-161)	3 (2-4)	1 (50.0)	0 (0)	-
	1-6	8	3	21 (0-22)	1 (0-7)	1 (33.3)	1 (33.3)	2 (66.7)
	1-7	14	2	0 (0-8)	1 (0-10)	1 (50.0)	1 (50.0)	1 (50.0)
	1-8	6	3	0 (0-2)	1.5 (0-3)	3 (100)	0 (0)	-
	1-9	4	1	-	8.5 (2-15)	1 (100)	0 (0)	-
2	2-1	9	5	4 (1-116)	0 (0-3)	1 (100)	2 (40.0)	-
	2-2	51	23	1 (0-655)	1 (0-3)	3 (13.0)	7 (30.4)	-
	2-3	11	2	0 (0-4)	1 (0-5)	1 (50.0)	1 (50.0)	-
	2-4	3	1	-	1 (0-1)	1 (100)	0 (0)	-
	2-5	2	1	-	0 (0-0)	1 (100)	0 (0)	-
	2-6	5	1	-	0 (0-0)	1 (100)	0 (0)	-
4	4-1	12	6	0 (0-2)	1 (0-3)	6 (100)	1 (16.7)	-
	4-2	9	5	0 (0-0)	0 (0-0)	0 (0)	4 (80.0)	-

499 Clusters were identified in the analysis including all samples, using hierBAPS [16]. Core SNPs within each Cluster were then identified by repeating the analysis,  
 500 but aligning the reads from isolates within each Cluster to a closely-related reference from the most represented sequence type in that Cluster. hierBAPS [16]  
 501 then run again for each Cluster using these new alignments as input (to obtain the highest resolution), and sub-groups of closely-related isolates were identified  
 502 within each Cluster. The number of core SNPs were tabulated within each sub-group and are comparable within, but not across, Clusters (as each Cluster has a  
 503 different core genome). Cluster 1 sub-groups were separated by a minimum of 34 core SNPs, Cluster 2 sub-groups were separated by a minimum of 150 core  
 504 SNPs, and Cluster 4 sub-groups were separated by a minimum of 100 core SNPs. Cluster 1 was predominantly ST8 (n=136/139); three non-ST8 isolates were  
 505 included based on hierBAPS analysis, including 1133\_2P (a single locus variant of ST8), 1105\_5P (a triple locus variant of ST8) and 1133\_2P (ST72, a triple  
 506 locus variant of ST8). These three non-ST8 isolates were included in sub-group 1-1, and were a maximum of 383 core SNPs to another other sample within this  
 507 sub-group. Cluster 2 was comprised of all ST5 isolates, while Cluster 4 was comprised of all ST87 isolates.

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

508 **Figure legends.**

509

510 **Figure 1. Number of isolates per trainee, by anatomical site of isolation.** 85 trainees had at

511 least one non-clinical sample positive for MRSA during the study. Includes samples that were

512 not available/sequenced.

513



RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

514 **Figure 2. Timing of first MRSA positive samples collected from each trainee, by company**  
515 **at Fort Benning, Georgia, 2015.** In total, 87 trainees were positive for MRSA from the two  
516 Companies under study. 27 trainees were positive at baseline; these samples were collected  
517 within 1-2 days of arrival to base, while all trainees were at one central location.  
518  
519

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

520  
521 **Figure 3. Maximum likelihood tree for colonization and clinical samples from trainees**  
522 **undergoing basic training at Fort Benning, Georgia in 2015.** One isolate (1134\_5N) was  
523 highly divergent from all others, at > 18,500 core SNPs compared to < 1000 SNPs for the next  
524 most divergent. Therefore, this isolate was excluded from this analysis. A concatenated SNP  
525 alignment of the remaining isolates was produced using snp-sites [28] and IQ-Tree [14] (v1.6.8)  
526 was used to produce the maximum likelihood tree, using the TVMe model with ascertainment  
527 bias correction. Model selection was based on the lowest Bayesian Information Criterion.  
528 Hierarchical Bayesian Analysis of Population Structure (hierBAPS) [16, 29] was used to identify  
529 clusters with the full alignment as input. The tree was visualized using the Interactive Tree of  
530 Life [30]. Samples from skin and soft tissue infection are indicated.

531

532

533

RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

534 **Figure 4. Recombination-adjusted core pairwise SNP distributions by company and**  
535 **platoon.**

536 **A. Pairwise core SNPs across and within companies.**

537 **B. Pairwise core SNPs across and within platoons.**

538 Core SNPs were identified compared to the USA300 reference (excluding isolate 1134\_5N due  
539 to its high divergence from all others isolates in the dataset). This resulted in 29,403 pairwise  
540 comparisons between trainees, including repeat samples from the same individuals. Retaining  
541 closely-related samples from the same trainees would be expected to artificially reduce the  
542 pairwise distances within companies and platoons. Therefore, where repeat samples from a  
543 single participant were within 5 SNPs of one another, only one of these was retained for this  
544 analysis. This threshold was chosen because it represented 90% of pairs from the same trainees.  
545 In total, 3,739 pairwise comparisons were therefore included in this analysis. Note that all  
546 trainees in the same platoon are, by default, in the same company.


547

548


RUNNING TITLE: GENOMICS OF MRSA IN MILITARY TRAINEES

549 **Figure 5. In-depth analysis of Cluster-1 and potential transmission within and across**  
550 **platoons.** There were 139 isolates in Cluster-1, as classified by hierBAPs [16, 29] in the overall  
551 analysis (Figure 3), 136 of which were ST8. A concatenated core SNP alignment was produced  
552 with these isolates. A maximum likelihood tree was produced using IQ-Tree [14] (v1.6.8) and  
553 visualized using ggtree [31] in R (v.3.5.2). hierBAPS was again used to identify sub-groups of  
554 closely-related isolates within this Cluster, which are indicated in colour. Note that this analysis  
555 includes repeat isolates from all trainees; sample names are indicated using the format  
556 *Participant Identifier + Visit Number + Anatomical Site of Isolation* (where N=nasal, I=Inguinal,  
557 P=Perianal and O=Oropharyngeal), such that repeat samples can be identified. The heatmap  
558 indicates epidemiological contact between trainees; the letters (A and B) represent the company  
559 and the numbers represent the specific platoon within that company.  
560  
561

**Figure 1.**










**Figure 2.**




**Figure 3.**

Tree scale: 0.001


Clusters	
	1
	2
	3
	4
	5

MLST	
	5
	8
	15
	30
	72
	87
	-




**Figure 4.**

A.



B.





**Figure 5.**

