

Distinct single cell gene expression in peripheral blood monocytes correlates with treatment response groups defined by type I interferon in rheumatoid arthritis

Theresa L. Wampler Muskardin^{1,2*}, Wei Fan^{3*}, Zhongbo Jin⁴, Mark A. Jensen^{1,2}, Jessica M. Dorschner⁵, Yogita Ghodke-Puranik^{1,2}, Betty Dicke⁵, Danielle Vsetecka⁶, Kerry Wright⁵, Thomas Mason⁵, Scott Persellin⁵, Clement J. Michet⁵, John M. Davis⁵, Eric Matteson⁵, Timothy B. Niewold¹

¹Colton Center for Autoimmunity – Department of Medicine, New York University School of Medicine, New York, NY

²Division of Rheumatology - Department of Medicine, New York University School of Medicine, New York, NY

³Department of Rheumatology, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, China

⁴Department of Pathology, Immunology and Laboratory Medicine, University of Florida College of Medicine, Gainesville, FL

⁵Department of Internal Medicine, Division of Rheumatology, Mayo Clinic, Rochester, MN

⁶UnitedHealth Group, Research & Development

*Co-first authors

Grant Numbers/Other Funding Source: *TWM: Arthritis National Research Foundation Arthritis Research Award; NYU Colton Center for Autoimmunity; NYU Department of Medicine Sapperstein Scholar Award; NYU CTSI Scholar Award; Central Society for Clinical and Translational Medicine Career Development Award; Mayo Clinic Career Development Award in Rheumatoid Arthritis Research; TBN: NIH (AR060861, AR057781, AR065964, DK107984), Rheumatology Research Foundation.*

Abstract

Objective: Previously, we have shown that the ratio of IFN β / α activity can predict response to tumor necrosis factor inhibitors (TNFi) in rheumatoid arthritis (RA). In this study, our objective was to better understand the impact of this type I IFN ratio in monocytes from RA patients.

Methods: We compared single cell gene expression in purified classical (CL) and non-classical (NC) monocytes from RA patients. Patients with high vs. low serum IFN β / α ratio were compared to each other, as determined by functional type I IFN assay.

Results: We found large differences in monocyte gene expression between patients with different IFN β / α ratios, which was apparent in principal component analysis. Cluster analysis also documented strong differences between these response groups. In CL monocytes, decreased expression of JAK1 and IFI27 were strongly associated with the non-response low IFN β / α ratio. In NC monocytes, increased expression of HLADRB1, TNFA, PDL1, TGFB, and others were associated with the non-response IFN ratio. Interestingly, JAK1 expression was absent in all monocytes from some patients. This pattern was strongly associated with the non-response type I IFN ratio, suggesting a major biological difference between JAK1 expressors and non-expressors.

Conclusions: Differences in the ratio of IFN β / α are associated with major changes in gene expression in RA patient monocytes, suggesting differential IFN pathway activation in RA patients who are either more or less likely to respond to TNFi. This work could suggest mechanisms for TNFi non-response, as well as potential therapeutic strategies for those patients who would not to respond to TNFi.

INTRODUCTION

Rheumatoid Arthritis (RA) is the most common inflammatory joint disease world-wide, characterized by a destructive arthritis and serious extra-articular manifestations, including accelerated vascular disease(1). Early, effective treatment prevents damage. Thus, remission or very low disease activity within the first 3 months is the goal (2, 3). New therapies have made remission possible for a greater number of patients. However, the current treatment strategy is one of trial-and-error, as we are not able to predict which medication will work for an individual patient. Tumor necrosis factor inhibitors (TNFi) are the most common biologic treatment employed(4). Responses are variable, with approximately 30% not responding and another 30% achieving only partial response. Insufficient treatment is associated with increased morbidity, mortality, and a heavy economic burden (5-8). Type I IFN levels are genetically determined to some degree (9, 10) and type I IFNs are pleiotropic biologic response modifiers, making them ideal candidate biomarkers for response to immunomodulatory therapies. Recently, we studied pre-treatment circulating IFN-alpha (IFN α), IFN-beta (IFN β), and total type I IFN activity in RA patients just prior to receiving a TNFi (11) in independent test and validation cohorts. The ratio of IFN β to IFN α activity (IFN β/α) >1.3 was strongly predictive of non-response to TNFi therapy (specificity=77% in the validation cohort). Remarkably, no patient with a ratio >1.3 achieved remission or low disease activity.

Monocytes are one of the major effector cells in RA (12, 13). By studying the effect of IFN β/α ratio on monocytes, we understand the functional impact of the IFN ratio on a key effector cell type. This may suggest cellular mechanisms that underlie response/non-response to TNFi therapy in RA. We chose to study gene expression in single cells, as effects of IFN on single immune cells or cell types may be masked in whole blood or mixed cell populations (14). We find that circulating type I IFN ratio corresponds to strikingly different gene expression patterns in RA patient monocytes, and that particular transcripts such as *JAK1* are highly informative and

could suggest alternate therapeutic avenues in patients who are predicted to be TNFi non-responders.

METHODS

Patient and Public Involvement. Patients/the public were not involved in the design of the study. The study design and plans to disseminate study results to participants were informed by patient priorities and preferences.

Subjects and Samples. Blood samples from 15 patients with RA were recruited from the Mayo Clinic in Rochester, Minnesota, USA. All of the patients fulfilled the 2010 American College of Rheumatology classification criteria for RA (15) and were seropositive. Exclusion criteria included overlap autoimmune connective tissue disease, pregnancy, active acute infection, chronic infection (e.g., hepatitis C, HIV, etc.), current intravenous therapy (e.g., methylprednisolone or cyclophosphamide), and history of biologic therapy. All samples were obtained prior to initiation of TNFi. All patients provided informed consent, and the study was approved by the institutional review board. Subjects were grouped by pre-TNFi serum type I IFN activity into two groups, those with detectable type I IFN activity but low IFN β/α ratio (IFN β/α > 0 and ≤ 1.3 , n=6, responder group), and those with either undetectable type I IFN activity or a high IFN β/α ratio (IFN β/α 0 or >1.3, n=9). Patients with undetectable type I IFN activity typically did not respond to TNFi therapy (11), and so these patients were grouped together with those who have an IFN β/α ratio >1.3 (non-responder group).

Determination of IFN β/α ratio. Type I IFN activity in serum was measured using a validated functional assay in which reporter cells are used to measure the ability of patient sera to cause type I IFN-induced gene expression (16). Reporter cells (WISH cells, ATCC #CCL-25) were

cultured with patient serum for 6 hours. Cells were then lysed, and cDNA was made from total cellular mRNA. Canonical type I IFN-induced gene expression (*MX1*, *PKR*, and *IFIT1*) (17), was measured using qPCR. The relative expression of these three genes was standardized to healthy donors and summed to generate a score reflecting the ability of sera to cause type I IFN-induced gene expression (serum type I IFN activity). This assay has been informative in a wide range of autoimmune diseases (11, 16, 18-20), and we have not found significant functional inhibitors in samples studied to date (21). Additional aliquots were tested following pre-incubation with polyclonal anti-IFN α (19.6 μ g/mL, PBL Assay Science) and anti-IFN β (10.1 μ g/mL, Chemicon) antibodies. The amount of inhibition of the observed type I IFN activity by anti-IFN α antibody allowed for quantitative assessment of IFN β activity, and that by anti β antibody allowed for quantitative assessment of IFN α activity. The ratio of IFN β activity to IFN α activity (IFN β/α activity ratio) was then calculated for each serum sample using these data. Those samples reading very low (<1 pg/mL) for total type I IFN activity were categorized as not having significant type I IFN present, and no ratio was calculated.

Purification of classical (CD14⁺⁺CD16⁻) and non-classical (CD14^{dim}CD16⁺) monocytes.

Classical (CL) and non-classical (NC) monocytes were isolated from peripheral blood using the protocol described in Jin et al(22). Briefly, CL (CD14⁺⁺CD16⁻) monocytes were purified using the Human Pan-Monocyte Isolation Kit (Miltenyi) with modification of adding anti-CD16-biotin (Miltenyi) into the biotin-antibody cocktail. CD14⁺ selection (Miltenyi) was used subsequently to further increase purity. Purified CL monocytes were stained with Molecular Probes CellTracker Green CMFDA Dye (Life Technologies). NC (CD14^{dim}CD16⁺) monocytes were purified similarly, using CD16 microbeads (Miltenyi) during positive selection. Purity checked by flow cytometry was very high (> 95%) for both CL and NC monocytes.

C1 single cell isolation and measurement of gene expression. Using the Fluidigm C1 Single-Cell Auto Prep System we isolated single cells from the bulk monocyte subsets. NC and

CL monocytes were loaded onto the C1 Integrated Fluidic Circuit (IFC) sequentially.

Determination of NC or CL lineage of individual cells was made by direct visualization using fluorescent microscopy. Pre-amplification was done using the C1 Single-Cell Auto Prep Array IFC, according to the manufacturer's protocol. Melt curves were inspected to ensure that all PCR products were uniform. Amplification curves were analyzed and those not following the expected log-growth curve were excluded. 87 target genes were analyzed (Supplemental Table 1). Target gene pre-amplified cDNAs were assayed using 96.96 IFCs on the BioMark HD System (Fluidigm) according to the manufacturer's protocol. Empty wells and wells that contained more than one cell after C1 automated single cell capture were identified by visual inspection using microscopy. These wells were excluded from the dataset. A cell was also removed from the dataset if failure score (total CT value) was greater than two standard deviations above the mean, as this indicated that the cell's overall expression level was too low to be trusted for downstream analysis.

Statistical Analysis. Principal component analysis was used to reduce dimensionality in the complex data sets, and compare overall trends between response groups and CL and NCL monocytes. Unsupervised hierarchical clustering was done to detect individual genes and gene sets that defined the response groups and to identify other possible strata within the data. Each gene was also tested individually for association with response group, using Mann Whitney U testing for the quantitative data and Fisher's exact test for the categorical expressed/not expressed analysis. For these analyses, we used the following strategy to account for multiple comparisons. We expected to find correlations between transcripts in the same cell, which would make a strict Bonferroni correction inappropriate, as each of the 87 tests is not independent. So we first calculated pairwise Spearman correlations (ρ) for each possible pairing of transcripts, resulting in 3741 pairwise correlations. The average correlation between

transcripts was then calculated and a threshold p-value was derived using the following modified Bonferroni method to account for between-transcript correlations:

$$p_{corr} = p_{obs} \times 87 \left[\frac{1 - \sum_{i=1}^{3741} |rho_i|}{3741} \right]$$

This resulted in a threshold p-value of <0.0008 for a corrected alpha of 0.05. We also examined correlations between transcripts in Mo from individual participants, to detect co-expressed genes using the same threshold p value (p<0.0008).

RESULTS

Circulating type I IFN ratio corresponds to large differences in monocyte gene expression

Among the participants in the groups, there were no significant differences in age or disease activity score (DAS), and treatments were comparable (Table 1).

Table 1. Patient characteristics.

	IFN-β/α 0 or > 1.3 (n=9)	IFN-β/α > 0 and ≤ 1.3 (n=6)
Age (mean,range)*	54 (27 - 75)	55 (42 - 70)
Gender (F, M)	4F, 5M	4F, 2M
CCP positivity (n, %)	8 (89)	6 (100)
RF positivity (n, %)	7 (78)	5 (83)
ESR (median, range)	7.5 (0 - 41)	21.5 (13 - 40)
DAS28-CRP (mean ± std.dev, range)*	2.92 ± 1.82	3.42 ± 0.62
	1.15 - 5.77	2.52 - 4.23
Medications (n, %)		
Prednisone	4 (44)	1 (17)
<i>median dose (mg/day)</i>		10
<i>dose range (mg/day)</i>	5 to 15	5
NSAIDs (prn)	0	3 (50)
Methotrexate	8 (89)	3 (50)
<i>median dose (mg/wk)</i>		20
<i>dose range (mg/wk)</i>	17.5 - 25	7.5 - 20
Sulfasalazine	2 (22)	0
Leflunomide	2 (22)	0
Hydroxychloroquine	5 (55)	2 (33)
Statin	3 (33)	1 (17)
ASA-81	2 (22)	0
Allopurinol	1 (11)	0

We were able to analyze results of 87 target genes from 701 individual monocytes (342 CL, 359 NC). Principal component analysis (PCA) showed large differences in the second principal component in CL Mo when cells are labeled by their response group category. Similar findings were observed in the NC monocytes, although the clustering by response group related to the second principal component on the PCA plot was not as strong in NC cells as in CL cells (Figure 1). We next performed unsupervised hierarchical clustering of the target genes to visualize the difference between groups with respect to individual transcripts (Figure 2). In this analysis, it was clear that certain genes strongly aligned with the type I IFN activity groups. In particular, *JAK1* appeared to be strongly predictive of type I IFN ratio group (Figure 2).

Different genes were associated with blood IFN ratio in CL vs. NC cells

In the categorical expressed/not expressed analysis, there were significant differences in the transcripts observed between IFN ratio response groups in CL as compared to NC monocytes. In CL monocytes, in addition to *JAK1*, *IFI27* was less likely to be expressed in the non-response IFN β/α 0 or >1.3 group (Table 2). Thus the two significant findings in this analysis in CL monocytes were both less likely to be expressed in the non-response group. In NC monocytes, a number of transcripts were more likely to be expressed in the non-response IFN β/α 0 or >1.3 group. These include *HLADRB1*, *TNFA*, *PDL1*, *TGFB*, *CD11c*, *IL8*, and *IFNAR1* (Table 2). One transcript, *JAK1*, was less likely to be expressed in the non-response IFN β/α 0 or >1.3 group (Table 2).

Table 2. Odds of being expressed in the IFN β / α 0 or > 1.3 group. P value by Fisher's Exact test.

Classical Monocytes			
Transcript	Odds ratio	95% CI	p value
<i>JAK1</i>	0.061	0.030 to 0.126	<0.0001
<i>IFI27</i>	0.380	0.229 to 0.637	0.0005
Non-classical Monocytes			
Transcript	Odds ratio	95% CI	p value
<i>HLADB1</i>	3.23	1.953 to 5.198	<0.0001
<i>TNFA</i>	2.96	1.780 to 4.985	<0.0001
<i>PDL1</i>	2.64	1.703 to 4.155	<0.0001
<i>TGFB</i>	2.61	1.674 to 4.126	<0.0001
<i>CD11c</i>	2.58	1.631 to 4.002	<0.0001
<i>IL8</i>	2.46	1.595 to 3.779	<0.0001
<i>JAK1</i>	0.39	0.248 to 0.617	<0.0001
<i>IFNAR1</i>	2.14	1.382 to 3.304	0.0008

Examining the quantitative data results in some similar findings and some additional findings are apparent. In CL monocytes, many genes were reduced in expression in the IFN β / α 0 or >1.3 group, including *JAK1* and *IFI27* which were significant in the expressed/not expressed analysis. Additional transcripts that were reduced in the IFN β / α 0 or >1.3 group in quantitative analysis were: *TLR7*, *TLR8*, *TLR2*, *MAVS*, *PKR*, *GMCSF*, *IRF8*, *IL4*, *IL1A*, *ILT7*, *CD127*, *CD16*, and *CCR4* (Supplemental Figure A). In NC Mo, a large number of genes were shown increased expression in the IFN β / α 0 or >1.3 group, including *HLADRB1*, *TNFA*, *PDL1*, *TGFB*, *CD11c*, *IL8*, and *IFNAR1* which were identified in the expressed/not expressed analysis. Additional genes that were significantly increased in the in the IFN β / α 0 or >1.3 group in quantitative analysis were: *TLR4*, *MYD88*, *IRF1*, *FCER1G* and *CD86* (Supplemental Figure B). Genes that differed between groups by non-parametric (Mann Whitney U) univariate analysis were tested in multivariate logistic regression models. In CL Mo, *JAK1*, *TLR2*, *IRF8*, *CD16*, and *IL1A* were retained as independent factors predictive of type I IFN activity group (Table 3).

Table 3. Multivariate analysis results.

Multivariate Logistic Regression			
	Transcript	Coefficient	P value
CL Mo	<i>JAK1</i>	-0.35391	<0.0001
	<i>TLR2</i>	-0.50455	0.0003
	<i>IRF8</i>	-0.64969	0.002
	<i>CD16</i>	-0.44492	0.029
	<i>IL1A</i>	-0.28175	0.035
NC Mo	<i>CD86</i>	0.22951	0.001
	<i>HLADRB1</i>	0.097105	0.016
	<i>IL8</i>	0.11685	0.002
	<i>PDL1</i>	0.099174	0.037
	<i>TGFB</i>	0.07669	0.042
	<i>FCER1G</i>	-0.25984	0.034

In NC Mo, *CD86*, *HLADRB1*, *IL8*, *PDL1*, *TGFB*, and *FCER1G* were retained in the model. (Table 3). ROC curve analysis of these transcripts demonstrated an area under the curve (AUC) of 0.89 (Std. error 0.0172, 95% CI 0.849-0.919) and 0.76 (Std. error 0.026, 95% CI 0.708-0.799), respectively.

JAK1 expression was completely suppressed in some RA patients, and this suppression correlated strongly with non-response

JAK1 was more likely to be not expressed in both CL and NC monocytes from patients in the IFN β / α 0 or > 1.3 group (Odds 0.06, p value < 0.0001, 95% CI 0.03-0.13 in CL; Odds 0.39, corrected p value < 0.0063, 95% CI 0.25-0.62 in NC). Ninety-one percent of CL monocytes and seventy-six percent of NC monocytes in the IFN β / α 0 or > 1.3 group did not express *JAK1*. Whereas in the other IFN group (IFN β / α > 0 \leq 1.3), the majority of CL monocytes (63%) and forty-five percent of NC monocytes expressed *JAK1*. Strikingly, only one participant in the

IFN β / α 0 or > 1.3 group expressed *JAK1* in CL monocytes (Figure 3). Interestingly, after enrollment into our study, this participant was found to have several pre-malignant melanoma lesions that were ultimately removed near the time she began TNFi. Melanocytes and melanoma cells produce IFN β and are capable of suppressing their own proliferation by secretion of endogenous IFN β (23). It is possible that the increased IFN β / α activity noted in this participant was significantly influenced by the pre-malignant melanoma and less informative as a physiologic immune phenotype predictive of response to TNFi.

The participants who did not express *JAK1* in CL monocytes also did not express *JAK1* in NC monocytes (Figure 3). This “on or off” expression pattern was not seen in any of the other transcripts studied, and was not observed in healthy controls (data not shown). In the *JAK1* expressors, most of the cells expressed *JAK1* (83% of cells in CL, 99% in NC). Given this observed distribution, even a subject with only 5 CL cells represented would have a 0.01% chance for miscategorization due to sampling error. The same primers were used to measure expression in all experiments, and also have been used to study healthy controls and lupus patients in other studies in which this pattern of *JAK1* expression was not observed (22). Each of the 96-well plates were run independently on different days, and it would be highly unlikely that the same one out of the 87 target genes would fail each time, and largely in those patients with a non-response IFN ratio. No other transcript shared this expression pattern in our study.

DISCUSSION

Using a novel single cell PCR approach, we used a panel of type I IFN and innate immune system related genes to define gene expression states in monocytes from RA patients with blood type I IFN ratios that would predict response or non-response to TNFi. Compared to RNA-seq data, PCR data is typically more robust and more quantitative, and we found that our

quantitative analyses both confirmed and extended the findings observed in the expressed/not expressed analyses. Also, by working with a more limited panel of carefully quantitated genes we were able to perform some innovative analyses, which led to novel insights in our study. We observed striking differences in gene expression patterns in circulating monocytes between those subjects likely to respond to TNFi as compared to those that are not likely to respond. Our data suggest differential IFN production and downstream pathway activation in in monocytes from patients who do not respond to TNFi. Production of type I IFNs depends on cell type and context (24-27). An RA patient's circulating IFN β/α activity may reflect different numbers of cells which produce IFN α and IFN β in the inflamed tissue. Also, steady-state levels of IFN β can be influenced by our microbiota (28), which raises the possibility that circulating IFN β activity could correlate with components of a patient's microbiome.

The outcome of type I IFN receptor activation depends on the pathway components present in the cell and the context (e.g., other cytokines can influence the outcome of IFN receptor ligation) (29, 30). Murine data has shown that JAK1 plays an essential and non-redundant role in promoting biologic responses induced by class II cytokine receptor family members, including the receptors for type I IFNs, type II IFN, and IL-10 (31). JAK1 is required for canonical type I IFN pathway signaling. In our data, it was striking that JAK1 expression was absent in some patients' monocytes, and that this was a strong predictor of IFN response group. Also, the density of the co-expression network was highly concordant with presence or absence of JAK1, supporting major biological differences between monocytes from individuals that express vs. do not express JAK1. This pattern in which none of the monocytes studied in a given subject expression JAK1 was only observed in patients and not in controls, suggesting that a disease related factor may be contributing to this pattern, such as a cytokine signal inducing a strong transcriptional repressor for example. It is possible that the type I IFNs could contribute to this

process, given that this pattern was associated with IFN ratio in our study.

In addition to JAK1, the IFNAR1, IFI27, PKR, and TNFA transcripts were differentially expressed between the response groups. IFNAR1 expression, but not IFNAR2, was enriched in NC Mo of participants in the IFN β/α 0 or >1.3 group. IFNAR1 functions in general as a heterodimer with IFNAR2, and, canonical signaling through the type I IFN receptor requires JAK1 and results in expression of interferon stimulated genes (ISGs), including IFI27 and PKR. Cell-surface IFNAR1 is a limiting factor for assembly of the functional type I IFN receptor complex. Intriguingly, IFNAR1 can form an active IFNB receptor by itself and activate signaling that does not involve activation of the Jak/STAT pathway (32). Among the transcripts uniquely upregulated by IFNAR1-IFNB signaling was *TNFA*, which in our study, together with IFNAR1, was increased in NC monocytes of participants in the IFN β/α 0 or >1.3 group. The pattern of gene expression that differed between the participant groups could suggest that canonical type I IFN pathway signaling is increased in peripheral blood CL monocytes of RA patients who are likely to respond to TNFi, whereas Jak/STAT-independent IFNB-IFNAR1 signaling is increased in NC monocytes of those who are not likely to respond to TNFi. It is possible that this pattern could be exploited therapeutically in those who would not be likely to respond to TNFi.

Key Messages

In this study we measured gene expression in single monocytes from seropositive RA patients prior to biologic treatment, in subjects with high vs. low IFN β/α ratio which would be predictive of treatment response. The data suggest large differences in monocyte gene expression between the two type I IFN ratio groups, supporting downstream effects upon a critical effector cell population. Interestingly, JAK1 expression was a major predictor of IFN ratio response group,

and also was observed to be completely lacking in some patient's monocytes. Our data suggest differences in IFN β / α ratio may skew canonical vs. non-canonical IFN pathway activation in RA patient monocytes.

Competing Interests

Dr. Niewold holds research grants from EMD Serono and Microdrop, Inc. which are not related to this study. Drs. Niewold and Wampler Muskardin have filed a patent on biomarker testing for the prediction of drug response in rheumatoid arthritis.

References

1. L. Carmona, M. Cross, B. Williams, M. Lassere, L. March, Rheumatoid arthritis. *Best Pract Res Clin Rheumatol* **24**, 733-745 (2010).
2. M. Boers, Understanding the window of opportunity concept in early rheumatoid arthritis. *Arthritis Rheum* **48**, 1771-1774 (2003).
3. J. S. Smolen *et al.*, Treating rheumatoid arthritis to target: recommendations of an international task force. *Ann Rheum Dis* **69**, 631-637 (2010).
4. K. S. Upchurch, J. Kay, Evolution of treatment for rheumatoid arthritis. *Rheumatology (Oxford)* **51 Suppl 6**, vi28-36 (2012).
5. J. M. Gwinnutt *et al.*, Twenty-Year Outcome and Association Between Early Treatment and Mortality and Disability in an Inception Cohort of Patients With Rheumatoid Arthritis: Results From the Norfolk Arthritis Register. *Arthritis Rheumatol* **69**, 1566-1575 (2017).
6. K. Puolakka *et al.*, Early suppression of disease activity is essential for maintenance of work capacity in patients with recent-onset rheumatoid arthritis: five-year experience from the FIN-RACo trial. *Arthritis Rheum* **52**, 36-41 (2005).
7. I. Filipovic, D. Walker, F. Forster, A. S. Curry, Quantifying the economic burden of productivity loss in rheumatoid arthritis. *Rheumatology (Oxford)* **50**, 1083-1090 (2011).
8. T. Sokka *et al.*, Work disability remains a major problem in rheumatoid arthritis in the 2000s: data from 32 countries in the QUEST-RA study. *Arthritis Res Ther* **12**, R42 (2010).
9. C. M. Lopez de Padilla, T. B. Niewold, The type I interferons: Basic concepts and clinical relevance in immune-mediated inflammatory diseases. *Gene* **576**, 14-21 (2016).
10. S. N. Kariuki *et al.*, Genetic analysis of the pathogenic molecular sub-phenotype interferon-alpha identifies multiple novel loci involved in systemic lupus erythematosus. *Genes and immunity* **16**, 15-23 (2015).
11. T. Wampler Muskardin *et al.*, Increased pretreatment serum IFN-beta/alpha ratio predicts non-response to tumour necrosis factor alpha inhibition in rheumatoid arthritis. *Annals of the rheumatic diseases* **75**, 1757-1762 (2016).
12. F. Liote, B. Boval-Boizard, D. Weill, D. Kuntz, J. L. Wautier, Blood monocyte activation in rheumatoid arthritis: increased monocyte adhesiveness, integrin expression, and cytokine release. *Clin Exp Immunol* **106**, 13-19 (1996).
13. F. M. Batliwalla *et al.*, Peripheral blood gene expression profiling in rheumatoid arthritis. *Genes Immun* **6**, 388-397 (2005).
14. S. Sharma *et al.*, Widely divergent transcriptional patterns between SLE patients of different ancestral backgrounds in sorted immune cell populations. *J Autoimmun* **60**, 51-58 (2015).
15. D. Aletaha *et al.*, 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* **69**, 1580-1588 (2010).
16. T. B. Niewold, T. L. Rivera, J. P. Buyon, M. K. Crow, Serum type I interferon activity is dependent on maternal diagnosis in anti-SSA/Ro-positive mothers of children with neonatal lupus. *Arthritis Rheum* **58**, 541-546 (2008).
17. K. A. Kirou *et al.*, Coordinate overexpression of interferon-alpha-induced genes in systemic lupus erythematosus. *Arthritis Rheum* **50**, 3958-3967 (2004).
18. C. E. Weckerle *et al.*, Network analysis of associations between serum interferon-alpha activity, autoantibodies, and clinical features in systemic lupus erythematosus. *Arthritis Rheum* **63**, 1044-1053 (2011).
19. T. B. Niewold, S. C. Wu, M. Smith, G. A. Morgan, L. M. Pachman, Familial aggregation of autoimmune disease in juvenile dermatomyositis. *Pediatrics* **127**, e1239-1246 (2011).

20. X. Feng *et al.*, Inhibition of interferon-beta responses in multiple sclerosis immune cells associated with high-dose statins. *Arch Neurol* **69**, 1303-1309 (2012).
21. T. B. Niewold, J. Hua, T. J. Lehman, J. B. Harley, M. K. Crow, High serum IFN-alpha activity is a heritable risk factor for systemic lupus erythematosus. *Genes Immun* **8**, 492-502 (2007).
22. Z. Jin *et al.*, Single-cell gene expression patterns in lupus monocytes independently indicate disease activity, interferon and therapy. *Lupus Sci Med* **4**, e000202 (2017).
23. H. Satomi, B. Wang, H. Fujisawa, F. Otsuka, Interferon-beta from melanoma cells suppresses the proliferations of melanoma cells in an autocrine manner. *Cytokine* **18**, 108-115 (2002).
24. T. Ito, H. Kanzler, O. Duramad, W. Cao, Y. J. Liu, Specialization, kinetics, and repertoire of type 1 interferon responses by human plasmacytoid dendritic cells. *Blood* **107**, 2423-2431 (2006).
25. A. Prakash, E. Smith, C. K. Lee, D. E. Levy, Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *J Biol Chem* **280**, 18651-18657 (2005).
26. M. Severa *et al.*, Sensitization to TLR7 agonist in IFN-beta-preactivated dendritic cells. *J Immunol* **178**, 6208-6216 (2007).
27. U. Nir, L. Maroteaux, B. Cohen, I. Mory, Priming affects the transcription rate of human interferon-beta 1 gene. *J Biol Chem* **260**, 14242-14247 (1985).
28. G. Weiss *et al.*, MyD88 drives the IFN-beta response to *Lactobacillus acidophilus* in dendritic cells through a mechanism involving IRF1, IRF3, and IRF7. *J Immunol* **189**, 2860-2868 (2012).
29. L. B. Ivashkiv, L. T. Donlin, Regulation of type I interferon responses. *Nat Rev Immunol* **14**, 36-49 (2014).
30. K. Fink *et al.*, IFNbeta/TNFalpha synergism induces a non-canonical STAT2/IRF9-dependent pathway triggering a novel DUOX2 NADPH oxidase-mediated airway antiviral response. *Cell Res* **23**, 673-690 (2013).
31. S. J. Rodig *et al.*, Disruption of the Jak1 gene demonstrates obligatory and nonredundant roles of the Jaks in cytokine-induced biologic responses. *Cell* **93**, 373-383 (1998).
32. N. A. de Weerd *et al.*, Structural basis of a unique interferon-beta signaling axis mediated via the receptor IFNAR1. *Nat Immunol* **14**, 901-907 (2013).

Figure Legends

Figure 1 Principal component analysis (PCA) of single monocyte (Mo) gene expression.

Plots depict the first two differentiating factors among the gene expression data of NCMo (A), and CLMo alone (B). Blue (●) denotes data from subjects in the “IFN β / α 0 or > 1.3” group. Green (●) denotes data from subjects in the “IFN β / α > 0 and \leq 1.3” group. In B, most of the monocytes from patients in the IFN β / α 0 or > 1.3 group are on one side of the Y axis; thus, the pretreatment type I IFN- β / α ratio appears to impact CLMo gene expression. Active variables (C) for CLMo PCA are shown.

Figure 2. Unsupervised hierarchical clustering of single cell pre-biologic gene expression in classical monocytes. Genes (Y-axis). Single classical monocytes (X-axis). Both genes and cells were selected for clustering. The bars under the heatmap indicate the IFN β / α group, subject, and width of data depicted that is from a single cell. Each subject is depicted by a different color. The bottom bar shows the width of data depicted that is from a single cell. See legends for relative expression level (yellow/black/blue) and IFN β / α group (black/light grey) color assignments.

Figure 3. Expression of JAK1 in single classical and non-classical monocytes. P-value by non-parametric Mann Whitney U. Top panels show each individual patient's cells in a separate column, bottom panels show cells from all patients in aggregate.

Figure 1




Figure 2




Figure 3

