Brain functional network integrity sustains cognitive function despite atrophy in presymptomatic genetic frontotemporal dementia.

Kamen A. Tsvetanov,[*,1,2] Stefano Gazzina,[*,1,3], P. Simon Jones,[1] John van Swieten,[3] Barbara Borroni,[4] Raquel Sanchez-Valle,[5] Fermin Moreno,[6, 7] Robert Laforce Jr,[8] Caroline Graff,[9] Matthis Synofzik,[10,11] Daniela Galimberti,[12,13] Mario Masellis,[14] Maria Carmela Tartaglia,[15] Elizabeth Finger,[16] Rik Vandenberghe,[17,18] Alexandre de Mendonça,[19] Fabrizio Tagliavini,[20] Isabel Santana,[21,22,23] Simon Ducharme,[24,25] Chris Butler,[26] Alexander Gerhard,[27,28] Adrian Danek,[29] Johannes Levin,[29] Markus Otto,[30] Giovanni Frisoni,[31,32] Roberta Ghidoni,[33] Sandro Sorbi,[34,35] Jonathan D. Rohrer,[36] and James B. Rowe,[1,2] on behalf of the Genetic FTD Initiative, GENFI.[#]

*Joint first authorship

Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK

Cambridge Centre for Ageing and Neuroscience (Cam-CAN), University of Cambridge and MRC Cognition and Brain Sciences Unit, Cambridge, UK

Department of Neurology, Erasmus Medical Center, Rotterdam, The Netherlands

Centre for Neurodegenerative Disorders, Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy

Alzheimer's disease and Other Cognitive Disorders Unit, Neurology Service, Hospital Clínic, Institut d'Investigacións Biomèdiques August Pi I Sunyer, University of Barcelona, Barcelona, Spain

Cognitive Disorders Unit, Department of Neurology, Hospital Universitario Donostia, San Sebastian, Gipuzkoa, Spain

Neuroscience Area, Biodonostia Health Research Insitute, San Sebastian, Gipuzkoa, Spain

Clinique Interdisciplinaire de Mémoire, Département des Sciences Neurologiques, CHU de Québec, and Faculté de Médecine, Université Laval, Québec, Canada

Karolinska Institutet, Department NVS, Center for Alzheimer Research, Division of Neurogenetics, Stockholm, Sweden

Department of Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research & Center of Neurology, University of Tübingen, Germany

German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany

University of Milan, Centro Dino Ferrari, Milan, Italy

13 Fondazione IRCSS Ca' Granda, Ospedale Maggiore Policlinico, Neurodegenerative Diseases Unit, Milan, Italy

14 LC Campbell Cognitive Neurology Research Unit, Sunnybrook Research Institute, Toronto, Ontario, Canada

15 Toronto Western Hospital, Tanz Centre for Research in Neurodegenerative Disease, Toronto, Ontario, Canada

16 Department of Clinical Neurological Sciences, University of Western Ontario, London, ON, Canada

17 Laboratory for Cognitive Neurology, Department of Neurosciences, KU Leuven, Leuven, Belgium

18 Neurology Service, University Hospitals Leuven, Belgium, Laboratory for Neurobiology, VIB-KU

19 Laboratory of Neurosciences, Institute of Molecular Medicine, Faculty of Medicine, University of Lisbon, Lisbon,
Portugal

20 Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy

21 Neurology Department, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal

22 Faculty of Medicine, University of Coimbra, Coimbra, Portugal

23 Centre of Neurosciences and Cell biology, Universidade de Coimbra, Coimbra, Portugal

24 Department of Psychiatry, McGill University Health Centre, McGill University, Montreal, Canada

25 McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University, Montreal, Canada

26 Nuffield Department of Clinical Neurosciences, Medical Sciences Division, University of Oxford, Oxford, UK

27 Division of Neuroscience and Experimental Psychology, Wolfson Molecular Imaging Centre, University
of Manchester, Manchester, UK

28 Departments of Geriatric Medicine and Nuclear Medicine, University of Duisburg-Essen, Germany

29 Neurologische Klinik und Poliklinik, Ludwig-Maximilians-Universität, Munich, German Center for Neurodegenerative
Diseases (DZNE), Munich, Germany

30 Department of Neurology, University Hospital Ulm, Ulm, Germany

31 Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia,
Italy

32 Memory Clinic and LANVIE-Laboratory of Neuroimaging of Aging, University Hospitals and University of Geneva,
Geneva, Switzerland

33 Molecular Markers Laboratory, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

34 *Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy*

35 *Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) "Don Gnocchi", Florence, Italy*

36 *Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK*

 # See Appendix for a list of GENFI consortium members.

Correspondence to:

Kamen A. Tsvetanov

e-mail: kat35@cam.ac.uk

Keywords (up to five): frontotemporal dementia (FTD), presymptomatic, functional magnetic resonance imaging (*f*MRI), network connectivity.

Abstract

INTRODUCTION: The presymptomatic phase of neurodegenerative disease can last many years, with sustained cognitive function despite progressive atrophy. We investigate this phenomenon in familial Frontotemporal dementia (FTD).

METHODS: We studied 121 presymptomatic FTD mutation carriers and 134 family members without mutations, using multivariate data-driven approach to link cognitive performance with both structural and functional magnetic resonance imaging. Atrophy and brain network connectivity were compared between groups, in relation to the time from expected symptom onset.

 RESULTS: There were group differences in brain structure and function, in the absence of differences in cognitive performance. Specifically, we identified behaviourally-relevant structural and functional network differences. Structure-function relationships were similar in both groups, but coupling between functional connectivity and cognition was stronger for carriers than for non-carriers, and increased with proximity to the expected onset of disease.

DISCUSSION: Our findings suggest that maintenance of functional network connectivity enables carriers to maintain cognitive performance.

1. Introduction

Across the adult healthy lifespan, the structural and functional properties of brain networks are coupled, and both are predictive of cognitive ability [1,2]. The connections between structure, function and performance have been influential in developing current models of ageing and neurodegeneration [3–5]. However, this work contrasts with the emerging evidence of neuropathological and structural changes many years before the onset of symptoms of Alzheimer's disease and frontotemporal dementia (FTD) [6–8]. Genetic FTD with highly-penetrant gene mutations provides the opportunity to examine the precursors of symptomatic disease. Three main genes account for 10-20% of FTD cases: chromosome 9 open reading frame 72 (*C9orf72*), granulin (*GRN*) and microtubule-associated protein tau (*MAPT*). These genes vary in their phenotypic expression and in the age of onset [9]. Despite pleiotropy [10], environmental and secondary genetic moderation [11,12] all three mutations cause significant structural brain changes in key regions over a decade before the expected age of disease onset [7,13], confirmed by longitudinal studies [14,15].

The divergence between early structural change and late cognitive decline begs the question: how do presymptomatic mutation carriers stay so well in the face of progressive atrophy? We propose that the answer lies in the maintenance of network dynamics and functional organisation [16]. Across the lifespan, functional brain network connectivity predicts cognitive status [17], and this connectivity-cognition relationship becomes stronger with age [18–20].

Our overarching hypothesis is that for those at genetic risk of dementia, the maintenance of network connectivity prevents the manifestation of symptoms despite progressive structural changes. A challenge is that neither the anatomical and functional substrates of cognition nor the targets of neurodegenerative disease are mediated by single brain regions: they are distributed across multi-level and interactive networks. We therefore used a multivariate data-driven approach to identify differences in the multidimensional brain-behaviour relationship between presymptomatic carriers and non-carriers of mutations in FTD genes. We identified key brain networks [21] from a large independent population-based age-matched dataset [22].

We tested three key hypotheses: (i) presymptomatic carriers differ from non-carriers in brain structure and brain function, but not in cognitive function, (ii) brain structure and function correlate with performance in

110 both groups, but functional network indices are stronger predictors of cognition in carriers, and (iii) the
111 dependence on network integrity for maintaining cognitive functioning increases as carriers approach the
112 onset of symptoms.

2. Methods

2.1. Participants

Thirteen research sites across Europe and Canada recruited participants as part of an international
multicentre partnership, the Genetic Frontotemporal Initiative (GENFI). 313 participants had usable
structural and resting state functional magnetic resonance imaging data (MRI) [7,13]. The study was
approved by the institutional review boards for each site, and participants providing written informed
consent. Inclusion criteria included anyone over the age of 18, who is symptomatic or a an asymptomatic
first-degree relative. Five participants were excluded due to excessive head motion (see below), resulting
in 308 datasets for further analysis.

Participants were genotyped based on whether they carried a pathogenic mutation in *MAPT, GRN*
and *C9or*f72. Mutation carriers were classified as either symptomatic or presymptomatic based on
clinician evaluation. Participants were only classified as symptomatic if the clinician judged that symptoms
were present, consistent with a diagnosis of a degenerative disorder, and progressive in nature. Additional
group of controls, termed non-carriers, comprised of mutation-negative family members. In this study,
we focus on non-carriers (NC, N=134) and presymptomatic carriers (PSC, N=121). Participants and site
investigators were blinded to the research genotyping, although a minority of participants had undergone
predictive testing outwith the GENFI study. See Table 1 for demographic information and Table 2 for
behavioural, cognitive and neuropsychological information of both groups. In keeping with other GENFI
reports, the years to expected onset (EYO) were calculated as the difference between age at assessment
and mean age at onset within the family [7].

2.2. Neurocognitive assessment

Each participant completed a standard clinical assessment consisting of medical history, family history,
functional status and physical examination, in complement with collateral history from a family member or
a close friend. In the current study 13 behavioural measures of cognitive function were correlated with
neuroimaging measures. These included the Uniform Data Set [23]: the Logical Memory subtest of the
Wechsler Memory Scale-Revised with Immediate and Delayed Recall scores, Digit Span forwards and

backwards from the Wechsler Memory Scale-Revised, a Digit Symbol Task, Parts A and B of the Trail Making

Test, the short version of the Boston Naming Test, and Category Fluency (animals). Additional tests included

Letter Fluency, Wechsler Abbreviated Scale of Intelligence Block Design task, and the Mini-Mental State

Examination. Latency measures for the Trail Making Test were inverted so that higher values across all tests

reflect better performance.

146   2.3. Neuroimaging assessment

Figure 1 provides a schematic representation of imaging data processing pipeline and the analysis strategy

for linking brain-behaviour data. MRI data were acquired using 3T scanners and 1.5T where no 3T scanning

was available from various vendors, with optimised scanning protocols to maximise synchronisation across

scanners and sites [7,13]. A 3D-structural MRI was acquired on each participant using T1-weighted

Magnetic Prepared Rapid Gradient Echo (MPRAGE) sequence over at least 283s (283-462s) and had a

median isotropic resolution of 1.1mm (1-1.3mm), repetition time of 2000ms (6.6-2400), echo time of 2.9ms

(2.6-3.5ms), inversion time of 8ms (8-9ms), and field of view 256x256x208mm (192-256x192-256x192-

208mm). The co-registered T1 images were segmented to extract probabilistic maps of 6 tissue classes:

grey matter (GM), white matter (WM), cerebrospinal fluid (CSF), bone, soft tissue, and residual noise. The

native-space GM and WM images were submitted to diffeomorphic registration to create equally

represented gene-group template images [DARTEL; 24]. The templates for all tissue types were normalised

to the Montreal Neurological Institute template using a 12-parameter affine transformation. The

normalised images were smoothed using an 8-mm Gaussian kernel.

For resting state fMRI measurements, Echo-Planar Imaging (EPI) data were acquired with at least six

minutes of scanning. Analogous imaging sequences were developed by the GENFI Imaging Core team, and

used at each GENFI study site to accommodate different scanner models and field strengths. EPI data were

acquired over at least 300s (inter-quartile range 309-440) and had a median repetition time of 2200ms

(2200-3000ms), echo time of 30ms, in-plane resolution of 2.75x2.75mm (2.75-3.31 x 2.75-3.31), and slice

thickness of 3.3mm (3.0-3.3).

The imaging data were analysed using Automatic Analysis [AA 4.0, 25] pipelines and modules which called

relevant functions from SPM12 [26]. To quantify the total motion for each participant, the root mean

square volume-to-volume displacement was computed using the approach of Jenkinson et al [27].

Participants with 3.5 or more standard deviations above the group mean motion displacement were

excluded from further analysis (N = 5). To further ensure that potential group bias in head motion did not

affect later analysis of connectivity, we took three further steps: i) fMRI data was further postprocessed

using whole-brain Independent Component Analysis (ICA) of single subject time-series denoising, with

noise components selected and removed automatically using *a priori* heuristics using the ICA-based

algorithm [28], ii) postprocessing of network node time-series (see below) and iii) a subject-specific

estimate of head movement for each participant [27] included as a covariate in group-level analysis [29].

2.4. Network definition

The location of the key cortical regions in each network was identified by spatial-ICA in an independent

dataset of 298 age-matched healthy individuals from a large population-based cohort [22]. Full details

about preprocessing and node definition are described previously [30]. Four networks commonly affected

by neurodegenerative diseases including FTD [21] were identified by spatially matching to pre-existing

templates [31]. The node time-series were defined as the first principal component resulting from the

singular value decomposition of voxels in an 8-mm radius sphere, which was centred on the peak voxel for

each node [18]. Visual representation of the spatial distribution of the nodes is shown in Figure 2.

We aimed to further reduce the effects of noise confounds on functional connectivity effects of node time-

series using general linear model (GLM) [29]. This model included linear trends, expansions of realignment

parameters, as well as average signal in WM and CSF, including their derivative and quadratic regressors

from the time-courses of each node. The WM and CSF signals were created by using the average signal

across all voxels with corresponding tissue probability larger than 0.7 in associated tissue probability maps

available in SPM12. A band-pass filter (0.0078-0.1 Hz) was implemented by including a discrete cosine

transform set in the GLM. Finally, the functional connectivity (FC) between each pair of nodes was

computed using Pearson's correlation on postprocessed time-series.

2.5. Statistical analysis

2.5.1. Group differences in brain structure, function and cognition

To assess the group-differences in neuroimaging and behavioural dataset we used multiple linear

regression with a well-conditioned shrinkage regularization [32,33] and 10-Fold Cross–Validation [34]. In

the analysis of brain structure we used as independent variables the mean grey matter volume (GMV) of

the 246 brain nodes in the Brainnetome atlas [35]. The Brainnetome atlas was developed to link functional

and structural characteristics of the human brain [35] and provides a fine-grained whole brain parcellation

with a superior representation of age-related differences in brain structure compared to other cortical

parcellation schemes [36,37]. In the analysis of brain function, we used the functional connectivity between

15 nodes, which were part of the four large-scale functional networks described above. In the analysis of

cognitive function, the independent variables comprised the performance measures on the 13

neuropsychological tests performed outside of the scanner. In all three analyses the dependent variable

was the genetic status (PSC vs NC) including age as a covariate of no interest. GENFI's large-sampled cohort

was created using harmonized multi-site neuroimaging data. Although, scanning protocols were optimised

to maximise comparability across scanners and sites [7,13], different scanning platforms can introduce

systematic differences which might confound true effects of interest [38]. Therefore, in the analysis of

neuroimaging data we included scanner site and head motion as additional covariates of no interest.

2.5.2. Brain-behaviour relationships

For the brain-behaviour analysis, we adopted a two-level procedure. In the first-level analysis, we assessed

the multidimensional brain-behaviour relationships using partial least squares [39]. This analysis described

the linear relationships between the two multivariate datasets, namely neuroimaging (either GMV or FC)

and behavioural performance, by providing pairs of latent variables (Brain-LVs and Cognition-LVs) as linear

combinations of the original variables which are optimised to maximise their covariance. Namely, dataset

1 consisted of a brain feature set, which could be either grey matter volume (GMV dataset) or functional

connectivity strength between pairs of regions for each individual (FC dataset). Dataset 2 included the

performance measures on the 13 tests (i.e. Cognition dataset), as considered in the multiple linear

regression analysis of group differences in cognition. Covariates of no interest included head motion,

scanner site, gender and handedness. In addition, we also included average GMV across all 15 nodes as a

covariate of no interest in the FC-behaviour analysis to ensure that the observed effects are over and above

differences in the level of atrophy.

Next, we tested whether the identified behaviourally-relevant LVs of brain structure and function were

differentially expressed by NC and PSC as a function of expected years to onset. To this end, we performed

a second-level analysis using multiple linear regression with robust fitting algorithm as implemented in

matlab's function "fitlm.m". Independent variables included subjects' brain scores from first level PLS

(either Structure-LV or Function-LV subject scores), group information, expected years to onset and their

interaction terms (e.g. brain scores x group, brain scores x years to expected onset, etc.). The dependent

variable was subjects' cognitive scores from the first level analysis in the corresponding PLS (Cognition-LV).

Given that the interaction effects were derived from continuous variables, we tested and interpreted
interactions based on simple slope analysis and slope difference tests [40–42]. Covariates of no interest
included gender, handedness, head movement and education (Figure 1). In addition, we included average
GMV across all 15 nodes as a covariate in the FC-behaviour analysis to ensure that the observed effects are
over and above differences in the level of atrophy.

3. Results

3.1. Group differences in neuroimaging and cognitive data

Brain structure

The multiple linear regression model testing for overall group differences in grey matter volume between
PSC and NC was significant (r=.14, p=.025), reflecting expected presymptomatic differences in brain-wide
atrophy. The frontal, parietal and subcortical regions had most atrophy in PSC (Figure 3). As expected, the
group difference in grey matter volume of these regions increased as EYO decreased, see Supplementary
Materials.

Brain Function

The multiple linear regression model testing for overall group differences in functional connectivity
between PSC and NC was marginally significant (r=.12, p=.049). The pattern of connectivity indicated mainly
increased connectivity between SN-DMN and SN-FPN in presymptomatic carriers, coupled with decreased
connectivity within the networks and DMN-FPN connectivity (Figure 3).

Cognitive Function

We did not identify group differences in cognition and behaviour (r=.002, p=.807), confirming the
impression of "healthy" status among presymptomatic carriers. However, in the next section, we consider
the relationships between structure, function and cognition that underlie this maintenance of cognitive
function.

3.2. Brain-behaviour relationships

Structure-cognition

Partial least squares analysis of grey matter volume and cognition identified one significant pair of latent
variables (r = .40, p = .019). This volumetric latent variable expressed negative loadings in frontal (superior
frontal gyrus, precentral gyrus, paracentral lobule), parietal (postcentral gyrus, precuneus, superior and
inferior parietal lobule) and occipital (lateral and medial occipital cortex) regions and positive loadings in

parahippocampal and hippocampal regions in addition to inferior temporal and insular cortex (Figure 4).

The Cognition-LV profile expressed positively a large array of cognitive tests, with strongest values on

delayed memory, Trail Making, Digit Symbol, Boston Naming and Fluency tests. The positive correlation

between volumetric and cognitive LV's confirms the expected relationship across the cohort as a whole,

between cortical grey matter volume and both executive, language and mnemonic function (Figure 4).

To understand the structure-cognition relationship in each group and in relation to the expected

years of onset, we performed a second-level interaction analysis using a regression model: we entered

Cognition-LV subject scores as dependent variable, and grey matter volume LV subject scores, genetic

status (i.e. mutation carrier or non-carrier), expected years to onset and their interactions as independent

variables in addition to covariates of no interest. The results indicated that the relationship between grey

matter volume and cognition could not be explained by genetic status, expected years to onset or their

interactions with grey matter volume LV subject scores. There was no evidence for genetic status- and

onset-dependent differences (over and above ageing and other covariates) in the associations between

grey matter volume and cognition in this analysis (Figure 4).

Connectivity-Cognition

PLS analysis of functional connectivity and cognition also identified one significant pair of LVs (Function-LV

and Cognition-LV, r=.32, p=.020), see Figure 5. This Function-LV reflected weak between-network

connectivity, coupled with strong within-network connectivity. This pattern indicates the segregation or

modularity of large-scale brain networks. The Cognition-LV expressed all tests, with positive loading values

indicating that higher performance on a wide range of cognitive tests is associated with stronger functional

network segregation. Cognitive deficits were associated with loss of segregation, with increased between-

network connectivity and decreased within-network connectivity.

To further test whether the observed behaviourally-relevant pattern of connectivity is differentially

expressed between genetic status groups and expected years of onset, we constructed a second-level

regression model with robust error estimates by including Function-LV subject scores, genetic status,

expected years of onset and their interaction terms as independent variables and Cognition-LV as

dependent variable in addition to covariates of no interest (Figure 5).

We found evidence for significant interaction between expected years of onset and Function-LV (r=.21,

p<.001) and between group and Function-LV (r=.16, p=.002) explaining unique variance in Cognition-LV.

We used simple slope analysis and slope difference tests [40–42] to test formally for differences in the

relationship between Function-LV and Cognition-LV for PSC and NC. The relationship between Function-LV

and Cognition-LV was stronger for PSC relative to NC (r=.16, p=.002), indicating the increasing importance

of functional connectivity between the large-scale networks for PSC participants to maintain performance

(Figure 5).

For ease of interpretation and illustration, we also computed the correlation between Cognition-LV and

Function-LV for high and low levels of expected years to onset (EYO) within each group separately, where

the levels were taken to be 1 standard deviation above and below the mean values of EYO following the

simple slopes approach [40–42]. The two EYO subgroups were labelled "near" and "far", with "near" for

EYO values close to zero (i.e. participant's age is "near" the age at which disease symptoms were

demonstrated in the family), and "far" for EYO being a largely negative value (i.e. participant's age is "far"

from the age at which disease symptoms were demonstrated in the family). The analysis indicated that as

the EYO decreases (i.e. participant's age is reaching the years of onset of symptoms) the relationship

between functional connectivity and performance becomes stronger. This effect was highly significant in

presymptomatic carriers (r=.31, p<.001) and tended towards significance in non-carriers (r=.12, p=.038,

one-sided). The differences in effects between presymptomatic carriers and non-carriers was qualified by

a significant interaction term (t=2.27, p=0.024, i.e. the effect in presymptomatic mutation carriers was

statistically stronger than the effect detected in non-carriers). These findings indicate that the relationship

between FC and cognition is stronger in PSC relative to NC, and that this relationship increases as a function

of EYO.

4. Discussion

In the present study, we confirmed previous findings of group differences in brain structure and function,

in the absence of differences in cognitive performance between non-carriers and presymptomatic carriers

of FTD-related genetic mutations. But, while the relationship between structure and cognition was similar

in both groups, the coupling between function and cognition was stronger for presymptomatic carriers,

and increased as they approached the expected onset of disease.

These results suggest that people can maintain good cognitive abilities and successful day-to-day

functioning despite significant neuronal loss and atrophy. This disjunction between structure and function

is a feature of healthy ageing, but we have shown that it also characterises presymptomatic FTD, over and

above the age effects in their other family members, despite widespread progressive atrophy. The
multivariate approach reveals two key findings: (i) presymptomatic carriers express stronger between-
network and weaker within-network functional connectivity than age-matched non-carriers, and (ii) as
carriers approach their estimated age of symptom onset, and atrophy becomes evident, the maintenance
of good cognition is increasingly associated with sustaining balance of within- and between-network
integration.

This balance of within- and between-network connectivity is characteristic of segregated and specialized
network organization of brain systems. Such functional segregation varies with physiological ageing
[17,18,43], with cognitive function [18] and in individuals at risk for Alzheimer's disease [44]. Graph-
theoretic quantification of network organisation confirms the relevance of modularity and efficiency to
function in FTD [16]. Conversely, the loss of neural systems' modularity mirrors the loss of functional
specialization with age [45] and dementia [44]. Here, we show the significance of the maintenance of this
functional network organisation, with a progressively stronger correlation with cognitive performance as
seemingly healthy adults approach the age of expected onset of FTD.

The uncoupling of brain function from brain structure indicates that there may be independent and
synergistic effects of multiple factors leading to cognitive preservation. This is consistent with a previous
work in healthy ageing where brain activity and connectivity provide independent and synergistic
predictions of performance across the lifespan [19]. Therefore, future studies need to consider the
independent and synergistic effects of many possible biomarkers, based on MRI, computed tomography,
positron-emission tomography, CSF, blood and brain histopathology. For example, functional network
impairment may be related to tau expression and tau pathology, amyloid load, or neurotransmitter deficits
in neurodegenerative diseases, independent of atrophy [30,46–48]. Importantly, studies need to recognise
the rich multivariate nature of cognition and of neuroimaging in order to improve stratification procedures,
e.g. based on integrative approaches that explain individual differences in cognitive impairment [30,49].
On a clinical level, this may facilitate future studies to establish whether presymptomatic carriers who
maintain such connectivity profiles and thereby neuropsychological function in the presence of atrophy
may have a lower risk of progression and better prognosis – information which will be important for future
triallists, patients and carers.

We also recognise the difficulty to determine a unique contribution of each factor (e.g. brain structure and
brain function), given the increasing interaction between factors in advanced stages of disease [50]. This is
further complicated by these alterations becoming irreversible with progression of neurodegeneration

[51]. This suggests that the critical interplay between multiple factors (including brain structure and
function) may be better studied in the asymptomatic and preclinical stages as well as across the healthy
lifespan, which could still be modifiable and their influences are likely to be more separable.

Our findings agree with the model of compensation in the presymptomatic and early phases of
Huntington's disease, where network coupling predicted better cognitive performance [52]. In a recent
longitudinal study a non-linear concave-down pattern of both brain activity and behaviour was present,
despite a linear decline in brain volume over time, [53]. Similar effects have been observed also in healthy
ageing and amnestic mild cognitive impairment, where greater connectivity with the default-mode network
and weaker connectivity between default-mode network and dorsal-attention network was associated with
higher cognitive status in both groups [54]. Network integrity may also play a role in compensatory
mechanisms in non-cognitive symptoms, such as motor impairment in Parkinson's disease [55].
Accordingly, increased network efficiency and connectivity has been shown in prodromal phases, followed
by decreased local connectivity in symptomatic phases, suggesting the emergence and dissipation of neural
compensation [56].

The current study has several limitations. First, despite the large size of the overall GENFI cohort, we did
not analyse each genetic group separately. The subdivision of each clinical group (PSC, NC) by three genes
would have led to small and unbalanced subgroups, lowering statistical power and robustness. Moreover,
genetic FTD is also characterised by multiple mutations within MAPT and GRN, and pleiotropy of clinical
phenotypes from the same mutation [10]. Pleiotropy of clinical phenotype is avoided by the study of
presymptomatic carriers, but we cannot rule out pleiotropy of intermediate phenotypes expressed as say
neural network diversity. In FTD as in other dementias, clinical heterogeneity is modified by environmental
factors such as education [which may be a surrogate of cognitive reserve, 12,57]. In addition, our analysis
included the estimated age of onset in some models, but we recognise that the precision of the estimated
years of onset (based on family history of onset) varies across mutations and families [7,58], being highest
for MAPT and low for C9ORF72 expansion. Genetic modifiers such as TMEM106B [59], APOE [60], have also
been identified. Further work, with larger cohorts is required to test for gene-specific effects, and the role
of environmental and genetic moderators on the relationships between brain structure, functional
networks and cognition. The harmonisation of sequences and data acquisition protocols in this multi-site
neuroimaging study aimed to reduce the susceptibility to systematic differences across scanning platforms,
but residual site variance cannot be ruled out [38,61]. The inclusion of study site as a covariate of no interest
[61] and the nature of our multivariate approach to identify shared signals between brain and behavioural

data reduce residual effects of scanner variance [38,62]. Future studies may use alternative brain measures
that reflect differences in cortical surface and thickness estimates [63,64], or which infer neural
connectivity directly from neurophysiology or from the separation of neurovascular from neuronal
contributors to BOLD fMRI variance [18,65], given the confounding effects of age, drug or disease on
neurovascular signals [66,67].

The current study is cross-sectional. Therefore, we cannot infer longitudinal progression within subjects as
the unambiguous cause of the effects we observe in relation to expected years of onset. Accumulating
evidence suggests that network integrity serves to maintain performance with either physiological ageing
or pathological conditions. However, longitudinal mediation studies and pharmacological or electroceutical
interventions would be needed to prove its causal role in cognitive preservation. Finally, our findings are
limited to autosomal dominant FTD, which represents a minority of FTD: generalisation to sporadic forms
of disease would be speculative.

In conclusion, we used a multivariate data-driven approach to demonstrate that brain functional integrity
may facilitate presymptomatic carriers to maintain cognitive performance in the presence of progressive
brain atrophy for years before the onset of symptoms. The multivariate approach to cognition and brain
function is well-suited to address the effects of multiple interacting risk factors on biomarkers of the
progression of neurodegeneration, ahead of clinical conversion to dementia. The approach and our findings
have implications for the design of presymptomatic disease-modifying therapy trials, which are likely to
rely initially on surrogate markers of brain health rather than clinical endpoints.

5. References

[1] Persson J, Nyberg L, Lind J, Larsson A, Nilsson LG, Ingvar M, et al. Structure-function correlates of cognitive decline in aging. CerebCortex 2006;16:907–15.

[2] Geerligs L, Cam-CAN, Henson RN. Functional connectivity and structural covariance between regions of interest can be measured more accurately using multivariate distance correlation. Neuroimage 2016;135:16–31. https://doi.org/10.1016/j.neuroimage.2016.04.047.

[3] Cope TE, Rittman T, Borchert RJ, Jones PS, Vatansever D, Allinson K, et al. Tau burden and the functional connectome in Alzheimer's disease and progressive supranuclear palsy. Brain 2018;141:550–67. https://doi.org/10.1093/brain/awx347.

[4] Seeley WW, Crawford RK, Zhou J, Miller BL, Greicius MD. Neurodegenerative diseases target large-scale human brain networks. Neuron 2009;62:42–52. https://doi.org/10.1016/j.neuron.2009.03.024.

[5] Raj A, Kuceyeski A, Weiner M. A network diffusion model of disease progression in dementia. Neuron 2012;73:1204–15. https://doi.org/10.1016/j.neuron.2011.12.040.

[6] Kinnunen KM, Cash DM, Poole T, Frost C, Benzinger TLS, Ahsan RL, et al. Presymptomatic atrophy in autosomal dominant Alzheimer's disease: A serial magnetic resonance imaging study. Alzheimer's Dement 2018;14:43–53. https://doi.org/10.1016/j.jalz.2017.06.2268.

[7] Rohrer JD, Nicholas JM, Cash DM, van Swieten J, Dopper E, Jiskoot L, et al. Presymptomatic cognitive and neuroanatomical changes in genetic frontotemporal dementia in the Genetic Frontotemporal dementia Initiative (GENFI) study: a cross-sectional analysis. Lancet Neurol 2015;14:253–62. https://doi.org/10.1016/S1474-4422(14)70324-2.

[8] Vatsavayai SC, Yoon SJ, Gardner RC, Gendron TF, Vargas JNS, Trujillo A, et al. Timing and significance of pathological features in *C9orf72* expansion-associated frontotemporal dementia. Brain 2016;139:3202–16. https://doi.org/10.1093/brain/aww250.

[9] Deleon J, Miller BL. Frontotemporal dementia, 2018, p. 409–30. https://doi.org/10.1016/B978-0-

427   444-64076-5.00027-2.

428   [10]   Snowden JS, Adams J, Harris J, Thompson JC, Rollinson S, Richardson A, et al. Distinct clinical and
429   pathological phenotypes in frontotemporal dementia associated with MAPT, PGRN and C9orf72
430   mutations. Amyotroph Lateral Scler Front Degener 2015;16:497–505.
431   https://doi.org/10.3109/21678421.2015.1074700.

432   [11]   Murphy NA, Arthur KC, Tienari PJ, Houlden H, Chiò A, Traynor BJ. Age-related penetrance of the
433   C9orf72 repeat expansion. Sci Rep 2017;7:2116. https://doi.org/10.1038/s41598-017-02364-1.

434   [12]   Premi E, Grassi M, Van Swieten J, Galimberti D, Graff C, Masellis M, et al. Cognitive reserve and
435   TMEM106B genotype modulate brain damage in presymptomatic frontotemporal dementia: a
436   GENFI study. Brain 2017;140:1784–91. https://doi.org/10.1093/brain/awx103.

437   [13]   Cash DM, Bocchetta M, Thomas DL, Dick KM, van Swieten JC, Borroni B, et al. Patterns of gray
438   matter atrophy in genetic frontotemporal dementia: results from the GENFI study. Neurobiol Aging
439   2018;62:191–6. https://doi.org/10.1016/j.neurobiolaging.2017.10.008.

440   [14]   Olm CA, McMillan CT, Irwin DJ, Van Deerlin VM, Cook PA, Gee JC, et al. Longitudinal structural gray
441   matter and white matter MRI changes in presymptomatic progranulin mutation carriers.
442   NeuroImage Clin 2018;19:497–506. https://doi.org/10.1016/J.NICL.2018.05.017.

443   [15]   Floeter MK, Danielian LE, Braun LE, Wu T. Longitudinal diffusion imaging across the *C9orf72* clinical
444   spectrum. J Neurol Neurosurg Psychiatry 2018;89:53–60. https://doi.org/10.1136/jnnp-2017-
445   316799.

446   [16]   Rittman DT, Borchert MR, Jones MS, van Swieten J, Borroni B, Galimberti D, et al. Functional
447   network resilience to pathology in presymptomatic genetic frontotemporal dementia. Neurobiol
448   Aging 2019;77:169–77. https://doi.org/10.1016/J.NEUROBIOLAGING.2018.12.009.

449   [17]   Chan MY, Park DC, Savalia NK, Petersen SE, Wig GS. Decreased segregation of brain systems across
450   the healthy adult lifespan. Proc Natl Acad Sci 2014;111:4997–5006.
451   https://doi.org/10.1073/pnas.1415122111.

452   [18]   Tsvetanov KA, Henson RNA, Tyler LK, Razi A, Geerligs L, Ham TE, et al. Extrinsic and intrinsic brain
453   network connectivity maintains cognition across the lifespan despite accelerated decay of regional
454   brain activation. J Neurosci 2016;36:3115–26. https://doi.org/10.1523/JNEUROSCI.2733-15.2016.

[19] Tsvetanov KA, Ye Z, Hughes L, Samu D, Treder MS, Wolpe N, et al. Activity and connectivity
differences underlying inhibitory control across the adult lifespan. J Neurosci 2018;38:7887–900.
https://doi.org/10.1523/JNEUROSCI.2919-17.2018.

[20] Bethlehem RAI, Paquola C, Seidlitz J, Ronan L, Bernhardt B, Consortium C-C, et al. Dispersion of
functional gradients across the adult lifespan. Neuroimage 2020:117299.
https://doi.org/10.1016/j.neuroimage.2020.117299.

[21] Chhatwal JP, Schultz AP, Johnson KA, Hedden T, Jaimes S, Benzinger TLS, et al. Preferential
degradation of cognitive networks differentiates Alzheimer's disease from ageing. Brain
2018;141:1486–500. https://doi.org/10.1093/brain/awy053.

[22] Shafto MA, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, et al. The Cambridge Centre for Ageing
and Neuroscience (Cam-CAN) study protocol: a cross-sectional, lifespan, multidisciplinary
examination of healthy cognitive ageing. BMC Neurol 2014;14:204.
https://doi.org/10.1186/s12883-014-0204-1.

[23] Morris JC, Weintraub S, Chui HC, Cummings J, DeCarli C, Ferris S, et al. The Uniform Data Set (UDS):
Clinical and Cognitive Variables and Descriptive Data From Alzheimer Disease Centers. Alzheimer
Dis Assoc Disord 2006;20:210–6. https://doi.org/10.1097/01.wad.0000213865.09806.92.

[24] Ashburner J. A fast diffeomorphic image registration algorithm. Neuroimage 2007;38:95–113.
https://doi.org/10.1016/j.neuroimage.2007.07.007.

[25] Cusack R, Vicente-Grabovetsky A, Mitchell DJ, Wild CJ, Auer T, Linke AC, et al. Automatic analysis
(aa): efficient neuroimaging workflows and parallel processing using Matlab and XML. Front
Neuroinform 2014;8:90. https://doi.org/10.3389/fninf.2014.00090.

[26] Friston KJ, Ashburner J, Kiebel S, Nichols T, Penny WD. Statistical parametric mapping : the analysis
of funtional brain images. Elsevier Academic Press; 2007.

[27] Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust and accurate
linear registration and motion correction of brain images. Neuroimage 2002;17:825–41.

[28] Pruim RHR, Mennes M, Buitelaar JK, Beckmann CF. Evaluation of ICA-AROMA and alternative
strategies for motion artifact removal in resting state fMRI. Neuroimage 2015;112:278–87.
https://doi.org/10.1016/j.neuroimage.2015.02.063.

[29] Geerligs L, Tsvetanov KA, Cam-Can, Henson RN. Challenges in measuring individual differences in
functional connectivity using fMRI: The case of healthy aging. Hum Brain Mapp 2017.
https://doi.org/10.1002/hbm.23653.

[30] Passamonti L, Tsvetanov KA, Jones PS, Bevan-Jones WR, Arnold R, Borchert RJ, et al.
Neuroinflammation and functional connectivity in Alzheimer's disease: interactive influences on
cognitive performance. J Neurosci 2019;39:2574–18. https://doi.org/10.1523/jneurosci.2574-
18.2019.

[31] Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding subject-driven cognitive states
with whole-brain connectivity patterns. Cereb Cortex 2012;22:158–65.
https://doi.org/10.1093/cercor/bhr099.

[32] Blankertz B, Lemm S, Treder M, Haufe S, Müller K-R. Single-trial analysis and classification of ERP
components--a tutorial. Neuroimage 2011;56:814–25.
https://doi.org/10.1016/j.neuroimage.2010.06.048.

[33] Ledoit O, Wolf M. A well-conditioned estimator for large-dimensional covariance matrices. J
Multivar Anal 2004;88:365–411. https://doi.org/10.1016/S0047-259X(03)00096-4.

[34] Lemm S, Blankertz B, Dickhaus T, Müller K-R. Introduction to machine learning for brain imaging.
Neuroimage 2011;56:387–99. https://doi.org/10.1016/j.neuroimage.2010.11.004.

[35] Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, et al. The Human Brainnetome Atlas: A New Brain Atlas
Based on Connectional Architecture. Cereb Cortex 2016;26:3508–26.
https://doi.org/10.1093/cercor/bhw157.

[36] Madan CR, Kensinger EA. Predicting age from cortical structure across the lifespan. Eur J Neurosci
2018;47:399–416. https://doi.org/10.1111/ejn.13835.

[37] Long Z, Huang J, Li B, Li Z, Li Z, Chen H, et al. A Comparative Atlas-Based Recognition of Mild
Cognitive Impairment With Voxel-Based Morphometry. Front Neurosci 2018;12:916.
https://doi.org/10.3389/fnins.2018.00916.

[38] Chen J, Liu J, Calhoun VD, Arias-Vasquez A, Zwiers MP, Gupta CN, et al. Exploration of scanning
effects in multi-site structural MRI studies. J Neurosci Methods 2014;230:37–50.
https://doi.org/10.1016/j.jneumeth.2014.04.023.

[39] Krishnan A, Williams LJ, McIntosh AR, Abdi H. Partial Least Squares (PLS) methods for
neuroimaging: A tutorial and review. Neuroimage 2011;56:455–75.
https://doi.org/10.1016/j.neuroimage.2010.07.034.

[40] Aiken LS, West SG. Multiple regression: Testing and interpreting interactions. Thousand Oaks, CA,
US: Sage Publications, Inc; 1991.

[41] Dawson JF, Richter AW. Probing three-way interactions in moderated multiple regression:
Development and application of a slope difference test. J Appl Psychol 2006;91:917–26.
https://doi.org/10.1037/0021-9010.91.4.917.

[42] Dawson JF. Moderation in Management Research: What, Why, When, and How. J Bus Psychol
2014;29:1–19. https://doi.org/10.1007/s10869-013-9308-7.

[43] Samu D, Campbell KL, Tsvetanov KA, Shafto MA, Consortium C-C, Brayne C, et al. Preserved
cognitive functions with age are determined by domain-dependent shifts in network responsivity.
Nat Commun 2017;8:ncomms14743. https://doi.org/10.1038/ncomms14743.

[44] Contreras JA, Goñi J, Risacher SL, Amico E, Yoder K, Dzemidzic M, et al. Cognitive complaints in
older adults at risk for Alzheimer's disease are associated with altered resting-state networks.
Alzheimer's Dement Diagnosis, Assess Dis Monit 2017;6:40–9.
https://doi.org/10.1016/J.DADM.2016.12.004.

[45] Cabeza R, Albert M, Belleville S, Craik FIM, Duarte A, Grady CL, et al. Maintenance, reserve and
compensation: the cognitive neuroscience of healthy ageing. Nat Rev Neurosci 2018;19:701–10.
https://doi.org/10.1038/s41583-018-0068-2.

[46] Hedden T, Van Dijk KRA, Becker JA, Mehta A, Sperling RA, Johnson KA, et al. Disruption of functional
connectivity in clinically normal older adults harboring amyloid burden. J Neurosci 2009;29:12686–
94. https://doi.org/10.1523/JNEUROSCI.3189-09.2009.

[47] Murley AG, Rowe JB. Neurotransmitter deficits from frontotemporal lobar degeneration. Brain
2018;141:1263–85. https://doi.org/10.1093/brain/awx327.

[48] Rittman T, Rubinov M, Vértes PE, Patel AX, Ginestet CE, Ghosh BCP, et al. Regional expression of
the MAPT gene is associated with loss of hubs in brain networks and cognitive impairment in
Parkinson disease and progressive supranuclear palsy. Neurobiol Aging 2016;48:153–60.

539 https://doi.org/10.1016/J.NEUROBIOLAGING.2016.09.001.

540 [49] Geerligs L, Tsvetanov KA. The use of resting state data in an integrative approach to studying
541 neurocognitive ageing – Commentary on Campbell and Schacter (2016). Lang Cogn Neurosci
542 2016;32. https://doi.org/http://dx.doi.org/10.1080/23273798.2016.1251600.

543 [50] Vemuri P, Lesnick TG, Przybelski SA, Knopman DS, Preboske GM, Kantarci K, et al. Vascular and
544 amyloid pathologies are independent predictors of cognitive decline in normal elderly. Brain
545 2015;138:761–71. https://doi.org/10.1093/brain/awu393.

546 [51] Rodrigue KM, Kennedy KM, Devous MD, Rieck JR, Hebrank AC, Diaz-Arrastia R, et al. β-Amyloid
547 burden in healthy aging: regional distribution and cognitive consequences. Neurology
548 2012;78:387–95. https://doi.org/10.1212/WNL.0b013e318245d295.

549 [52] Klöppel S, Gregory S, Scheller E, Minkova L, Razi A, Durr A, et al. Compensation in Preclinical
550 Huntington's Disease: Evidence From the Track-On HD Study. EBioMedicine 2015;2:1420–9.
551 https://doi.org/10.1016/j.ebiom.2015.08.002.

552 [53] Gregory S, Long JD, Klöppel S, Razi A, Scheller E, Minkova L, et al. Operationalizing compensation
553 over time in neurodegenerative disease. Brain 2017;140:1158–65.
554 https://doi.org/10.1093/brain/awx022.

555 [54] Sullivan MD, Anderson JAE, Turner GR, Spreng RN. Intrinsic neurocognitive network connectivity
556 differences between normal aging and mild cognitive impairment are associated with cognitive
557 status and age. Neurobiol Aging 2019;73:219–28.
558 https://doi.org/10.1016/J.NEUROBIOLAGING.2018.10.001.

559 [55] Blesa J, Trigo-Damas I, Dileone M, del Rey NL-G, Hernandez LF, Obeso JA. Compensatory
560 mechanisms in Parkinson's disease: Circuits adaptations and role in disease modification. Exp
561 Neurol 2017;298:148–61. https://doi.org/10.1016/J.EXPNEUROL.2017.10.002.

562 [56] Wen M-C, Heng HSE, Hsu J-L, Xu Z, Liew GM, Au WL, et al. Structural connectome alterations in
563 prodromal and de novo Parkinson's disease patients. Parkinsonism Relat Disord 2017;45:21–7.
564 https://doi.org/10.1016/j.parkreldis.2017.09.019.

565 [57] Stern Y. Cognitive reserve in ageing and Alzheimer's disease. Lancet Neurol 2012;11:1006–12.
566 https://doi.org/10.1016/S1474-4422(12)70191-6.

[58] Moore KM, Nicholas J, Grossman M, McMillan CT, Irwin DJ, Massimo L, et al. Age at symptom onset
and death and disease duration in genetic frontotemporal dementia: an international
retrospective cohort study. Lancet Neurol 2020;19:145–56. https://doi.org/10.1016/S1474-
4422(19)30394-1.

[59] Lattante S, Le Ber I, Galimberti D, Serpente M, Rivaud-Péchoux S, Camuzat A, et al. Defining the
association of TMEM106B variants among frontotemporal lobar degeneration patients with GRN
mutations and C9orf72 repeat expansions. Neurobiol Aging 2014;35:2658.e1-2658.e5.
https://doi.org/10.1016/J.NEUROBIOLAGING.2014.06.023.

[60] Koriath C, Kenny J, Adamson G, Druyeh R, Taylor W, Beck J, et al. Predictors for a dementia gene
mutation based on gene-panel next-generation sequencing of a large dementia referral series. Mol
Psychiatry 2018. https://doi.org/10.1038/s41380-018-0224-0.

[61] Alfaro-Almagro F, McCarthy P, Afyouni S, Andersson JLR, Bastiani M, Miller KL, et al. Confound
modelling in UK Biobank brain imaging. Neuroimage 2020:117002.
https://doi.org/10.1016/j.neuroimage.2020.117002.

[62] Li H, Smith SM, Gruber S, Lukas SE, Silveri MM, Hill KP, et al. Denoising scanner effects from
multimodal MRI data using linked independent component analysis. Neuroimage
2020;208:116388. https://doi.org/10.1016/j.neuroimage.2019.116388.

[63] Brodoehl S, Gaser C, Dahnke R, Witte OW, Klingner CM. Surface-based analysis increases the
specificity of cortical activation patterns and connectivity results. Sci Rep 2020;10:1–13.
https://doi.org/10.1038/s41598-020-62832-z.

[64] Coalson TS, Van Essen DC, Glasser MF. The impact of traditional neuroimaging methods on the
spatial localization of cortical areas. Proc Natl Acad Sci U S A 2018;115:E6356–E6365.
https://doi.org/10.1073/pnas.1801582115.

[65] Sami S, Williams N, Hughes LE, Cope TE, Rittman T, Coyle-Gilchrist ITS, et al. Neurophysiological
signatures of Alzheimer's disease and frontotemporal lobar degeneration: pathology versus
phenotype. Brain 2018;141:2500–10. https://doi.org/10.1093/brain/awy180.

[66] Campbell KL, Shafto MA, Wright P, Tsvetanov KA, Geerligs L, Cusack R, et al. Idiosyncratic
responding during movie-watching predicted by age differences in attentional control. Neurobiol

Aging 2015;36:3045–55. https://doi.org/10.1016/j.neurobiolaging.2015.07.028.

[67]   Tsvetanov KA, Henson RNA, Rowe JB. Separating vascular and neuronal effects of age on fMRI

BOLD signals. Philos Trans R Soc B Biol Sci 2020. https://doi.org/10.1098/rstb.2019.0631.

6. Acknowledgements

K.A.T. is supported by the British Academy Postdoctoral Fellowship (PF160048) and the
Guarantors of Brain (101149). J.B.R. is supported by the Wellcome Trust (103838) the Medical Research
Council (SUAG/051 G101400) and the Cambridge NIHR Biomedical Research Centre. R. S.-V. is supported
by the Instituto de Salud Carlos III and the JPND network PreFrontAls (01ED1512/AC14/0013) and the
Fundació Marató de TV3 (20143810). M.M and E.F are supported by the UK Medical Research Council, the
Italian Ministry of Health and the Canadian Institutes of Health Research as part of a Centres of Excellence
in Neurodegeneration grant, and also a Canadian Institutes of Health Research operating grant (MOP
327387) and funding from the Weston Brain Institute. J.D.R., D.C. and K.M.M. are supported by the NIHR
Queen Square Dementia Biomedical Research Unit, the NIHR UCL/H Biomedical Research Centre and the
Leonard Wolfson Experimental Neurology Centre (LWENC) Clinical Research Facility. J.D.R. is supported
by an MRC Clinician Scientist Fellowship (MR/M008525/1) and has received funding from the NIHR Rare
Disease Translational Research Collaboration (BRC149/NS/MH), the MRC UK GENFI grant (MR/
M023664/1) and The Bluefield Project. F.T. is supported by the Italian Ministry of Health (Grant NET-2011-
02346784). L.C.J. and J.V.S. are supported by the Association for Frontotemporal Dementias Research
Grant 2009, ZonMw Memorabel project number 733050103 and 733050813, and the Bluefield project.
R.G. supported by Italian Ministry of Health, Ricerca Corrente. The Swedish contributors C.G., L.O. and
C.A. were supported by grants from JPND Prefrontals Swedish Research Council (VR) 529-2014-7504,
Swedish Research Council (VR) 2015- 02926, Swedish Research Council (VR) 2018-02754, Swedish FTD
Initiative-Schorling Foundation, Swedish Brain Foundation, Swedish Alzheimer Foundation, Stockholm
County Council ALF, Karolinska Institutet Doctoral Funding and StratNeuro, Swedish Demensfonden,
during the conduct of the study.

7. Tables

Table 1. Demographics of participants included in the analysis, grouped by genetic status as non-carriers (NC) and presymptomatic carriers (PSC). * denotes whether demographics vary between NC and PSC groups.

	Gene Status Group		Statistical tests*	
	NC	PSC	X^2 or F-test	*P*-value
N	134	121		
Mutated gene, n (%)			0.86	0.649
MAPT	17 (12.7)	19 (15.7)		
GRN	77 (57.5)	63 (52.1)		
C9Orf72	40 (29.9)	39 (32.2)		
Gender, n (%)			0.01	0.908
Male	53 (39.6)	47 (38.8)		
Handedness, n (%)			0.06	0.806
Right-handed	122 (91)	107 (88.4)		
Age (Years)			2.68	0.103
Mean / SD	49 / 14	46 / 11		
Range [Min/Max]	19 / 86	20 / 70		
Expected Years to Onset			0.23	0.631
Mean / SD	-10 / 12	-11 / 11		
Range [Min/Max]	-25 / 10	-25 / 10		
Education (Years)			0.05	0.826
Mean / SD	14 / 3	14 / 3		
Range [Min/Max]	5 / 24	5 / 22		

* Statistical test to indicate whether demographics vary between NC and PSC groups

632 *Table 2. Behavioural, cognitive and neuropsychological estimates in presymptomatic carriers and non-carriers*

	Gene Status Group		Statistical tests*	
	NC	PSC	χ^2	*P*-value
Behavioural				
Cambridge Behavioural Inventory—Revised (/180)	3.5 (5.4)	4.7 (10)	0.03	0.864
Cognitive				
Mini-Mental State Examination	29.3 (1.1)	29.2 (1.3)	< 0.01	0.963
Neuropsychological				
Logical Memory—Immediate Recall	15.2 (5.6)	15.7 (5.6)	0.47	0.495
Logical Memory—Delayed Recall	14.1 (4.7)	14 (5)	0.97	0.356
Digit Span - Forwards	6.4 (1.2)	6.3 (1.3)	0.52	0.470
Digit Span - Backwards	4.9 (1.2)	4.8 (1.2)	1.62	0.203
Digit Symbol Task	32 (14.1)	35 (14)	0.35	0.556
Trail Making Test Part A	28.9 (17.2)	28.9 (11.5)	0.97	0.325
Trail Making Test Part B	72.5 (43.7)	72.3 (45.5)	0.02	0.895
Verbal Fluency - Letter	42 (12.2)	40.7 (15.1)	0.95	0.330
Verbal Fluency - Animal	23.3 (6)	23.7 (5.8)	0.58	0.445
Boston Naming Test	28.1 (2.1)	27.6 (2.7)	0.58	0.446
Block Design	41.8 (16.1)	42.5 (17.1)	0.17	0.683

633   * Statistical test to indicate whether scores vary between NC and PSC groups

8. Figures

Figure 1. Schematic representation of data processing and analysis pipeline to test for brain-behaviour differences between presymptomatic carriers (PSC) and non-carriers (NC) as a function of expected years to onset (EYO) of symptoms, while controlling for covariates of no interest (Covs). Brain structural measures were based on the mean grey matter volume (GMV) in 246 nodes, as defined in the Brainnetome atlas [35]. Brain functional measures were based on the functional connectivity between 15 nodes as part of four large-scale networks, which were defined in an independent cohort of 298 age-matched individuals part of the Cam-CAN dataset.

Figure 2. Visualisation of spatial localisation of the nodes part of the four large-scale networks and their mean functional connectivity (circular plot) across all participants in this study. Nodes and networks were defined in an independent cohort of 298 age-matched individuals part of the Cam-CAN dataset [30].*The default mode network (DMN) contained five nodes: the ventral anterior cingulate cortex (vACC), dorsal and ventral posterior cingulate cortex (vPCC and dPCC), and right and left inferior parietal lobes (rIPL and lIPL). The salience network (SN) was defined using right and left anterior insular (rAI and lAI) and dorsal anterior cingulate cortex (dACC). The frontoparietal network (FPN) was defined using right and left anterior superior frontal gyrus (raSFG and laSFG), and right and left angular gyrus (rAG and lAG). The dorsal attention Network (DAN) was defined using right and left intraparietal sulcus (rIPS and lIPS).*

659　　　　*Figure 3. Group differences between PSC and NC in grey matter volume (left panel) and*
660　*functional connectivity between nodes within four large scale networks (right panel). Hot colour*
661　*scheme indicates the strength of effect size of PSC showing higher GMV and FC than NC, while cold*
662　*colour scheme indicates the opposite effect (i.e. NC > PSC).*

Figure 4. PLS analysis of grey matter volume (GMV) and cognition indicating the spatial distribution of GMV loading values (a), where hot and cold colour schemes are used for the strength of positive and negative correlations with the profile of Cognitive LV (b). (c) The scatter plot on the left represents the relationship between subjects scores of GMV LV and Cognition LV for presymptomatic carriers (PSC) and non-carriers (NC). The scatter plots in the middle and right hand-side represent GMV-Cognition LV relationship as a function of expected years to onset (EYO, split in two groups, Near and Far, see text) in each genetic status group separately.

Figure 5. *PLS analysis of functional connectivity and cognition indicating the connectivity pattern of loading values (a), where hot and cold colour schemes are used for the strength of positive and negative correlations with the profile of Cognitive LV (b). (c) The scatter plot on the left represents the relationship between subjects scores of Function LV and Cognition LV for presymptomatic carriers (PSC) and non-carriers (NC). The scatter plots in the middle and right hand-side represents Function-Cognition LV relationship* as a function of expected years to onset *(EYO split in two groups, Near and Far, see text) in each genetic status group separately. This is also represented using a bar chart in (d), where continuous and dashed lines indicate significance of effect differences and difference in differences, respectively. † and * denote significant tests at p-value < 0.05 (one- and two-sided, respectively).*

9. APPENDIX

List of other GENFI consortium members

Sónia Afonso - Instituto Ciencias Nucleares Aplicadas a Saude, Universidade de Coimbra, Coimbra, Portugal

Maria Rosario Almeida - Centre of Neurosciences and Cell Biology, Universidade de Coimbra, Coimbra, Portugal

Sarah Anderl-Straub – Department of Neurology, Ulm University, Ulm, GermanyChristin Andersson - Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

Anna Antonell - Alzheimer's disease and other cognitive disorders unit, Neurology Department, Hospital Clinic, Institut d'Investigacions Biomèdiques, Barcelona, Spain

Silvana Archetti - Biotechnology Laboratory, Department of Diagnostics, Spedali Civili Hospital, Brescia, Italy

Andrea Arighi - Fondazione IRCSS Ca' Granda, Ospedale Maggiore Policlinico, Neurodegenerative Diseases Unit, Milan, Italy

Mircea Balasa - Alzheimer's disease and other cognitive disorders unit, Neurology Department, Hospital Clinic, Institut d'Investigacions Biomèdiques, Barcelona, Spain

Myriam Barandiaran - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San Sebastian, Gipuzkoa, Spain

Nuria Bargalló - Radiology Department, Image Diagnosis Center, Hospital Clínic and Magnetic Resonance Image core facility, IDIBAPS, Barcelona, Spain

Robart Bartha - Department of Medical Biophysics, Robarts Research Institute, University of Western Ontario, London, Ontario, Canada

Benjamin Bender - Department of Diagnostic and Interventional Neuroradiology, University of Tuebingen, Tuebingen, Germany

Luisa Benussi - Istituto di Ricovero e Cura a Carattere Scientifico Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

Valentina Bessi - Department of Neuroscience, Psychology, Drug Research, and Child Health, University of Florence, Florence, Italy

Giuliano Binetti - Istituto di Ricovero e Cura a Carattere Scientifico Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

Sandra Black - LC Campbell Cognitive Neurology Research Unit, Sunnybrook Research Institute, Toronto, Canada

Martina Bocchetta – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square London, UK

Sergi Borrego-Ecija - Alzheimer's disease and other cognitive disorders unit, Neurology Department, Hospital Clinic, Institut d'Investigacions Biomèdiques, Barcelona, Spain

Jose Bras – Dementia Research Institute, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Rose Bruffaerts - Laboratory for Cognitive Neurology, Department of Neurosciences, KU Leuven, Leuven, Belgium

Paola Caroppo - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy

David Cash – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Miguel Castelo-Branco - Neurology Department, Centro Hospitalar e Universitário de Coimbra, Instituto de Ciências Nucleares Aplicadas à Saúde (ICNAS), Coimbra, Portugal

Rhian Convery – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Thomas Cope – Department of Clinical Neuroscience, University of Cambridge, Cambridge, UK

Maura Cosseddu – Centre for Neurodegenerative Disorders, Neurology Unit, Spedali Civili Hospital, Brescia, Italy

María de Arriba - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San Sebastian, Gipuzkoa, Spain

Giuseppe Di Fede - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy

Zigor Díaz - CITA Alzheimer, San Sebastian, Spain

Katrina M Moore – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Diana Duro - Faculty of Medicine, Universidade de Coimbra, Coimbra, Portugal

Chiara Fenoglio - University of Milan, Centro Dino Ferrari, Milan, Italy

Camilla Ferrari - Department of Neuroscience, Psychology, Drug Research, and Child Health, University of Florence, Florence, Italy

Carlos Ferreira - Instituto Ciências Nucleares Aplicadas à Saúde, Universidade de Coimbra, Coimbra, Portugal

Catarina B. Ferreira - Faculty of Medicine, University of Lisbon, Lisbon, Portugal

Toby Flanagan – Faculty of Biology, Medicine and Health, Division of Neuroscience and Experimental Psychology, University of Manchester, Manchester, UK

Nick Fox – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Morris Freedman - Division of Neurology, Baycrest Centre for Geriatric Care, University of Toronto, Toronto, Canada

Giorgio Fumagalli - Fondazione IRCSS Ca' Granda, Ospedale Maggiore Policlinico, Neurodegenerative Diseases Unit, Milan, Italy; Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy

Alazne Gabilondo - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San Sebastian, Gipuzkoa, Spain

Roberto Gasparotti - Neuroradiology Unit, University of Brescia, Brescia, Italy

Serge Gauthier - Department of Neurology and Neurosurgery, McGill University, Montreal, Québec, Canada

Stefano Gazzina - Centre for Neurodegenerative Disorders, Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy

Giorgio Giaccone - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy

Ana Gorostidi - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San Sebastian, Gipuzkoa, Spain

Caroline Greaves – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square London, UK

Rita Guerreiro – Dementia Research Institute, Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK

Carolin Heller – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Tobias Hoegen - Department of Neurology, Ludwig-Maximilians-University of Munich, Munich, Germany

Begoña Indakoetxea - Cognitive Disorders Unit, Department of Neurology, Donostia University Hospital, Paseo Dr Begiristain sn, CP 20014, San Sebastian, Gipuzkoa, Spain

Vesna Jelic - Division of Clinical Geriatrics, Karolinska Institutet, Stockholm, Sweden

Lize Jiskoot - Department of Neurology, Erasmus Medical Center, Rotterdam, The Netherlands

Hans-Otto Karnath - Section of Neuropsychology, Department of Cognitive Neurology, Center for Neurology & Hertie-Institute for Clinical Brain Research, Tübingen, Germany

Ron Keren - University Health Network Memory Clinic, Toronto Western Hospital, Toronto, Canada

Maria João Leitão - Centre of Neurosciences and Cell Biology, Universidade de Coimbra, Coimbra, Portugal

Albert Lladó - Alzheimer's disease and other cognitive disorders unit, Neurology Department, Hospital Clinic, Institut d'Investigacions Biomèdiques, Barcelona, Spain

Gemma Lombardi - Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Florence, Italy

Sandra Loosli - Department of Neurology, Ludwig-Maximilians-University of Munich, Munich, Germany

Carolina Maruta - Lisbon Faculty of Medicine, Language Research Laboratory, Lisbon, Portugal

Simon Mead - MRC Prion Unit, Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

Lieke Meeter - Department of Neurology, Erasmus Medical Center, Rotterdam, Netherlands

Gabriel Miltenberger - Faculty of Medicine, University of Lisbon, Lisbon, Portugal

Rick van Minkelen - Department of Clinical Genetics, Erasmus Medical Center, Rotterdam, The Netherlands

Sara Mitchell - LC Campbell Cognitive Neurology Research Unit, Sunnybrook Research Institute, Toronto, Canada

Benedetta Nacmias - Department of Neuroscience, Psychology, Drug Research and Child Health, University of
Florence, Florence, Italy
Mollie Neason - Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology,
Queen Square, London, UK
Jennifer Nicholas – Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK
Linn Öijerstedt - Department of Geriatric Medicine, Karolinska Institutet, Stockholm, Sweden
Jaume Olives - Alzheimer's disease and other cognitive disorders unit, Neurology Department, Hospital Clinic, Institut
d'Investigacions Biomèdiques, Barcelona, Spain
Alessandro Padovani - Centre for Neurodegenerative Disorders, Neurology Unit, Department of Clinical and
Experimental Sciences, University of Brescia, Brescia, Italy
Jessica Panman – Department of Neurology, Erasmus Medical Center, Rotterdam, The Netherlands
Janne Papma - Department of Neurology, Erasmus Medical Center, Rotterdam, The Netherlands
Irene Piaceri - Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence,
Florence
Michela Pievani - Istituto di Ricovero e Cura a Carattere Scientifico Istituto Centro San Giovanni di Dio
Fatebenefratelli, Brescia, Italy
Yolande Pijnenburg - VUMC, Amsterdam, The Netherlands
Cristina Polito - Department of Biomedical, Experimental and Clinical Sciences "Mario Serio", Nuclear Medicine Unit,
University of Florence, Florence, Italy
Enrico Premi - Stroke Unit, Neurology Unit, Spedali Civili Hospital, Brescia, Italy
Sara Prioni - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan,
Italy
Catharina Prix - Department of Neurology, Ludwig-Maximilians-University Munich, Germany
Rosa Rademakers - Department of Neurosciences, Mayo Clinic, Jacksonville, Florida, USA
Veronica Redaelli - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta,
Milan, Italy
Tim Rittman – Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK
Ekaterina Rogaeva - Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto,
Canada
Pedro Rosa-Neto - Translational Neuroimaging Laboratory, McGill University Montreal, Québec, Canada
Giacomina Rossi - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta,
Milan, Italy
Martin Rossor – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology,
Queen Square, London, UK
Beatriz Santiago - Neurology Department, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal
Elio Scarpini - University of Milan, Centro Dino Ferrari, Milan, Italy; Fondazione IRCSS Ca' Granda, Ospedale Maggiore
Policlinico, Neurodegenerative Diseases Unit, Milan, Italy
Sonja Schönecker - Neurologische Klinik, Ludwig-Maximilians-Universität München, Munich, Germany
Elisa Semler – Department of Neurology, Ulm University, Ulm, Germany
Rachelle Shafei – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology,
Queen Square, London, UK
Christen Shoesmith - Department of Clinical Neurological Sciences, University of Western Ontario, London, Ontario,
Canada
Miguel Tábuas-Pereira - Centre of Neurosciences and Cell Biology, Universidade de Coimbra, Coimbra, Portugal
Mikel Tainta - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San
Sebastian, Gipuzkoa, Spain
Ricardo Taipa - Neuropathology Unit and Department of Neurology, Centro Hospitalar do Porto - Hospital de Santo
António, Oporto, Portugal
David Tang-Wai - University Health Network Memory Clinic, Toronto Western Hospital, Toronto, Canada
David L Thomas - Neuroradiological Academic Unit, UCL Institute of Neurology, London, UK
Hakan Thonberg - Center for Alzheimer Research, Division of Neurogeriatrics, Karolinska Institutet, Stockholm,
Sweden
Carolyn Timberlake - University of Cambridge, Cambridge, UK

Pietro Tiraboschi - Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta,
Milano, Italy
Philip Vandamme - Neurology Service, University Hospitals Leuven, Belgium; Laboratory for Neurobiology, VIB-KU
Leuven Centre for Brain Research, Leuven, Belgium
Mathieu Vandenbulcke - Geriatric Psychiatry Service, University Hospitals Leuven, Belgium; Neuropsychiatry,
Department of Neurosciences, KU Leuven, Leuven, Belgium
Michele Veldsman - University of Oxford, UK
Ana Verdelho - Department of Neurosciences, Santa Maria Hospital, University of Lisbon, Portugal
Jorge Villanua - OSATEK Unidad de Donostia, San Sebastian, Gipuzkoa, Spain
Jason Warren – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of Neurology,
Queen Square, London, UK
Carlo Wilke - Hertie Institute for Clinical Brain Research, University of Tuebingen, Tuebingen, Germany
Ione Woollacott – Dementia Research Centre, Department of Neurodegenerative Disease, UCL Institute of
Neurology, Queen Square, London, UK
Elisabeth Wlasich - Neurologische Klinik, Ludwig-Maximilians-Universität München, Munich, Germany
Henrik Zetterberg - Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK
Miren Zulaica - Neuroscience Area, Biodonostia Health Research Institute, Paseo Dr Begiristain sn, CP 20014, San
Sebastian, Gipuzkoa, Spain.
860