Title: Projected Impact of Concurrently Available Long-Acting Injectable and Daily-Oral HIV Pre-Exposure

Prophylaxis: A Mathematical Model

Running Head: Projected Impact of Injectable PrEP

Authors: Kevin M. Maloney[1*], Adrien Le Guillou[1,2], Robert A. Driggers[1§], Supriya Sarkar[1,3], Emeli J.

Anderson[1], Amyn A. Malik[1,4,5], Samuel M. Jenness[1]

[1]Department of Epidemiology, Emory University, Atlanta, GA, USA

[2]Department of Research and Public Health, Reims Teaching Hospitals, Robert Debré Hospital, Reims,

France

[3]ViiV Healthcare, Brentford, UK

[4]Yale Institute for Global Health, New Haven, CT

[5]Yale School of Medicine, New Haven, CT

§ *Deceased*

Word Count: Main text, **3,498** words (3,500 limit); Abstract, **200** words (200 limit)

ABSTRACT

Background: Long-acting injectable HIV pre-exposure prophylaxis (LAI-PrEP) is reportedly efficacious,

although full trial results have not been published. We used a dynamic network model of HIV transmission

among men who have sex with men (MSM) to assess the population impact of LAI-PrEP when available

concurrently with daily-oral (DO) PrEP.

Methods: We calibrated our reference model to the current HIV epidemiology and DO-PrEP coverage

(15% among indicated) among MSM in the southeastern US. Primary analyses investigated varied PrEP

uptake and proportion selecting LAI-PrEP. Secondary analyses evaluated uncertainty in pharmacokinetic

efficacy and LAI-PrEP persistence relative to DO-PrEP.

Results: Compared to the reference scenario, if 50% chose LAI-PrEP, 4.3% (95% SI: -7.3%, 14.5%) of

infections would be averted over 10 years. LAI-PrEP impact is slightly greater than the DO-PrEP only

regimen from higher adherence and partial protection after discontinuation. If the total PrEP initiation rate

doubled, 17.1% (95% SI: 6.7%, 26.4%) of infections would be averted. The highest population-level

impact occurred when LAI-PrEP uptake and persistence improved.

Conclusions: If LAI-PrEP replaces DO-PrEP, its availability will modestly improve the population impact.

LAI-PrEP will make a more substantial impact if its availability drives higher total PrEP coverage, or if

persistence if greater among LAI-PrEP users.

KEY WORDS

HIV; Men who have sex with men; Mathematical model; Preexposure prophylaxis; Sexual network

BACKGROUND

Men who have sex with men (MSM) remain at increased risk for HIV infection in the United

States, accounting for two-thirds of all new infections in 2018 [1]. Pre-exposure prophylaxis

(PrEP) is a highly effective method to reduce HIV acquisition [2, 3]. However, implementation

challenges undermine the potential for PrEP to reduce incidence at the population level. For

example, uptake of PrEP remains low [4-7]. The US Centers for Disease Control and Prevention

(CDC) estimate that 1.1 million US adults were behaviorally indicated to use PrEP in 2017 [8],

but only 100,000 accessed PrEP [4]. Efforts to increase PrEP use are undermined by poor

persistence, resulting in early discontinuation among individuals with ongoing HIV risk [9-11].

Previous PrEP models have predicted that current coverage levels are insufficient to reach

many HIV prevention targets [12, 13]. Among MSM accessing PrEP, inadequate adherence

further reduces its population-level prevention benefits [10, 14].

Alternative PrEP products that are more acceptable and easier to use for some may

address these implementation challenges. Daily-oral PrEP (DO-PrEP) is currently the only

dosing strategy recommended by the CDC and the US Food and Drug Administration [15].

However, alternative dosing regimens [16] have proven efficacious; novel formulations are in

various stages of development, including long-acting injectable [17] and topical formulations

[18], implants and other devices [19]. Cabotegravir is an investigational antiretroviral (strand-

transfer integrase inhibitor) [17]. When formulated as a long-acting injectable suspension (LAI-

PrEP), cabotegravir is maintained with periodic dosing at 8-week intervals [17]. The Phase III

clinical trial for LAI-PrEP efficacy among MSM (HPTN-083) has been discontinued early with

fewer incident infections observed among MSM randomized to LAI-PrEP compared to DO-PrEP

[20], although full trial outcomes have yet to be released.

LAI-PrEP may address adherence challenges with DO-PrEP because the pill burden

would be eliminated [14, 17]. However, the clinical burden for LAI-PrEP may be greater for

some because of the need for intramuscular injections administered by a healthcare

professional at 8-week intervals (compared to 12-week intervals for clinical monitoring of DO-

PrEP) [15]. LAI-PrEP may be more acceptable for users unwilling or unable to take a daily pill

[21-25]. Estimates of preference for LAI-PrEP range from 30% to 67% [21, 23, 25-28], but real-

world selection of LAI-PrEP over DO-PrEP is unknown. Inferring population-level benefits from

these hypothetical LAI-PrEP preferences depends on the interaction of uptake and persistence.

Mathematical modeling provides insights into the mechanisms by which LAI-PrEP may

differ from DO-PrEP under these conditions of uncertainty. A recent study [29] simulated the

population impact when either DO-PrEP or LAI-PrEP were allocated to MSM. The study found

that in scenarios with LAI-PrEP there was a greater decline in incidence than in scenarios with

DO-PrEP. However, the impact when LAI-PrEP is available as an additional option to DO-PrEP

remains unclear. It was also unclear how challenges to persistence will limit the population

impact of LAI-PrEP which, unlike DO-PrEP, provides continued but waning protection even after

discontinuation [17]. Since publication of this model, pharmacokinetic data from a Phase II

clinical trial of LAI-PrEP has become available [17], which can be used to specify model

parameters.

To estimate the potential population-level impact of LAI-PrEP, we used a network-based

mathematical model of HIV transmission dynamics among MSM in the southeastern US. To

understand how concurrent availability of LAI-PrEP and DO-PrEP could maximize HIV

prevention, we simulated scenarios varying overall PrEP uptake and the proportion choosing

LAI-PrEP versus DO-PrEP. We also conducted sensitivity analyses of the LAI-PrEP biological

parameters (pharmacokinetic and efficacy) and behavioral parameters (discontinuation rate

compared to DO-PrEP).

METHODS

We used a mathematical model of HIV transmission dynamics among black, white, and

Hispanic MSM, aged 15–65 years, in the Atlanta area over a 10-year period. The model was

simulated using *EpiModel* [30], a software platform for modeling disease dynamics in sexual

networks using temporal exponential random graph models (TERGMs) [31]. Building on past

PrEP models [12], we developed the model framework and parameterization for LAI-PrEP to

explore differences with DO-PrEP. Full methodological details can be found in the Web

Appendix.

Sexual Network Model. All network and behavioral parameters were drawn from the ARTnet

study, a 2017–2019 egocentric network study of 4,904 MSM reporting on 16,198 sexual

partnerships, including duration, concurrency and other attributes [32]. The model was initialized

with 10,000 MSM, but the network size varied based on entry (sexual debut) and exit (mortality

or aging out at 65). Summary statistics from ARTnet were fit to statistical models under the

TERGM framework to simulate one-off, casual, and main partnerships. Partnership formation

was determined stochastically by partnership type, age and race mixing, number of ongoing

partnerships, and sorting by receptive and insertive positioning. Relational dissolution was

modeled with a constant hazard based on the median duration for partnership type.

For MSM with HIV, rates were assigned for diagnosis, initiation of antiretroviral therapy

(ART), and viral suppression [33, 34]. HIV disease progressed through the acute, chronic and

AIDS stages. HIV viral loads were modeled continuously, in the absence of ART. ART

decreased HIV viral loads, with a corresponding decrease in the probability of HIV transmission

[35, 36]. Within serodiscordant sexual partnerships, HIV transmission occurred stochastically

with a per-act probability which was modified based on PrEP use [14], condom use [37], sexual

positioning [38], and circumcision status of the insertive partner [39].

PrEP Uptake, Adherence, and Persistence. Uptake of PrEP was determined using a two-step

process. Parameters for LAI-PrEP and DO-PrEP are shown in **Table 1**. First, we specified the

overall probability of initiating PrEP. Second, we specified the probability of choosing LAI-PrEP

versus DO-PrEP. By simulating uptake and selection using separate probabilities, we were able

to explore counterfactual scenarios regarding how user decisions interact. Eligibility to initiate

PrEP of either formulation was determined based on CDC guidelines for DO-PrEP indications

[12, 15]. For MSM using DO-PrEP, we assigned a probability distribution for medication

adherence corresponding to low (< 2 doses/week, 8.9%), medium (2–3 doses/week, 12.7%),

and high adherence (≥ 4 doses/week, 78.4%) [40], which modified the level of protection against

HIV acquisition (hazard ratios: 0.69, 0.19, and 0.02, respectively) [14]. PrEP users were

screened for HIV every 90 days according to CDC guidelines [15].

Based on CDC guidelines, behavioral indications were reassessed annually; if

indications had lapsed, PrEP would be discontinued for both DO-PrEP and LAI-PrEP [15]. We

also allowed for spontaneous discontinuation of DO-PrEP to occur based on an observational

study showing 57% of PrEP patients remained in care 6 months after initiation [41]. We

translated this into a constant probability of discontinuation assuming a geometric distribution to

calculate the median time to stoppage (224 days) [41]. It is unknown whether MSM using LAI-

PrEP would have the same persistence as has been observed among MSM using DO-PrEP, so

our base LAI-PrEP conditions assumed that the rate of spontaneous discontinuation of LAI-

PrEP would be the same as the rate assigned to DO-PrEP.

Injectable PrEP Pharmacokinetics and Efficacy. Based on studies of nonhuman primates, the

target cabotegravir plasma concentration in clinical trials was set to 4-times the protein-adjusted

90% inhibitory concentration (PA-IC$_{90}$) [42-44]. A Phase IIa clinical trial showed that 600-mg

intramuscular injections at 8-week intervals were sufficient to maintain the target plasma

concentration [17]. For simplicity, we assumed that men achieved the target peak plasma

concentration following the first injection (in clinical trials, the second injection was administered

after 4-weeks to initially achieve the target concentration) and without an initial oral safety

phase.

Using data from the Phase IIa trial to model the pharmacokinetics of injectable

cabotegravir [17], we estimated the median peak plasma concentration (3.59 µg/ml) following

each injection after and including the second injection, as well as the rate of decay in

longitudinal follow-up. We then estimated a continuous half-life function (35 days) of drug

elimination based on these two parameters. This estimate shows that plasma concentrations

above PA-IC_{90} and 4-times PA-IC_{90} are maintained for 22 and 12 weeks, respectively, with

limited drug still present 52 weeks after the final injection.

The Phase III trial results for cabotegravir as LAI-PrEP have not been published. Thus,

we used the results of a macaque study to model the relative reduction in the per-exposure

probability of HIV infection [42]. We used logistic regression to estimate the probability of

infection given the plasma concentration of cabotegravir at the time of exposure. We then

transformed the results to the human equivalent of the PA-IC_{90} to obtain a single parameter for

the relative reduction in infection probability given the plasma concentration of cabotegravir. We

estimated that LAI-PrEP is greater than 99% effective at 4-times PA-IC_{90} (HR = 0.002).

Reference Model and Counterfactual Scenarios. The reference model represents current uptake

of DO-PrEP, in which approximately 15% of indicated MSM in the southeastern US use PrEP

[45-47]. We calibrated the model to maintain a steady state prevalence of DO-PrEP use among

indicated MSM. To reach 15% DO-PrEP, the probability of initiating PrEP given indications was

set to 0.411% per time-step, with LAI-PrEP selection set to 0%.

Our primary analysis explored scenarios of overall PrEP uptake by simultaneously

varying the probability of PrEP initiation (supplementing current DO-PrEP uptake by relative

factors of 0 to 6 times the base parameter) and the probability of selecting LAI-PrEP (replacing

current DO-PrEP uptake from 0% to 100%). The probability of selecting LAI-PrEP was fixed at

50% for all secondary analyses, while the probability of initiating PrEP was fixed at the

reference model value. Second, to better understand the pharmacological properties of LAI-

PrEP, we varied the pharmacokinetic and efficacy parameters. We varied the LAI-PrEP

pharmacokinetics by simultaneously modifying the peak plasma concentration after injection

(1.5 to 6.5 µg/ml) and the half-life of drug elimination (15 to 60 days). Separately, we varied the

efficacy of LAI-PrEP by scaling the per-exposure hazard ratio for transmission (5%, 10%, 25%,

and 50% increase or decrease in efficacy compared to the base parameter). Finally, to

understand how the effectiveness of LAI-PrEP could depend on continued use during periods of

sexual risk, we varied the rate of spontaneous discontinuation. For this, we modified the median

time using LAI-PrEP from 33% (approximately 75 days) to 300% (672 days) and calculated new

spontaneous discontinuation rates.

Calibration, Simulation, and Analysis. The network was initialized with 10,000 MSM and

calibrated to the estimated prevalence of diagnosed HIV (33% for black MSM; 27% for Hispanic

MSM; and 8.4% for white MSM) [48]. Each model scenario was simulated over 10 years and for

250 times. The outcomes reported were HIV prevalence and incidence per 100 person-years at

risk at the end of 10 years, and the percent of infections averted (PIA). The PIA was calculated

by comparing the cumulative incidence in each counterfactual scenario to the reference model.

We summarized the results using the median values of all simulations and 95% simulation

intervals (SI).

RESULTS

In the reference scenario, 15.1% (95% SI: 14.4%, 15.9%) of indicated MSM were using DO-

PrEP (0% LAI-PrEP). At the end of the 10-year simulation, the HIV incidence was 1.19 (95% SI:

0.93, 1.50) per 100 person-years and prevalence was 22.4% (95% SI: 21.5%, 23.4%).

Increasing both the probability of initiating PrEP (supplementing current DO-PrEP use) and the

probability of selecting LAI-PrEP (replacing current DO-PrEP use) would result in fewer

infections over 10 years, compared to the reference scenario (**Figure 1**). The benefit of

introducing LAI-PrEP is greater in scenarios where overall uptake of PrEP also increased. Small

improvements, however, would still be realized at lower proportions of MSM initiating PrEP and

selecting LAI-PrEP.

Increasing either or both the initiation and the LAI-PrEP selection probabilities will

increase the proportion of indicated MSM protected from PrEP, which in turn causes HIV

incidence to decline (**Table 2**, with additional scenarios in **Supplemental Table 14**). With PrEP

limited to DO-PrEP, increasing the probability of initiating PrEP by two-fold resulted in a greater

proportion of indicated MSM using DO-PrEP (26.7%; 95% SI: 25.8%, 27.5%) and 11.3% (95%

SI: 0.1%, 21.1%) of infections averted. Alternatively, when the probability of selecting LAI-PrEP

was increased to 50% and the PrEP initiation rate was held constant, the proportion of MSM

using DO-PrEP dropped to 7.6% (95% SI: 7.1%, 8.2%) while LAI-PrEP use increased to 8.4%

(95% SI: 7.9%, 9.0%). In addition to the MSM actively using LAI-PrEP, 7.3% (95% SI: 6.8%,

7.7%) of indicated MSM had waning plasma concentration and thus partial protection after

discontinuing LAI-PrEP. This scenario resulted in a modest decrease in incidence (4.3% of

infections averted; 95% SI: -7.3%, 14.5%). While the median result indicates an improvement,

the lower bound of the 95% SI shows that incidence increased for many of the scenarios with

LAI-PrEP, relative to the median value of the reference model; similarly, the upper bound of the

interval shows that a larger decrease in incidence is possible.

If 100% of the MSM accessing PrEP were to select LAI-PrEP, and initiation remained at

the reference level, then 16.8% (95% SI: 16.1%, 17.5%) would have full protection from LAI-

PrEP and 13.9% (95% SI: 13.2%, 14.6%) would have waning protection, resulting in 7.5% of

infections averted (95% SI: -3.5%, 18.7%). Finally, if the probability of initiating PrEP increased

two-fold and the proportion choosing LAI-PrEP was 50%, then incidence would decrease with

17.1% (95% SI: 6.7%, 26.4%) of infections averted.

The underlying estimates for LAI-PrEP pharmacokinetics are sensitive to uncertain

parameter values (**Figure 2**, **Table 3**, and **Supplemental Table 15**). If the peak plasma

concentration were reduced to 2 µg/ml and the half-life were reduced to 15 days (an extreme

scenario, well below the *a priori* clinical trial targets), then the modest effect of introducing LAI-

PrEP at 50% selection probability (4.3% of infections averted; 95% SI: -7.3%, 14.5%) would be

almost eliminated (1.4% of infections averted; 95% SI: -11.5%, 11.9%). When parameters were

modified within a smaller range around our estimates (e.g., 3 µg/ml peak plasma concentration

and 25 days half-life) then the impact would be limited (3.1% of infections averted; 95% SI: -

10.4%, 13.5%). Similarly, the population impact of LAI-PrEP is sensitive to our estimate of the

drug efficacy (**Figure 3**, **Table 3**, and **Supplemental Table 16**). For example, a 50% decrease

in LAI-PrEP efficacy resulted in increased incidence with -4.4% (95% SI: -17.4%, 8.4%) of

infections averted. However, a 10% decrease in efficacy would result in a more modest decline

in the population benefit of introducing LAI-PrEP (2.4% of infections averted; 95% SI: -9.7%,

13.8%).

Changes to duration of persistence on LAI-PrEP, relative to the base model duration

based on current estimates of DO-PrEP persistence, also impacted population-level impact

(**Figure 4**, **Table 3**, and **Supplementary Table 17**). In scenarios where persistence improved

for MSM accessing LAI-PrEP, the prevention benefit of introducing LAI-PrEP increased. For

example, if the median time to discontinuation were 448 days (versus 224 days for the base

parameter), then active LAI-PrEP use would increase from 8.4% (95% SI: 7.9%, 9.0%) to 13.6%

(SI: 12.9%, 14.4%) with a predicted 7.9% (95% SI: -3.0%, 19.6%) of infections averted

(compared to 4.3% in the base model; 95% SI: -7.3%, 14.5%). Alternatively, as persistence

decreased, the prevalence of LAI-PrEP use among indicated MSM also decreased; the

proportion with waning protection from LAI-PrEP increased negligibly because overall uptake of

PrEP was fixed. When the rate of discontinuation was doubled (median time = 112 days), the

predicted prevention benefit declined to 1.8% of infections averted, with the 95% SI (-11.7%,

13.6%) showing that many simulations represented null findings or increased incidence.

DISCUSSION

In this study, we projected that LAI-PrEP concurrently available with DO-PrEP would modestly

reduce HIV incidence among MSM compared to offering only DO-PrEP. The benefit of replacing

some DO-PrEP use with LAI-PrEP is due to the elimination of challenges associated with daily

adherence to DO-PrEP and the partial protection provided by the long metabolic half-life of LAI-

PrEP even after discontinuation. The benefit of LAI-PrEP would increase if uptake supplements

current DO-PrEP uptake. To our knowledge, this is the first modeling study to predict the

population impact of LAI-PrEP when available concurrently with DO-PrEP.

249   The introduction of LAI-PrEP could yield small improvements to the prevention benefit of

PrEP if uptake replaces current DO-PrEP use, but our model predicts even greater benefit if

LAI-PrEP acts to supplement current uptake of DO-PrEP. We designed our model to explore

PrEP uptake using two components: total PrEP coverage and cross-sectional mix of DO-PrEP

and LAI-PrEP. It is unknown how many MSM currently using DO-PrEP will switch to LAI-PrEP

once available; however, future preference estimates range from 35–67% would switch to LAI-

PrEP [25, 28]. Low uptake has been a limiting factor to the population benefit of DO-PrEP [4-6].

Similar to other studies [12, 13], we found that large-scale reductions in HIV incidence will

require greater uptake of PrEP among MSM than has currently been observed in the US [4-6].

This may be achieved if the availability of LAI-PrEP increases the proportion of indicated

individuals accessing PrEP. Although 66% of PrEP inexperienced MSM indicate interest in LAI-

PrEP [27], translating interest to actual uptake has proven difficult for DO-PrEP.

261   In our model, the prevention benefits of LAI-PrEP were greater when persistence

increased above the levels currently observed for DO-PrEP. Our finding is in direct contrast to

an earlier LAI-PrEP modeling study, which counterintuitively reported lower HIV incidence as

LAI-PrEP persistence decreased [29]. However, that model assumed PrEP was allocated to

maintain a constant prevalence of active users in the population so that a new LAI-PrEP user

immediately replaced one who had discontinued. As a result, the total person-time on LAI-PrEP

would artificially increase in scenarios with lower persistence, through the increased number of

individuals with waning protection after discontinuation. This model assumption is unrealistic,

and the corresponding relationship between discontinuation and LAI-PrEP impact implausible.

In our model, we disentangled these two components of PrEP prevalence (initiation and

persistence) in order to evaluate the individual contribution of each factor. We therefore varied

LAI-PrEP persistence without artificially inflating the person-time on LAI-PrEP, isolating the

effects of initiation and persistence. Our model shows that improving persistence would

increase total PrEP coverage in the population and thus decrease HIV incidence, even if the

overall PrEP initiation rate remained constant.

Persistence will likely remain a challenge for all persons accessing PrEP. The reasons

for discontinuation off DO-PrEP in practice include decreased HIV risk perception, non-

adherence to the clinical protocol for PrEP, insurance barriers and experiencing side-effects [9,

49]. The clinical procedures of LAI-PrEP may be even more challenging for some to maintain,

given more frequent clinical care visits (i.e., six annual visits for LAI-PrEP compared to four for

DO-PrEP). However, the history of contraceptive medication could serve as a guide: increased

choices improved uptake, persistence, and adherence, as women selected the modality which

most closely aligned with their preferences [50]. As formulation options increase for PrEP,

uptake, persistence and adherence may similarly improve across the board. However, when

LAI-PrEP becomes available, clinical practice guidelines should be developed to counsel

potential users on the formulation that best meets their sexual health needs.

Limitations. Our model has a number of important limitations. The first is uncertainty of the LAI-

PrEP parameters, since Phase III clinical trials have not been published. Our model parameters

may be misspecified which would cause our primary results to be biased. To accommodate this,

we tested a range of plausible parameters for LAI-PrEP pharmacokinetics and efficacy, which

had only a small effect on the model results. Our model was also a simplification of the protocol

292 for LAI-PrEP used in the Phase III clinical trial [17]. Increased complexity and clinical burden of

293 the protocol may result in lower persistence. Specifically, we did not include the 4-week oral

294 dosing phase of cabotegravir that is used to ensure safety. We also assumed that the peak

295 plasma concentration for cabotegravir would be achieved after a single intramuscular dose,

296 instead of the two initial doses at a 4-week interval. Similarly, we did not model an oral ramp-

297 down phase for individuals discontinuing LAI-PrEP under the supervision of their clinician

298 (opposed to spontaneous discontinuation). This latter phase of LAI-PrEP is used to prevent

299 infection during the waning tail of LAI-PrEP protection. Finally, we did not model resistance to

300 cabotegravir among individuals that acquire HIV infection following discontinuation of LAI-PrEP,

301 while the drug remains present but provides only partial protection. This phenomenon presents

302 a new challenge compared to DO-PrEP. Future studies will need to evaluate resistance and

303 what impact this may have on transmission dynamics, if clinical trial and observational data

304 demonstrate the occurrence of resistance.

305 *Conclusions.* The introduction of LAI-PrEP will modestly improve the prevention benefit of DO-

306 PrEP if LAI-PrEP only serves as a replacement to DO-PrEP. This is due to the non-inferiority in

307 the biological benefit of LAI-PrEP over DO-PrEP. However, LAI-PrEP could make a more

308 substantial impact on HIV incidence if the availability of LAI-PrEP increases the likelihood that

309 indicated users will start on PrEP, or decreases the likelihood that current PrEP users will

310 discontinue despite ongoing risk. Prior to the FDA approval of LAI-PrEP, public health policy

311 and implementation science on LAI-PrEP should anticipate these user behaviors as critical to

312 maximizing the benefits of this new HIV prevention tool.

ACKNOWLEDGEMENTS

We would like to acknowledge our friend and colleague, Robert A. Driggers, who contributed

substantially to the design and conduct of this study, and in authoring this paper, prior to his

unexpected death. Rob was an important member of our team and he will be missed.

DISCLOSURES

S.S. is an employee of ViiV Healthcare, which owns the patent for Cabotegravir. K.M., R.D.,

E.A., A.M., and S.J. have no conflicts to report.

AUTHORSHIP

K.M., R.D., S.S., E.A., and A.M. conceived the analysis and assisted in coding the model. K.M.

conducted the statistical analyses and drafted the manuscript. R.D. provided critical input with

K.M. to specify the pharmacokinetic properties of the LAI-PrEP model. A.L.G. assisted in coding

the model and provided input to the analysis plan. S.J. provided direction to the development of

the analysis plan, led the coding of the model, and provided critical input to the manuscript and

technical appendix. All authors reviewed and provided substantial feedback to drafts of the

manuscript.

FUNDING

This work was supported by National Institutes of Health grants R21 MH112449, R01 AI138783,

and P30 AI050409 (Emory University Center for AIDS Research).

CORRESPONDING AUTHOR:

Kevin Maloney, MPH

Department of Epidemiology

Emory University

1518 Clifton Rd.

Atlanta, GA 30322

Email: kevin.maloney@emory.edu

Phone: (404) 727-0483

Fax: (404) 712-8392

MEETINGS

A previous version of this model was presented as a poster at the International AIDS Society's

10th Conference on HIV Science (Abstract ID: TUPEC390). A pre-print of this manuscript has

been posted to medRxiv (medrxiv.org/content/10.1101/19012443v1).

SUPPLEMENTAL DIGITAL CONTACT

Maloney_Projected_Impact_of_Injectable_PrEP_Appendix_version_1.docx

REFERENCES

1. Centers for Disease Control and Prevention. Estimated HIV incidence and prevalence in the United States, 2014-2018. HIV Surveillance Supplemental Report. Vol. 25, **2020**.

2. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. N Engl J Med **2010**; 363:2587-99.

3. Spinner CD, Brunetta J, Shalit P, et al. DISCOVER study for HIV pre-exposure prophylaxis (PrEP): F/TAF has a more rapid onset and longer sustained duration of HIV protection compared with F/TDF. 10th IAS Conference on HIV Science. Mexico City, Mexico, **2019**.

4. Sullivan PS, Giler RM, Mouhanna F, et al. Trends in the use of oral emtricitabine/tenofovir disoproxil fumarate for pre-exposure prophylaxis against HIV infection, United States, 2012-2017. Ann Epidemiol **2018**; 28:833-40.

5. Finlayson T, Cha S, Xia M, et al. Changes in HIV Preexposure Prophylaxis Awareness and Use Among Men Who Have Sex with Men - 20 Urban Areas, 2014 and 2017. MMWR Morb Mortal Wkly Rep **2019**; 68:597-603.

6. Kamitani E, Wichser ME, Adegbite AH, et al. Increasing prevalence of self-reported HIV preexposure prophylaxis use in published surveys: a systematic review and meta-analysis. AIDS **2018**; 32:2633-5.

7. Parsons JT, Rendina HJ, Lassiter JM, Whitfield TH, Starks TJ, Grov C. Uptake of HIV Pre-Exposure Prophylaxis (PrEP) in a National Cohort of Gay and Bisexual Men in the United States. J Acquir Immune Defic Syndr **2017**; 74:285-92.

8. Smith DK, Van Handel M, Grey J. Estimates of adults with indications for HIV pre-exposure prophylaxis by jurisdiction, transmission risk group, and race/ethnicity, United States, 2015. Ann Epidemiol **2018**; 28:850-7 e9.

9. Krakower D, Maloney KM, Powell VE, et al. Patterns and clinical consequences of discontinuing HIV preexposure prophylaxis during primary care. J Int AIDS Soc **2019**; 22:e25250.

10. Marcus JL, Hurley LB, Hare CB, et al. Preexposure Prophylaxis for HIV Prevention in a Large Integrated Health Care System: Adherence, Renal Safety, and Discontinuation. J Acquir Immune Defic Syndr **2016**; 73:540-6.

11. Marcus JL, Hurley LB, Nguyen DP, Silverberg MJ, Volk JE. Redefining Human Immunodeficiency Virus (HIV) Preexposure Prophylaxis Failures. Clin Infect Dis **2017**; 65:1768-9.

12. Jenness SM, Goodreau SM, Rosenberg E, et al. Impact of the Centers for Disease Control's HIV Preexposure Prophylaxis Guidelines for Men Who Have Sex With Men in the United States. J Infect Dis **2016**; 214:1800-7.

13. Sullivan PS, Carballo-Dieguez A, Coates T, et al. Successes and challenges of HIV prevention in men who have sex with men. Lancet **2012**; 380:388-99.

14. Grant RM, Anderson PL, McMahan V, et al. Uptake of pre-exposure prophylaxis, sexual practices, and HIV incidence in men and transgender women who have sex with men: a cohort study. Lancet Infect Dis **2014**; 14:820-9.

15. Centers for Disease Control and Prevention. Preexposure prophylaxis for the prevention of HIV infection in the United States - 2017 update: a clinical practice guideline, **2018**.

16. Molina JM, Capitant C, Spire B, et al. On-Demand Preexposure Prophylaxis in Men at High Risk for HIV-1 Infection. N Engl J Med **2015**; 373:2237-46.

17. Landovitz RJ, Li S, Grinsztejn B, et al. Safety, tolerability, and pharmacokinetics of long-acting injectable cabotegravir in low-risk HIV-uninfected individuals: HPTN 077, a phase 2a randomized controlled trial. PLoS Med **2018**; 15:e1002690.

18. Baeten J. Topical and on-demand PrEP. 10th IAS Conference on HIV Science. Mexico City, Mexico: International AIDS Society, **2019**.

19. Flexner CW. Implants and transdermal drug delivery systems for HIV prevention. 10th IAS Conference on HIV Science. Mexico City, Mexico: International AIDS Society, **2019**.

20. Bokoch K. HIV Prevention Trials Network (HPTN). Long-acting injectable cabotegravir is highly effective for the prevention of HIV infection in cisgender men and transgender women who have sex with men. . Available at: https://www.hptn.org/news-and-events/press-releases/long-acting-injectable-cabotegravir-highly-effective-prevention-hiv.

21. Parsons JT, Rendina HJ, Whitfield TH, Grov C. Familiarity with and Preferences for Oral and Long-Acting Injectable HIV Pre-exposure Prophylaxis (PrEP) in a National Sample of Gay and Bisexual Men in the U.S. AIDS Behav **2016**; 20:1390-9.

22. Biello K, Coffey-Esquivel J, Hosek S, et al. Development of a mobile-based application to increase uptake of HIV testing among young US men who have sex with men. J Int Aids Soc **2016**; 19.

23. Greene GJ, Swann G, Fought AJ, et al. Preferences for Long-Acting Pre-exposure Prophylaxis (PrEP), Daily Oral PrEP, or Condoms for HIV Prevention Among U.S. Men Who Have Sex with Men. AIDS Behav **2017**; 21:1336-49.

24. Biello KB, Hosek S, Drucker MT, et al. Preferences for Injectable PrEP Among Young U.S. Cisgender Men and Transgender Women and Men Who Have Sex with Men. Arch Sex Behav **2018**; 47:2101-7.

25. Meyers K, Wu Y, Brill A, Sandfort T, Golub SA. To switch or not to switch: Intentions to switch to injectable PrEP among gay and bisexual men with at least twelve months oral PrEP experience. PLoS One **2018**; 13:e0200296.

26. Levy ME, Patrick R, Gamble J, et al. Willingness of community-recruited men who have sex with men in Washington, DC to use long-acting injectable HIV pre-exposure prophylaxis. PLoS One **2017**; 12:e0183521.

27. Biello KB, Mimiaga MJ, Santostefano CM, Novak DS, Mayer KH. MSM at Highest Risk for HIV Acquisition Express Greatest Interest and Preference for Injectable Antiretroviral PrEP Compared to Daily, Oral Medication. AIDS Behav **2018**; 22:1158-64.

28. John SA, Whitfield THF, Rendina HJ, Parsons JT, Grov C. Will Gay and Bisexual Men Taking Oral Pre-exposure Prophylaxis (PrEP) Switch to Long-Acting Injectable PrEP Should It Become Available? AIDS Behav **2018**; 22:1184-9.

29. Marshall BDL, Goedel WC, King MRF, et al. Potential effectiveness of long-acting injectable pre-exposure prophylaxis for HIV prevention in men who have sex with men: a modelling study. Lancet HIV **2018**; 5:e498-e505.

30. Jenness SM, Goodreau SM, Morris M. EpiModel: An R Package for Mathematical Modeling of Infectious Disease over Networks. J Stat Softw **2018**; 84.

31. Krivitsky PN, Handcock MS. A Separable Model for Dynamic Networks. J R Stat Soc Series B Stat Methodol **2014**; 76:29-46.

32. Weiss KM, Goodreau SM, Morris M, et al. Egocentric sexual networks of men who have sex with men in the United States: Results from the ARTnet study. Epidemics **2020**; 30:100386.

33. Beer L, Mattson CL, Bradley H, Shouse RL, Medical Monitoring P. Trends in ART Prescription and Viral Suppression Among HIV-Positive Young Adults in Care in the United States, 2009-2013. J Acquir Immune Defic Syndr **2017**; 76:e1-e6.

34. Beer L, Oster AM, Mattson CL, Skarbinski J, Medical Monitoring P. Disparities in HIV transmission risk among HIV-infected black and white men who have sex with men, United States, 2009. AIDS **2014**; 28:105-14.

35. Hughes JP, Baeten JM, Lingappa JR, et al. Determinants of per-coital-act HIV-1 infectivity among African HIV-1-serodiscordant couples. J Infect Dis **2012**; 205:358-65.

36. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. N Engl J Med **2011**; 365:493-505.

37. Smith DK, Herbst JH, Zhang X, Rose CE. Condom effectiveness for HIV prevention by consistency of use among men who have sex with men in the United States. J Acquir Immune Defic Syndr **2015**; 68:337-44.

38. Goodreau SM, Carnegie NB, Vittinghoff E, et al. What drives the US and Peruvian HIV epidemics in men who have sex with men (MSM)? PLoS One **2012**; 7:e50522.

39. Wiysonge CS, Kongnyuy EJ, Shey M, et al. Male circumcision for prevention of homosexual acquisition of HIV in men. Cochrane Database Syst Rev **2011**:CD007496.

40. Liu AY, Cohen SE, Vittinghoff E, et al. Preexposure Prophylaxis for HIV Infection Integrated With Municipal- and Community-Based Sexual Health Services. JAMA Intern Med **2016**; 176:75-84.

41. Chan PA, Mena L, Patel R, et al. Retention in care outcomes for HIV pre-exposure prophylaxis implementation programmes among men who have sex with men in three US cities. J Int AIDS Soc **2016**; 19:20903.

42. Andrews CD, Spreen WR, Mohri H, et al. Long-acting integrase inhibitor protects macaques from intrarectal simian/human immunodeficiency virus. Science **2014**; 343:1151-4.

43. Andrews CD, Yueh YL, Spreen WR, et al. A long-acting integrase inhibitor protects female macaques from repeated high-dose intravaginal SHIV challenge. Sci Transl Med **2015**; 7:270ra4.

44. Radzio J, Spreen W, Yueh YL, et al. The long-acting integrase inhibitor GSK744 protects macaques from repeated intravaginal SHIV challenge. Sci Transl Med **2015**; 7:270ra5.

45. Kelley CF, Kahle E, Siegler A, et al. Applying a PrEP Continuum of Care for Men Who Have Sex With Men in Atlanta, Georgia. Clin Infect Dis **2015**; 61:1590-7.

46. Goedel WC, Halkitis PN, Greene RE, Hickson DA, Duncan DT. HIV Risk Behaviors, Perceptions, and Testing and Preexposure Prophylaxis (PrEP) Awareness/Use in Grindr-Using Men Who Have Sex With Men in Atlanta, Georgia. J Assoc Nurses AIDS Care **2016**; 27:133-42.

47. Harris NS, Johnson AS, Huang YA, et al. Vital Signs: Status of Human Immunodeficiency Virus Testing, Viral Suppression, and HIV Preexposure Prophylaxis - United States, 2013-2018. MMWR Morb Mortal Wkly Rep **2019**; 68:1117-23.

48. Rosenberg ES, Purcell DW, Grey JA, Hankin-Wei A, Hall E, Sullivan PS. Rates of prevalent and new HIV diagnoses by race and ethnicity among men who have sex with men, U.S. states, 2013-2014. Ann Epidemiol **2018**; 28:865-73.

49. Lankowski AJ, Bien-Gund CH, Patel VV, Felsen UR, Silvera R, Blackstock OJ. PrEP in the Real World: Predictors of 6-Month Retention in a Diverse Urban Cohort. AIDS Behav **2019**; 23:1797-802.

50. Gray AL, Smit JA, Manzini N, Beksinska M. Systematic review of contraceptive medicines: "Does choice make a difference?". WHO RHRU, **2006**.

TABLES

Table 1. Pre-exposure prophylaxis parameters in reference model, base LAI-PrEP parameters and LAI-PrEP values varied in counterfactual scenarios.

Parameter	Reference Scenario	Base LAI-PrEP Scenario	Counterfactual LAI-PrEP Scenarios	Reference
Initiation and Persistence				
Prevalence of overall PrEP use given indication	15.1%	15.1%	Varies	[45-47]
Probability of initiating PrEP per time-step	0.411%	0.411%	0 – 100%	
Probability of selecting LAI-PrEP	0%	50%	0 – 100%	[21, 23, 25-28]
Spontaneous discontinuation rate, median time	224 days	224 days	75 – 672 days	[41]
DO-PrEP				
Adherence level, % of DO-PrEP users				[40]
Low (< 2 doses/week)	8.9%	---	---	
Medium (2-3 doses/week)	12.7%	---	---	
High (4+ doses/week)	78.4%	---	---	
Efficacy, per-exposure hazard-ratio				[14]
Low adherence	0.69	---	---	
Medium adherence	0.19	---	---	
High adherence	0.02	---	---	
LAI-PrEP				
Peak plasma concentration (P)	---	3.59 µg/ml	1.5 – 6.5 µg/ml	[17]
Half-life ($T_{1/2}$)	---	35 days	15 – 60 days	[17]
Days since last injection (t)	---	Varies	Varies	
Plasma concentration (P_t)	---	$P_t = P * \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}}$	---	[17]
Efficacy, per-exposure hazard-ratio	---	$HR = e^{P_t * B}$	± 5% – 50%	
Beta (B)	---	-9.06	---	[42]
Hazard-ratio at 4 x PA-IC$_{90}$	---	0.002	Varies	[42]

Table 2. Varying rate of pre-exposure prophylaxis initiation and proportion choosing long-acting injectable pre-exposure prophylaxis (LAI-PrEP) versus daily-oral (DO) PrEP compared to the reference scenario of current uptake of DO-PrEP.

PrEP Initiation Probability[1]	LAI-PrEP Selection Probability[2]	% DO-PrEP[3] (95% SI[4])	% LAI-PrEP[5] (95% SI[4])	% Waning LAI-PrEP[6] (95% SI[4])	HIV Prevalence[7] (95% SI[4])	HIV Incidence[8] (95% SI[4])	PIA[9] (95% SI[4])
	0%	**15.1% (14.4, 15.9)**	**0.0% (0.0, 0.0)**	**0.0% (0.0, 0.0)**	**22.4% (21.5, 23.4)**	**1.19 (0.93, 1.50)**	**Reference Model**
Base Probability	25%	11.4% (10.7, 12.0)	4.2% (3.8, 4.6)	3.7% (3.4, 4.1)	22.2% (21.5, 23.1)	1.15 (0.90, 1.42)	2.0% (-9.6, 13.1)
	50%	7.6% (7.1, 8.2)	8.4% (7.9, 9.0)	7.3% (6.8, 7.7)	22.1% (21.3, 22.9)	1.14 (0.87, 1.40)	4.3% (-7.3, 14.5)
	75%	3.8% (3.4, 4.2)	12.6% (11.9, 13.3)	10.7% (10.1, 11.3)	21.9% (21.0, 22.9)	1.09 (0.84, 1.38)	6.3% (-7.0, 16.7)
	100%	0.0% (0.0, 0.0)	16.8% (16.1, 17.5)	13.9% (13.2, 14.6)	21.8% (20.9, 22.8)	1.07 (0.84, 1.36)	7.5 (-3.5, 18.7)
	0%	26.7% (25.8, 27.5)	0.0% (0.0, 0.0)	0.0% (0.0, 0.0)	21.5% (20.6, 22.3)	1.02 (0.78, 1.31)	11.3% (0.1, 21.1)
	25%	20.0% (19.1, 20.8)	7.4% (6.9, 7.9)	6.4% (5.9, 6.9)	21.2% (20.5, 22.1)	0.99 (0.74, 1.25)	14.6% (2.6, 24.4)
2x Base Probability	50%	13.4% (12.7, 14.0)	14.8% (14.0, 15.5)	12.3% (11.7, 12.9)	21.0% (20.2, 21.8)	0.95 (0.72, 1.20)	17.1% (6.7, 26.4)
	75%	6.7% (6.2, 7.1)	22.2% (21.3, 23.0)	17.6% (16.8, 18.4)	20.8% (20.0, 21.7)	0.90 (0.68, 1.17)	19.8% (9.4, 29.1)
	100%	0.0% (0.0, 0.0)	29.5% (28.5, 30.4)	22.3% (21.6, 23.1)	20.5% (19.7, 21.4)	0.87 (0.68, 1.11)	22.4% (13.3, 31.8)
	0%	43.1% (42.0, 44.0)	0.0% (0.0, 0.0)	0.0% (0.0, 0.0)	20.4% (19.7, 21.3)	0.87 (0.60, 1.10)	24.4% (13.9, 33.0)
	25%	32.3% (31.3, 33.1)	11.9% (11.2, 12.6)	10.0% (9.4, 10.5)	20.1% (19.3, 20.8)	0.81 (0.60, 1.03)	28.5% (20.0, 36.0)
4x Base Probability	50%	21.5% (20.7, 22.3)	23.8% (23.0, 24.6)	18.4% (17.6, 19.1)	19.7% (18.9, 20.4)	0.74 (0.55, 0.98)	32.0% (23.6, 40.6)
	75%	10.8% (10.1, 11.4)	35.6% (34.7, 36.6)	25.2% (24.4, 25.9)	19.4% (18.6, 20.1)	0.70 (0.53, 0.90)	36.2% (27.5, 43.9)
	100%	0.0% (0.0, 0.0)	47.5% (46.5, 48.6)	30.5% (29.7, 31.4)	19.1% (18.4, 19.9)	0.66 (0.49, 0.85)	39.5% (31.6, 46.3)

[1]Weekly probability of initiating PrEP given behavioral indications. [2]Probability of selecting LAI-PrEP (versus DO-PrEP) given initiation. [3]Median proportion using daily-oral pre-exposure prophylaxis among HIV-negative MSM with a behavioral indication for PrEP. [4]95% simulation interval. [5]Median proportion using long-acting injectable pre-exposure prophylaxis among HIV-negative MSM with a behavioral indication for PrEP. [6]Median proportion with waning protection from LAI-PrEP (>8 weeks and ≤ 52 weeks since last dose) among HIV-negative MSM with a behavioral indication for PrEP. [7]HIV prevalence among all MSM. [8]HIV incidence per 100 person-years at risk. [9]Percent of infections averted over 10 years compared to the reference model scenario.

Table 3. Varying base parameters for long-acting injectable pre-exposure prophylaxis compared to the reference scenario of current uptake of DO-PrEP.

Scenario	% DO-PrEP[1] (95% SI[3])	% LAI-PrEP[2] (95% SI[3])	% Waning LAI-PrEP[2] (95% SI[3])	HIV Prevalence (95% SI[3])	HIV Incidence[4] (95% SI[3])	PIA[5] (95% SI[3])
Reference	15.1% (14.4, 15.9)	0.0% (0.0, 0.0)	0.0% (0.0, 0.0)	22.4% (21.5, 23.4)	1.19 (0.93, 1.50)	---
Base Parameters	7.6% (7.1, 8.2)	8.4% (7.9, 9.0)	7.3% (6.8, 7.7)	22.1% (21.3, 22.9)	1.14 (0.87, 1.40)	4.3% (-7.3%, 14.5)
Pharmacokinetics (base parameters: peak plasma concentration = 3.59 μg/ml; half-life = 35 days)						
2 μg/ml 15 days	7.6% (7.1, 8.1)	8.4% (7.8, 8.9)	7.3% (6.8, 7.8)	22.3% (21.4, 23.3)	1.17 (0.89, 1.49)	1.4% (-11.5, 11.9)
3 μg/ml 25 days	7.6% (7.0, 8.1)	8.4% (7.9, 8.9)	7.3% (6.8, 7.8)	22.2% (21.3, 23.1)	1.15 (0.90, 1.44)	3.1% (-10.4, 13.5)
LAI-PrEP Efficacy (base parameter: continuous function; hazard ratio at PA-IC$_{90}$ = 0.002)						
50% Decrease	7.6% (7.1, 8.2)	8.4% (7.8, 9.0)	7.3% (6.8, 7.8)	22.8% (22.0, 23.7)	1.32 (1.05, 1.66)	-4.4% (-17.4, 8.4)
10% Decrease	7.6% (7.1, 8.1)	8.4% (7.8, 9.0)	7.3% (6.8, 7.8)	22.2% (21.3, 23.1)	1.17 (0.89, 1.45)	2.4% (-9.7, 13.8)
10% Increase	7.6% (7.1, 8.2)	8.4% (7.8, 9.0)	7.3% (6.8, 7.8)	21.9% (21.1, 22.8)	1.08 (0.82, 1.40)	6.0% (-6.4, 16.6)
50% Increase	7.6% (7.1, 8.1)	8.4% (7.8, 9.0)	7.3% (6.8, 7.8)	21.5% (20.6, 22.3)	0.98 (0.73, 1.23)	10.9% (0.4, 21.4)
LAI-PrEP Spontaneous Discontinuation (base parameter: median time to discontinuation = 224 days)						
1/3x = 74.7 days	8.0% (7.4, 8.5)	3.8% (3.5, 4.1)	7.7% (7.2, 8.2)	22.3% (21.4, 23.2)	1.16 (0.89, 1.47)	1.1% (-12.3, 12.9)
1/2x = 112 days	7.9% (7.3, 8.4)	5.0% (4.6, 5.4)	7.6% (7.0, 8.1)	22.3% (21.3, 23.2)	1.17 (0.92, 1.49)	1.8% (-11.7, 13.6)
2x = 448 days	7.2% (6.7, 7.7)	13.6% (12.9, 14.4)	6.8% (6.3, 7.3)	21.7% (20.8, 22.6)	1.08 (0.83, 1.34)	7.9% (-3.6, 18.6)
3x = 672 days	6.9% (6.4, 7.4)	17.5% (16.6, 18.3)	6.5% (6.0, 6.9)	21.6% (20.7, 22.5)	1.02 (0.77, 1.28)	10.5% (-2.2, 20.5)

[1]Constant discontinuation rate with a median time to discontinuation of 224 days in the base parameter. [2]Median proportion using daily-oral pre-exposure prophylaxis among HIV-negative MSM with a behavioral indication for PrEP. [3]95% simulation interval. [4]Median proportion using long-acting injectable pre-exposure prophylaxis among HIV-negative MSM with a behavioral indication for PrEP. [5]Median proportion with waning protection from LAI-PrEP (>8 weeks and ≤ 52 weeks since last dose) among HIV-negative MSM with a behavioral indication for PrEP. [6]HIV prevalence among all MSM. [7]HIV incidence per 100 person-years at risk. [8]Percent of infections averted over 10 years compared to the reference model scenario.

FIGURES

Figure 1. Population benefit of varying overall PrEP uptake and proportion using LAI-PrEP.

Legend: Percent of infections averted (PIA) over 10 years, compared to the reference scenario of 15% (dashed line) of behaviorally indicated men who have sex with men using pre-exposure prophylaxis and 0% choosing long-acting injectable pre-exposure prophylaxis (LAI-PrEP) versus daily-oral pre-exposure prophylaxis, across varying overall PrEP uptake and proportion using LAI-PrEP.

Figure 2. Population benefit of long-acting injectable pre-exposure prophylaxis (LAI-PrEP) across varying pharmacokinetic parameters.

Legend: Percent of infections averted (PIA) over 10 years, compared to the reference scenario of 15% (dashed line) of behaviorally indicated men who have sex with men using pre-exposure prophylaxis and 0% choosing long-acting injectable pre-exposure prophylaxis (LAI-PrEP) versus daily-oral pre-exposure prophylaxis, if 50% of PrEP users choose LAI-PrEP when available, across varying parameters for LAI-PrEP pharmacokinetics. Dashed lines indicate peak plasma concentration (3.59 µg/ml) and half-life (5 weeks) used in the base model parameters.

Figure 3. Population benefit of long-acting injectable pre-exposure prophylaxis (LAI-PrEP) across varying efficacy.

Legend: HIV outcomes among all men who have sex with men (MSM) in a population with 15% of behaviorally indicated men using PrEP and 50% choosing long-acting injectable pre-exposure prophylaxis (LAI-PrEP) versus daily-oral PrEP, across varying scenarios of the relative change in per-exposure LAI-PrEP efficacy. Panel A: HIV incidence rate per 100 person-years at risk. Panel B: Percent of infections averted compared to the reference scenario of 15% of behaviorally indicated MSM using PrEP and 0% choosing LAI-PrEP.

Figure 4. Population benefit of long-acting injectable pre-exposure prophylaxis (LAI-PrEP) across varying rates of spontaneous discontinuation.

Legend: Long-acting injectable pre-exposure prophylaxis (LAI-PrEP) use and HIV outcomes among all men who have sex with men (MSM) across scenarios with fixed overall PrEP initiation rate (calibrated to 15% of behaviorally indicated MSM in the reference scenario), 50% choosing LAI-PrEP, and varying LAI-PrEP spontaneous discontinuation rates, compared to a reference scenario of 0% choosing LAI-PrEP.

Panel A: Percent of behaviorally indicated MSM actively using LAI-PrEP (≤ 8 weeks since last dosage).

Panel B: Percent of behaviorally indicated MSM with waning protection from LAI-PrEP after discontinuation (> 8 weeks and ≤ 52 weeks since last dosage). **Panel C**: HIV incidence rate per 100 person-years at risk. **Panel D**: Percent of infections averted compared to the reference scenario.