Current practice and challenges in screening for visual perception deficits after stroke: a qualitative study

Kathleen Vancleef,

Department of Experimental Psychology, University of Oxford, Anna Watts Building,

Radcliffe Observatory Quarter, Woodstock Road, University of Oxford, OX2 6GG,

United Kingdom

Email: kathleen.vancleef@psy.ox.ac.uk

Twitter: @katvancleef

Michael J Colwell,

Department of Experimental Psychology, University of Oxford, Anna Watts Building,

Radcliffe Observatory Quarter, Woodstock Road, University of Oxford, OX2 6GG,

United Kingdom

Email: michael.colwell@psy.ox.ac.uk

Twitter: @MichaelJColwell

Olivia Hewitt,

The Oxford Institute of Clinical Psychology Training and Research, University of

Oxford, Isis Education Centre, Warneford Hospital, Oxford, OX3 7JX, United Kingdom

Email: olivia.hewitt@hmc.ox.ac.uk

Nele Demeyere,

20   Department of Experimental Psychology, University of Oxford, Anna Watts Building,

21   Radcliffe Observatory Quarter, Woodstock Road, University of Oxford, OX2 6GG,

22   United Kingdom

23   Email: nele.demeyere@psy.ox.ac.uk

24   Twitter: @NeleDemeyere

Abstract

AIM: We aimed to document current clinical practice and needs in screening for visual

perception problems after stroke to inform development of new screening tools.

METHODS: We interviewed 25 health care professionals (12 occupational therapists, 13

orthoptists) from 16 organisations in England. Interviews were transcribed and coded in

NVivo Software. Data were thematically analysed using the Value Proposition Canvas, a

model which establishes what people want to achieve, the challenges they face and what

facilitates their jobs.

RESULTS: Participants' understanding of visual perception varied and often included sensory

and cognitive deficits. Occupational therapists commonly screened for visual field deficits

and hemispatial neglect, while other aspects of visual cognition were rarely assessed. They

decided on referrals to orthoptists for further assessment. Screening generally occurred

during functional assessments and/or with in-house developed tools. Challenges to practice

were: lack of time, lack of training, environmental and stroke survivor factors (e.g. aphasia),

insufficient continuation of care, and test characteristics (e.g. not evidence-based).

Facilitators to practice were: quick and practical tools, experienced staff or tools with

minimal training requirements, a streamlined care pathway between a stroke unit and eye

hospital supported by occupational therapists and orthoptists.

CONCLUSION: Screening employs non-standardised assessments and rarely covers visual

perceptual deficits in higher order perception. Our service evaluation demonstrates the

need for a standardised visual perception screen, which should ideally be 15 minutes or less,

be portable, and require minimal equipment. The screen should be suitable for bedside

testing in noisy environments, inclusive for participants with aphasia and evidence-based.

Introduction

Lilian Kallir, a concert pianist, could no longer read music from one day to the next. Brain damage left her with the inability to read although she was perfectly capable of writing and had not gone blind. This case described by Oliver Sacks illustrates the importance of our brain in processing visual information to make sense of the world around us.[1]

Visual perception is the dynamic process of perceiving the environment through sensory inputs and translating the sensory input into meaningful concepts associated with visual knowledge of the environment.[2] Visual perception problems are therefore distinct from sensory visual impairments such as reduced visual acuity, visual field and eye movements.[3] Where sensory visual impairments result from damage to the eye or early visual pathways from the eye to the primary visual cortex, visual perception deficits are attributed to impaired function in later visual processing areas in the occipital, parietal and temporal cortex.[4] Examples of visual perceptual deficits include apperceptive and associative agnosia, prosopagnosia, akinetopsia, achromatopsia, problems in visual memory, and in visuospatial abilitities.[4] Visual inattention or hemispatial neglect is sometimes considered to be part of visual perception,[2,5] though neuropsychology research attributes this to an attentional deficit.[6] In particular the presence of preserved perception when attention is stretched to focus on the stimuli, the existence of cross-modal neglect and manipulations of stimulus density on the extent of neglect support the classification of hemispatial neglect as a disorder of attention. [7,8]

Lilian is not an isolated case: Rowe and colleagues reported that 20% of stroke survivors with a suspected visual difficulty have visual perception deficits.[9] This study made use of reports by stroke survivors and carers rather than formal assessment. With systematic

screening with the Rivermead Perceptual Assessment Battery, Edmans and Lincoln

identified visual perception problems in 76% of hemiplegic stroke survivors.[10] The

discrepancy between prevalence with self-reports compare to neuropsychological

assessment suggests that not all visual perceptual problems are picked up based on self-

report. This means many stroke survivors are discharged without the appropriate

rehabilitation or adjustments in their home environment or care packages.

Under-diagnosis of visual perception problems can severely impact stroke survivors' quality

of life, functional outcome, participation in the community, independence and pose

substantial risk. [11–15] For instance, participation in traffic with visual perception problems

can be dangerous and even life-threatening. Risks are also heightened indoors when

preparing a meal or in perceiving trip hazards. Better diagnosis of visual perception

difficulties after stroke will allow better care planning and substantially impact stroke

survivors' life.

The first step toward better diagnosis is an in-depth understanding of the clinical reality and

the reasons behind under-diagnosis. We need to understand what the challenges are in

screening for visual perceptual difficulties after stroke. This can then inform the

development of solutions.

In the current study, we aimed to conduct an in-depth exploration of current clinical

practice, challenges and facilitators of screening for visual perception problems after stroke.

To achieve a rich understanding of these issues, we performed semi-structured interviews

with orthoptists and occupational therapists. These professionals are most commonly

involved in visual perception screening after stroke in the United Kingdom's National Health

Service.

Methods

Participants

All participants were recruited via opportunistic and snowball sampling. Invitations for the interviews were sent out through (i) email to the British and Irish Orthoptic Society Stroke and Neuro Rehab Special Interest Group, (ii) Twitter, (iii) a sign-up sheet at a conference poster and (iv) informal face-to-face conversations at the United Kingdom Stroke Forum 2018. Participants met inclusion criteria if they were working in the National Health Service as an occupational therapist or an orthoptist, and were involved in the assessment of visual perceptual problems after stroke. The participants were informed about the aims of the study, the organisations running the study (University of Oxford and North Bristol NHS Trust), and the planned outcomes and dissemination. The latter made explicit that the results would be used to design a national survey on the topic to inform the development of a new visual perception screen for stroke survivors. The project was reviewed and approved by the Patient Safety Assurance & Audit Service at NHS North Bristol Trust as a Clinical Effectiveness study (CE45999). All participants gave verbal consent.

Interviewer's training

All interviews were conducted over the phone by the first author, KV, who holds an MSc in Clinical Psychology and a PhD in Experimental Psychology. At the time of the research study KV was working as a postdoctoral research fellow at the University of Oxford. KV had nine years' experience as a researcher, including one year as a qualitative researcher. Her formal training in qualitative research included an undergraduate course in qualitative research methods at the University of Leuven which included a research project. In preparation for the current project, she was trained and supervised by senior author, OH. OH is a clinical

psychologist who has worked predominantly as a qualitative researcher and health care

professional within the field of intellectual disabilities. OH has numerous publications within

this field, teaches qualitative research methods at the University of Oxford, and has

supervised a number of doctoral projects using various qualitative methodologies. KV's

training included reading and reflecting on reference works in qualitative research,[16,17] a

question and answer session with an experienced moderator, and guidance during coding

and analyses via regular meetings with OH.

Model

The interview process and qualitative analysis was guided by the Value Proposition Canvas.

This model was originally designed to guide product development, but can be applied to

different contexts.[18] Central to this model is the principle that development of products or

services are best informed by the users. A good product or service takes into account the

goals of the user, it reduces the pains a user experiences while trying to achieve their goals,

and it provides gains in achieving their goals. For instance a good electric car helps a person

to get to work (goal to achieve) without having to fear that the battery will run out (reduce

pains) while at the same time provides a beautiful design to impress colleagues and family

(increase gains). We can apply this model in trying to understand current clinical practice in

visual perception screening after stroke (goal), the challenges (pains) it brings, and what

facilitates the process (gains).

Data collection

A semi-structured interview guide was developed by KV in collaboration with an

occupational therapist and an orthoptist (Table 1). The interview guide was reviewed by an

experienced moderator and piloted with one orthoptist and three occupational therapists.

The interviews consisted of open-ended questions about current practice in visual

perception screening after stroke, and the challenges and facilitators of practice. The

interviewer conducted interviews at the participant's workplace or over the phone in a quiet

office with no other people present. The participants took the call in a location of their

choice. The location and presence of others at the participant's side was not recorded. After

obtaining participant consent, the interview was audio recorded. All participants were

interviewed once. Interviews took place between October 2018 and January 2019.

Recruitment was ongoing during data collection and interviews were conducted until data

saturation was reached.

Table 1. Interview structure

Section	Interview route
Introduction	*Q.* Which duties are central to your job role?
Definition of visual perception	*Q.* What comes to your mind when I say visual perception problems in stroke survivors?
Current practice	*Q.* Do you do visual perception screening as a routine on your stroke survivors? (Orthoptists only) *Q.* What did you do the last time you suspected visual perception problems in a stroke survivor? *Cues: standardised tests, unstandardized tests, observations, conversation with stroke survivor, family members, other staff, medical notes* *Q.* What is the care pathway for these stroke survivor? *Cues: Which health care professional (occupational therapist,, neuropsychologist, orthoptist, consultant, physiotherapist) does what (clinical judgement, systematic screen, in depth assessment) and where (A&E, Acute stroke ward, rehab ward, care home, at home, outpatient appointment at eye clinic), referral process*
Facilitators to practice	*Q.* What does an initial visual perception screen need to be like for you? *Cues: time it takes, format, what to include, what result, who should do it*
Challenges to practice	*Q.* Can you name three things that frustrate you about visual perception screening on the stroke ward? *Cues: assessment methods, time, training, other staff, care pathway,* *Q.* Which of the three challenges you have mentioned is the most important?

Ending	*Q.* If you could give one piece of advice to someone making a visual perception screen, what would it be?
	Q. My goal was to have an idea of how visual perception screening is done and what the clinical needs are. Do you think we've talked about everything or is there anything we've missed?

Qualitative Analyses

As a method of data analysis we used thematic analysis.[19] Interviews were transcribed

verbatim by a professional typist via an external service provider. The transcripts were not

returned to the participants for feedback. Two researchers (KV and MC) were involved in

data analysis. KV was familiar with the data from conducting the interviews (Phase 1 in

thematic analysis). The first 15 interviews were checked and coded by KV in NVivo Software

(Version 12.0). She developed an initial coding tree (Phase 2 in thematic analysis). Overall

themes were set in advance in alignment with the Value Proposition Model: Jobs,

Challenges, and Facilitators. The coding tree was reviewed and substantially adapted after

coding of five interviews. MC familiarised himself with the data by checking transcripts and

reading through interviews coded by KV. Two of these interview transcripts were checked,

coded and discussed by both KV and MC to align their coding. MC checked and coded the

remaining 10 interview transcripts. The frequency of text extracts coded under each node

was visualised with a hierarchical chart of nodes in NVivo to explore overall patterns in the

data describing current practice. This chart shows the frequency of references for each node

when participants were talking about current practice in routine visual and visual perception

screening. Only nodes with at least ten references were included. Overall themes and

subthemes were derived from the data by KV (Phase 3 of thematic analysis). This was done

in two stages: (1) we performed search queries of coded text for challenges and facilitators

of visual perception screening in current practice and (2) read through the resulting text.

The themes were subsequently reviewed by reading all answers to the questions under the

headings Challenges and Facilitators (see Table 2). The information was then synthesised by

KV (Phase 4 of thematic analysis). Lastly, themes and subthemes were reviewed, refined and

defined by KV (Phase 5 of thematic analysis) and a report was prepared (Phase 6 of thematic

analysis). This report was sent out to the participants for validation. Ten out of 25

participants replied and nine agreed with the thematic analysis. One participant pointed out

that participants' understanding of visual perception was underexplored. We added a

results section on this topic posthoc.

Results

12 occupational therapists and 13 orthoptists from 16 health care organisations in England

took part in the study. Characteristics of each participant are reported in Table 2. All

participants worked in England, and were based across the East Midlands (n=1), East of

England (n=1), Greater London (n=2), North East (n=3), North West (n=4), South East (n=6),

South West (n=4), and Yorkshire and the Humber (n=4). The interviewer had no personal

relationship with any of the participants. Two orthoptists received referrals from the same

acute stroke unit where the first author recruits stroke survivors as participants for other

studies (pseudonyms cannot be disclosed to ensure anonymity). One of the participants was

also involved in the development of the interview guide. All but one participant (Jessica)

agreed to be audio recorded. For Jessica, notes were made during and immediately after the

interview. Interviews lasted between 16 and 46 minutes with an average of 27 minutes. For

Christopher, the interview was split over two consecutive days.

 Table 2. Participant characteristics

Pseudonym	Gender	Profession	Clinical setting	Years of experience in stroke care
Amanda	Female	Orthoptist	Eye clinic	10+ years
Amber	Female	Occupational therapist	Rehabilitation service	10+ years
Amy	Female	Orthoptist	Eye clinic	unknown
Brittany	Female	Occupational therapist	Rehabilitation service	10+ years
Christopher	Male	Occupational therapist	Acute stroke service	unknown
Crystal[a]	Female	Occupational therapist	Rehabilitation service	5-10 years
Danielle	Female	Occupational therapist	Acute stroke service	unknown
Elizabeth	Female	Orthoptist	Eye clinic	10+ years
Emily[a]	Female	Occupational therapist	Rehabilitation service	unknown
Erin	Female	Occupational therapist	Acute stroke service	5-10 years
Heather[b]	Female	Orthoptist	Eye clinic	10+ years
Jamie[b]	Female	Orthoptist	Eye clinic	2-5 years
Jessica[c]	Female	Orthoptist	Eye clinic	10+ years
Joshua	Male	Orthoptist	Eye clinic	2-5 years
Kimberley	Female	Occupational therapist	Acute stroke service	unknown
Matthew	Male	Orthoptist	Eye clinic	5-10 years
Megan	Female	Occupational therapist	Rehabilitation service	10+ years
Melissa	Female	Occupational therapist	Acute stroke service	10+ years
Michelle	Female	Occupational therapist	Rehabilitation service	2-5 years
Nicole	Female	Orthoptist	Eye clinic	unknown
Rachel	Female	Orthoptist	Eye clinic	unknown
Rebecca	Female	Occupational therapist	Acute stroke service	5-10 years
Sarah[c]	Female	Orthoptist	Eye clinic	10+ years
Stephanie	Female	Orthoptist	Eye clinic	unknown
Tiffany	Female	Orthoptist	Eye clinic	2-5 years

 [a,b,c] Participants working in the same clinical teams A, B or C.

Participants understanding of visual perception

The interviews highlighted that participants' understanding of visual perception differed from textbook definitions of visual perception. Visual perception is defined as a dynamic process of translating sensory visual information into meaning percepts[2]. These percepts can subsequently be linked to higher cognitive functions like visual memory for recognition (ventral of what stream) and praxis for visually guided actions (dorsal or where stream)[4]. Visual perception must be differentiated from low level sensory visual functions like acuity, visual fields, and ocular movements, and from higher cognitive processes like (visual) memory or (visual) inattention, although some authors consider visual inattention to be part of visual perception. Our participants seemed to have a broader understanding of visual perception and often did not differentiate between sensory and higher cognitive functions.

Participants typically mentioned the following examples when asked about what they

consider visual perceptual problems (in descending order of frequency): visual inattention,

agnosia's for objects, faces, letters, or shapes, hallucinations or Charles Bonnet syndrome,

difficulties in depth perception, visual field deficits like hemianopia, and spatial vision (e.g.

navigating and perceiving space around them).

Participants' understanding seemed to vary widely but we observed general trends in the

different professions. Orthoptists seemed to differentiate visual perception difficulties from

low level sensory visual impairments, but considered visual inattention the prime (and only)

example of visual perception difficulties after stroke.

"[Interviewer: What comes to your mind, when I say visual perception problems in stroke

patients?] So I would probably say visual neglect, ignoring one side of their world when the

other side is presented to them as well, is the main definition of it. That it's decreased

awareness of that side of their vision and their external world as well as themselves as well,

on that side. *[Interviewer: Anything else, that falls under that term for you?]* No, not that I

can think of, off the top of my head, no." (Tiffany, Orthoptist in an eye clinic)

In addition to visual inattention, occupational therapists mentioned impairments that are

typically (but not exclusively) related to low level sensory vision like contrast sensitivity,

hemianopia, and depth perception. They seemed to interpret visual perception more

broadly.

"A range really. So we would look at visual perception as things like depth perception, …

recognition of objects, … and any kind of visual inattention or neglect, that sort of thing…

But yes, that's it basically." (Danielle, Occupation Therapist in an acute stroke service)

These observations are relevant in interpreting the subsequent results. Although the

interview questions asked for experiences with visual perception assessment after stroke,

the results below should be interpreted as potentially referring to all or any of the following:

sensory visual impairments (for occupational therapist participants), visual perception

impairments (for all participants), and visual inattention (for all participants).

Current Practice

A hierarchical chart of nodes is presented in Figure 1. The figure illustrates that participants

most often mentioned occupational therapists working in acute stroke or rehabilitation

services when discussing screening for visual neglect, visual field deficits, and ocular

movements. For screening they referred to in-house developed screening tools,

standardised tests or observations in functional tasks like dressing and making a cup of tea.

"In terms of the visual field, we do the standard, you know, wiggle your fingers

test to see how people respond. We do the tracking tests. Otherwise it's mainly

through observing people in functional tasks what they're doing. ... Through that

you can often realise ... if they're having problems with their visual perception

because they're missing things or not seeing things." (Crystal, occupational

therapist in a stroke rehabilitation service)

"I'd rather do it in function. So it's more meaningful especially with stroke

patients because it makes more sense. Rather than getting them to balance

cubes on each other or, you know, I can say 'can you, can you find the

toothpaste?' " (Megan, occupational therapist in an acute stroke service)

Participants mentioned both positive and negative aspects of the process, referring to the

conditions of the stroke survivor and the physical environment, time limitations, staff

training and relationships with other staff members. These themes are explored in greater

detail in subsequent paragraphs on Challenges and Facilitators.

"I think for us, it's a multitude of things, it's an acute unit, through-put is quite

quick, it's an environment that often doesn't lend itself to a nice quiet space

where patients can attend … from an OT [occupational therapist] point of view,

there's not a lot of OTs who are actually for [delivering] the therapy, so time-wise

we haven't got a lot of capacity at times, to do that." (Danielle, occupational

therapist in an acute stroke service)

"It can be a bit frustrating … because lots of the occupational therapists are on a

rotational job. So getting them trained up [is difficult]. I think they do find vision

scary." (Stephanie, orthoptist in an eye clinic)

Figure 1. Hierarchical chart of nodes showing the frequency of references for each node. Text is filtered for

extracts where participants were talking about current practice in routine visual and visual perceptions

screening. Only nodes with at least ten references are shown. (colour version available online)

Most participants mentioned that the service they work within has a referral pathway in

place: if problems are suspected following screening by the occupational therapist, a

referral is made to orthoptics. A stroke orthoptist will then see the stroke survivor as an

inpatient on the unit or as an outpatient after discharge from the hospital.

"It's really as and how other health professionals have identified them, then an

orthoptist will go up to try and assert a diagnosis and then give them the

rehabilitation options." (Amy, orthoptist in an eye clinic)

Screening by an occupational therapist followed by referral (if suitable) to orthoptics is seen

as an efficient system.

"Whereas since we've set up a proper vision screening it just means that things

have been a lot more efficient and that's made a much better service because

they're not having to see a lot of people that don't need to be seen. ... And we've

got our waiting lists down substantially so it's just helped things a long quite a

lot." (Nicole, orthoptist in an eye clinic)

Challenges

284 Themes around challenges in visual perception screening are presented in Table 3. We

285 identified five major themes and 13 subthemes.

Lack of time

Occupational therapists and orthoptists both reported that time pressure makes visual

perception screening difficult. They mentioned that because of understaffing they spend

little time with stroke survivors or experience delays in seeing them following referral. In

addition, occupational therapists report that stroke survivors are discharged from an acute

stroke unit as soon as medically safe and a suitable discharge location has been organised.

For survivors with milder strokes, discharge is reported to happen within 1-2 days, giving

very little time for therapists to complete their assessments.

"Well one [challenge] is just logistically and organisationally, just in terms of

caseload and time to hand. We are having to prioritise more direct functional

assessments in terms of establishing readiness for discharge." (Christopher,

occupational therapist in an acute stroke service)

"Often we don't get the chance to use it [assessment instruments] that much

because they're either too unwell for us to do them with or they get to the point

where they're well enough and then they get taken off to a rehab facility."

(Danielle, occupational therapist in an acute stroke service)

"There's not a lot of OTs who are actually for [delivering] the therapy, so time-

wise we haven't got a lot of capacity at times." (Danielle, occupational therapist

in an acute stroke service)

"Lack of time is always going to be frustrating knowing that you've got a clinic

here and you can't get up to the ward and there's an urgent stroke survivor."

(Heather, orthoptist in an eye clinic)

"Erm… it sounds awful doesn't it when you're constantly having to slim down

what you do." (Melissa, occupational therapist in an acute stroke service)

310 Lack of training

Training is another theme that frequently emerged across both professions. Visual

perception difficulties following stroke appear to be only briefly covered in the

undergraduate education of orthoptists and occupational therapists, and is mostly learned

about post-qualification through on-the-job-training. Many occupational therapists

mentioned that they rotate between services, leaving little time for them to be trained in

many aspects of stroke care. In particular, they said that this lack of experience made them

feel uncertain when it comes to vision and visual perception deficits following stroke.

Inexperience and limited training seemed to be an important hurdle for occupational

therapists in visual perception screening. Because vision and visual perception was not well

understood by the staff (see above), participants reported that mistakes in assessments are

commonly made, leading to a potential under-diagnosis of visual perception deficits.

"[It] can be a bit frustrating ... lots of the occupational therapists are on a

rotational job. So getting them trained up, I think they do find vision scary....

Vision, it's the scariest thing about stroke." (Stephanie, orthoptists in an eye

clinic)

"[Visual perception is] one of those things that is perhaps not widely

understood.... think unless you've got a unit with a special interest in stroke and

you've taken the time to look into that... then it's perhaps under-diagnosed as

well, so yeah I think it's recognition that it's a potential problem." (Amanda,

orthoptist in an eye clinic)

"Erm, yeah because visual and cognitive *[problems]* post stroke ... are not well

known. Everybody thinks you know the FAST but nobody thinks of vision as part

of that. ... And sometimes vision is the only thing that's affected post stroke."

(Rachel, orthoptist in an eye clinic)

"The key is really, somebody in the hospital to be able to deliver the training and

the therapy staff to actually take on board the training and to actually deliver it."

(Sarah, orthoptist in an eye clinic)

"I think it probably is underreported; the amount of incidents of perceptual

problems post stroke." (Rachel, orthoptist in an eye clinic)

Environmental and stroke survivor factors

Participants reported that visual perception screening is limited by the environment in

which they take place. They said that in acute stroke care, most assessments are done in a

hospital bay with several beds. Even if curtains can be drawn around the stroke survivor's

bed, distractions were reported to be numerous: sounds from neighbouring beds,

interruptions for timed caring needs, distractions from medical equipment around the bed

and presence of personal objects on a small table in front of the stroke survivor (spectacles,

drinks, food, tissues, personal care items, newspaper, get well cards, crossword book, pen,

etc.) can be visually distracting and interfere with visual perceptual assessment.

"Frustrating for us is ... that we .. depending on which unit we are [on] or where

we are, it can be quite difficult to do [our assessment at] the bedside if there is a

lot [of] distractions going on; if there is no quiet space to take the patient.

Obviously, yes, we can draw the curtains but it's sometimes more the noise levels

than everything. They get easily distracted. That makes it quite hard actually to

do an assessment." (Amber, occupational therapist in a stroke rehabilitation

service)

"It's *[the]* environment. Often *[it]* doesn't lend itself to a nice quiet space where

patients can attend." (Danielle, occupational therapist in an acute stroke service)

Participants also mentioned the condition of the stroke survivors as a limiting factor to

visual perception screening. Both professions reported that tests and tools are often not

suitable for stroke survivors with poor alertness, poor cognition, aphasia or severe weakness

in their upper limbs. For instance, stroke survivors may not able to answer a question on

whether their vision has changed because they developed aphasia following stroke.

Similarly, copying and cancellation tasks to assess visual neglect were reported to be

unsuitable for a stroke survivor who can no longer hold a pen.

"Another component would be around the client or the patient themselves in

terms of factors that might work against screening: so attention, concentration,

fatigue, the ability to deliver a verbal response in terms of aphasia or dysarthria

difficulties. ... That has significant impact on our ability to undertake formal

testing of whatever nature." (Christopher, occupational therapist in an acute

stroke service)

"Our patients who either are fatigued quite quickly, or are ... medically not well

enough to do a lot of taking down to the therapy department ..., or their

attention span is quite limited, or they've got a weakness in their upper limb that

makes it more difficult to complete the tests." (Danielle, occupational therapist in

an acute stroke service)

Continuation of care

A further concern of the participants was the follow-up process after screening for visual

perception deficits. They highlighted that quite often the information from screening is not

passed on to the next care team. Therefore, even if visual perception problems are picked

up in an acute setting, no follow-up assessments are carried out or no rehabilitation is

provided in their experience. With respect to rehabilitation, occupational therapists found it

frustrating that few treatment options are available. They reported that little information

and few guidelines are available on how visual perception problems can be treated or

managed. Some noted that the lack of options for rehabilitation reduced their motivation to

screen for visual perception problems.

"Yes, and it [*the discharge summary*] is very medical because the occupational

therapists or myself don't like the discharge summaries. ... We do all this

information and then it gets lost ....Yeah, it's not part of our discharge summary

in enough detail, so when they go somewhere new which could be two or three

weeks later ... they don't have that information." (Matthew, orthoptist in an eye

clinic)

"It might be because it [*visual perception deficits*] is limited in treatment ... that's

why it's not the focus. So ... it would be more useful to spend time during my

assessment speaking to them about adapting to their field loss and giving them a

prism or occlusion for their double vision." (Rachel, orthoptist in an eye clinic)

Test characteristics

Both professions highlighted limitations in the characteristics of the available tests for visual

perception. Occupational therapists more frequently reported to be hindered by practical

limitations and the unknown impact on daily life. With respect to the practical limitations,

they referred to a large number of test materials like stimulus booklets, stopwatch, cubes,

etc. needed for assessment or too many loose pages. With respect to the unknown impact

on daily life, they referred to stimuli and tests that were thought to be limited in ecological

validity. They considered the tests too abstract with no clear link to the implications for

everyday tasks.

"We have our little screening tool. It's a few pages, just to prompt us …. That's all

right but then … everything else you are … searching together from different kits.

… That's a little bit frustrating." (Amber, occupational therapist in a stroke

rehabilitation service)

"There's an awful lot of bits of paper and a bit of faffing about and preparing to

get it all ready before we actually go out to see the patients." (Melissa,

occupational therapist in an acute stroke service)

"We use Rivermead Perceptual Battery. But to be fair, I don't tend to use it as

often because I'd rather do it in function. …It's more meaningful, especially with

stroke patients because it makes more sense. Rather than getting them to

balance cubes on each other or, you know I can say "Can you find the

toothpaste? Can you take the lid off? Can you brush your own teeth?" and then

it just makes more sense. And then … it's easier for them to identify *[a]* goal."

(Megan, occupational therapist in a stroke rehabilitation service)

Orthoptists, on the other hand, seemed to emphasize more the lack of evidence and

consequently the lack of specific clinical guidelines on how to assess visual perception.

"If you see the national guidelines for visual problems after stroke, they have a

really small section on vision with orthoptics but we don't really have anything in

it that's specific to what we should be screening them with and what we should

be treating them with. So even though we're using evidence-based practice, the

evidence sometimes isn't great and I think that's probably why it's not in the

guidelines because there's not really good evidence to support it. But sometimes

you're seeing a patient and you're not entirely sure if you're doing the right thing

because you're maybe giving them information based on clinical experience,

what other people have been doing but you're not really sure if that is the best

thing to be doing or could you be doing something else with them." (Amy,

orthoptist in an eye clinic)

Table 3. Challenges in visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Lack of time	Limited time with stroke survivor due to discharge pressure	x	
	Limited time with stroke survivor due to staff shortage	x	x
Lack of training	Inexperienced staff with limited training	x	
	Vision and visual perception is not well understood	x	x
	Mistakes made in assessment	x	x
	Vision and visual perception deficits are underdiagnosed	x	x
Environmental and stroke survivor factors	Condition of stroke survivor (e.g. low alertness, poor cognition, aphasia, not mobile)	x	x
	Distractions in test environment (often in hospital bay)	x	x
Continuation of care	Information on vision and visual perception problems gets lost after discharge	x	x
	No information on treatment or management of vision or visual perception problems	x	
Test characteristics	Test materials not readily available or making use clumsy	x	
	Lack of evidence-based screening instruments		x
	No specific recommendations in clinical guidelines		x
	Tests are not meaningful for everyday life	x	

Facilitators

435   The facilitators to practice that participants reported, largely mirrored the challenges

436   reported above (Table 4). We identified three major themes and 12 subthemes.

Practical tool

On a practical level, participants preferred a tool that can be administered in approximately

10-20 minutes, so it can fit in with busy every-day practice and current staff levels. They also

favoured a tool that consists of few test materials and pages to prevent test materials

getting lost or mixed up during testing. According to participating orthoptists, the tool

should have a 1 to 2 page proforma. They reported that it should include questions to ask

the stroke survivor and simple quick tests for a range of functions. They suggested that the

proforma could subsequently be used in a referral for orthoptic input. To allow bedside

testing or testing in the community, participants preferred portable tools. Lastly, many

participants highlighted that a tool should ideally be suitable for all stroke survivors, such as

those with aphasia, poor concentration, poor cognition or severe fatigue.

"It needs to be something that's fairly quick and easy to administer." (Danielle,

occupational therapist in an acute stroke service)

"Just pen and paper, quite simple or a few things to hand that will fit in

something that's quite portable and mobile, I think [*that*] would be handy, yes."

(Brittany, occupational therapist in a stroke rehabilitation service)

"Reasonably short ... for logistical [*reasons*]: ... our time and patient

concentration time, fatigue in the more acute stages. I think really half an hour is

the maximum time." (Christopher, occupational therapist in an acute stroke

service)

"It would have to be a list of standardised questions: Are they able to look

between two points specifically? ... I guess the visual perception question would

be if they're following a moving object? ... You know are they able to identify

their friends or their family? Can they recognise themselves in a photo?" (Rachel,

orthoptist in an eye clinic)

462 Training and experience

Orthoptists and occupational therapists agreed on the importance of training and

experience for robust visual perception screening. The presence of experienced colleagues

or training by other staff members was seen as very advantageous. Given the limited

training in visual perception highlighted above, a tool that is easy to administer and score

was preferred by our participants. Similarly, they reported that results should be easy to

interpret, and should provide clear guidance for management of the deficit (if present).

"If it was something that was quite self-explanatory then anybody could do it and

therefore anybody can interpret it, then that would be much clearer." (Amy,

orthoptist in an eye clinic)

"Just easy to administer and easy to score. … So making instructions clear and

easy to understand would be good." (Emily, occupational therapist in a stroke

rehabilitation service)

Staff roles and interactions

Occupational therapists reported that their preferred practice is to screen stroke survivors

for vision and visual perception problems during their functional assessments (e.g. washing,

making a cup of tea) or via a screening tool. Both professions mention that if a problem is

flagged, more in-depth assessment can then take place.

"Whereas since we've set up a proper vision screening it just means that things

have been a lot more efficient and that's made a much better service because

they're not having to see a lot of people that don't need to be seen." (Nicole,

orthoptists in an eye clinic)

"So, I'm happy with the main basic screen to begin with, and if it digs something

up then we'll dig it in some more detail. I think that works quite well."

(Matthew, orthoptists in an eye clinic)

If the problem concerns visual perception difficulties, an occupational therapist is seen by

our participants as the most appropriate professional to follow it up. If the problem

concerns vision, a referral to an orthoptist was the preferred practice in our sample. Good

relationships between stroke occupational therapists and orthoptists in eye clinics were

frequently mentioned as invaluable.

"I think the key is having the staff, the occupational therapist and the physio staff

actually take that on board." (Sarah, orthoptist in an eye clinic)

Table 4. Facilitators to visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Practical tool	Quick to administer (10-20 min)	x	x
	Paper and pen based without too many test materials or pages	x	x
	Proforma with questions and simple tests to be used in referral		x
	Portable	x	
	Suitable for stroke survivors with poor concentration, communication and fatigue	x	
Training and experience	Easy to administer and score	x	x
	Easy to interpret with management advice included	x	x
	Trained and experienced staff	x	x
Staff roles and interactions	Screening for visual and visual perception problems by occupational therapists during functional assessment or with screening tool	x	x
	In depth assessment with range of complimentary tests	x	x
	Specialised staff for visual (orthoptists) and visual perceptual (occupational therapists) problems		x
	Good relationship between stroke and eye departments		x

Discussion

The aim of this qualitative research was to gain a deeper understanding of current practice

in visual perception screening after stroke, the challenges faced by health care

professionals. This information can then inform what an ideal solution to visual perception

screening might look like. In our participants' stroke care services, occupational therapists

most often screened for visual and visual perceptual difficulties. If visual difficulties like

hemianopia, double vision or reduced acuity, are found, a referral to an orthoptist is made

for an in depth assessment. If visual neglect is suspected, occupational therapists will often

follow this up themselves with a range of tasks like cancellations, copying, line bisection,

etc. The biggest challenges faced by our participants were lack of time and training.

Facilitating factors mentioned were simple, easy to use instruments and good relationships

between stroke and eye departments.

The qualitative approach of our study allowed us to gather views directly and in depth.

Instead of reporting frequency of views (e.g. how many health care professionals report

staff shortage), we focus on how central certain views are to a person's experience. Our

large sample size covers the two main professions involved in visual perception screening

after stroke.

National Clinical Guidelines recommend that every stroke survivor who appears to have

perceptual difficulties should have an assessment with standardised measures.[3] As is

evident from our interviews, only a small proportion of stroke survivors receive an

assessment with standardised measures for visual perceptions difficulties, and screening is

not routinely done. The means of assessment of visual perception deficits varied widely

between participants. Orthoptists will often use standardised tools in their assessments,

though these are not necessarily validated with stroke survivors.[20,21] Overall, standardised

assessments were rarely mentioned by our participants Occupational therapists expressed

preferring assessment during functional tasks like washing and dressing and assessment

with in-house developed screening checklists. This may be due to a lack of a validated

screening tool for a wide range of visual and visual perceptual deficits that can be used by

any health care professional.[20] The diversity in tools that we observed in our group of

English participants contrasts with the reported use of standardised tools in an Australian

population where the Occupational Therapy Adult Perceptual Screening Test and the

Loewenstein Occupational Therapy Cognitive Assessment are most commonly used to

evaluate visual perceptual difficulties.[21]

McClusky and colleagues investigated the barriers and enablers for following recommended

practice in stroke care. In line with our findings the occupational therapists they interviewed

reported the condition of the stroke survivors as a barrier for standardised assessments.

Many tests are not designed and not suitable for stroke survivors with aphasia and/or

dysarthria. Occupational therapists in their study also mentioned the lack of training and

knowledge, specifically for visual neglect rehabilitation. Time pressure and fluctuating staff

levels meant occupational therapists in their study prioritised assessments and

interventions which would produce the best clinical outcome. A lack of time to see stroke

survivors was also mentioned by their orthoptists. However, in our study this referred to the

lack of time to see all referred stroke survivors before they were discharged, while in the

study by McClusky, the orthoptists referred to the lack of time to treat stroke survivors. The

standardised assessments used by Australian occupation therapists, Occupational Therapy

Adult Perceptual Screening Test and the Loewenstein Occupational Therapy Cognitive

Assessment, require training and take 20-45 minutes to complete. Given the frequently

reported lack of training and time, it may appear that the standardised tools recommended

by clinical guidelines are not adapted to the clinical reality of the National Health Service in

England. Although McClusky's study was limited to one Australian hospital and only involved

5 occupational therapists and 2 orthoptists, all working in the same service and

circumstances, the barriers are remarkably similar to what our 12 occupation therapists and

13 orthoptists reported experiencing in 16 English healthcare services. The stroke survivor's

condition, lack of time and staff seem to be universal barriers for providing evidence-based

stroke care across domains, not just visual perception screening. Also, the lack of

information on treatment or management of vision or visual perception problems that

occupational therapists mention as a barrier is not surprising given the very limited evidence

for treatment and management options.[22]

The first challenge exposed by our study is the variation between participants in their

understanding of constitute visual perception problems after stroke. This led to participants

answering interview questions with different reference frameworks. When sharing the

challenges of visual perception screening in their clinical practice, some might have been

considering challenges in just screening for visual inattention difficulties while others

considered a broad range of sensory and cognitive visual impairments. To maintain an equal

relationship between participant and interviewer, and an open non-judgemental

environment to freely share their experiences, participants' definition of visual perception

was not challenged during the interview. A disadvantage of this approach is that it is unclear

whether certain themes apply specifically to one aspect of visual and visual perception

screening or to the whole range of vision related difficulties after stroke.

The second challenge lies in our sample size. Our sample, though the largest for a qualitative

study in this topic, is not representative for the whole population of health care

professionals involved in visual perception screening after stroke. We limited ourselves to

occupational therapists and orthoptists as they are most commonly involved in visual

perception screening in England. Experiences might be very different in clinical settings

without an orthoptic department or with more involvement of neuropsychologists in stroke

care. In addition, voluntary participation and asking our participants to commit 30 minutes

of their time might have induced a bias for allied health professionals with a keen interest in

the topic. We have possibly missed opinions from people with a more negative or neutral

attitude towards visual perception screening. In addition, newly qualified health care

professionasl with limited experience in stroke care might not have felt confident enough to

share their opinions on the topic. Lack of confidence on this topic with junior staff was

brought up by several of our participants. However, the aim of this qualitative research was

not to generalise findings to the population, but rather to provide an in-depth exploration of

the topic to generate hypotheses. We are following this research up with a large scale

survey for all health care professionals involved in visual perception screening in the United

Kingdom and Republic of Ireland. In the recruitment to this survey study, we are

emphasizing that we welcome the opinions of heath care professionals with all levels of

experience.

A third potential limitation is that the interviewer was the principal investigator of the study

and was therefore not independent. Unconscious bias of the interviewer might have guided

the participants' answers. We have tried to minimize this by using a detailed interview guide

that included a list of cues that could be given. In addition, all interviews were recorded and

a second coder was involved.

Our study limits itself to health care professionals' perceptive on visual perception screening

after stroke. We have not explored experiences by stroke survivors, although they can be

substantially different. Also, we have not investigated the management of visual perceptual

problems after stroke of their impact in daily life as is previously reported for sensory visual

problems. [23,24]

The current research highlighted that a clear and consistent definition of visual perception

should be provided to orthoptists and occupational therapists. More training is needed on

the assessment (and management) of visual perceptual screening. This is preferably

provided via professional organisations to ensure consistency. This will raise awareness and

reduce insecurity experienced by junior staff members. Establishing good relationships

between staff at stroke and eye departments is strongly recommended. We suggest this

could be achieved may in the form of knowledge exchanges or via a re-evaluation of the

existing referral system. Lastly, a quick and portable visual perception screening tool that is

easy to administer, score and interpret would highly benefit both staff and stroke survivors

if it is inclusive for stroke survivors with aphasia, motor impairments and cognitive

problems. The tool should be evidence-based and self-explanatory to use

605 Clinical Messages

• There is a high demand for training on what visual perception deficits are and how to

screen for them.

• Building local relationships between orthoptists and occupational therapists is

perceived as highly beneficial for providing good vision and visual perception care for

stroke survivors.

• There is a need for a visual perception screening tool that is self-explanatory, quick,

portable, paper and pen based, accessible to stroke survivors with aphasia or low

concentration, and includes management advice.

614 Acknowledgments

615 We would like to express our sincere gratitude to all the participating allied health

616 professionals who gifted their time. In addition we would like to thank Philip Clatworthy and

617 Claire Murray for facilitating this project.

 Author contributions

#	Role	Definition	KV	MC	OH	ND
1	**Conceptualization**	Ideas; formulation or evolution of overarching research goals and aims.	●			
2	**Data curation**	Management activities to annotate (produce metadata), scrub data and maintain research data (including software code, where it is necessary for interpreting the data itself) for initial use and later re-use.	●	●		
3	**Formal analysis**	Application of statistical, mathematical, computational, or other formal techniques to analyse or synthesize study data.	●	●	●	
4	**Funding acquisition**	Acquisition of the financial support for the project leading to this publication.	●			●
5	**Investigation**	Conducting a research and investigation process, specifically performing the experiments, or data/evidence collection.	●			
6	**Methodology**	Development or design of methodology; creation of models.	●		●	
7	**Project administration**	Management and coordination responsibility for the research activity planning and execution.	●			
8	**Resources**	Provision of study materials, reagents, materials, patients, laboratory samples, animals, instrumentation, computing resources, or other analysis tools.	●			●
9	**Software**	Programming, software development; designing computer programs; implementation of the computer code and supporting algorithms;	●	●		

		testing of existing code components.				
10	**Supervision**	Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team.	●		●	●
11	**Validation**	Verification, whether as a part of the activity or separate, of the overall replication/reproducibility of results/experiments and other research outputs.			●	
12	**Visualization**	Preparation, creation and/or presentation of the published work, specifically visualization/data presentation.		●		
13	**Writing – original draft**	Preparation, creation and/or presentation of the published work, specifically writing the initial draft (including substantive translation).	●	●	●	●
14	**Writing – review & editing**	Preparation, creation and/or presentation of the published work by those from the original research group, specifically critical review, commentary or revision – including pre- or post-publication stages.	●	●	●	●

619   CRediT (Contributor Roles Taxonomy) – see www.carai.org – taxonomy from Allen L, Scott J,

620   Brand A, Hlava M, Altman M. Publishing: Credit where credit is due. Nature News. 2014 Apr

621   17;508(7496):312.

622   Competing interests

623   The authors declare that there is no conflict of interest.

Funding support

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Stroke Association [grant number TSA PDF 2017/03, TSA LECT 2015/02].

References

1.   Sacks O. The Mind's Eye. Alfred A. Knopf; 2010.

2.   Bouska MJ, Kauffman NA, Marcus SE. Disorders of the visual perceptual system. In: Umphred DA, editor. Neurological Rehabilitation. 2nd editio. St. Louis, MO: The C. V. Mosby Company; 1990. p. 705–40.

3.   Intercollegiate Stroke Working Party. National clinical guideline for stroke. 2016.

4.   Kolb B, Whishaw IQ. Fundamentals of human neuropsychology. 7th editio. New York, NY: Worth Publishers; 2003.

5.   Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence and prevalence of visual problems after acute stroke: An epidemiology study with implications for service delivery. PLoS One [Internet]. 2019 [cited 2019 Jul 2];14(3):e0213035. Available from: http://www.ncbi.nlm.nih.gov/pubmed/30840662

6.   Làdavas E. The Role of Visual Attention in Neglect: A Dissociation between Perceptual and Directional Motor Neglect. Neuropsychol Rehabil. 1994;4(2):155–9.

7.   Mancini F, Bricolo E, Mattioli FC, Vallar G. Visuo-haptic interactions in unilateral spatial neglect: The cross modal Judd illusion. Front Psychol. 2011;2(NOV).

8.   Chatterjee A, Thompson KA, Ricci R. Quantitative analysis of cancellation tasks in

neglect. Cortex. 1999;35(2):253–62.

9.   Rowe FJ, Brand D, Jackson CA, Price A, Walker L, Harrison S, et al. Visual impairment

following stroke: Do stroke patients require vision assessment? Age Ageing.

2009;38(2):188–93.

10.   Edmans J, Lincoln N. The frequency of perceptual deficits after stroke. Clin Rehabil

[Internet]. 1987 Nov 1;1(4):273–81. Available from:

http://cre.sagepub.com/cgi/doi/10.1177/026921558700100403

11.   Kerkhoff G. Neurovisual rehabilitation: Recent developments and future directions. J

Neurol Neurosurg Psychiatry. 2000;68(6):691–706.

12.   Jehkonen M, Ahonen JP, Dastidar P, Koivisto AM, Laippala P, Vilkki J, et al. Visual

neglect as a predictor of functional outcome one year after stroke. Acta Neurol Scand

[Internet]. 2000 Mar [cited 2016 Dec 8];101(3):195–201. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/10705943

13.   Plante M, Demers L, Swaine B, Desrosiers J. Association Between Daily Activities

Following Stroke Rehabilitation and Social Role Functioning Upon Return to the

Community. Top Stroke Rehabil. 2010;17(1):47–57.

14.   Mercier L, Audet T, Hébert R, Rochette a, Dubois MF. Impact of motor, cognitive, and

perceptual disorders on ability to perform activities of daily living after stroke. Stroke.

2001;32(11):2602–8.

15.   Kalra L, Smith DH, Crome P. Stroke in patients aged over 75 years: outcome and

predictors. Postgrad Med J [Internet]. 1993 Jan 1;69(807):33–6. Available from:

http://pmj.bmj.com/cgi/doi/10.1136/pgmj.69.807.33

16.   Bazeley P. Qualitative data analysis with NVivo. SAGE; 2007. 217 p.

17.   Krueger RA, Casey MA (Mary AW. Focus groups⬚: a practical guide for applied

research. SAGE; 2009. 219 p.

18.   Strategyzer | Value Proposition Canvas [Internet]. [cited 2019 Jul 26]. Available from:

https://www.strategyzer.com/canvas/value-proposition-canvas

19.   Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol [Internet].

2006 Jan;3(2):77–101. Available from:

http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa

20.   Hanna KL, Hepworth LR, Rowe FJ. Screening methods for post-stroke visual

impairment: a systematic review. Disabil Rehabil [Internet]. 2016 Sep 26;1–13.

Available from:

https://www.tandfonline.com/doi/full/10.1080/09638288.2016.1231846

21.   Kitsos G, Harris D, Pollack M, Hubbard IJ. Assessments in Australian stroke

rehabilitation units: a systematic review of the post-stroke validity of the most

frequently used. Disabil Rehabil [Internet]. 2011 Jan 9 [cited 2019 Aug 5];33(25–

26):2620–32. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21554012

22.   Bouwmeester L, Heutink J, Lucas C. The effect of visual training for patients with

visual field defects due to brain damage: A systematic review. J Neurol Neurosurg

Psychiatry. 2007;78(6):555–64.

23.   Hepworth LR, Rowe FJ. Visual Impairment Following Stroke - The Impact on Quality of

Life: A Systematic Review. Ophthalmol Res An Int J [Internet]. 2016;5(2):1–15.

Available from: http://sciencedomain.org/abstract/12864

24.   Pollock A, Hazelton C, Brady M. Visual Problems After Stroke: A Survey of Current

690   Practice by Occupational Therapists Working in UK Stroke Inpatient Settings. Top

691   Stroke Rehabil. 2011;18(sup1):643–51.

692