Current Practice and Challenges in Screening for Visual Perception Deficits after

Stroke: a Qualitative Study

Kathleen Vancleef, (corresponding author)

Department of Experimental Psychology, University of Oxford, Anna Watts Building,

Radcliffe Observatory Quarter, Woodstock Road, University of Oxford, OX2 6GG,

United Kingdom

Email: kathleen.vancleef@psy.ox.ac.uk

Twitter: @katvancleef

ORCiD: 0000-0002-9943-9341

Michael J Colwell,

Department of Experimental Psychology, University of Oxford

Twitter: @MichaelJColwell

Olivia Hewitt,

The Oxford Institute of Clinical Psychology Training and Research, University of

Oxford

Nele Demeyere,

Department of Experimental Psychology, University of Oxford

Twitter: @NeleDemeyere

ORCiD: 0000-0003-0416-5147

Abstract

PURPOSE: We document current clinical practice and needs in screening for visual

perception problems after stroke to inform development of new screening tools.

MATERIALS AND METHODS: We interviewed 12 occupational therapists and 13

orthoptists. Interviews were thematically analysed using the Value Proposition Canvas, a

model which establishes challenges and facilitators in what people want to achieve.

RESULTS: Participants' understanding of visual perception varied and often included

sensory and cognitive deficits. Occupational therapists commonly screened for visual field

deficits and hemispatial neglect, while other aspects of visual cognition were rarely assessed.

A positive screening result triggered an orthoptic referral. Screening generally occurred

during functional assessments and/or with in-house developed tools. Challenges to practice

were: lack of time, lack of training, environmental and stroke survivor factors (e.g. aphasia),

insufficient continuation of care, and test characteristics (e.g. not evidence-based).

Facilitators were: quick and practical tools, experienced staff or tools with minimal training

requirements, a streamlined care pathway.

CONCLUSIONS: Screening employs non-standardised assessments and rarely covers higher

visual perceptual deficits. We demonstrates the need for an evidence-based visual perception

screen, which should ideally be 15 minutes or less, be portable, and require minimal

equipment. The screen should be suitable for bedside testing and aphasia-friendly.

Keywords

Visual perception, screening, stroke, assessment, clinical needs, facilitators, barriers

Introduction

Visual perception is the dynamic process of perceiving the environment through sensory inputs and translating the sensory input into meaningful concepts associated with visual knowledge of the environment.(1) Visual perception problems are therefore distinct from sensory visual impairments such as reduced visual acuity, visual field and eye movements.(2) Where sensory visual impairments result from damage to the eye or early visual pathways from the eye to the primary visual cortex, visual perception deficits are attributed to impaired function in later visual processing areas in the occipital, parietal and temporal cortex.(3) Examples of visual perceptual deficits include apperceptive and associative agnosia (object recognition difficulties), prosopagnosia (face recognition difficulties), akinetopsia (difficulties in perceiving motion), achromatopsia (difficulties in perceiving motion), problems in visual memory (remembering what you have seen before), and in visuospatial abilities (e.g. judging distances or spatial relations between objects).(3) Visual inattention or hemispatial neglect is sometimes considered to be part of visual perception,(1)'(4) though neuropsychology research attributes this to an attentional deficit.(5) In particular, the presence of preserved perception when attention is stretched to focus on the stimuli, the existence of cross-modal neglect and manipulations of stimulus density on the extent of neglect support the classification of hemispatial neglect as a disorder of attention. (6,7)

Rowe and colleagues reported that 20.5% of stroke survivors with a suspected visual difficulty have visual perception deficits such as visual inattention (14%), visual hallucinations (2.5%), and object agnosia (2.2.%)(8) This study made use of reports by stroke survivors and carers rather than formal assessment (except for visual inattention). With systematic screening with the Rivermead Perceptual Assessment Battery, Edmans and Lincoln identified visual perception problems in 76% of hemiplegic stroke survivors.(9) The discrepancy between prevalence with self-reports compare to neuropsychological assessment

suggests that not all visual perceptual problems are picked up based on self-report. This

means many stroke survivors are discharged without the appropriate rehabilitation or

adjustments in their home environment or care packages.

Under-diagnosis of visual perception problems can severely impact stroke survivors' quality

of life, functional outcome, participation in the community, independence and pose

substantial risk. (10–14) For instance, participation in traffic with visual perception problems

can be dangerous and even life-threatening. Risks are also heightened indoors when

preparing a meal or in perceiving trip hazards. Better diagnosis of visual perception

difficulties after stroke will allow better care planning and substantially impact stroke

survivors' life.

The first step toward better diagnosis is an in-depth understanding of the clinical reality and

the reasons behind under-diagnosis. We need to understand what the challenges are in

screening for visual perceptual difficulties after stroke. This can then inform the development

of solutions in the future.

In the current study, we aimed to conduct an in-depth exploration of current clinical practice,

challenges and facilitators of screening for visual perception problems after stroke. To

achieve a rich understanding of these issues, we performed semi-structured interviews with

orthoptists and occupational therapists. These professionals are most commonly involved in

visual perception screening after stroke in the United Kingdom's National Health Service.

Materials and methods

Participants

All participants were recruited via opportunistic and snowball sampling. Invitations for the

interviews were sent out through (i) email to the British and Irish Orthoptic Society Stroke

and Neuro Rehab Clinical Advisory Group, (ii) Twitter, (iii) a sign-up sheet at a conference

poster and (iv) informal face-to-face conversations at the United Kingdom Stroke Forum

2018. Participants met inclusion criteria if they were working in the National Health Service

as an occupational therapist or an orthoptist, and were involved in the assessment of visual

perceptual problems after stroke. The participants were informed about the aims of the study,

the organisations running the study (University of Oxford and North Bristol NHS Trust), and

the planned outcomes and dissemination. The project was reviewed and approved by the

Patient Safety Assurance & Audit Service at NHS North Bristol Trust as a Clinical

Effectiveness study (CE45999). All participants gave verbal consent.

Interviewer's training

All interviews were conducted over the phone by the first author, KV, who holds an MSc in

Clinical Psychology and a PhD in Experimental Psychology. At the time of the research

study KV was working as a postdoctoral research fellow at the University of Oxford. KV had

nine years' experience as a researcher, including one year as a qualitative researcher. Her

formal training in qualitative research included an undergraduate course in qualitative

research methods at the University of Leuven which included a research project. In

preparation for the current project, she was trained and supervised by senior author, OH. OH

is a clinical psychologist who has worked predominantly as a qualitative researcher and

health care professional within the field of intellectual disabilities. OH has numerous

publications within this field, teaches qualitative research methods at the University of

Oxford, and has supervised a number of doctoral projects using various qualitative

methodologies. KV's training included reading and reflecting on reference works in

qualitative research,(15,16) a question and answer session with an experienced moderator,

and guidance during coding and analyses via regular meetings with OH.

Model

The interview process and qualitative analysis was guided by the Value Proposition Canvas.

This model was originally designed to guide product development, but can be applied to

different contexts.(17) Central to this model is the principle that development of products or

services are best informed by the users. A good product or service takes into account the

goals of the user, it reduces the pains a user experiences while trying to achieve their goals,

and it provides gains in achieving their goals. For instance a good electric car helps a person

to get to work (goal to achieve) without having to fear that the battery will run out (reduce

pains) while at the same time provides an attractive design to impress colleagues and family

(increase gains). We can apply this model in trying to understand current clinical practice in

visual perception screening after stroke (goal), the challenges (pains) it brings, and what

facilitates the process (gains).

Data collection

A semi-structured interview guide was developed by KV in collaboration with an

occupational therapist and an orthoptist (table 1). The interview guide was reviewed by an

experienced moderator and piloted with one orthoptist and three occupational therapists. The

interviews consisted of open-ended questions about current practice in visual perception

screening after stroke, and the challenges and facilitators of practice. The interviewer

conducted interviews at the participant's workplace or over the phone in a quiet office with

no other people present. The participants took the call in a location of their choice. The

location and presence of others at the participant's side was not recorded. After obtaining

participant consent, the interview was audio recorded. All participants were interviewed once.

Recruitment was ongoing during data collection and interviews were conducted until data

saturation was reached.

Table 1. Interview structure

Section	Interview route
Introduction	*Q.* Which duties are central to your job role?
Definition of visual perception	*Q.* What comes to your mind when I say visual perception problems in stroke survivors?
Current practice	*Q.* Do you do visual perception screening as a routine on your stroke survivors? (Orthoptists only) *Q.* What did you do the last time you suspected visual perception problems in a stroke survivor? *Cues: standardised tests, unstandardized tests, observations, conversation with stroke survivor, family members, other staff, medical notes* *Q.* What is the care pathway for these stroke survivor? *Cues: Which health care professional (occupational therapist,, neuropsychologist, orthoptist, consultant, physiotherapist) does what (clinical judgement, systematic screen, in depth assessment) and where (A&E, Acute stroke ward, rehab ward, care home, at home, outpatient appointment at eye clinic), referral process*
Facilitators to practice	*Q.* What does an initial visual perception screen need to be like for you? *Cues: time it takes, format, what to include, what result, who should do it*
Challenges to practice	*Q.* Can you name three things that frustrate you about visual perception screening on the stroke ward? *Cues: assessment methods, time, training, other staff, care pathway,* *Q.* Which of the three challenges you have mentioned is the most important?
Ending	*Q.* If you could give one piece of advice to someone making a visual perception screen, what would it be? *Q.* My goal was to have an idea of how visual perception screening is done and what the clinical needs are. Do you think we've talked about everything or is there anything we've missed?

Qualitative Analyses

As a method of data analysis we used thematic analysis.(18) Interviews were transcribed

verbatim by a professional typist. The transcripts were not returned to the participants for

feedback. Two researchers (KV and MC) were involved in data analysis. KV was familiar

with the data from conducting the interviews (Phase 1 in thematic analysis). The first 15

interviews were checked and coded by KV in NVivo Software (Version 12.0). She developed

an initial coding tree (Phase 2 in thematic analysis). Overall themes were set in advance in

alignment with the Value Proposition Model: Jobs, Challenges, and Facilitators. The coding

tree was reviewed and substantially adapted after coding of five interviews. MC familiarised

himself with the data by checking transcripts and reading through interviews coded by KV.

Two of these interview transcripts were checked, coded and discussed by both KV and MC to

align their coding. MC checked and coded the remaining 10 interview transcripts. The

frequency of text extracts coded under each node was visualised with a hierarchical chart of

nodes in NVivo to explore overall patterns in the data describing current practice. The use of

frequencies in qualitative analysis can be a valuable strategy when used as a complement to

an overall process orientation to the research. They can provide evidence for generalizability

of the findings between participants, highlight diversity, reveal larger patterns beyond a

participant's immediate experience, minimise cherry picking of the data.(19) This chart

shows the frequency of references for each node when participants were talking about current

practice in routine visual and visual perception screening. If a participant referred to a node

multiple times during one interview, each reference was counted. Only nodes with at least ten

references were included in the hierarchical chart of nodes. Overall themes and subthemes

were derived from all data by KV (Phase 3 of thematic analysis). This was done in two

stages: (1) we performed search queries of coded text for challenges and facilitators of visual

perception screening in current practice and (2) read through the resulting text. The themes

were subsequently reviewed by reading all answers to the questions under the headings

Challenges and Facilitators (see table 1). The information was then synthesised by KV (Phase

4 of thematic analysis). Lastly, themes and subthemes were reviewed, refined and defined by

KV (Phase 5 of thematic analysis) and a report was prepared (Phase 6 of thematic analysis).

This report was sent out to the participants for validation. Ten out of 25 participants replied

and nine agreed with the thematic analysis. One participant pointed out that participants'

understanding of visual perception was underexplored. We added a results section on this

topic posthoc.

Results

12 occupational therapists and 13 orthoptists from 16 health care organisations in England

took part in the study. Characteristics such as gender, years of experience and main clinical

setting of each participant are reported in table 2. All participants worked in England, and

were based across the East Midlands (n=1), East of England (n=1), Greater London (n=2),

North East (n=3), North West (n=4), South East (n=6), South West (n=4), and Yorkshire and

the Humber (n=4). The interviewer had no personal relationship with any of the participants.

Two orthoptists received referrals from the same acute stroke unit where the first author

recruits stroke survivors as participants for other studies (pseudonyms cannot be disclosed to

ensure anonymity). One of the participants was also involved in the development of the

interview guide. All but one participant (Jessica) agreed to be audio recorded. For Jessica,

notes were made during and immediately after the interview. Interviews lasted between 16

and 46 minutes with an average of 27 minutes. For Christopher, the interview was split over

two consecutive days.

Table 2. Participant characteristics

Pseudonym	Gender	Profession	Clinical setting[†]	Years of experience in stroke care
Amanda	Female	Orthoptist	Eye clinic	10+ years
Amber	Female	Occupational therapist	Rehabilitation service	10+ years
Amy	Female	Orthoptist	Eye clinic	unknown
Brittany	Female	Occupational therapist	Rehabilitation service	10+ years
Christopher	Male	Occupational therapist	Acute stroke service	unknown
Crystal[a]	Female	Occupational therapist	Rehabilitation service	5-10 years
Danielle	Female	Occupational therapist	Acute stroke service	unknown
Elizabeth	Female	Orthoptist	Eye clinic	10+ years
Emily[a]	Female	Occupational therapist	Rehabilitation service	unknown
Erin	Female	Occupational therapist	Acute stroke service	5-10 years
Heather[b]	Female	Orthoptist	Eye clinic	10+ years
Jamie[b]	Female	Orthoptist	Eye clinic	2-5 years
Jessica[c]	Female	Orthoptist	Eye clinic	10+ years
Joshua	Male	Orthoptist	Eye clinic	2-5 years
Kimberley	Female	Occupational therapist	Acute stroke service	2-5 years
Matthew	Male	Orthoptist	Eye clinic	5-10 years
Megan	Female	Occupational therapist	Rehabilitation service	10+ years
Melissa	Female	Occupational therapist	Acute stroke service	10+ years
Michelle	Female	Occupational therapist	Rehabilitation service	2-5 years
Nicole	Female	Orthoptist	Eye clinic	unknown
Rachel	Female	Orthoptist	Eye clinic	unknown
Rebecca	Female	Occupational therapist	Acute stroke service	5-10 years
Sarah[c]	Female	Orthoptist	Eye clinic	10+ years
Stephanie	Female	Orthoptist	Eye clinic	unknown
Tiffany	Female	Orthoptist	Eye clinic	2-5 years

[a,b,c] Participants working in the same clinical teams A, B or C. [†]All orthoptists were based in eye clinics but

could offer inpatient appointments on the stroke unit, inpatient appointments in the eye clinic and outpatient

appointments in the eye clinic.

Lack of clarity around visual perception

The interviews highlighted that participants' understanding of visual perception differed from

textbook definitions of visual perception. Visual perception is defined as a dynamic process

of translating sensory visual information into meaning percepts(1). These percepts can

subsequently be linked to higher cognitive functions like visual memory for recognition

(ventral or what stream) and praxis for visually guided actions (dorsal or where stream)(3).

Visual perception must be differentiated from low level sensory visual functions like acuity,

visual fields, and ocular movements, and from higher cognitive processes like (visual)

memory or (visual) inattention, although some authors consider visual inattention to be part

of visual perception. Our participants seemed to have a broader understanding of visual

perception and often did not differentiate between sensory and higher cognitive functions.

Participants typically mentioned the following examples when asked about what they

consider visual perceptual problems (in descending order of frequency): visual inattention,

agnosia's for objects, faces, letters, or shapes, hallucinations or Charles Bonnet syndrome,

difficulties in depth perception, visual field deficits like hemianopia, and spatial vision (e.g.

navigating and perceiving space around them).

Participants' understanding seemed to vary widely but we observed general trends in the

different professions. Orthoptists seemed to differentiate visual perception difficulties from

low level sensory visual impairments, but considered visual inattention the prime (and only)

example of visual perception difficulties after stroke.

"[*Interviewer: What comes to your mind, when I say visual perception problems in stroke*

patients?] So I would probably say visual neglect, ignoring one side of their world when the

other side is presented to them as well, is the main definition of it. That it's decreased

awareness of that side of their vision and their external world as well as themselves as well,

on that side. *[Interviewer: Anything else, that falls under that term for you?]* No, not that I

can think of, off the top of my head, no." (Tiffany, Orthoptist in an eye clinic)

In addition to visual inattention, occupational therapists mentioned impairments that are

typically (but not exclusively) related to low level sensory vision like contrast sensitivity,

hemianopia, and depth perception. They seemed to interpret visual perception more broadly.

"A range really. So we would look at visual perception as things like depth perception, …

recognition of objects, … and any kind of visual inattention or neglect, that sort of thing…

But yes, that's it basically." (Danielle, Occupation Therapist in an acute stroke service)

These observations are relevant in interpreting the subsequent results. Although the interview

questions asked for experiences with visual perception assessment after stroke, the results

below should be interpreted as potentially referring to all or any of the following: sensory

visual impairments (for occupational therapist participants), visual perception impairments

(for all participants), and visual inattention (for all participants).

Current Practice

A hierarchical chart of nodes is presented in figure 1. The figure illustrates that participants

most often mentioned occupational therapists working in acute stroke or rehabilitation

services when discussing screening for visual neglect, visual field deficits, and ocular

movements. For screening they referred to in-house developed screening tools, standardised

tests or observations in functional tasks like dressing and making a cup of tea.

"In terms of the visual field, we do the standard, you know, wiggle your fingers

test to see how people respond. We do the tracking tests. Otherwise it's mainly

through observing people in functional tasks what they're doing. ... Through that

you can often realise ... if they're having problems with their visual perception

because they're missing things or not seeing things." (Crystal, occupational

therapist in a stroke rehabilitation service)

"I'd rather do it in function. It's more meaningful especially with stroke patients

because it makes more sense. Rather than getting them to balance cubes on each

other or, you know, I can say 'Can you find the toothpaste?' " (Megan,

occupational therapist in an acute stroke service)

248
Figure 1. Hierarchical chart of nodes showing the frequency of references for each node. The size of a box

reflects the frequency of references for each node with larger boxes demonstrating more frequency mentioned

nodes. Parent nodes are shown in lightly shaded boxes, child nodes are shown in darker shaded boxes within

their parent's box. The colours are added for easy distinction between different parent boxes but have no further

meaning. Text is filtered for extracts where participants were talking about current practice in routine visual and

visual perceptions screening. Only nodes with at least ten references are shown.

Most participants mentioned that the service they work within has a referral pathway in place:

if problems are suspected following screening by the occupational therapist, a referral is

made to orthoptics. A stroke orthoptist will then see the stroke survivor as an inpatient on the

unit or as an outpatient after discharge from the hospital.

260   "It's really as and how other health professionals have identified them, then an

261   orthoptist will go up to try and assert a diagnosis and then give them the

262   rehabilitation options." (Amy, orthoptist in an eye clinic)

Challenges

Themes around challenges in visual perception screening are presented in table 3. We

identified five major themes and 13 subthemes.

Lack of Time

Occupational therapists and orthoptists both reported that time pressure makes visual

perception screening difficult. They mentioned that because of understaffing they spend little

time with stroke survivors or experience delays in seeing them following referral. In addition,

occupational therapists report that stroke survivors are discharged from an acute stroke unit

as soon as medically safe and a suitable discharge location has been organised. For survivors

with milder strokes, discharge is reported to happen within 1-2 days, giving very little time

for therapists to complete their assessments.

"Well one *[challenge]* is just logistically and organisationally, just in terms of

caseload and time to hand. We are having to prioritise more direct functional

assessments in terms of establishing readiness for discharge." (Christopher,

occupational therapist in an acute stroke service)

"There's not a lot of OTs who are actually for *[delivering]* the therapy, so time-

wise we haven't got a lot of capacity at times." (Danielle, occupational therapist in

an acute stroke service)

"Lack of time is always going to be frustrating knowing that you've got a clinic

here and you can't get up to the ward and there's an urgent stroke survivor."

(Heather, orthoptist in an eye clinic)

"Erm… it sounds awful doesn't it when you're constantly having to slim down

what you do." (Melissa, occupational therapist in an acute stroke service)

Lack of Training

Training is another theme that frequently emerged across both professions. Visual perception

difficulties following stroke appear to be only briefly covered in the undergraduate education

of orthoptists and occupational therapists, and is mostly learned about post-qualification

through on-the-job-training. Many occupational therapists mentioned that they rotate between

services, leaving little time for them to be trained in many aspects of stroke care. In

particular, they said that this lack of experience made them feel uncertain when it comes to

vision and visual perception deficits following stroke. Inexperience and limited training

seemed to be an important hurdle for occupational therapists in visual perception screening.

Because vision and visual perception was not well understood by the staff (see section 'Lack

of clarity around visual perception'), participants reported that mistakes in assessments are

commonly made, leading to a potential under-diagnosis of visual perception deficits.

"[It] can be a bit frustrating ... lots of the occupational therapists are on a

rotational job. So getting them [occupational therapists] trained up [is difficult], I

think they [occupational therapists] do find vision scary.... Vision, it's the scariest

thing about stroke." (Stephanie, orthoptists in an eye clinic, using 'vision' as a

general term to cover a wide range of sensory and perceptual visual functions)

"[Visual perception is] one of those things that is perhaps not widely

understood.... think unless you've got a unit with a special interest in stroke and

you've taken the time to look into that... then it's perhaps under-diagnosed as well,

so yeah I think it's recognition that it's a potential problem." (Amanda, orthoptist

in an eye clinic)

"The key is really, somebody in the hospital to be able to deliver the training and

the therapy staff to actually take on board the training and to actually deliver it."

(Sarah, orthoptist in an eye clinic)

311   "I think it probably is underreported; the amount of incidents of perceptual

312    problems post stroke." (Rachel, orthoptist in an eye clinic)

313 *Environmental and Stroke Survivor Factors*

314 Participants reported that visual perception screening is limited by the environment in which

315 they take place. They said that in acute stroke care, most assessments are done in a hospital

316 bay with several beds. Even if curtains can be drawn around the stroke survivor's bed,

317 distractions were reported to be numerous: sounds from neighbouring beds, interruptions for

318 timed caring needs, distractions from medical equipment around the bed and presence of

319 personal objects on a small table in front of the stroke survivor (spectacles, drinks, food,

320 tissues, personal care items, newspaper, get well cards, crossword book, pen, etc.) can be

321 visually distracting and interfere with visual perceptual assessment.

322   "Frustrating for us is ... that we .. depending on which unit we are *[on]* or where

323    we are, it can be quite difficult to do *[our assessment at]* the bedside if there is a

324    lot *[of]* distractions going on; if there is no quiet space to take the patient.

325    Obviously, yes, we can draw the curtains but it's sometimes more the noise levels

326    than everything. They get easily distracted. That makes it quite hard actually to do

327    an assessment." (Amber, occupational therapist in a stroke rehabilitation service)

328   "It's *[the]* environment. Often *[it]* doesn't lend itself to a nice quiet space where

329    patients can attend." (Danielle, occupational therapist in an acute stroke service)

330 Participants also mentioned the condition of the stroke survivors as a limiting factor to visual

331 perception screening. Both professions reported that tests and tools are often not suitable for

332 stroke survivors with poor alertness, poor cognition, aphasia or severe weakness in their

333 upper limbs. For instance, stroke survivors may not able to answer a question on whether

334 their vision has changed because they developed aphasia following stroke. Similarly, copying

and cancellation tasks to assess visual neglect were reported to be unsuitable for a stroke

survivor who can no longer hold a pen.

"Another component would be around the client or the patient themselves in terms

of factors that might work against screening: so attention, concentration, fatigue,

the ability to deliver a verbal response in terms of aphasia or dysarthria

difficulties. ... That has significant impact on our ability to undertake formal

testing of whatever nature." (Christopher, occupational therapist in an acute stroke

service)

"Our patients who either are fatigued quite quickly, or are ... medically not well

enough to do a lot of taking down to the therapy department ..., or their attention

span is quite limited, or they've got a weakness in their upper limb that makes it

more difficult to complete the tests." (Danielle, occupational therapist in an acute

stroke service)

Continuation of Care

A further concern of the participants was the follow-up process after screening for visual

perception deficits. They highlighted that quite often the information from screening is not

passed on to the next care team. Therefore, even if visual perception problems are picked up

in an acute setting, no follow-up assessments are carried out or no rehabilitation is provided

in their experience. With respect to rehabilitation, occupational therapists found it frustrating

that few treatment options are available. They reported that little information and few

guidelines are available on how visual perception problems can be treated or managed. Some

noted that the lack of options for rehabilitation reduced their motivation to screen for visual

perception problems.

"Yes, and it [*the discharge summary*] is very medical because the occupational

therapists or myself don't like the discharge summaries. ... We do all this

information and then it gets lost ….Yeah, it's not part of our discharge summary

in enough detail, so when they go somewhere new which could be two or three

weeks later ... they don't have that information." (Matthew, orthoptist in an eye

clinic)

"It might be because it [*visual perception deficits*] is limited in treatment … that's

why it's not the focus. So … it would be more useful to spend time during my

assessment speaking to them about adapting to their field loss and giving them a

prism or occlusion for their double vision." (Rachel, orthoptist in an eye clinic)

Test Characteristics

Both professions highlighted limitations in the characteristics of the available tests for visual

perception. Occupational therapists more frequently reported to be hindered by practical

limitations and the unknown impact on daily life. With respect to the practical limitations,

they referred to a large number of test materials like stimulus booklets, stopwatch, cubes, etc.

needed for assessment or too many loose pages. With respect to the unknown impact on daily

life, they referred to stimuli and tests that were thought to be limited in ecological validity.

They considered the tests too abstract with no clear link to the implications for everyday

tasks.

"We have our little screening tool. It's a few pages, just to prompt us .... That's all

right but then ... everything else you are ... searching together from different kits.

... That's a little bit frustrating." (Amber, occupational therapist in a stroke

rehabilitation service)

"There's an awful lot of bits of paper and a bit of faffing about and preparing to

get it all ready before we actually go out to see the patients." (Melissa,

occupational therapist in an acute stroke service)

384   "We use Rivermead Perceptual Battery. But to be fair, I don't tend to use it as

385   often because I'd rather do it in function. ...It's more meaningful, especially with

386   stroke patients because it makes more sense. Rather than getting them to balance

387   cubes on each other or, you know I can say "Can you find the toothpaste? Can

388   you take the lid off? Can you brush your own teeth?" and then it just makes more

389   sense. And then ... it's easier for them to identify *[a]* goal." (Megan, occupational

390   therapist in a stroke rehabilitation service)

391   Orthoptists, on the other hand, seemed to emphasize more the lack of evidence and

392   consequently the lack of specific clinical guidelines on how to assess visual perception.

393   "If you see the national guidelines for visual problems after stroke, they have a

394   really small section on vision with orthoptics but we don't really have anything in

395   it that's specific to what we should be screening them with and what we should be

396   treating them with. So even though we're using evidence-based practice, the

397   evidence sometimes isn't great and I think that's probably why it's not in the

398   guidelines because there's not really good evidence to support it. But sometimes

399   you're seeing a patient and you're not entirely sure if you're doing the right thing

400   because you're maybe giving them information based on clinical experience, what

401   other people have been doing but you're not really sure if that is the best thing to

402   be doing or could you be doing something else with them." (Amy, orthoptist in an

403   eye clinic)

405   Table 3. Challenges in visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Lack of time	Limited time with stroke survivor due to discharge pressure	x	
	Limited time with stroke survivor due to staff shortage	x	x
Lack of training	Inexperienced staff with limited training	x	
	Vision and visual perception is not well understood	x	x
	Mistakes made in assessment	x	x
	Vision and visual perception deficits are underdiagnosed	x	x
Environmental and stroke survivor factors	Condition of stroke survivor (e.g. low alertness, poor cognition, aphasia, not mobile)	x	x
	Distractions in test environment (often in hospital bay)	x	x
Continuation of care	Information on vision and visual perception problems gets lost after discharge	x	x
	No information on treatment or management of vision or visual perception problems	x	
Test characteristics	Test materials not readily available or making use clumsy	x	
	Lack of evidence-based screening instruments		x
	No specific recommendations in clinical guidelines		x
	Tests are not meaningful for everyday life	x	

Facilitators

408   The facilitators to practice that participants reported, largely mirrored the challenges reported

409   above (table 4). We identified three major themes and 12 subthemes.

Practical Tool

On a practical level, participants preferred a tool that can be administered in approximately

10-20 minutes, so it can fit in with busy every-day practice and current staff levels. They also

favoured a tool that consists of few test materials and pages to prevent test materials getting

lost or mixed up during testing. According to participating orthoptists, the tool should have a

1 to 2 page proforma. They reported that it should include questions to ask the stroke survivor

and simple quick tests for a range of functions. They suggested that the proforma could

subsequently be used in a referral for orthoptic input. To allow bedside testing or testing in

the community, participants preferred portable tools. Lastly, many participants highlighted

that a tool should ideally be suitable for all stroke survivors, such as those with aphasia, poor

concentration, poor cognition or severe fatigue.

"It needs to be something that's fairly quick and easy to administer." (Danielle,

occupational therapist in an acute stroke service)

"Just pen and paper, quite simple or a few things to hand that will fit in something

that's quite portable and mobile, I think [*that*] would be handy, yes." (Brittany,

occupational therapist in a stroke rehabilitation service)

"Reasonably short … for logistical [*reasons*]: … our time and patient

concentration time, fatigue in the more acute stages. I think really half an hour is

the maximum time." (Christopher, occupational therapist in an acute stroke

service)

"It would have to be a list of standardised questions: Are they able to look

between two points specifically? … I guess the visual perception question would

be if they're following a moving object? … You know are they able to identify

their friends or their family? Can they recognise themselves in a photo?" (Rachel,

orthoptist in an eye clinic)

Training and Experience

Orthoptists and occupational therapists agreed on the importance of training and experience

for robust visual perception screening. The presence of experienced colleagues or training by

other staff members was seen as very advantageous. Given the limited training in visual

perception highlighted above, a tool that is easy to administer and score was preferred by our

participants. Similarly, they reported that results should be easy to interpret, and should

provide clear guidance for management of the deficit (if present).

"If it was something that was quite self-explanatory then anybody could do it and

therefore anybody can interpret it, then that would be much clearer." (Amy,

orthoptist in an eye clinic)

"Just easy to administer and easy to score. … So making instructions clear and

easy to understand would be good." (Emily, occupational therapist in a stroke

rehabilitation service)

Staff Roles and Interactions

Occupational therapists reported that their preferred practice is to screen stroke survivors for

vision and visual perception problems during their functional assessments (e.g. washing,

making a cup of tea) or via a screening tool. Both professions mention that if a problem is

flagged, more in-depth assessment can then take place.

"Whereas since we've set up a proper vision screening it just means that things

have been a lot more efficient and that's made a much better service because

they're not having to see a lot of people that don't need to be seen." (Nicole,

orthoptists in an eye clinic)

"So, I'm happy with the main basic screen to begin with, and if it digs something

up then we'll dig it in some more detail. I think that works quite well." (Matthew,

orthoptists in an eye clinic)

If the problem concerns visual perception difficulties, an occupational therapist is seen by our

participants as the most appropriate professional to follow it up. If the problem concerns

sensory vision, a referral to an orthoptist was the preferred practice in our sample. Good

relationships between stroke occupational therapists and orthoptists in eye clinics were

frequently mentioned as invaluable.

"I think the key is having the staff, the occupational therapist and the physio staff

actually take that on board." (Sarah, orthoptist in an eye clinic)

Table 4. Facilitators to visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Practical tool	Quick to administer (10-20 min)	x	x
	Paper and pen based without too many test materials or pages	x	x
	Proforma with questions and simple tests to be used in referral		x
	Portable	x	
	Suitable for stroke survivors with poor concentration, communication and fatigue	x	
Training and experience	Easy to administer and score	x	x
	Easy to interpret with management advice included	x	x
	Trained and experienced staff	x	x
Staff roles and interactions	Screening for visual and visual perception problems by occupational therapists during functional assessment or with screening tool	x	x
	In depth assessment with range of complimentary tests	x	x
	Specialised staff for visual (orthoptists) and visual perceptual (occupational therapists) problems		x
	Good relationship between stroke and eye departments		x

Discussion

The aim of this qualitative research was to gain a deeper understanding of current practice in

visual perception screening after stroke, the challenges faced by health care professionals. In

our participants' stroke care services, occupational therapists most often screened for visual

and visual perceptual difficulties. If visual difficulties like hemianopia, double vision or

reduced acuity, are found, a referral to an orthoptist is made for an in depth assessment. This

referral strategy corresponds to an earlier report of 76.7% of stroke units having access to an

orthoptic stroke service.(20) If visual neglect is suspected, occupational therapists will often

follow this up themselves with a range of tasks like cancellations, copying, line bisection, etc.

The biggest challenges faced by our participants were lack of time and training. Facilitating

factors mentioned were simple, easy to use instruments and good relationships between

stroke and eye departments.

The qualitative approach of our study allowed us to gather views directly and in depth.

Instead of reporting frequency of views (e.g. how many health care professionals report staff

shortage), we focus on how central certain views are to a person's experience. Our sample

covers the two main professions involved in visual perception screening after stroke.

National Clinical Guidelines recommend that every stroke survivor who appears to have

perceptual difficulties should have an assessment with standardised measures.(2) As is

evident from our interviews, only a small proportion of stroke survivors receive an

assessment with standardised measures for visual perceptions difficulties, and screening is

not routinely done. The means of assessment of visual perception deficits varied widely

between participants. Orthoptists will often use standardised tools in their assessments,

though these are not necessarily validated with stroke survivors.(21,22) Overall, standardised

assessments were rarely mentioned by our participants. Occupational therapists expressed

preferring assessment during functional tasks like washing and dressing and assessment with

in-house developed screening checklists. This may be due to a lack of a validated screening

tool for a wide range of visual and visual perceptual deficits that can be used by any health

care professional.(21) The diversity in tools that we observed in our group of English

participants contrasts with the reported use of standardised tools in an Australian population

where the Occupational Therapy Adult Perceptual Screening Test and the Loewenstein

Occupational Therapy Cognitive Assessment are most commonly used to evaluate visual

perceptual difficulties.(22)

McCluskey and colleagues investigated the barriers and enablers for following recommended

practice in stroke care.(23) In line with our findings the occupational therapists they

interviewed reported the condition of the stroke survivors as a barrier for standardised

assessments. Many tests are not designed and not suitable for stroke survivors with aphasia

and/or dysarthria. Occupational therapists in their study also mentioned the lack of training

and knowledge, specifically for visual neglect rehabilitation. Time pressure and fluctuating

staff levels meant occupational therapists in their study prioritised assessments and

interventions which would produce the best clinical outcome. A lack of time to see stroke

survivors was also mentioned by their orthoptists. However, in our study this referred to the

lack of time to see all referred stroke survivors before they were discharged, while in the

study by McCluskey, the orthoptists referred to the lack of time to treat stroke survivors. The

standardised assessments used by Australian occupational therapists, Occupational Therapy

Adult Perceptual Screening Test and the Loewenstein Occupational Therapy Cognitive

Assessment, require training and take 20-45 minutes to complete. Given the frequently

reported lack of training and time, it may appear that the standardised tools recommended by

clinical guidelines are not adapted to the clinical reality of the National Health Service in

England. Although McCluskey's study was limited to one Australian hospital and only

involved 5 occupational therapists and 2 orthoptists, all working in the same service and

circumstances, the barriers are remarkably similar to what our 12 occupation therapists and

13 orthoptists reported experiencing in 16 English healthcare services. The stroke survivor's

condition, lack of time and staff seem to be universal barriers for providing evidence-based

stroke care across domains, not just visual perception screening. Also, the lack of information

on treatment or management of vision or visual perception problems that occupational

therapists mention as a barrier is not surprising given the very limited evidence for treatment

and management options.(24)

The first challenge exposed by our study is the variation between participants in their

understanding of constitute visual perception problems after stroke. This led to participants

answering interview questions with different reference frameworks. When sharing the

challenges of visual perception screening in their clinical practice, some might have been

considering challenges in just screening for visual inattention difficulties while others

considered a broad range of sensory and cognitive visual impairments. To maintain an equal

relationship between participant and interviewer, and an open non-judgemental environment

to freely share their experiences, participants' definition of visual perception was not

challenged during the interview. A disadvantage of this approach is that it is unclear whether

certain themes apply specifically to one aspect of visual and visual perception screening or to

the whole range of vision related difficulties after stroke.

The second challenge lies in our sample size. Our sample, though the largest for a qualitative

study in this topic, is not representative for the whole population of health care professionals

involved in visual perception screening after stroke. We limited ourselves to occupational

therapists and orthoptists as they are most commonly involved in visual perception screening

in England. Experiences might be very different in clinical settings without an orthoptic

department or with more involvement of neuropsychologists in stroke care. In addition,

voluntary participation and asking our participants to commit 30 minutes of their time might

have induced a bias for allied health professionals with a keen interest in the topic. We have

possibly missed opinions from people with a more negative or neutral attitude towards visual

perception screening. In addition, newly qualified health care professionals with limited

experience in stroke care might not have felt confident enough to share their opinions on the

topic. Lack of confidence on this topic with junior staff was brought up by several of our

participants. However, the aim of this qualitative research was not to generalise findings to

the population, but rather to provide an in-depth exploration of the topic to generate

hypotheses. We are following this research up with a large scale survey for all health care

professionals involved in visual perception screening in the United Kingdom and Republic of

Ireland. In the recruitment to this survey study, we are emphasizing that we welcome the

opinions of heath care professionals with all levels of experience.

A third potential limitation is that the interviewer was the principal investigator of the study

and was therefore not independent. Unconscious bias of the interviewer might have guided

the participants' answers. We have tried to minimize this by using a detailed interview guide

that included a list of cues that could be given. In addition, all interviews were recorded and a

second coder was involved.

Our study limits itself to health care professionals' perceptive on visual perception screening

after stroke. We have not explored experiences by stroke survivors, although they can be

substantially different. Also, we have not investigated the management of visual perceptual

problems after stroke of their impact in daily life as is previously reported for sensory visual

problems. (25–27)

The current research highlighted that a clear and consistent definition of visual perception

should be provided to orthoptists and occupational therapists. More training is needed on the

assessment (and management) of visual perceptual screening. This is preferably provided via

professional organisations to ensure consistency and in line with national guidelines(2,28).

This will raise awareness and reduce insecurity experienced by junior staff members.

Establishing good relationships between staff at stroke and eye departments is strongly

recommended. We suggest this could be achieved in the form of knowledge exchanges or via

a re-evaluation of the existing referral system. Lastly, a quick and portable visual perception

screening tool that is easy to administer, score and interpret would highly benefit both staff

and stroke survivors if it is inclusive for stroke survivors with aphasia, motor impairments

and cognitive problems. The tool should be evidence-based and self-explanatory to use. Such

a tool is currently not available. Our research provides test developers with the requirements

of occupational therapists and orthoptists for such a screening tool to be clinically useful and

thereby increase the likelihood of adoption.

Acknowledgments

We would like to express our sincere gratitude to all the participating allied health

professionals who gifted their time. In addition we would like to thank Philip Clatworthy and

Claire Murray for facilitating this project. This work was supported by the Stroke Association

under TSA PDF 2017/03; and under TSA LECT 2015/02.

Declaration of interest statement

The authors report no conflicts of interest.

Data availability statement

Given the small group of stroke orthoptists and stroke occupational therapists in England,

sharing the interview transcripts would potentially disclose participants' identity.

References

1.   Bouska MJ, Kauffman NA, Marcus SE. Disorders of the visual perceptual system. In:

592   Umphred DA, editor. Neurological Rehabilitation. 2nd editio. St. Louis, MO: The C.

593   V. Mosby Company; 1990. p. 705–40.

594   2.   Intercollegiate Stroke Working Party. National clinical guideline for stroke. 2016.

595   3.   Kolb B, Whishaw IQ. Fundamentals of human neuropsychology. 7th editio. New

596   York, NY: Worth Publishers; 2003.

597   4.   Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence

598   and prevalence of visual problems after acute stroke: An epidemiology study with

599   implications for service delivery. PLoS One [Internet]. 2019 [cited 2019 Jul

600   2];14(3):e0213035. Available from: http://www.ncbi.nlm.nih.gov/pubmed/30840662

601   5.   Làdavas E. The Role of Visual Attention in Neglect: A Dissociation between

602   Perceptual and Directional Motor Neglect. Neuropsychol Rehabil. 1994;4(2):155–9.

603   6.   Mancini F, Bricolo E, Mattioli FC, Vallar G. Visuo-haptic interactions in unilateral

604   spatial neglect: The cross modal Judd illusion. Front Psychol. 2011;2(NOV).

605   7.   Chatterjee A, Thompson KA, Ricci R. Quantitative analysis of cancellation tasks in

606   neglect. Cortex. 1999;35(2):253–62.

607   8.   Rowe FJ, Brand D, Jackson CA, Price A, Walker L, Harrison S, et al. Visual

608   impairment following stroke: Do stroke patients require vision assessment? Age

609   Ageing. 2009;38(2):188–93.

610   9.   Edmans J, Lincoln N. The frequency of perceptual deficits after stroke. Clin Rehabil

611   [Internet]. 1987 Nov 1;1(4):273–81. Available from:

612   http://cre.sagepub.com/cgi/doi/10.1177/026921558700100403

613   10.   Kerkhoff G. Neurovisual rehabilitation: Recent developments and future directions. J

614   Neurol Neurosurg Psychiatry. 2000;68(6):691–706.

615   11.   Jehkonen M, Ahonen JP, Dastidar P, Koivisto AM, Laippala P, Vilkki J, et al. Visual

616   neglect as a predictor of functional outcome one year after stroke. Acta Neurol Scand

617   [Internet]. 2000 Mar [cited 2016 Dec 8];101(3):195–201. Available from:

618   http://www.ncbi.nlm.nih.gov/pubmed/10705943

619   12.   Plante M, Demers L, Swaine B, Desrosiers J. Association Between Daily Activities

620   Following Stroke Rehabilitation and Social Role Functioning Upon Return to the

621   Community. Top Stroke Rehabil. 2010;17(1):47–57.

622   13.   Mercier L, Audet T, Hébert R, Rochette a, Dubois MF. Impact of motor, cognitive,

623   and perceptual disorders on ability to perform activities of daily living after stroke.

624   Stroke. 2001;32(11):2602–8.

625   14.   Kalra L, Smith DH, Crome P. Stroke in patients aged over 75 years: outcome and

626   predictors. Postgrad Med J [Internet]. 1993 Jan 1;69(807):33–6. Available from:

627   http://pmj.bmj.com/cgi/doi/10.1136/pgmj.69.807.33

628   15.   Bazeley P. Qualitative data analysis with NVivo. SAGE; 2007. 217 p.

629   16.   Krueger RA, Casey MA (Mary AW. Focus groups⬚: a practical guide for applied

630   research. SAGE; 2009. 219 p.

631   17.   Strategyzer | Value Proposition Canvas [Internet]. [cited 2019 Jul 26]. Available from:

632   https://www.strategyzer.com/canvas/value-proposition-canvas

633   18.   Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol

634   [Internet]. 2006 Jan;3(2):77–101. Available from:

635   http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa

636   19.   Maxwell JA. Using Numbers in Qualitative Research. Qual Inq [Internet]. 2010 Jul

637   15;16(6):475–82. Available from:

638   http://journals.sagepub.com/doi/10.1177/1077800410364740

639   20.   Hepworth L, Rowe F. Ten Years On – A Survey of Orthoptic Stroke Services in the

640   UK and Ireland. Br Ir Orthopt J [Internet]. 2019 May 8;15(1):89–95. Available from:

641   http://www.bioj-online.com/articles/10.22599/bioj.135/

642   21.   Hanna KL, Hepworth LR, Rowe FJ. Screening methods for post-stroke visual

643   impairment: a systematic review. Disabil Rehabil [Internet]. 2016 Sep 26;1–13.

644   Available from:

645   https://www.tandfonline.com/doi/full/10.1080/09638288.2016.1231846

646   22.   Kitsos G, Harris D, Pollack M, Hubbard IJ. Assessments in Australian stroke

647   rehabilitation units: a systematic review of the post-stroke validity of the most

648   frequently used. Disabil Rehabil [Internet]. 2011 Jan 9 [cited 2019 Aug 5];33(25–

649   26):2620–32. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21554012

650   23.   McCluskey A, Vratsistas-Curto A, Schurr K. Barriers and enablers to implementing

651   multiple stroke guideline recommendations: a qualitative study. BMC Health Serv Res

652   [Internet]. 2013 Dec 19 [cited 2019 Aug 5];13(1):323. Available from:

653   http://www.ncbi.nlm.nih.gov/pubmed/23958136

654   24.   Bouwmeester L, Heutink J, Lucas C. The effect of visual training for patients with

655   visual field defects due to brain damage: A systematic review. J Neurol Neurosurg

656   Psychiatry. 2007;78(6):555–64.

657   25.   Hepworth LR, Rowe FJ. Visual Impairment Following Stroke - The Impact on Quality

658   of Life: A Systematic Review. Ophthalmol Res An Int J [Internet]. 2016;5(2):1–15.

659   Available from: http://sciencedomain.org/abstract/12864

660   26.   Pollock A, Hazelton C, Brady M. Visual Problems After Stroke: A Survey of Current

661   Practice by Occupational Therapists Working in UK Stroke Inpatient Settings. Top

662   Stroke Rehabil. 2011;18(sup1):643–51.

663   27.   Rowe FJ. Stroke survivors' views and experiences on impact of visual impairment.

664   Brain Behav [Internet]. 2017 Sep;7(9):e00778. Available from:

665   http://doi.wiley.com/10.1002/brb3.778

666   28.   NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE (NICE) .

667   Stroke Rehabilitation: Long term rehabilitation after stroke [Internet]. NICE. Londo;

668   2013. Available from: http://www.nice.org.uk/nicemedia/live/14182/64094/64094.pdf

669