

Out of hospital cardiac arrests in the Gulf Region: a scoping review

Authors

Alan M. Batt¹, Chelsea Lanos², Dalal Al-Hasan³, Assim Alhmoudi⁴, Megan Anderson⁵, Shane Knox⁶, Shannon Delpont⁷, Saleh Fares⁸, Fergal H. Cummins⁹

Affiliations

1. Fanshawe College, 1001 Fanshawe College Blvd, London, ON N5Y5R6, Canada; CQUniversity, 554-700 Yaamba Rd, Norman Gardens, QLD 4701, Australia; Monash University, VIC, Australia; REDSPOT Medical International, Cork, Ireland
2. County of Renfrew Paramedic Service, 9 International Dr, Pembroke, ON K8A 6W5, Canada; Cardiff University, Cardiff, CF10 3AT, United Kingdom.
3. Department of Applied Medical Sciences, Health Sciences College, Public Authority of Applied Education and Training, State of Kuwait.
4. Emergency and Public Safety, Abu Dhabi Police, Sheikh Zayed Road, Al Muroor, Abu Dhabi, UAE.
5. Fanshawe College, 1001 Fanshawe College Blvd, London, ON N5Y5R6, Canada.
6. National Ambulance Service College, Creagh Road, Ballinasloe, Ireland; College of Medicine and Health, University College Cork, Brookfield House, College Road, Cork, Ireland.
7. CQUniversity, 554-700 Yaamba Rd, Norman Gardens, QLD 4701, Australia
8. Zayed Military Hospital, Al Khaleej Al Arabi Street, Abu Dhabi, UAE; Emirates Society of Emergency Medicine, Abu Dhabi, UAE.
9. University Hospital Limerick, Raheen, Limerick, Ireland; University of Limerick, Castletroy, Limerick, Ireland; Charles Sturt University, Panorama Ave, Bathurst, NSW 2795, Australia, Australia; REDSPOT Medical International, Cork, Ireland

Corresponding author: Alan M. Batt, Fanshawe College, 1001 Fanshawe College Blvd, London, ON N5Y5R6, Canada. Email: abatt@fanshawec.ca

Author Declarations

The authors declare that this work has not been published elsewhere. Data from this study were presented at the Emirates Society of Emergency Medicine scientific meeting on December 13th 2019 in Abu Dhabi, UAE. The authors declare that they are responsible and accountable for the accuracy and integrity of all aspects of this work.

Author Contributions

AB, CL, SD, SK, MA, SF and FC designed the study. MA and AB designed and conducted the literature searches. CL and AB performed screening and data abstraction. DA and AA contributed additional information to data analysis. AB authored the first draft of the

Batt et al. Out of hospital cardiac arrests in the Gulf Region: a scoping review. Pre-print 07/12/2019

manuscript. All authors contributed to the authoring and editing of the manuscript. All authors have approved the final version for publication.

Funding

This research was supported by a research grant from the ZOLL Foundation, who had no role in the design or conduct of the study; the collection or analysis of the data; or the drafting or submission of the manuscript.

Conflicts of interest

None declared

Abstract

Background

Out-of-hospital cardiac arrest (OHCA) is a major cause of mortality worldwide. Recent studies demonstrated low survival rates in Middle Eastern countries. Anecdotally there are unique demographic, cultural and logistical challenges in this region. However, there remains a paucity of data published on OHCA in the Middle East. In order to address OHCA in a meaningful manner in the region, we first need to quantify the issue.

Methods

We conducted a scoping review of published and grey literature on OHCA in the Gulf Cooperative Council region that utilised Arksey and O'Malley's framework. Electronic databases and grey literature sources were identified and searched. Subject matter experts in the region were consulted. All types of studies in English and Arabic were included.

Results

A total of 24 studies were included from Saudi Arabia, the UAE, Oman, Kuwait, and Qatar. No literature was identified from the state of Bahrain. OHCA victims in the region are younger, predominantly male, and more co-morbid than those in Western studies. We observed low Emergency Medical Service utilisation, low bystander cardiopulmonary resuscitation, return of spontaneous circulation, and survival to discharge rates across the region. There are differences in characteristics of OHCA among ethnic groups.

Discussion and conclusions

We identified unique characteristics associated with OHCA in the region, variances in processes and outcomes when compared to Western studies, and a lack of coordinated effort to research and address OHCA. We recommend creating lead agencies responsible for coordinating and developing strategies such as community response, public education, and reporting databases.

Introduction

Out-of-hospital cardiac arrest (OHCA) is a major cause of mortality worldwide, with variable survival reported across countries and systems. In particular, Middle-Eastern and South Asian countries report low survival rates¹. Reported reasons for low OHCA survival in these contexts include unique demographic, cultural, and logistical challenges related to the management of OHCA. Such contexts may also present a challenge when we attempt to implement evidence produced for the most part in Western systems. Characteristics and outcomes of OHCA in Middle Eastern countries remains poorly researched, compounding these challenges. The resulting lack of knowledge prevents us from understanding the issue, and thereby prevents us from improving the issue. Thus, to meaningfully address OHCA in the Middle East, we first need to understand the issue within its context.

Within the Middle East, the Cooperation Council for the Arab States of the Gulf, also known as the Gulf Cooperation Council (GCC, مجلس التعاون الخليجي) was established in 1981, and comprises the governments of the states of the Kingdom of Bahrain, Kuwait, the Sultanate of Oman, Qatar, the Kingdom of Saudi Arabia, and the United Arab Emirates (UAE). The countries in the GCC are predominantly Islamic, and they share a common cultural and historical background². Another shared trait is their economic reliance on expatriate workforces³, estimated at 48% of the total regional population of 53 million^{4,5}. Within this regional population life expectancy has risen dramatically since the 1960s to 77 years (2017)⁵. However, increased population numbers, the healthcare needs of different demographics and increased life expectancy have led to a rise in healthcare demands. This rise is predominantly due to high rates of non-communicable diseases such as diabetes, cardiovascular disease, and obesity, attributable to unhealthy lifestyles and a general lack of coordinated public health programmes^{4,6}. In order to reduce the risk associated with such morbidity, the majority of GCC member countries have initiated public health strategies which include the recording and reporting of non-communicable diseases

Sasser et al. first reported on out-of-hospital emergency care in Abu Dhabi in 2004⁷, when emergency medicine in the GCC was rudimentary at best. They observed no real medical control, no treatment policies, a lack of training and education, and no quality assurance policies, which echoed the situation across the region. However, emergency medicine and prehospital care have developed across the GCC in the last 15 years. Reporting of OHCA rates, response, and outcomes are also considered benchmarks of emergency medical response readiness. As a result, we elected to conduct a review of OHCA studies in the GCC in order to collectively summarise their characteristics and properties, as well as to identify any gaps and areas for further research.

Methods and analysis

We conducted a scoping review, which enabled us to identify, map and present an overview of a heterogeneous body of literature^{8,9}. We deemed a scoping review to be appropriate given the predicted lack of published data, the anticipated heterogeneity among studies and the

expected variability in reporting. We performed a systematic literature search for studies that investigated OHCA in the GCC region. We utilised Arksey and O'Malley's five-stage framework which included (1) identifying the research question, (2) identifying relevant studies, (3) refining the study selection criteria, (4) collecting relevant data from each article, and (5) collating, summarizing, reporting, and interpreting the results. We also implemented a recommended additional stage, (6) consultation with experts. We reported our process according to the PRISMA Extension for Scoping Reviews¹⁰.

1. Identify the research questions

1. What are the characteristics of out-of-hospital cardiac arrest cases in the GCC region?
2. How do these studies differ from cases in Western countries which are predominantly used to inform resuscitation guidelines?
3. What insights can be gained from these findings in order to better inform policy and practice in the GCC region?

Ethics and dissemination

This is a scoping study of completed studies and hence no ethics approval was required.

2. Identify relevant studies

Systematic search

An information scientist (MA) developed the search strategy. We used this to search the electronic databases MEDLINE, CINAHL, Web of Science and EMBASE from 1990 until June 2019. The search strategy was modified accordingly for each database. Relevant non-indexed journals were identified and searched, and we performed multiple Google web and Google Scholar searches. We searched the grey literature with guidance from the Grey Matters checklist¹¹. References of included studies were scanned for relevant articles. See Appendix I for search terms and a list of non-indexed sources.

3. Select the studies

Eligibility criteria

Studies in English and Arabic were eligible for inclusion if they reported OHCA in the GCC. We excluded studies that were not related to OHCA (e.g. in-hospital cardiac arrest), case-reports, studies that reported only on traumatic OHCA, studies that were related to cardiopulmonary resuscitation (CPR) training, and those not conducted in GCC countries.

Title and abstract screening

Initial screening comprised of a review of title and abstracts by two reviewers (AB and CL). We resolved disagreements through discussion until consensus was achieved. Where disagreement remained or there was insufficient evidence to make a decision, we included the citation for full text review and subsequent decision.

Batt et al. Out of hospital cardiac arrests in the Gulf Region: a scoping review. Pre-print 07/12/2019

4. Chart the data

To support the full-text review, we developed a standardised data extraction form to organize information, confirm relevance, and to extract study characteristics (See Appendix II)¹². The information we collected included general study characteristics such as authorship, year of publication, country, study design, sample size, outcomes measured, and OHCA characteristics. We compiled all data into a single spreadsheet in Microsoft Excel 2013 (Microsoft, Redmond, WA) for analysis.

5. Collate, summarize, report and interpret

Data summary and synthesis

Due to variations in study designs, terminology used, and reporting conventions, it was necessary to identify common areas for reporting summarised results. We selected a number of demographic, process and outcome measures that reflect key components of effective OHCA response and management. We abstracted and reported the findings for each measure per study where possible, and we report on them under the headings of characteristics, process and outcomes measures.

Critical appraisal

In line with the scoping review framework, we did not conduct a critical appraisal⁸.

6. Consultation (optional stage)


In addition to our own experience and expertise with OHCA in the region, we conferred with a number of OHCA subject matter experts (SMEs) in the GCC, and invited them to contribute additional materials and unpublished studies. Two of these individuals subsequently joined the research team (DA and AA).

Results

Search results and study selection

The search yielded 590 citations. We identified an additional 57 citations through searches of grey literature and hand searching. After elimination of duplicates, we screened 284 citations at the title and abstract level. This led to the exclusion of 236 citations. After full-text review of 48 citations, we included 24 full-texts for data extraction and analysis. See Fig 1 for an illustration of these findings using PRISMA Diagram, Appendix III for details of adult OHCA studies, and Appendix IV for details of paediatric OHCA studies.

Fig 1. PRISMA diagram of search results.



Characteristics of included studies

Included studies were published between 1999 and 2019. Studies comprised full-text articles (both published and unpublished manuscripts) (n=16), technical reports or theses (n=3), conference abstracts (n=3), and letters to the editor (n=2). Studies were from the UAE (n=10), Qatar (n=6), Saudi Arabia (n=4), Kuwait (n=3), Oman (n=1), and the region (n=1). We did not identify any literature from the state of Bahrain despite consultation with an SME in the country. A total of 20 studies described adult cases or all cases of OHCA. Four studies specifically or independently described paediatric OHCA and we reported these separately in our results; Conroy (1999)¹³ is reported as both an adult and a paediatric OHCA study.

Characteristics of adult OHCA studies

Batt et al. Out of hospital cardiac arrests in the Gulf Region: a scoping review. Pre-print 07/12/2019

Adult OHCA victims in the GCC region were consistently younger, predominantly male, and more co-morbid than their counterparts in Western studies. Most OHCA cases occurred at home (ranging from 54% to 85% of all OHCA cases), consistent with international findings¹⁴. See Table 1, Fig 2 and Fig 3 for further characteristics, and Appendix III for full

Table 1. Key characteristics, process and outcome measures of included OHCA studies

Study	Characteristics							Process measures			Outcome measures	
	Country	n	Median age	Mean age	% of males	% of females	OHCA at home %	bCPR rate %	baED rate %	ROSC rate %	Survival to discharge %	Survival at 30 days %
Al Hasan, 2019a ¹⁵	Kuwait	314		54.7 (GP); 61 (EMS)	70 (GP); 67 (EMS)	30 (GP); 33 (EMS)				7 (GP); 3 (EMS)		7 (GP); 1 (EMS)
Al Hasan, 2019b ¹⁶	Kuwait	332		59 (pre); 63 (post)	74 (pre); 61.7(post)	26 (pre); 38.3 (post)	75 (pre); 84.7 (post)	14 (pre); 12 (post)		6 (pre); 1.6 (post)	2 (pre); 0.8 (post)	0 (pre); 0.8 (post)
Al Hasan, 2019c*	Kuwait	286	63	61.1	67.1	32.9	76	8.7		3.5	0.3	
Alqahtani, 2019a ¹⁷	UAE	715	50		77	23	69.9	27.5	1.8	9.2		
Arabi, 2013 ¹⁸	Qatar	987		49 (ME); 62 (SA)	63 (ME); 88 (SA)	37 (ME); 12 (SA)						
Batt, 2016a ¹⁹	UAE	384	53	50.9	76	24	54.2	30	0.5	3.1		
Batt, 2017a ²⁰	UAE	514	52		75	25	62.6	28.6	1	6.6		
bin Salleeh, 2015 ²¹	KSA	96		62.9	62.5	37.5	74	20.8		4.8		
Conroy, 1999 ¹³	KSA	39		55.4	57.3	42.7		5.1			5.1	
Irfan, 2016 ²²	Qatar	447	51		80.5	19.5	63.3	20.6	2.7	13	8.1	
Irfan, 2018 ²³	Qatar	397	71 (GCC); 52 (NA)		60 (GCC); 84.8 (NA)	40 (GCC); 15.2 (NA)	88.8 (GCC); 72.3 (NA)	17.2 (GCC); 16.1 (NA)		10.2 (GCC); 9.9 (NA)	4.9 (GCC); 10.7 (NA)	
Nadar, 2018 ²⁴	Oman	216		55	69.4	30.6				39.4	13	
Ong, 2015 ²⁵	UAE	405	50	49.7	82.7	17.3	54.3	10.5	0.8	14	3	
Patel, 2014 ²⁶	Qatar	987		57	72.7	27.3					40.2	
Salam, 2013 ²⁷	Qatar	801		56								
Salam, 2018 ²⁸	Regional	89									13	
Paediatric OHCA												
Batt, 2017b ²⁹	UAE	14	4		43	57	36	26		14		
bin Salleeh, 2016 ³⁰	KSA	23		4	69.6	30.4	87	4.3				
Conroy, 1999 ¹³	KSA	27		5.3	58.8	41.2					5.8	
Tham, 2018 ³¹	UAE	17	7		76.5	23.5	52.9	11.8		5.9	5.9	

details.

Table 1. Key characteristics, process and outcome measures of abstracted OHCA studies.

OHCA=out of hospital cardiac arrest; bCPR=bystander CPR; baED=bystander AED; EMS=Emergency Medical Services; ROSC=Return of spontaneous circulation; GP=General Practitioner; UAE=United Arab Emirates; KSA=Kingdom of Saudi Arabia; ED=Emergency Department; GCC=Gulf Cooperative Council; NA=North Africa (defined in Appendix III); ME=Middle East (defined in Appendix III); SA=South Asian (defined in Appendix III)

Abstracted adult OHCA studies: Al Hasan, 2019a¹⁵; Al Hasan, 2019b¹⁶; Al Hasan, 2019c*; Alqahtani, 2019a¹⁷; Arabi, 2013¹⁸; Batt, 2016a¹⁹; Batt, 2017a²⁰; bin Salleeh, 2015²¹; Conroy, 1999¹³; Irfan, 2016²²; Irfan, 2018²³; Nadar, 2018²⁴; Ong, 2015²⁵; Patel, 2014²⁶; Salam, 2013²⁷; Salam, 2018²⁸

Abstracted paediatric OHCA studies: Batt, 2017b²⁹; bin Salleeh, 2016³⁰; Conroy, 1999¹³; Tham, 2018³¹

*note: Al Hasan 2019c is an unpublished manuscript.

Fig 2. Age profile of included adult OHCA studies.





Fig 3. Gender and location distribution of included adult OHCA studies.




We observed differences in the characteristics of OHCA among ethnic groups in the region (e.g. Arab, South Asian and North African). For example, compared to GCC patients, South Asian and North African patients were more likely to smoke, but had lower levels of comorbidities^{18,23}. Despite this, South Asian and North African patients were generally younger, and accounted for a large percentage of total OHCA cases in a number of studies^{17,19,22}, possibly attributable to population demographics.

Process and outcome characteristics

We observed low levels of EMS utilisation, low bystander CPR rates, low return of spontaneous circulation (ROSC) rates, and lower survival to hospital discharge rates across the region when compared to Western studies. EMS utilization rates were rarely reported, and ranged from 1.4%²⁴ to 31%²¹. Bystander CPR (bCPR) rates ranged from 5.1%¹³ to 30%¹⁹. Bystander AED (bAED) rates were reported at under 2% for any study which reported such data. ROSC rates ranged from 1.6%¹⁶ to 39.4%²⁴ while most studies reported ROSC rates under 10%. Survival to hospital discharge rates were often not reported, and ranged from 2%¹⁶ to 40.2%²⁴; however, most studies reported rates under 10%. See Fig 4 for information on process and outcome results.

Fig 4. Process and outcome results for included adult OHCA studies



Characteristics of paediatric OHCA cases

A small number (4) of paediatric OHCA (POHCA) studies were identified. Despite the limitations associated with such a small number of studies, we report them here nonetheless in order to provide some collective insight into POHCA in the GCC. POHCA victims in the GCC region were predominantly male, and a significant percentage were due to non-medical aetiologies (23.5%-45%), including drowning. Most POHCA cases occurred at home, ranging from 36%²⁹ to 87%³⁰ of all POHCA cases. Despite this, and similar to adult OHCA findings, we observed low levels of bystander CPR. See Table 1, Fig 5 and Fig 6 for further characteristics, and Appendix IV for full details.

Fig 5. Age profile of included paediatric OHCA studies.





Fig 6. Gender and location distribution of included paediatric OHCA studies.




Process and outcome characteristics

We observed low bCPR rates, low ROSC rates, and low survival to hospital discharge rates across POHCA studies in the region. Bystander CPR (bCPR) ranged from 0%¹³ to 26%²⁹. No bAED use was reported for any POHCA case in the region. Two studies reported ROSC,

5.9%³¹ and 14%²⁹. Two studies reported a survival to discharge rate of approximately 6%^{13,31}. Pre-existing medical conditions were reported for POHCA patients in several studies and varied from 23.5%³¹ to almost 70%³⁰. See Fig 7 for information on process and outcome results for POHCA.

Fig 7. Process and outcome results for included paediatric OHCA studies.



Discussion

The causes of lower OHCA survival in the GCC are anecdotally related to unique demographic, cultural and logistical contexts of the region. Given these challenges, and the lack of synthesized literature, we sought to describe and explore OHCA in the GCC via a scoping review methodology. Included studies reported low out-of-hospital cardiac arrest survival to discharge rates (1-13%) when compared to Western studies (e.g. 10-23% reported in Gräsner³²). After we examined studies from multiple countries in the GCC region, we suggest several unique characteristics are associated with OHCA in the region; variances exist in both OHCA processes and outcomes when compared to Western studies; and, despite the mortality associated with OHCA in the region it remains poorly researched, with an observed lack of coordinated effort to improve process and outcomes.

Unique characteristics of OHCA in the region

OHCA victims in the GCC are younger, predominantly male, and more co-morbid than their counterparts in Western studies. We suggest a number of reasons for these findings. One proposed cause for the observed increase in cardiac conditions is oil wealth-induced “obesogenic urbanization” that has occurred rapidly in the region³³. A number of studies in our review reported high rates of cardiac risk factors including diabetes mellitus,

hypertension, hyperlipidaemia, chronic renal failure, and acute coronary syndrome^{18,19,23,24,26,28,34}. Additionally, the high rate of expatriate populations living in some GCC countries^{17,19,22}, and their unique lifestyle risk factors (such as high rates of smoking) may have contributed to the observed young age and male dominance of OHCA cases in our review. We also observed differences in OHCA demographics in these populations compared to the national populations. For example, expatriates were normally younger, and more likely to smoke compared to locals^{18,23}.

Variances in pre-hospital resuscitation processes and outcomes

Cardiac arrest cases in public places have generally demonstrated better outcomes in Western studies, likely due to an increased rate of witnessed arrests, early activation of EMS and early provision of bystander CPR and AED. Existing evidence acknowledges this portion of care prior to EMS arrival is crucial in OHCA outcomes, i.e. recognising OHCA, calling for help and providing bystander CPR and AED^{35,36}. We observed lower levels of EMS utilisation (reported in 2 studies; range 1.4-35%), lower bystander CPR rates (reported in 10 studies; range 3-30%), and very low bystander AED rates in the studies included in our review. Variation in response times, and a noteworthy absence of integrated community response systems means that mortality from OHCA in the GCC likely occurs on scene. These findings highlight a significant weakness in the early access and public response to OHCA in the region. Anecdotal reasons for such challenges in the GCC include a lack of defibrillators in public places, “Good Samaritan” laws that stipulate only certified people should provide medical assistance, lack of governing bodies to oversee all aspects of medical emergency response, rurality, and cultural norms¹⁹. Cultural challenges include a lack of knowledge of CPR, a lack of confidence to intervene, and barriers to CPR and AED use, especially on female OHCA patients³⁷. We also observed over-reliance on private transport rather than EMS to transport OHCA patients, mirroring results in non-OHCA studies in the region³⁸. Attempts to address such challenges are yielding results, as evidenced in the reported increased use of EMS for OHCA^{17,19,20}; however, public perceptions related to recognising and intervening in OHCA remain unstudied in the GCC.

Elsewhere, developing integrated community response models has yielded significant improvements in OHCA measures. The Republic of Ireland is a good exemplar, and it shares characteristics of rurality in common with GCC countries^{39,40}, despite their obvious geographical differences. In 2018 national figures for Ireland (OHCA n=2,442) were reported as: 81% bCPR (excluding EMS witnessed cases), 22% bAED, 26% ROSC and 7.2% (n=176) survival to discharge, the majority with cerebral performance score of 1 or 2⁴¹. These results represent significant improvements when compared to national figures from only five years prior (n=1,885): 69% bCPR (excluding EMS witnessed cases), 16% bAED, 23% ROSC and 6% survival to discharge (n=120)⁴². In that five year period, Ireland further developed a national network of integrated community response schemes, focused on the provision of early CPR, including dispatcher-assisted CPR (DA-CPR). As a result, continued improvements in OHCA measures are likely to be realised in the future. Due to the challenges we observed in this review in relation to access to EMS, bCPR, bAED, and

cultural and logistical barriers, we observed low ROSC rates, and subsequent low levels of survival to hospital discharge across the region. In other words, lower survival as an outcome is reflective of process issues in the resuscitation attempt, particularly in the pre-arrival phase.

Lack of research and coordinated effort

The processes and outcomes related to OHCA are well-researched in Western countries, in part due to established, coordinated cardiac arrest registries. These may be national (e.g. OHCAR in Ireland⁴⁰) or regional (e.g. EuReCa in Europe³²), and they allow researchers to examine trends, correlations and concerns related to OHCA with large prospectively-collected datasets. Another example is the Pan-Asian Resuscitation Outcomes Study (PAROS²⁵), an international cardiac arrest registry study of which the UAE is a reporting member. Our review also demonstrated that OHCA research efforts are sporadic across the GCC, and even within countries, such research efforts are not strategically coordinated. For example, none of the existing UAE PAROS-member studies reported on OHCA in the Emirate of Abu Dhabi, and thus these publications do not present a complete account of OHCA in the UAE. The heterogeneity of populations within PAROS across Asia and the Middle East presents a challenge when we attempt to utilise it as a solution for the GCC³⁴. In some GCC countries, OHCA outcomes remain unknown due to a complete lack of reporting. Despite a previous call for a regional initiative for data collection and sharing⁴³, there has been no concerted effort to create OHCA reporting databases in the GCC. As a result of this lack of coordinated effort, we cannot provide scientific answer to research questions, which creates a barrier to evidence-informed policy and decision making in the GCC. Additionally, the provision of public education, coordination of public information campaigns, bystander and community response to OHCA, strategic planning of emergency medical response, and the coordination of research efforts are (where they exist) spread across multiple local or national agencies within each country. This may lead to a lack of coordinated effort in relation to the management of OHCA. Without such strategic oversight, there is potential for duplication of effort, missed opportunities, and lack of compatibility, both within and between countries in the region.

Recommendations

The low OHCA survival rates and characteristics observed in our review highlight the regional challenges when we attempt to implement existing OHCA management strategies produced by Western systems. Regional challenges include pre-existing morbidity of patients, immigrant populations, unclear “Good Samaritan” laws, low levels of EMS utilisation, lack of AED response, low bystander CPR rates, cultural challenges, and an unknown OHCA recognition rate. As a result, the traditional cardiac arrest ‘chain of survival’ faces challenges as a ‘one size-fits all’ strategy within the region due to such differences in context. Therefore countries within the GCC region may need to customise their implementation of accepted OHCA management strategies to reflect cultural, logistical and demographical variations. To inform such approaches, it is first necessary to highlight regional OHCA characteristics, identify potential OHCA management challenges, and subsequently propose areas for consideration and improvement. Based on our review, we

provide several recommendations to reduce the OHCA burden in the region (see Textbox 1), largely reflective of the recommendations of the Global Resuscitation Alliance⁴⁴. In the first instance, and essential in order to implement additional recommendations, we recommend that a lead agency should be established within each country to coordinate and manage OHCA response within the community. This agency should advocate for the passing of "Good Samaritan Law" equivalents within the region to encourage and protect community responders and bystanders, and may provide leadership on the integration of trained responder dispatch systems^{45,46}. This agency would continue to develop EMS response, in particular through the creation of integrated community response models, and the provision of DA-CPR to improve OHCA process and outcome measures. Community response models empower the local community, and may work to alleviate cultural barriers and the increased response times associated with rurality. Additionally, advocating for uniform emergency numbers, coordinated dispatch centres, and compatible EMS system designs are essential components of early recognition and response in OHCA^{47,48}.

Public health initiatives should be developed and implemented (or continued where already present) to reduce risk factors for cardiovascular disease, respiratory conditions, and preventable trauma in paediatrics, including drowning. We also recommend targeted public health interventions aimed at improving public health concerns highlighted in expatriate populations. Additionally, in light of the proportion of OHCA that occurred at home in the region (similar to Western studies), the lead agency should implement widespread information campaigns through various media, that inform the public on the existence of "Good Samaritan Law" equivalents, the importance of early OHCA recognition, how to access EMS, and the importance of early CPR (which can be further supported via provision of DA-CPR). Furthermore, the lead agency should promote improved access to AEDs in public places that have high incidence of OHCA⁴⁹, as well as develop and maintain a national registry of publicly accessible AEDs⁵⁰. This focus on improved care **prior** to EMS arrival is crucial in improving OHCA outcomes in GCC countries³⁵. Finally, in order to improve our understanding of OHCA in the region, we advocate for the creation of a reporting database within each GCC country, and a further cooperative database within the region. The lead agency for each country should oversee the development of such a registry, which will enable continued research into the characteristics and management of OHCA in the region. Such databases should align with international data collection and reporting conventions to facilitate data sharing and collaboration⁵¹.

Limitations

Our findings need to be interpreted in the context of certain limitations. We may not have identified all relevant studies despite our attempts to be comprehensive. Our search and review was restricted to articles published in English and Arabic, but this does not inherently bias a review⁵². The Google Scholar searches were limited to the first 300 results which is considered adequate for grey literature searches⁵³. Further grey literature sources may exist that we did not identify; however, we attempted to minimize this risk by using the Grey-Matters Checklist¹¹, contacting SMEs in the region, and searching non-indexed sources. A

Batt et al. Out of hospital cardiac arrests in the Gulf Region: a scoping review. Pre-print 07/12/2019

number of studies included data on presumed cardiac aetiology as well as traumatic OHCA. We attempted to identify and report only non-traumatic OHCA data where possible, but this was not always possible. We noted in our data where we were able to perform this. Finally, there are a limited number of POHCA studies in our review. Despite these limitations, we propose that our review offers the first comprehensive overview of OHCA in the GCC region.

Textbox 1. Summary of recommendations.

1. Create a lead agency in each country for strategic oversight of OHCA preparedness, response, management and research
2. Develop community response models, including dispatcher-assisted CPR, and community responder dispatch models
3. Advocate for the legislation of ‘Good Samaritan Law’ equivalents
4. Implement (or continue) public health initiatives to reduce risk factors, including cardiac related health checks for immigrant populations
5. Coordinate widespread public information campaigns related to recognition and management of OHCA, early EMS access, and early CPR
6. Improve access to AEDs in high-risk locations, and create national AED registries
7. Establish and maintain OHCA registries in each country and the GCC region which align with international conventions

Conclusion

Our review is the first to collate and comprehensively report data on OHCA in the GCC region. We highlighted unique demographic traits, and challenges related to OHCA processes and outcomes in the region. We also acknowledged where countries have implemented strategies to address OHCA, and process and outcome measures are improving. Finally, despite the significant burden associated with OHCA in the region, we observed a concerning lack of research and a lack of coordinated effort. In order to address these concerns, we recommend the creation of lead agencies responsible for coordinating and developing OHCA strategies. Such strategies may include implementing community response models including dispatcher-assisted CPR, advocating for “Good Samaritan Law” equivalents, coordinating mass public education campaigns targeted at cultural and logistical issues, supporting public health risk-reduction initiatives, and establishing OHCA reporting databases.

Acknowledgments

This research was supported by a research grant from the ZOLL Foundation, who had no role in the design or conduct of the study; the collection or analysis of the data; or, the drafting or submission of the manuscript.

References

1. Berdowski J, Berg RA, Tijssen JGP, Koster RW. Global incidences of out-of-hospital cardiac arrest and survival rates: Systematic review of 67 prospective studies. *Resuscitation*. 2010;
2. Kirk D, Napier D. The Transformation of Higher Education in the UAE: Issues, Implications and Intercultural Dimensions. In: Zajda J, Daun H, Saha L, editors. *Nation-Building , Identity and Citizenship Education: Cross-cultural Perspectives* [Internet]. 2008. p. 131–42. Available from: <http://books.google.com/books?id=WXLtU14ggwQC>
3. Naithani P, Jha AN. Challenges Faced by Expatriate Workers in Gulf Cooperation Council Countries. *Int J Bus Manag*. 2009;5(1).
4. Khoja T, Rawaf S, Qidwai W, Rawaf D, Nanji K, Hamad A. Health Care in Gulf Cooperation Council Countries: A Review of Challenges and Opportunities. *Cureus*. 2017;9(8):1–11.
5. World Bank. Data for Saudi Arabia, United Arab Emirates, Kuwait, Bahrain, Oman, Qatar [Internet]. World Bank Data. 2019 [cited 2019 Dec 2]. Available from: <https://data.worldbank.org/?locations=SA-AE-KW-BH-OM-QA>
6. Razzak H, Harbi A, Shelpai W, Qawas A. Prevalence and risk factors of cardiovascular disease in the United Arab Emirates. *Hamdan Med J* [Internet]. 2018;11(3):105. Available from: <http://www.hamdanjournal.org/text.asp?2018/11/3/105/236269>
7. Sasser S, Gibbs M, Blackwell T. Prehospital emergency care in Abu Dhabi, United Arab Emirates. *Prehospital Emerg Care* [Internet]. 2004 Jan 28;8(1):51–7. Available from: <http://www.tandfonline.com/doi/full/10.1080/312703002818>
8. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* [Internet]. 2005 Feb;8(1):19–32. Available from: <https://doi.org/10.1080/1364557032000119616>
9. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. 2018;1–7.
10. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* [Internet]. 2018; Available from: <http://annals.org/article.aspx?doi=10.7326/M18-0850>
11. Canadian Agency for Drugs and Technologies in Health. *Grey Matters: a practical tool for searching health-related grey literature* [Internet]. Ottawa; 2019. Available from: <https://www.cadth.ca/resources/finding-evidence/grey-matters>
12. Ritchie J, Spencer L. Qualitative data analysis for applied policy research. In: Huberman A, Miles M, editors. *The Qualitative Researcher's Companion*. Thousand Oaks, CA: SAGE Publications Inc.; 2002. p. 305–29.
13. Conroy KM, Jolin SW. Cardiac arrest in Saudi Arabia: A 7-year experience in Riyadh. *J Emerg Med*. 1999;17(4):617–23.
14. Masterson S, Teljeur C, Cullinan J, Murphy AW, Deasy C, Vellinga A. Out-of-hospital cardiac arrest in the home: Can area characteristics identify at-risk communities in the Republic of Ireland? *Int J Health Geogr* [Internet]. 2018;17(1):1–11. Available from: <https://doi.org/10.1186/s12942-018-0126-z>
15. Al Hasan D, Brightwell R. General Practitioner Pre-Hospital Resuscitation Contribution to out of Hospital Cardiac Arrest Survival; A Retrospective Study *Emergency Medicine*: Open Access. *Emerg Med Open Access*. 2019;9(3):397.

16. Al Hasan D, Drennan J, Monger E, Mahmud S Al, Ahmad H, Ameen M, et al. Dispatcher assisted cardiopulmonary resuscitation implementation in Kuwait. *Medicine (Baltimore)*. 2019;98(44):e17752.
17. Alqahtani S, Alhajeri A, Ahmed A, Mashal S. Characteristics of out of hospital cardiac arrest in the United Arab Emirates. *Heart Views [Internet]*. 2019;20(4):146. Available from: <http://www.heartviews.org/text.asp?2019/20/4/146/271031>
18. Arabi A, Patel A, Alsuwaidi J, Singh R, Albinali H. Ethnic Differences in Patients With Out-of-Hospital Cardiac Arrest: Insight From a 20-Year Registry in the State of Qatar. *J Am Coll Cardiol [Internet]*. 2013;61(10):E15. Available from: [http://dx.doi.org/10.1016/S0735-1097\(13\)60016-9](http://dx.doi.org/10.1016/S0735-1097(13)60016-9)
19. Batt AM, Al-Hajeri AS, Cummins FH. A profile of out-of-hospital cardiac arrests in northern Emirates, United Arab Emirates. *Saudi Med J*. 2016;37(11).
20. Batt AM, Al-Hajeri AS, Cummins FH. Urgent Need to Strengthen the Chain of Survival in the United Arab Emirates; a Letter to the Editor. *Emergency*. 2017;5(1):5968600.
21. Bin Salleh HM, Gabralla KA, Leggio WJ, Al Aseri ZA. Out-of-hospital adult cardiac arrests in a university hospital in central Saudi Arabia. *Saudi Med J*. 2015;36(9):1071–5.
22. Irfan FB, Bhutta ZA, Castren M, Straney L, Djarv T, Tariq T, et al. Epidemiology and outcomes of out-of-hospital cardiac arrest in Qatar: A nationwide observational study. *Int J Cardiol [Internet]*. 2016;223:1007–13. Available from: <http://dx.doi.org/10.1016/j.ijcard.2016.08.299>
23. Irfan FB, Castren M, Bhutta ZA, George P, Qureshi I, Thomas SH, et al. Ethnic differences in out-of-hospital cardiac arrest among Middle Eastern Arabs and North African populations living in Qatar. *Ethn Heal*. 2018;(December).
24. Nadar SK, Mujtaba M, Al-Hadi H, Sadiq M, Al-Riyami A, Ali M, et al. Epidemiology, outcomes and coronary angiography findings of patients following out-of-hospital cardiac arrest: A single-centre experience from Oman. *Sultan Qaboos Univ Med J*. 2018;18(2):e155–60.
25. Ong MEH, Shin S Do, De Souza NNA, Tanaka H, Nishiuchi T, Song KJ, et al. Outcomes for out-of-hospital cardiac arrests across 7 countries in Asia: The Pan Asian Resuscitation Outcomes Study (PAROS). *Resuscitation*. 2015;96:100–8.
26. Patel AA, Arabi AR, Alzaeem H, Al Suwaidi J, Singh R, Al Binali HA. Clinical profile, management, and outcome in patients with out of hospital cardiac arrest: Insights from a 20-year registry. *Int J Gen Med*. 2014;7:373–81.
27. Salam AM, AlBinali HA, Singh R, Al Suwaidi J. Impact of the 2005 resuscitation guidelines on patient survival after out-of-hospital cardiac arrest: Experience from a 20-year registry in a middle-eastern country. *Resuscitation [Internet]*. 2013;84(8):e97–8. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2013.04.008>
28. Salam AM, AlHabib K, Almahmeed W, Alsheikh-Ali A, Sulaiman K, Amin H, et al. Abstract 11181: Presentation With Cardiac Arrest in Patients With Acute Coronary Syndromes: Insights From the Second Gulf Registry of Acute Coronary Events. *Circulation [Internet]*. 2018 Nov 25;130(suppl_2):A11181–A11181. Available from: https://doi.org/10.1161/circ.130.suppl_2.11181
29. Batt AM, Al-Hajeri AS, Ward G, Pilapil CS, Delport S, Cummins FH. Paediatric cardiac arrests in the northern emirates, United Arab Emirates. *Mediterr J Emerg Med*. 2017;(25):30–4.
30. Bin Salleh HM, Al Tom M, Ahmed Y, Leggio WJ, Abdulqader NF. Out of Hospital Pediatric Cardiac Arrest: Prospective Study from Riyadh, Saudi Arabia. *Biosci Biotechnol Res Asia*. 2016;13(1):569–72.

31. Tham LP, Wah W, Phillips R, Shahidah N, Ng YY, Shin S Do, et al. Epidemiology and outcome of paediatric out-of-hospital cardiac arrests: A paediatric sub-study of the Pan-Asian resuscitation outcomes study (PAROS). *Resuscitation* [Internet]. 2018;125:111–7. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2018.01.040>
32. Gräsner JT, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, et al. EuReCa ONE—27 Nations, ONE Europe, ONE Registry: A prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation*. 2016;105:188–95.
33. AlHabib KF, Hersi A, AlFaleh H, AlNemer K, AlSaif S, Taraben A, et al. Baseline characteristics, management practices, and in-hospital outcomes of patients with acute coronary syndromes: Results of the Saudi project for assessment of coronary events (SPACE) registry. *J Saudi Hear Assoc* [Internet]. 2011;23(4):233–9. Available from: <http://dx.doi.org/10.1016/j.jsha.2011.05.004>
34. Batt AM, Al-Hajeri AS, Cummins FH, Bin Salleeh H. Out-of-hospital adult cardiac arrests in a university hospital in Central Saudi Arabia. *Saudi Med J*. 2015;37(6):714.
35. Deakin CD. The chain of survival: Not all links are equal. *Resuscitation*. 2018;126(January):80–2.
36. Glass GF, Brady WJ. Bystander Intervention in Out-of-Hospital Cardiac Arrest. *JAMA Netw Open* [Internet]. 2019 Mar 29;2(3):e191008. Available from: <http://jamanetworkopen.jamanetwork.com/article.aspx?doi=10.1001/jamanetworkopen.2019.1008>
37. Orla M. Abstract 229: Are There Any Cultural Barriers to Automated External Defibrillator Use in Middle Eastern Countries? *Circulation* [Internet]. 2011 Nov 22;124(suppl_21):A229–A229. Available from: https://doi.org/10.1161/circ.124.suppl_21.A229
38. Fares S, Zubaid M, Al-Mahmeed W, Ciottone G, Sayah A, Al Suwaidi J, et al. Utilization of emergency medical services by patients with acute coronary syndromes in the Arab Gulf States. *J Emerg Med* [Internet]. 2011;41(3):310–6. Available from: <http://dx.doi.org/10.1016/j.jemermed.2010.05.002>
39. Masterson S, Teljeur C, Cullinan J, Murphy AW, Deasy C, Vellinga A. The Effect of Rurality on Out-of-Hospital Cardiac Arrest Resuscitation Incidence: An Exploratory Study of a National Registry Utilizing a Categorical Approach. *J Rural Heal*. 2019;35(1):78–86.
40. Masterson S, Wright P, O'Donnell C, Vellinga A, Murphy AW, Hennelly D, et al. Urban and rural differences in out-of-hospital cardiac arrest in Ireland. *Resuscitation* [Internet]. 2015;91:42–7. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2015.03.012>
41. OHCAR Steering Group. Out-of-hospital Cardiac Arrest (OHCAR) Annual Report 2018 [Internet]. Galway, Ireland; 2018. Available from: <https://www.nuigalway.ie/media/collegeofmedicinenuresinghealthsciences/disciplines/generalpractice/files/OHCAR-Annual-Report-2018.pdf>
42. OHCAR Steering Group. Out-of-hospital Cardiac Arrest (OHCAR) Annual Report 2014 [Internet]. Health Policy. 2014. Available from: <https://www.phecit.ie/Images/PHECC/Publications and Media/Other Publications/5.1 OHCAR 6th Annual Report.pdf>
43. Batt AM, Al-Hajeri AS, Cummins FH. Re: A profile of out-of-hospital cardiac arrests in northern emirates, United Arab Emirates. *Saudi Med J*. 2017;38(6).
44. Eisenberg M, Lippert FK, Castren M, Moore F, Ong M, Rea T. Improving Survival from Out-of-Hospital Cardiac Arrest: Acting on the Call [Internet]. 2018. Available from:

- https://www.cercp.org/images/stories/recursos/articulos_docs_interes/doc_GRA_Acting_on_the_call_1.2018.pdf
45. Smith CM, Wilson MH, Hartley-Sharpe C, Gwinnutt C, Dicker B, Perkins GD. The use of trained volunteers in the response to out-of-hospital cardiac arrest – the GoodSAM experience. *Resuscitation* [Internet]. 2017;121:123–6. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2017.10.020>
 46. Brooks SC, Worthington H, Gonedalles T, Bobrow B, Morrison LJ. Implementation of the PulsePoint smartphone application for crowd-sourcing bystander resuscitation. *Crit Care* [Internet]. 2014;18(1):P484. Available from: <https://doi.org/10.1186/cc13674>
 47. Morais DA, Carvalho DV, Correa A dos R. Out-of-hospital cardiac arrest: Determinant factors for immediate survival after cardiopulmonary resuscitation. *Rev Lat Am Enfermagem*. 2014;
 48. Ng YY, Leong SHB, Ong MEH. The role of dispatch in resuscitation. *Singapore Med J*. 2017;
 49. Moran PS, Teljeur C, Masterson S, O’Neill M, Harrington P, Ryan M. Cost-effectiveness of a national public access defibrillation programme. *Resuscitation* [Internet]. 2015;91:48–55. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2015.03.017>
 50. Lijovic M, Bernard S, Nehme Z, Walker T, Smith K. Public access defibrillation-results from the victorian ambulance cardiac arrest registry. *Resuscitation* [Internet]. 2014;85(12):1739–44. Available from: <http://dx.doi.org/10.1016/j.resuscitation.2014.10.005>
 51. Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: The utstein style: A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, and Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation*. 1991;
 52. Morrison A, Polisena J, Husereau D, Moulton K, Clark M, Fiander M, et al. The effect of English-language restriction on systematic review-based meta-analyses: a systematic review of empirical studies. *Int J Technol Assess Health Care*. 2012 Apr;28(2):138–44.
 53. Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of google scholar in evidence reviews and its applicability to grey literature searching. *PLoS One*. 2015;10(9):1–17.