

Manuscript

1 **Title** [Knowledge and attitudes regarding infant BLS among babysitters and nursemaids in
2 north rural Jordan]

3 **Author(s) and Affiliation(s)**

4 **Liqaa Raffee¹, Shereen Hamadneh², Manal Kassab³, Fidaa Almomani⁴, Jehan Hamadneh⁵,**
5 **Mohammed AlBashtawy⁶, Rawan A. Obeidat⁵, Othman Almazloom¹**

6
7 1. Department of Accident and Emergency Medicine, Faculty of Medicine, Jordan
8 University of Science and Technology, Irbid, Jordan

9
10 2. Department of Maternal and Child Health, Faculty of Nursing, Al al-Bayt University,
11 Mafraq, Jordan

12
13 3. Department of Maternal and Child Health, Faculty of Nursing, Jordan University of
14 Science and Technology, Irbid, Jordan; University of Technology, Sydney, Australia

15 4. Department of Rehabilitation Sciences, Faculty of Applied Medical Sciences, Jordan
16 University of Science and technology, Irbid, Jordan

17 5. Department of Obstetrics and Gynaecology, Faculty of Medicine, Jordan
18 University of Science and technology, Irbid, Jordan

19
20 6. Department of Community Health Nursing, Faculty of Nursing, Al al-Bayt University,
21 Mafraq, Jordan

22
23
24 **Corresponding Author:**

25 * **Liqaa Raffee**

26
27 **Disclose of interest:** There is no conflict of interest to disclose

28
29 **Ethics:** The study was approved by the institutional research board and ethics committees at
30 Al al-Bayt University and Jordan University of Science & Technology.

31
32 **Acknowledgment:** Acknowledge for AABU and JUST universities for supporting this project.
33 Thanks to caregivers who accepted to take part to participate in the study survey.

34
35 **Contributions:**

36 **Shereen Hamadneh; Liqaa Raffee**

37 *Study conception and design; Acquisition of data; Analysis and interpretation of data;*
38 *Drafting of manuscript; Critical revision; supervised whole study*

39 **Manal Kassab; Jehan Hamadneh; Mohammed AlBashtawy; Rawan A. Obeidat, Fidaa**
40 **Almomani**

Manuscript

41 *Acquisition of data; Analysis and interpretation of data; Drafting of manuscript; Critical*
42 *revision*
43 **Othman Almazloom**
44 *Acquisition of data; Analysis and interpretation of data;*
45
46
47

Manuscript

48 **Title**

49 **[Knowledge and attitudes regarding infant BLS among babysitters and nursemaids in**
50 **north rural Jordan]**

51

52 **Abstract**

53 **Background:** There is a lack of national data related to the public knowledge and attitudes
54 concerning Basic Life Support (BLS) for an infant. This study is the first to assess the
55 knowledge and attitude regarding infant BLS among babysitters and nursemaids who are
56 working in registered nurseries in North Jordan.

57 **Methods:** A semi-structured questionnaire was conducted among a convenience sample of 77
58 child caretakers (babysitters and nursemaids) who were looking after infants in registered
59 nurseries in north of Jordan.

60 **Results:** The findings show that the majority of participants were not aware of skills for
61 managing foreign body aspiration (or CPR) among infants, while more than two-thirds of
62 them never attended training and or education sessions regarding managing foreign body
63 aspiration among infants.

64 **Conclusions:** This research strongly recommends training for babysitters and care center
65 staff. The findings of this study indicated that caregivers in Jordanian kindergartens and
66 nurseries lack awareness regarding safe infant care. As such, there is a lack of basic skills for
67 resuscitation in urgent cases; which can increase the risk of sudden and unexpected child
68 death. Accordingly, the infant life support course is highly recommended for babysitters and
69 nursemaids. Health care professionals could help to implement infant basic life support
70 training courses through local community basic service centers, non-governmental
71 organizations, local associations, universities and health colleges through providing

Manuscript

72 workshops that aim at enhancing knowledge and skills for infant BLS among caregivers and
73 babysitters.

74

75 **Keywords**

76 *Nursing; Emergency; CPR; Aspiration; infant death; safety; nursery; kindergarten*

77

78 **Introduction**

79 Approximately 11% of children die each year as a result of a variety of causes including road
80 traffic accidents (RTAs), falls, blast injuries, burns, insect bites, drowning, suffocations, and
81 aspiration.(1) Pediatric cardiac arrest is a life-threatening event, which often is required a fast
82 effective intervention.(2,3) There are more than 5,000 cases of pediatric cardiac-arrests out-
83 hospital annually.(2,3) In particular, one-fifth of total child deaths were due to mechanical
84 suffocation where the child was younger than four years.(3, 4) Dealing with infants' urgent
85 life-threatening situations and providing life support measures may improve infant survival
86 rate. Moreover, the quality of the CPR administered is of paramount importance because it
87 directly affects the infants' hemodynamics, their survival, and the neurological outcomes after
88 the arrest.(3)

89

90 The aspiration of the foreign body is a potentially life-threatening event especially in those
91 under the age of 5 years.(5) An aspirated child can be presented to the emergency as a
92 spectrum, ranging from an event that has no symptoms to more severe events including
93 suffocation, cyanosis, respiratory distress, or even death.(5-7) However, in some cases,
94 previous foreign objects that were topical in the lung can travel back to the mouth and are
95 swallowed, after some severe cough episodes.(7)

96

Manuscript

97 The American Academy of Pediatrics has recommended that children's caregivers should
98 have life support skills and be trained to deal with life-threatening situations.(8) Survival after
99 cardiopulmonary arrest is depending on the effectiveness of the intervention which includes
100 the early management, CPR quality and the time to defibrillation.(9) Babysitters and the local
101 community in Jordan lack awareness regarding safe infant care and basic skills for urgent
102 cases which can increase the risk of sudden and unexpected child death.(10)

103

104 Background

105 Choking and foreign body aspiration remain among the major causes of mortality and
106 morbidity in young children, however, it could be preventable.(4, 5, 6, 11-14) Anyone can be
107 a lifesaver for a cardiac arrest victim with the proper awareness and training.(10, 14, 15)

108 Primary and secondary infant caregivers including parents, babysitters, and nursemaids
109 should have the basic skills for life support to be able to use it when the baby's life is at stake.
110 (10-14) Young children aging between 0 and 3 years, are at the highest risk for accidental
111 suffocation, due to several factors; young children tend to put small things in their mouths and
112 often crying, screaming, running, and playing while such objects are still in their mouths; all
113 of which could be accidentally ingested and cause suffocation. (4) In addition, they do not
114 have molar teeth to chew some foods completely. (4) The most common reported cases of
115 foreign body aspiration are organic or food related, especially when the child begins to eat
116 solid food. (4)

117

118 CPR accurate application depends on experiences, skills, training, and confidence of the
119 rescuer.(12) The American Heart Association released revised CPR recommendations in 2010
120 which state that the compression to breathing ratio is 30:2 for children with one rescuer, and
121 15:2 for children with two rescuers.(12)

Manuscript

122

123 Public awareness campaigns and the strict guidelines for toy manufacturers to be within
124 safety requirements in the developed countries have contributed to decreasing the number of
125 childhood deaths due to foreign body aspiration over the past decade.(11) However, in
126 developing countries like Jordan, the public knowledge and skills related to infant basic life
127 support and safety care are still not well established.(1, 10, 13)

128

129 **Study significance**

130 In Jordan, there is a lack of data related to the public knowledge and attitudes towards Basic
131 Life Support (BLS) for an infant. Therefore, assessing the knowledge and attitudes regarding
132 infant BLS among babysitters and nursemaids who are working in registered kindergarten and
133 nurseries in Jordan is crucial to providing and improving care.This study was the first of its
134 kind in the region that investigated babysitters' and nursemaids' knowledge about Infant BLS
135 and provided basic information which can be used to construct more appropriate and effective
136 intervention programs.

137

138 **Aims**

139 This study was designed to assess the level of awareness and the attitudes caregivers held
140 towards the practice of BLS for an infant, among registered nurseries in north rural Jordan.

141 **METHODS**

142 A cross-sectional study conducted during February 2017, child caretakers (babysitters and
143 nursemaids) who were looking after infants in registered nurseries in north state of Jordan. A
144 semi-structured questionnaire was distributed among babysitters and nursemaids who agreed
145 to participate in the study. The questionnaire explored their knowledge, attitudes and
146 perceptions regarding Infant BLS.

Manuscript

147

148 **Instruments**

149 A semi-structured questionnaire was used, which was adapted from a study conducted by
150 Chia and Lian (2014) in Singapore to assess parental knowledge, attitudes, and perceptions
151 towards IBLs.(15) The questionnaire is to collect information about the demographic profile
152 of the participants; knowledge for participants on IBLs skill, and questions of perceptions and
153 attitudes about infant BLS. The questionnaire contains multiple options, True or False
154 questions, and open ended questions. (15)

155

156 **Ethical Considerations**

157 This study was approved by the concerned institutional review board. All participants signed
158 a consent form after had information about the study and they have been notified that their
159 participation was voluntary and they could withdraw at any time. They were guaranteed
160 anonymity and confidentiality

161

162 **Data Analysis**

163 Data were analyzed using SPSS 19 (SPSS, Inc., Chicago IL, USA). Descriptive statistical
164 analysis was used to identify frequencies and percentages of studied categories, and
165 correlation between demographical profile and knowledge regarding BLS.

166

167 **RESULTS**

168 A total of 77 babysitters and nursemaids from an eighteen registered nurseries were recruited
169 to participate in the study. The demographic characteristics are presented in Table-1. As seen
170 in the table, approximately half of the participants have a diploma degree (52%), while
171 another third (34%) has a bachelor's degree, with only 14% having a secondary or less degree

Manuscript

172 of education. In terms of age groups, around half of the participants (47%) were between 26
173 to 35 years old, while a quarter (26%) were between 36 to 45 years old, followed by (21%) of
174 25 years or less, and only (6%) were more than 45 years old. Lastly, concerning nationality,
175 the majority (70%) were of Jordanian nationality, while the rest (30%) were refugees or
176 migrants.

177

178 **Babysitters' and Nursemaids' Knowledge of IBLS**

179 The study investigated babysitters' and nursemaids' knowledge of the guidelines for IBLS,
180 according to the American Academy of Pediatrics and the American Heart Association;
181 including foreign body aspiration and CPR. Table-2 presents the responses related to
182 participants' knowledge regarding IBLS and foreign body aspiration. Based on babysitters'
183 and nursemaids' responses, more than a third of them (38%, n= 29) never heard about CPR
184 for children. Furthermore, the vast majority of participants (70%, n= 54) did not attend the
185 training/education session regarding managing foreign body aspiration. However, 84 percent
186 (n= 65) indicated that they would welcome participating in IBLS if it is offered to them.

187

188 **The Proportion of Participants Who Obtained Passing Marks on General Information 189 for BLS**

190 As shown in Table-3, more than one-quarter (27%) of the participants were unaware of the
191 Jordanian ambulance access number. Furthermore, only (66%) of nursemaids and babysitters
192 answered correctly in relation to the child's brain survival without oxygenation (breathing).
193 When asked about the location to check a pulse in a child aged less than one year old, only
194 22% could identify the correct answer.

195

Manuscript

196 Additionally, when asked the nursemaids and babysitters about the methods they may use to
197 wake an unresponsive infant, only 44% answered correctly and indicated that they need to tap
198 or flick the infant's feet soles gently. Finally, participants were asked about how they can
199 confirm if the baby is not breathing, about 48% of the participants answered the first item
200 correctly, while correct answers composed 29% for the second item, and 12% for the third
201 item (see Table-3).

202

203 With regards to their knowledge of the correct ratio of chest compression to breath rescue
204 during BLS, only 13% of the participants were aware of the correct ratio compression to
205 breath to provide an effective CPR for an infant with only one rescuer, and 39% indicated the
206 correct answer for providing effective CPR for an infant when two rescuers are available (see
207 Table-3).

208

209 As shown in Table-3, the questionnaire further investigated participants' knowledge about
210 proper interventions to be done in case of infant foreign body aspiration. Four alternative
211 answers were provided with the right answer being “Provide infant of five back blows and
212 five chest thrusts until the object is expelled,” only 21% participants were able to provide
213 correct answer. Another 43% indicated that they should remove the piece aspirated if only
214 were able to see it.

215 **The Main Source for the Information about IBLS**

216 The majority of study participants, (52%, n=40), indicated that social media was their main
217 source for information regarding BLS. While less than a quarter (21%, n=16) indicated that
218 they learned about IBLS from conferences and training workshops, and only (16%, n=12)
219 learned about it from friends and colleagues. Of the participants, few (12%, n= 9) reported
220 that they access professional journals.

Manuscript

221

222 **Perception for Confidence Level among Babysitters and Nursemaids Regarding IBLS** 223 **Skills**

224 The study sample varied in their responses regarding their confidence in providing IBLS and
225 dealing with foreign body aspiration situations. A high proportion of babysitters and
226 nursemaids reported being not confident (42%) or not confident at all (22%). While only 30%
227 of babysitters and nursemaids reported being somewhat confident, and just 6% were very well
228 confident of being able to provide IBLS.

229

230 Contrarily, when they were asked about their ability to manage foreign body aspiration in
231 infants, 35% of participants reported being not confident, and another 13% reported being not
232 confident at all. Those who were somewhat confident composed 42% of the babysitters and
233 nursemaids, while only 10% were very confident that they could provide proper intervention
234 if such emergent event happened (see Table-4).

235

236 Furthermore the level of awareness among the babysitters/nursemaid regarding infant BLS
237 was significantly ($p<0.01$) associated with level of education and age. Babysitters/nursemaid
238 with no university/institution degrees were have less knowledge than babysitters/nursemaid
239 holding college/university degrees. As well babysitters/nursemaid of ageing more than age
240 of 35 have better knowledge, while there is no relation with residency status ($p>0.01$).
241 (Table-5)

242 **Challenges for Attending IBLS Courses**

243 The study participants were asked to indicate their most significant challenge for attending
244 such BLS training courses. The study participants indicated that the major challenge they face
245 for attending IBLS courses was a time limitation due to overload work and busy schedule at

Manuscript

246 nurseries. They also indicated that the long day shift prevent them from engaging in such
247 course, some participants indicated that they only have a one day off work per week and so it
248 difficult to attend these training events. In addition, participants indicated financial
249 difficulties, and transportation difficulties were a challenging factor for attending the IBL
250 course.

251 **Suggestion for getting the training**

252 The participants were also asked to choose their more significant way they prefer to receive
253 such training or information about BLS. Less than quarter (21%) preferred the online training
254 courses if it offered. Another 33% chosen face-to-face training, while 53% indicated that they
255 prefer both ways to get the knowledge about BLS, another 3% were unsure about the method
256 to receive such training.

257

258 **DISCUSSION**

259 This study was conducted among 77 babysitters and nursemaids with the majority having a
260 bachelor's or diploma degree. The majority of babysitters and nursemaids were not aware of
261 skills for managing foreign body aspiration among infants, and more than two-thirds of them
262 never attended training or education sessions regarding managing foreign body aspiration and
263 CPR practicing among children. However, the majority indicated that they are motivated to
264 participate in such an event if it is offered to them. Significant differences were found in a
265 previous randomized controlled trial study conducted in a tertiary pediatrics' hospital in UK,
266 which aimed to compare the infant CPR technical skill performance among several groups—
267 one which participated in a self-motivated CPR refresher course using a competitive
268 leaderboard, another with a CPR training feedback device, and a final group who did not
269 participate in the course. The results showed that those who chose to participate in the

Manuscript

270 refresher course by self-motivation scored significantly higher than the other groups. The
271 study concluded that self-motivated participation in infant CPR training would help to
272 improve knowledge and skills to manage infants in a critical situation.(16) Furthermore, the
273 level of awareness among the babysitters/nursemaid regarding infant BLS was associated
274 with the level of education and age, while there is no relation with residency status.
275 Babysitters/nursemaid with college or university education level could have a chance of
276 getting information about life support skills through meetings or studying. Also,
277 babysitters/nursemaid with higher ages may have more life experiences.

278

279 One of the alarming findings of the current Jordanian study is that almost two-thirds of the
280 proportion of babysitters and nursemaids were not confident in performing CPR for the
281 infants in case of emergency. Furthermore, they lacked knowledge about the guidelines that
282 they should follow in managing foreign body aspiration among infants. Babysitters and
283 nursemaids were also unaware of how to confirm that a child is not breathing when is
284 unconscious. There is insufficient training or qualification for kindergarten staff, especially in
285 rural areas, where factors such as distance from the city centre may play a role in reducing the
286 opportunities for continuous training of nursery staff. The economic factor could be a
287 challenge, transport costs and training fees may influence the opportunity to get babysitters
288 involved in such training courses.

289

290 A descriptive study was conducted by Chia and Lain (2014), utilizing a survey questionnaire
291 for knowledge, attitudes, and perceptions of parents in Singapore about IBLS.(15) The results
292 of the study showed that IBLS training, as part of basic cardiac life support training, is
293 important given that CPR can significantly alter the outcome in children with
294 cardiopulmonary arrest. The study revealed that knowledge gaps can be filled through official

Manuscript

295 training. It recommended a refreshing course regularly update ' knowledge and skills among
296 child caregivers.(15) Developing such training intervention for babysitters and nursemaids is
297 highly needed in Jordan. Education campaigns have succeeded in part, and primary health
298 care providers can play a crucial role in increasing education efforts about the risk of
299 suffocation and management measures.(17)

300

301 Collaboration between the ministry of health, the ministry of education, the ministry of social
302 development and civil defense in creating programs and policies for enhancing awareness of
303 basic life support skills would be beneficial in accomplishing the mission. More than half of
304 babysitters and nursemaids in this study indicated they use social media to gain information
305 regarding basic life support skills. Using the Internet and adapting online training courses to
306 address this need would be beneficial for targeting babysitters and nursemaids providing a
307 suitable time for them to access the information, as well as being cost-effective by eliminating
308 transportation needs. Value is added when thinking about further benefits of online training
309 courses, such as increasing candidate independence, lowering teacher burden, and
310 maintaining better standardization of course materials; additionally, it would be effective in
311 terms cost.(18) However, first aid online courses may be useful for acquiring knowledge, but
312 maybe not effective for performing BLS skills.(19) Nonetheless, pre-training evaluation of
313 care providers knowledge and skills may be particularly beneficial in improvement BLS skills
314 acquisition immediately after the course of the training.(20) Officials, governmental decision-
315 makers, and stakeholders should support such initiative programs

316

317 The present study had a limitation with regard to the small sample size which affects general
318 claims about all of the babysitters and nursemaids in Jordan. However, conducting this
319 research was highly needed in Jordan, as far it was the first of its kind in the region that

Manuscript

320 investigated caregivers' (babysitters and nursemaids) knowledge about infant BLS, and
321 provided basic information that used in constructing such an intervention program.

322

323 **CONCLUSION AND RECOMMENDATIONS**

324 A majority of the study participants had not attended any CPR or BLS courses. However, it
325 was encouraging that most of them expressed an interest in attending such a training session
326 in the future. It is highly needed that administrators run continuous education and training
327 programs on IBLS for babysitters and nursemaids and to provide them with updated
328 information and keep them aware of best practices for dealing with an urgent situation, such
329 as choking that may often occur among young children. Policy changes are needed so that
330 each kindergarten will include a child nurse at each setting; so nurses can lead the
331 development of regular in-service training to improve caregivers' skills through their
332 participation in IBLS courses. This regular and continuous scheduling will minimize any
333 challenges that prevent caregivers from attending IBLS course including time limitation,
334 transportation issues, and financial difficulties. Further, providing such Internet-based
335 resources would help eliminate barriers like cost and transportation. Officials, governmental
336 decision-makers, and stakeholders should support such initiative programs.

337 A special effort could be made to engage the newly graduated students by including them in
338 such repairing courses before they begin working in the nurseries and kindergartens. At the
339 community level, local community basic service centers, non-governmental organizations,
340 local associations, universities and health colleges could support awareness through providing
341 courses that aimed at enhancing knowledge and skills for basic life support among child
342 caregivers and babysitters

343

Manuscript

344 REFERENCES

1. Hamadneh SM, Kassab M, Abu-Hammad S, Al-bayyari NS, Hamadneh J, Obeidat NA, et al. Common Factors Associated with Child Deaths, at a Tertiary Care Centre in Irbid Government, Jordan. *Int J Pediatr* 2018; 6(8): 8021-30.
2. Atkins DL, Everson-Stewart S, Sears GK, Daya M, Osmond MH, Warden CR, Berg RA. Epidemiology and outcomes from out-of-hospital cardiac arrest in children: the Resuscitation Outcomes Consortium Epistry-Cardiac Arrest. *Circulation*. 2009;119:1484-1491.
<https://doi.org/10.1161/CIRCULATIONAHA.108.802678>
PMid:19273724 PMCID:PMC2679169
3. Smereka J, Szarpak L, Smereka A, Leung S, Ruetzler K. Evaluation of new two-thumb chest compression technique for infant CPR performed by novice physicians. A randomized, crossover, manikin trial. *The American journal of emergency medicine*. 2017; 35(4): 604-609.
<https://doi.org/10.1016/j.ajem.2016.12.045>
PMid:28040386
4. Rovin J D, Rodgers B M. Pediatric foreign body aspiration. *Pediatrics in review*. 2000; 21(3): 86-90.
<https://doi.org/10.1542/pir.21-3-86>
PMid:10702322
5. Lluna J, Olabarri M, Domènech A, Rubio B, Yagüe F, Bentez MT, et al. Recommendations for the prevention of foreign body aspiration. *Anales de Pediatría*. 2017;86: 50-56
<https://doi.org/10.1016/j.anpede.2016.04.003>
PMid:27234822
6. Domènech AB, Gutiérrez C, Ibáñez V, Barrios JE, Lluna J, Vila JJ. Foreign body aspiration, a potentially lethal underestimated problem. *Rev Pediatr Aten Primaria*. 2014;16:211.81-86
<https://doi.org/10.4321/S1139-76322014000400003>
7. Sayedi SJ, Gharavi M, Khakshour A, Davoudi Kang M, Kashani I, Jafarzadeh Esfehiani R. Case Report of a Vanished Aspirated Foreign Body; Cough may be more Helpful. *Int J Pediatr*. 2019; 7(5): 9363-67. DOI: 10.22038/ijp.2018.36214.3160
8. Callahan J. M., Fuchs, S. M. Advocating for life support training of children, parents, caregivers, school personnel, and the public. *Pediatrics*. 2018; 114(6):1676
<https://doi.org/10.1542/peds.2018-0704>
PMid:29793987
9. Roshana S, Batajoo K, Piryani R, Sharma M. Basic life support: knowledge and attitude of medical/paramedical professionals. *World journal of emergency medicine*. 2012; 3(2): 141.
<https://doi.org/10.5847/wjem.j.issn.1920-8642.2012.02.011>
PMid:25215053 PMCID:PMC4129799
10. Hamadneh S, Kassab M, Hamadneh J, Amarin Z. Sudden unexpected infant death in Jordan and the home environment. *Pediatrics international*. 2016; 58(12), 1333-1336.
<https://doi.org/10.1111/ped.13016>
PMid:27110690
11. Green, S. Ingested and Aspirated Foreign Bodies. *Pediatrics in review*. 2015. 36(10), 430-436

Manuscript

<https://doi.org/10.1542/pir.36-10-430>

PMid:26430203

12. Berg MD, Schexnayder SM, Chameides L, Terry M, Donoghue A, Hickey RW, Berg RA, Sutton RM, Hazinski MF. Part 13: pediatric basic life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010; 2;122):S862-75.

<https://doi.org/10.1161/CIRCULATIONAHA.110.971085>

PMid:20956229 PMCID:PMC3717258

13. Hamadneh S, Al-Shdayfat N, Al-Omari O, Hamadneh J, Bashtawi M, Alkhatib A., et al. Sudden Infant Death Syndrome in the Middle East: An Exploration of the Literature on Rates, Risk Factors, High Risk Groups and Intervention Programs. *Research Journal of Medical Sciences*. 2016;10(4):199-204

14. American Heart Association. 2005 American Heart Association (AHA) guidelines for cardiopulmonary resuscitation (CPR) and emergency cardiovascular care (ECC) of pediatric and neonatal patients: pediatric basic life support. *Pediatrics*. 2006 ;117(5):e989-1004.

<https://doi.org/10.1542/peds.2006-0219>

PMid:16651298

15. Chia, P., & Lian, W. B. Parental knowledge, attitudes, and perceptions regarding infant basic life support. *Singapore medical journal*. 2014; 55(3): 137-45

<https://doi.org/10.11622/smedj.2014033>

PMid:24664380 PMCID:PMC4293985

16. MacKinnon R, Stoeter R, Doherty C, Fullwood C, Cheng A, Nadkarni V, Stenfors-Hayes T, Chang T. Self-motivated learning with gamification improves infant CPR performance, a randomised controlled trial. *BMJ Simulation and Technology Enhanced Learning*. 2015;1(3)

<https://doi.org/10.1136/bmjstel-2015-000061>

17. Hayes NM, Chidekel A. Pediatric choking. *Del Med J*. 2004; 76(9):335-40.

18. Thorne CJ, Lockey AS, Bullock I, Hampshire S, Begum-Ali S, Perkins GD E-learning in advanced life support--an evaluation by the Resuscitation Council (UK). *Advanced Life Support Subcommittee of the Resuscitation Council (UK)*. *Resuscitation*. 2015 May; 90:79-84

<https://doi.org/10.1016/j.resuscitation.2015.02.026>

PMid:25766092

19. Teague G, Riley RH. Online resuscitation training. Does it improve high school students' ability to perform cardiopulmonary resuscitation in a simulated environment? *Resuscitation*. 2006;71(3):352-7

<https://doi.org/10.1016/j.resuscitation.2006.05.007>

PMid:17069951

20. Li Q, Ma EL, Liu J, Fang LQ, Xia T. Pre-training evaluation and feedback improve medical students' skills in basic life support. *Med Teach*. 2011; 33(10):e549-55.

<https://doi.org/10.3109/0142159X.2011.600360>

PMid:21942491

Manuscript

346 **Table 1:**

347 *The demographic profile of the participants (N=77)*

Characteristic	Sub-factors	n	%
education	Do not hold college/university degree	28	36%
	Hold college/university degree	49	64%
Age group	20-35 years	52	68%
	36-50	25	32%
Nationality	Residents	54	70%
	Not resedent (Refugees/Migrant)	23	30%

348

349

350

Manuscript

351

352 **Table 2:**

353 *Knowledge regarding Infant basic life support/ foreign body aspiration*

Questions	Sub-factors	n	%
1- Have you ever heard about CPR for children?	Yes	48	62%
	No	29	38%
2- Have you ever attended a training/education session on the management of foreign body aspiration for infants?	Yes	23	30%
	No	54	70%
3- Will you attend an IBLS course if it is offered to you?	Yes	65	84%
	No	12	16%

354

355

Manuscript

356 **Table 3:**

357 *The proportion of participants who obtained pass marks*

Question	Correct answer	% of correct responses
1. What is the Jordanian Ambulance access number?	199	73%
2. How long can a child’s brain live without oxygen (breathing)?	4-6 minutes	67%
3. Where is the location to check a pulse in a child aged less than1 year old?	On the brachial artery (inside of the upper arm between the elbow and the shoulder)	22%
4. I can try to wake an unresponsive infant by:	Gently tap or flick the infant's feet soles	44%
5. I can confirm that an unconscious child is not breathing by:		
	Looking for rising and fall of the chest.	48%
	Listening for air escaping during exhalation.	29%
	Feeling for the flow of air from the infant's mouth and nose using my cheek.	12%
6. the correct ratio compression to breath to provide effective CPR for an infant		
	30 compressions: 2 breaths if you are alone as a rescuer	13%
	15 compressions: 2 breaths if there are two rescuers	39%
7. If you witness a conscious infant unable to breathe and choking from a foreign body aspiration, what you should do to save his life?	Provide infant of 5 back blows and 5 chest thrusts until the object is expelled	21%
	Remove the piece aspirated by my fingers, only if I can see it	43%

358

359

Manuscript

360 **Table 4:**

361 *Perception about confidence in providing IBLs/foreign body aspiration management*

Questions	Count (%)				Total
	Not confident at all	Not confident	Somewhat confident	Very well confident	
Do you feel confident to provide a CPR for an infant in such an urgent situation?	17 (22%)	32 (42%)	23 (30%)	5 (6%)	77 (100%)
Do you feel confident about your current knowledge of managing foreign body aspiration among infants?	10 (13%)	27 (35%)	32 (42%)	8 (10%)	77 (100%)

362

363

364

365 Table-5: **Correlations**

366

	knowledge
	Pearson Correlation Sig. (2-tailed)
Education	.468(**) .000
Age	.451 (**) .000
Residency	.092 .734
N	77

367

368

** Correlation is significant at the 0.01 level (2-tailed).