Malignant pleural effusions as an alternative to biopsies for the detection of EGFR TKI sensitizing mutations

George D'Souza[1,2*], Chirag Dhar[1,2*^], Vishal Kyalanoor[3], Lokendra Yadav[3], Mugdha Sharmra[1,4], Mohammad Nawaz S[2], Sweta Srivastava[3]

[1]St. John's Research Institute, Bangalore, India

[2]Department of Pulmonary Medicine, St. John's Medical College and Hospital, Bangalore, India

[3]Department of Transfusion Medicine and Immunohaematology, St. John's Medical College and Hospital, Bangalore, India

[4]Department of General Medicine, St. John's Medical College and Hospital, Bangalore, India

* Equal contribution of authors

^ Present affiliation: University of California, San Diego, USA

Corresponding authors: Dr. Sweta Srivastava- sweta.s@stjohns.in (Lead contact), Dr. George D'Souza- george.dsouza@stjohns.in and Dr. Chirag Dhar- cdhar@health.ucsd.edu

Abstract

Lung cancer is the cause of a fourth of all cancer related deaths. About a third of all lung adenocarcinoma tumours harbour mutations on exons 18, 19, 20 and 21 of the epidermal growth factor receptor (EGFR) gene. Patients having these mutations are treated with EGFR Tyrosine kinase inhibitors and show better clinical outcomes. Detection of these mutations must be both accurate and timely. We utilized malignant pleural effusions for EGFR mutation detection. We started with a direct sequencing approach and detected a common exon 19 deletion. We then attempted to test the sensitivity of this method by sequencing this sample diluted with wildtype DNA. While signatures of this large deletion were detectable even in a 1 part mutant and 99 part wildtype mixture, this method was going to limit our ability to detect single base pair mutations confidently from pleural effusions. We then tried a EGFR mutant-specific quantitative polymerase chain reaction (PCR) based assay on n=10 pleural effusion samples (1 non-malignant pleural effusion as an internal negative control). 5/9 (55.55%) samples harboured EGFR mutations with 2/9 (22.22%) being exon 19 deletions and 3/9 (33.33%) had the S768I exon 20 mutation. These data suggest malignant pleural effusions may be a utilized to detect mutations even before a biopsy is made available. The high rate of S768I mutations seen in our study suggests that cancer cells harbouring these mutations may be superior in their ability to migrate, home or reside in pleural fluid.

Introduction

Lung cancer is a major contributor to death due to cancer. Amongst the various types of lung cancer, the non-small cell lung cancer accounts for approximately 80% of all cases[1]. In India, a significant rise has been seen in the time trends of lung cancer in Delhi, Chennai and Bangalore. Lung cancer is the cause of 6.9 percent of all new cancer cases and 9.3 percent of all cancer related deaths in both men and women in India[2]. Mutations seen on Exons 18, 19, 20 and 21 of the Epidermal growth factor receptor (EGFR) gene are frequently observed in these cases and have been shown to be present in nearly a third of all lung adenocarcinoma cases[3-5]. Epidermal growth factor receptor (EGFR) is a family member of receptor tyrosine kinases and play a central role in cellular signalling which promote cell growth and proliferation. Some mutations in this protein strongly predict the efficacy of EGFR inhibitors used in the treatment of these cases. There are reportedly more than 32 different mutations that have been detected in EGFR gene, and are distributed across the 4 exons namely, 18, 19, 20, and 21[6, 7]. More than 20 different in-frame deletions on exon 19 have been reported accounting for nearly half of the cases of EGFR mutant lung adenocarcinoma. Importantly, tumours harbouring these deletions are found to be sensitive to EGFR TKIs such as erlotinib and gefitinib[8]. This targeted therapy has been found to increase clinical outcomes in these patients[9]. It is imperative that such mutations are diagnosed accurately and early for the benefit of patients.

Response rates of more than 70 percent have been observed in patients on EGFR targeted therapy[10]. Patients with EGFR mutant positive advanced NSCLC treated with Erlotinib, an EGFR Tyrosine kinase inhibitor (TKI) have shown better response rates and Progression-free survival (PFS) as compared to those on first line chemotherapeutic agents[11-14].

Diagnostic tests for the detection of these EGFR mutations are now part of the routine management

of lung adenocarcinoma[15]. In spite of the invasiveness and morbidity associated with lung

biopsies[16], the primary tumour is preferred for detection of mutations[17]. Most often, an

amplification-refractory mutation system (ARMS) is used for the detection of these mutations

from the biopsy. The utility of pleural effusion (PE) in the detection of EGFR mutations has been

well established over the years in the field of lung cancer diagnosis[18, 19]. In this study, we utilized

malignant pleural effusions in an attempt to detect EGFR mutations at our tertiary care centre.

Ethics

Ethical clearance was obtained from the Institutional Ethics Committee of St. John's National

Academy of Health as per IERB guidelines (Study No. 1/2016). Appropriate consents for carrying

out experimental genetic tests were obtained.

Methods

Pleural fluid processing and DNA isolation

Pleural fluid was tapped from patients with malignant pleural effusions and centrifuged at 3500

rpm for 5 mins at 4°C to collect the cell pellet. The cell pellets were stored in -80°C until further

processing. The DNA was extracted from the samples using the QIAamp DNA Mini Kit (Qiagen)

as per the manufacturer's instructions. Samples were analyzed for their quality using BioSpec Nano

UV-VIS Spectrophotometer (Shimadzu Corp.) and electrophoresed on an agarose gel to ascertain

its integrity.

Direct sequencing

Appropriate primers for exons 18, 19, 20 and 21 of the EGFR gene were designed and obtained

based on the reference sequence available on NCBI (Table 1). Genomic DNA obtained was

subjected to a gradient PCR to optimize the annealing temperature and was visualized on a 1.2%

agarose gel (Fig.1). The PCR products were subjected to Direct sequencing.

Table 1

Region of Interest	Forward Primer 5' to 3'	Reverse Primer 5' to 3'
Exon 18	GTCCTTCCAAATGAGCTGGCAAG	ACAAAGAGTAAAGTAGATGATGG
Exon 19	TGTCCCTCACCTTCGGGGTGCAT	ACATTTAGGATGTGGAGATGAGCA
Exon 20	CTCAAGATCGCATTCATGCGTC	GACAGGCACTGATTTGTGCAC
Exon 21	AGTAGTCACTAACGTTCGCCAG	TCCCAGCAAGTACTGTTCCC

Real-time PCR based detection of EGFR mutations

DNA samples were subjected for testing a total of 10 detectable EGFR mutations in a 96-well Real

Time PCR format using the TRUPCR® EGFR Kit (3B BlackBio Biotech India Ltd.) as per the

manufacturer's instructions. Real-time PCR data analysis and identification of EGFR mutant

samples was done. To the best of our knowledge, this is the first time this kit is being using to

detect mutations from PEs.

Results

Optimization of a polymerase chain reaction protocol to amplify EGFR exon 18, 19, 20 and

21 from malignant pleural effusion (MPE).

Initially, we attempted a PCR amplification coupled with a direct sequencing approach to detect

EGFR mutations from malignant pleural effusions. Appropriate primers (sequences listed in table

1) were designed and optimized on sample S0. Successful amplification of exons 18-21 by PCR

on S0 are shown in figure 1. The primers for exon 18 and 21 had 55 °C as the annealing temperature

while exon 19 and 20 were best amplified at an annealing temperature of 57.8 °C.

Figure 2: 1.2% Agarose gel imaged under ultra-violet light: 100bp ladder, A: GAPDH, B: Exon

20, lanes C-G: Exon 19 at annealing temperatures of 55 C, 57.8 C, 60.5 C, 63.1 C and 65 C lanes

H-L: Exon 18 at annealing temperatures of 55 C, 57.8 C, 60.5 C, 63.1 C and 65 C, lanes M-Q:

Exon 21 at annealing temperatures of 55 C, 57.8 C, 60.5 C, 63.1 C and 65 C Interestingly, two

distinct bands are visible for Exon 19 (lanes C-G)

130 **Detection of a common exon 19 deletion detected from MPE by direct sequencing.**

131 The amplicon sequences were compared with the reference EGFR sequence available on the

132 National Center for Biotechnology Information (https://www.ncbi.nlm.nih.gov/) database and

```
Query  11         CCAGATCACTGGGCAGCATGTGGCACCATCTCACAATTGACAGTTAACGTCTTCCTTCTC  70
                  |||||||||||||||||||||||||||||||||||||| |||||||||||||||||||||
Sbjct  55174650   CCAGATCACTGGGCAGCATGTGGCACCATCTCACAATTGCCAGTTAACGTCTTCCTTCTC
       55174709

Query  71         TCTCTGTCATAGGGACTCTGGATCCCAGAAGGTGAGAAAGTTAAAATTCCCGTCGCTATC  130
                  ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  55174710   TCTCTGTCATAGGGACTCTGGATCCCAGAAGGTGAGAAAGTTAAAATTCCCGTCGCTATC
       55174769

                                    <----------------->
Query  131        AAGGAATTA------------------AAAGCCAACAAGGAAATCCTCGATGTGAGTTTC  172
                  |||||||||                 |||||||||||||||||||||||||||||||||
Sbjct  55174770   AAGGAATTAAGAGAAGCAACATCTCCGAAAGCCAACAAGGAAATCCTCGATGTGAGTTTC
       55174829

Query  173        TGCTTTGATGTGTGGTC-TGCATGGCTCTGAACCTCAGGCCCACCTTTTCTCATGTCTGG  231
                  ||||||| |||||||  |  ||||||||||||||||||||||||||||||||||||||||
Sbjct  55174830   TGCTTTGCTGTGTGGGGGTCCATGGCTCTGAACCTCAGGCCCACCTTTTCTCATGTCTGG
       55174889
```

exons 18, 20 and 21 were found to be identical. However, exon 19 was found to have an 18 bp

deletion (Fig. 3). On comparing with the wildtype EGFR cDNA sequence, this sequence was found

to have a deletion (c.2240_2258del18, p.L747_P753delinsS). Searches on cBioPortal and

COSMIC yielded multiple other studies that have reported the same mutation.

Figure 3: Exon 19 sequence of S0 when aligned with the wildtype sequence indicated that this

was the c.2240_2258del18, p.L747_P753delinsS mutation.

The analysis of the sequence using FinchTV has been shown in figure 4. We observed that this

patient's sequences for exon 18, 20 and 21 were of good quality with no underlying peaks.

However, exon 19 sequence had variations in the sequence.

Figure 4: Chromatogram of exon 19 sequence detecting a deletion in sample S0.

Common exon 19 deletion detectable even in diluted DNA samples but base calling is error

prone.

The PCR product for exon 19 was then diluted in DNA obtained from a control subject to evaluate

the efficiency of Direct sequencing. These varying dilutions (1 in 10, 1 in 25, 1 in 50 and 1 in 100)

were then subjected to PCR and subsequent Direct sequencing. We observed that though the

mutated sequence could be detected even at a 1 in 100 dilution (Fig 5) there were significant

limitations of this sequencing-based approach.

```
Query  19         ATCACTGCTGCAGCATGTGTGCACGCATCTCACAATTGCCAGTTAACGTCTTCCTTCTCT  78
                  |||||| |||||||||| ||||  ||||||||||||||||||||||||||||||||||||
Sbjct  55174654   ATCACTG-GGCAGCATGTG-GCAC-CATCTCACAATTGCCAGTTAACGTCTTCCTTCTCT  55174710

Query  79         CTCTGTCATAGGGACTCTGGATCCCAGAAGGTGAGAAAGTTAAAATTCCCGTCGCTATCA  138
                  ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  55174711   CTCTGTCATAGGGACTCTGGATCCCAGAAGGTGAGAAAGTTAAAATTCCCGTCGCTATCA  55174770

Query  139        AGGAATC--GA-AAG-AA-------G-------ACAAGGAAATCCTCGATGTGAGTTTCT  180
                  |||||||  || ||| ||       |       |||||||||||||||||||||||||||
Sbjct  55174771   AGGAATTAAGAGAAGCAACATCTCCGAAAGCCAACAAGGAAATCCTCGATGTGAGTTTCT  55174830

Query  181        GCTTTGATGTGAGTTTCTGCTTTGCTCTGAACCTCAGGCCCACCTTTTCTCATGTCTGGC  240
                  |||| ||||| | ||||| ||||||||||||||||||||||||||||||||||||||||||
Sbjct  55174831   GCTTTGCTGTGTGGGGGTCCATGGCTCTGAACCTCAGGCCCACCTTTTCTCATGTCTGGC  55174890

Query  241        AGCTTTCTCATG--CTAGACCCTGCTCATCTCTCAC  274
                  |||| ||| || ||||||||||||||||| |||
Sbjct  55174891   AGCTG-CTC-TGCTCTAGACCCTGCTCATCTC-CAC  55174923
```

Figure 5: Sequencing of diluted S0 sample (1-part S0 DNA and 99 parts wildtype control DNA)

showed that while signatures of the deletion were still coming through, there was significant error

in base calls.

Malignant pleural effusions often contain wildtype stromal and hematopoietic cells along with the

metastatic lung cancer cells. In light of the above results, detection of single base part mutations

by direct sequencing of this heterogenous mix was obviously going to be a challenge. To overcome

this limitation of direct sequencing, we decided to use a mutant-specific PCR amplification assay

instead.

Mutant-specific quantitative PCR based detection of EGFR mutations from MPE suggests a

high frequency of S768I exon 20 mutation.

An established EGFR mutation real-time PCR kit was used to identify EGFR mutations in

malignant pleural effusions. The kit is optimized to detect mutations in a background of wild type

DNA with starting material as less as 3-5 ng of purified genomic DNA. The resultant qPCR was

analyzed for reaction controls, DNA assessment done based on an EGFR Reference gene and the

samples thus passed were analyzed for mutation as described in the manual according to the

defined ΔCt parameters. Our results showed that 5/9 (55.55%) of pleural effusions probed had

EGFR mutations. Of these, 2/9 (22.22%) were the exon 19 deletion E746_S750del18 and 3/9

(33.33%) were the exon 20 S768I mutations. Table 2 describes these findings and Figure 6 presents

these findings in the form of a doughnut chart. The amplification plots and ΔCt values are provided

in supplementary figures 1-7 and supplementary tables 1 and 2.

Table 2

Sample. No.	Sex	Mutation detected	Exon where mutation is present
S1	M	WT	-
S2	M	E746_S750DEL	Exon 19
S3	M	WT	-
S4	F	E746_S750DEL	Exon 19
S5	F	WT	-
S6	M	WT	-
S7	M	S768I	Exon 20
S8	M	WT	-
S9	M	S768I	Exon 20
S10	M	S768I	Exon 20

184   M= Male, F= Female, WT= wildtype EGFR, S5 was a non-malignant pleural fluid

185   sample that served as an internal assay negative control

Figure 6: Doughnut chart representing the various mutations detected in this study. Percentages are embedded in the slices and number of cases observed are indicated in parenthesis beneath the genotype.

Discussion

The prevalence of somatic mutations in the EGFR gene is high (~ 33%) in lung adenocarcinomas

[21, 22]. The detection of any EGFR mutation and the resultant treatment with EGFR TKIs (Tyrosine

Kinase Inhibitors) aids in sensitizing the tumours and thereby enhance the prognosis of patients.

EGFR mutation testing on tissue biopsies is the gold standard for diagnosis; however, it has its

own drawbacks, mainly resulting in uncontrolled bleeding and morbidity [23]. Utility of pleural

effusion fluid for mutation testing along with a plethora of other secondary specimens (pericardial

effusion, fine needle aspirates, CSF, cfDNA, etc) is reported in the quest of development of

alternate strategies for reliable testing [24]. We attempted to detect EGFR mutations from the pleural

effusion collected from patients as a part of routine treatment/diagnostic processes. As it involves

no exclusive sampling procedures, we proposed to develop a minimally invasive and inexpensive

method for EGFR mutation diagnostics.

In our study we first used PCR coupled to direct sequencing to detect mutation in pleural effusion derived malignant cells. We were able to detect a common mutation using this method. One of the biggest drawbacks of Direct sequencing versus allele specific detection lies in its limit of detection. Pleural fluid consists of a heterogeneous pool of stromal and inflammatory cells apart from the metastatic cells. Since the non-tumor cells harbour wildtype genomic sequence, the signal to noise ratio in electropherogram is lower for mutated sequence. The mutation sequence identification may be compromised in the event of suboptimal malignant cells numbers. We showed that at dilution of 1:100 of a mutated DNA with wildtype genomic DNA signatures of the mutation continue but the wildtype sequence bases provide conflicting results in certain positions of the gene. These inadequate results are more likely to be seen in single base pair mutations. We then utilized an EGFR mutation detection kit. Commercially available diagnostic kits are designed for testing EGFR mutation on FFPE sections. To the best of our knowledge, this is the first report of EGFR mutation testing by Taqman technology platform using pleural effusions in India. Jong Sik Lee and colleagues have used PNA clamping technology to detect EGFR mutations in the supernatant derived from the pleural effusion where in the objective was to detect mutations using cell free DNA [25]. Jie Lin et al have shown that EGFR mutations can be detected in the supernatant, cell pellets of the pleural effusion using High Resolution Melting (HRM) analysis and Sanger sequencing [26].

When implemented on a small pilot scale study (n=9), we were successful in detecting EGFR mutations in 5/9 (55%) subjects. The mutations that were detected in these samples were either S768I or E746_S750del. Importantly, a study by Leventakos and group have shown that S768I

mutation in exon 20 of EGFR is rare and is usually found in combination with sensitizing mutations

and they have concluded that the predictive and prognostic role of this mutation is yet to be fully

explored [27]. However, in another study it has been shown that S768I conferred reduced sensitivity

to gefitinib in vitro as compared to other mutations [28, 29]. In our study we have found that 33.3%

of the samples exhibited this mutation in isolation with absence of any other mutation. Some of

the reasons for this large difference are discussed here. Most studies utilize the primary tissue and

may miss this mutation if it appears later in the pathogenesis of the disease. Another reason could

be that the mutation detection panels used in these studies did not include this mutation. One may

also simply come to the conclusion that our cohort has a larger number of patients with this

mutation, we think otherwise. We hypothesize that S768I is a secondary mutation that develops

later in the course of the disease and cells harbouring this mutation are able to migrate to, home in

or reside within the pleural cavity more easily that cells without this mutation. One could further

this speculation that cells having this mutation have a greater metastatic potential (figure 6).

Further studies will be needed to prove these hypotheses.

Figure 6: Hypothesized role of the S768I mutation in the pathogenesis of metastasis in lung cancer (Elements of images sourced from http://smart.servier.com/).

The need to develop minimally invasive diagnostic tests in the management of cancers is gaining momentum, thus upon further evaluation and validation, EGFR mutation testing in lung adenocarcinoma patients could be potentially done using pleural effusion by exploiting Taqman based RQ-PCR in a dynamic fashion paving way for a reliable, robust, minimally invasive clinical diagnostic platform. However, this study is limited in its impact as it evaluates only 11 patients and further study needs to be undertaken to evaluate and validate this method in a larger cohort. Additionally, the detection of circulating tumour DNA should also be explored as an alternative to repeated biopsies is one of the techniques being studied [30]. This detection is being done using highly sensitive techniques such as droplet digital polymerase chain reaction. Resources must be directed towards the development of such minimally/non-invasive techniques, especially if these are inexpensive as well.

Acknowledgements

This work is supported by a grant from the Advanced Research Wing of the Rajiv Gandhi University of Health Sciences, Bangalore. We also thank Prof. Sudhir Krishna, Ms. Pranatharthi Annapurna and the sequencing division of NCBS, Bangalore for their support. Some of these data were presented at the National Lung Cancer Conference 2016 in Bhubaneshwar and at the American Society for Cellular Biology-European Molecular Biology Organization joint conference 2018 at San Diego[31-33]. A special mention to Dr. Paul Kalanithi for his battle against EGFR mutant lung cancer. His memoir When Breath becomes Air is humbling and yet, inspiring.

Conflict of Interest

The authors declare no conflict of interest.

Author contributions

G.D.S and C.D contributed equally this work which was conceived, planned, supervised and interpreted by G.D.S, C.D and S.S. Funding acquisition was by G.D.S. Experimental assays were performed and interpreted by C.D., V.K., L.Y. and M.S. Clinical sample acquisition was by C.D., M.N.S. and G.D.S. Manuscript was written and edited by C.D. and V.K. All authors approved the final manuscript.

References

1. Zhang X, Zhao Y, Wang M, Yap WS, Chang AY. Detection and comparison of epidermal growth factor receptor mutations in cells and fluid of malignant pleural effusion in non-small cell lung cancer. Lung Cancer. 2008 May; 60(2): 175-82.

2. Malik PS, Raina V. Lung cancer: prevalent trends & emerging concepts. Indian J Med Res. 2015, 141: 5-7.

3. Doval D, Prabhash K, Patil S, Chaturvedi H, Goswami C, Vaid A, Desai S, Dutt S, Veldore V, Jambhekar N, Mehta A, Hazarika D, Azam S, Gawande S, Gupta S. Clinical and epidemiological study of EGFR mutations and EML4-ALK fusion genes among Indian patients with adenocarcinoma of the lung. Onco Targets Ther. 2015; 8: 117-23.

4. Ding L, Getz G, Wheeler DA, Mardis ER, McLellan MD, Cibulskis K, Sougnez C, Greulich H, Muzny DM, Morgan MB, Fulton L, Fulton RS, Zhang Q, Wendl MC, Lawrence MS, Larson DE, Chen K, Dooling DJ, Sabo A, Hawes AC, Shen H, Jhangiani SN, Lewis LR, Hall O, Zhu Y, Mathew T, Ren Y, Yao J, Scherer SE, Clerc K, Metcalf GA, Ng B, Milosavljevic A, Gonzalez-Garay ML, Osborne JR, Meyer R, Shi X, Tang Y, Koboldt DC, Lin L, Abbott R, Miner TL, Pohl C, Fewell G, Haipek C, Schmidt H, Dunford-Shore BH, Kraja A, Crosby SD, Sawyer CS, Vickery T, Sander S, Robinson J, Winckler W, Baldwin J, Chirieac LR, Dutt A, Fennell T, Hanna M, Johnson BE, Onofrio RC, Thomas RK, Tonon G, Weir BA, Zhao X, Ziaugra L, Zody MC, Giordano T, Orringer MB, Roth JA, Spitz MR, Wistuba, II, Ozenberger B, Good PJ, Chang AC, Beer DG, Watson MA, Ladanyi M, Broderick S, Yoshizawa A, Travis WD, Pao W, Province MA, Weinstock GM, Varmus HE, Gabriel SB, Lander ES, Gibbs RA, Meyerson M, Wilson RK. Somatic mutations affect key pathways in lung adenocarcinoma.

Nature. 2008; 455: 1069-75.

5. Pao W, Miller V, Zakowski M, Doherty J, Politi K, Sarkaria I, Singh B, Heelan R, Rusch V,

Fulton L, Mardis E, Kupfer D, Wilson R, Kris M, Varmus H. EGF receptor gene mutations are

common in lung cancers from "never smokers" and are associated with sensitivity of tumors

to gefitinib and erlotinib. Proc Natl Acad Sci U S A. 2004; 101: 13306-11.

6. Sharma SV, Bell DW, Settleman J, Haber DA. Epidermal growth factor receptor mutations in

lung cancer. Nat Rev Cancer. 2007; 7: 169-181.

7. Catalogue   of   Somatic   Mutations   in   Cancer   (COSMIC),   2011,   v.51.

http://www.sanger.ac.uk/genetics/CGP/cosmic.

8. Cheng L, Alexander RE, Maclennan GT, Cummings OW, Montironi R, Lopez-Beltran A,

Cramer HM, Davidson DD, Zhang S. Molecular pathology of lung cancer: key to personalized

medicine. Mod Pathol. 2012; 25: 347-69.

9. Rocha-Lima CM, Raez LE. Erlotinib (tarceva) for the treatment of non-small-cell lung cancer

and pancreatic cancer. P T. 2009; 34: 554-64.

10. Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, Sunpaweravong P, Han B,

Margono B, Ichinose Y, Nishiwaki Y, Ohe Y, Yang JJ, Chewaskulyong B, Jiang H, Duffield

EL, Watkins CL, Armour AA, Fukuoka M. Gefitinib or carboplatin-paclitaxel in pulmonary

adenocarcinoma. N Engl J Med. 2009; 361: 947-57.

11. Zhou C, Wu YL, Chen G, Feng J, Liu XQ, Wang C, Zhang S, Wang J, Zhou S, Ren S, Lu S,

Zhang L, Hu C, Luo Y, Chen L, Ye M, Huang J, Zhi X, Zhang Y, Xiu Q, Ma J, You C. Erlotinib

versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-

positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label,

randomised, phase 3 study. Lancet Oncol. 2011, 12: 735-42.

341 12. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia-

342   Gomez R, Pallares C, Sanchez JM, Porta R, Cobo M, Garrido P, Longo F, Moran T, Insa A,

343   De Marinis F, Corre R, Bover I, Illiano A, Dansin E, de Castro J, Milella M, Reguart N,

344   Altavilla G, Jimenez U, Provencio M, Moreno MA, Terrasa J, Munoz-Langa J, Valdivia J, Isla

345   D, Domine M, Molinier O, Mazieres J, Baize N, Garcia-Campelo R, Robinet G, Rodriguez-

346   Abreu D, Lopez-Vivanco G, Gebbia V, Ferrera-Delgado L, Bombaron P, Bernabe R, Bearz A,

347   Artal A, Cortesi E, Rolfo C, Sanchez-Ronco M, Drozdowskyj A, Queralt C, de Aguirre I,

348   Ramirez JL, Sanchez JJ, Molina MA, Taron M, Paz-Ares L. Erlotinib versus standard

349   chemotherapy as first-line treatment for European patients with advanced EGFR mutation-

350   positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase

351   3 trial. Lancet Oncol. 2012 13: 239-46.

352 13. Sequist LV, Martins RG, Spigel D, Grunberg SM, Spira A, Janne PA, Joshi VA, McCollum

353   D, Evans TL, Muzikansky A, Kuhlmann GL, Han M, Goldberg JS, Settleman J, Iafrate AJ,

354   Engelman JA, Haber DA, Johnson BE, Lynch TJ. First-line gefitinib in patients with advanced

355   non-small-cell lung cancer harboring somatic EGFR mutations. J Clin Oncol. 2008; 26: 2442-

356   9.

357 14. Shigematsu H, Lin L, Takahashi T, Nomura M, Suzuki M, Wistuba, II, Fong KM, Lee H,

358   Toyooka S, Shimizu N, Fujisawa T, Feng Z, Roth JA, Herz J, Minna JD, Gazdar AF. Clinical

359   and biological features associated with epidermal growth factor receptor gene mutations in

360   lung cancers. J Natl Cancer Inst. 2005; 97: 339-46.

361 15. Peters S, Adjei AA, Gridelli C, Reck M, Kerr K, Felip E. Metastatic non-small-cell lung cancer

362   (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann

363   Oncol. 2012 23 Suppl 7: vii56-64.

16. Wiener RS, Schwartz LM, Woloshin S, Welch HG. Population-based risk for complications

after transthoracic needle lung biopsy of a pulmonary nodule: an analysis of discharge records.

Ann Intern Med. 2011 155: 137-44.

17. Pirker R, Herth FJ, Kerr KM, Filipits M, Taron M, Gandara D, Hirsch FR, Grunenwald D,

Popper H, Smit E, Dietel M, Marchetti A, Manegold C, Schirmacher P, Thomas M, Rosell R,

Cappuzzo F, Stahel R. Consensus for EGFR mutation testing in non-small cell lung cancer:

results from a European workshop. J Thorac Oncol. 2010; 5: 1706-13.

18. Dan Liu ,Yachao Lu , Zhenli Hu , Ning Wu, Xiaomeng Nie, Yang Xia, Yiping Han, Qiang Li,

Guanshan Zhu, Chong Bai Malignant Pleural Effusion Supernatants Are Substitutes for

Metastatic Pleural Tumor Tissues in EGFR Mutation Test in Patients with Advanced Lung

Adenocarcinoma. Plos ONE February 2014; 9 (2): e89946.

19. H Kimura, Y Fujiwara, T Sone, H Kunitoh, T Tamura, K Kasahara, K Nishio. *EGFR* mutation

status in tumour-derived DNA from pleural effusion fluid is a practical basis for predicting the

response to gefitinib. Molecular Diagnostics. 2006.

20. TRUPCR® EGFR PCR Kit, https://www.3bblackbio.com/trupcr-oncology-egfr.html

21. Yue-Lun Zhang, Jin-Qiu Yuan, Kai-Feng Wang, Xiao-Hong Fu, Xiao-Ran Han, Diane

Threapleton, Zu-Yao Yang, Chen Mao, Jin-Ling Tang. The prevalence of *EGFR* mutation in

patients with non-small cell lung cancer: a systematic review and meta-analysis. Oncotarget.

2016; 7 (48): 78985–78993.

22. Han B, Tjulandin S, Hagiwara K, Normanno N, Wulandari L, Laktionov K, Hudoyo A, He

Y, Zhang YP, Wang MZ, Liu CY, Ratcliffe M, McCormack R, Reck M. EGFR mutation

prevalence in Asia-Pacific and Russian patients with advanced NSCLC of adenocarcinoma

and non-adenocarcinoma histology: The IGNITE study. Lung Cancer. 2017; 113: 37-44.

23. Lim, Minji, Kim, Chi-Ju, Sunkara, Vijaya, Kim, Mi-Hyun, Cho, Yoon-Kyoung. Liquid Biopsy

in Lung Cancer: Clinical Applications of Circulating Biomarkers (CTCs and ctDNA).

Micromachines. 2018; 9, 100.

24. Gillian Ellison, Guanshan Zhu, Alexandros Moulis, Simon Dearden, Georgina Speake, Rose

McCormack. EGFR mutation testing in lung cancer: a review of available methods and their

use for analysis of tumour tissue and cytology samples. J Clin Pathol 2013; 66: 79–89.

25. Lee, J.S., Hur, J.Y., Kim, I.A. *et al.* Liquid biopsy using the supernatant of a pleural effusion

for EGFR genotyping in pulmonary adenocarcinoma patients: a comparison between cell-free

DNA and extracellular vesicle-derived DNA. BMC Cancer. 2018; 18**,** 1236.

26. Jie Lin, Ye Gu, Rui Du, Min Deng, Yaodan Lu, Yanqing Ding. Detection of *EGFR* mutation

in supernatant, cell pellets of pleural effusion and tumor tissues from non-small cell lung cancer

patients by high resolution melting analysis and sequencing. Int J Clin Exp Pathol. 2014; 7(12):

8813–8822.

27. Leventakos K, Kipp BR, Rumilla KM, Winters JL, Yi ES, Mansfield AS. S768I Mutation in

EGFR in Patients with Lung Cancer. J Thorac Oncol. 2016; 11(10): 1798-801.

28. Kancha RK, von Bubnoff N, Peschel C, Duyster J. Functional analysis of epidermal growth

factor receptor (EGFR) mutations and potential implications for EGFR targeted therapy. Clin

Cancer Res. 2009; 15(2): 460-7.

29. Chen YR, Fu YN, Lin CH, Yang ST, Hu SF, Chen YT, Tsai SF, Huang SF. Distinctive

activation patterns in constitutively active and gefitinib-sensitive EGFR

mutants. Oncogene. 2006; 25: 1205–1215.

30. Suraj, S., Dhar, C., Srivastava, S. Circulating nucleic acids: An analysis of their occurrence in

malignancies (Review). Biomedical Reports 2017; 6.1: 8-14.

31. Dhar C, Sharma M, Nawaz M, D'Souza G, Srivastava S. Sanger sequencing of metastatic

pleural effusion: An inexpensive, minimally invasive tool for the detection of EGFR mutations

in lung adenocarcinoma- Presented at the National Lung Cancer Conference, Bhubaneshwar

2016

(https://www.researchgate.net/publication/313108485_Sanger_sequencing_of_metastatic_ple

ural_effusion_An_inexpensive_minimally_invasive_tool_for_the_detection_of_EGFR_muta

tions_in_lung_adenocarcinoma-

_Presented_at_the_National_Lung_Cancer_Conference_201)

32. Dhar C, Sharma M, Nawaz M, D'Souza G, Srivastava S. Detection of EGFR TKI sensitizing

mutations from metastatic pleural fluid secondary to lung adenocarcinoma: a perspective from

Southern India. F1000Research. 2018 Dec 17;7.

33. 16. American Society for Cell Biology. 2018 ASCB Annual Meeting abstracts. Molecular

biology of the cell. 2018 Dec 15;29(26):3063.

Supplemental figure 1 shows the amplification curves for different mutations tested in Sample S2

Supplemental figure 2 shows the amplification curves for different mutations tested in Sample S4

Supplemental figure 3 shows the amplification curves for different mutations tested in Sample S7

Supplemental figure 5 shows the amplification curves for different mutations tested in Sample S9

**Supplemental figure 6 shows the amplification curves for different mutations tested in
Sample S10**

Supplemental figure 7 shows the representative amplification curves for different mutations tested and classified as a Wild Type sample or below LOD

Sample No.	ΔCt								
	G719X	T790M	S768I	EX20INS	C797S	L858R	L861Q	E746_S750DEL	EX19DEL
S1	11.261	12.919	12.393	15.518	13.704	21.527	19.990	14.589	8.697
S2	11.595	12.284	12.148	13.915	12.470	20.835	ND	**6.327**	9.159
S3	10.997	12.154	11.452	13.572	13.223	19.834	19.788	14.197	8.416
S4	10.981	12.106	12.276	15.961	12.879	18.905	ND	**5.597**	8.640
S5	12.438	11.107	11.385	14.670	19.424	ND	ND	ND	8.957
S6	11.314	12.079	11.776	17.363	12.901	ND	ND	14.173	8.642

**Supplemental table 1 shows the ΔCt values of
Samples S1- S6**

Sample No.	ΔCt								
	G719X	T790M	S768I	EX20INS	C797S	L858R	L861Q	E746_S750DEL	EX19DEL
S7	12.212	20.890	**4.077**	ND	ND	ND	ND	ND	ND
S8	15.164	ND	13.393	ND	ND	ND	ND	ND	ND
S9	13.942	ND	**8.558**	ND	ND	ND	ND	ND	ND
S10	12.041	ND	**6.280**	ND	ND	ND	ND	ND	ND

**Supplemental table 2 shows the ΔCt values of
Samples S7-S11**