Post-Exertional Malaise in People with Chronic Cancer-Related Fatigue

Rosie Twomey[1], Samuel T. Yeung[1], James G. Wrightson[2], Guillaume Y. Millet[1,3*], S. Nicole

Culos-Reed[1, 4, 5]

[1]Faculty of Kinesiology, University of Calgary, Calgary, Canada; [2]Department of Clinical

Neurosciences, University of Calgary, Canada; [3]Univ Lyon, UJM Saint-Etienne, Inter-

university Laboratory of Human Movement Biology, EA 7424, F-42023, Saint-Etienne,

France; [4]Department of Oncology, Cumming School of Medicine, Calgary, Canada;

[5]Department of Psychosocial Resources, Tom Baker Cancer Centre, Alberta Health Services,

Calgary, Canada

Corresponding Author:

Rosie Twomey, Ph.D.

Faculty of Kinesiology, University of Calgary, 2500 University Dr. NW, Calgary, Alberta,

T2N 1N4, Canada

T: +1 403-210-8688

Email: Rosemary.Twomey@ucalgary.ca

Abstract

Introduction: Cancer-related fatigue (CRF) is a distressing and persistent sense of tiredness or exhaustion that interferes with usual functioning. Chronic CRF continues for months after curative cancer treatment is complete. Post-exertional malaise (PEM) is a worsening of symptoms after physical or mental activity, with limited investigations in people with chronic CRF. The purpose of this study was to identify and describe self-reported incidences of PEM in people with chronic CRF.

Methods: Participants (n=18) were eligible if they scored ≤34 on the Functional Assessment of Chronic Illness Therapy-Fatigue scale and had a cancer-related onset of fatigue. Participants completed a brief questionnaire to assess PEM over a 6-month time-frame (the DePaul Symptom Questionnaire – Post-Exertional Malaise; DSQ-PEM). In addition, a maximal exercise test was used to investigate self-reported symptom exacerbation (via an open-ended questionnaire) after strenuous physical exertion.

Results: On the DSQ-PEM, three participants met objective scoring criteria, which included experiencing moderate to very severe symptoms at least half of the time, worsening of fatigue after minimal effort, plus a recovery duration of >24 h. Content analysis of responses to open-ended questionnaires identified five people who experienced a delayed recovery and symptoms of PEM after the exercise test.

Discussion: While only a minority (up to 29% in this sample) may experience PEM, exercise specialists and health care professionals working with people with chronic CRF must be aware that PEM may be an issue. Symptom exacerbation after exercise should be monitored, and exercise should be tailored and adapted to limit the potential for harm.

Keywords: exercise oncology; maximal exercise test; cancer rehabilitation

Introduction

Cancer-related fatigue (CRF) has been defined as a distressing, persistent sense of physical, emotional, and/or cognitive tiredness or exhaustion that is not proportional to recent activity and interferes with usual functioning [1]. The majority of people with cancer will experience CRF during active treatment, but this will resolve in the weeks after treatment. However, 20-30% of people will continue to experience CRF for months or years following treatment [2–4], Variations in the estimates of prevalence may reflect variability in the measurement of CRF [2–4]. Previous research has shown that people who have severe CRF at least 2 months after cancer treatment often do not recover and remain fatigued [3]. CRF that continues for months or years has been called chronic CRF [4] or post-cancer fatigue [5]. Although there is no strict consensus on when CRF can be described as chronic, we and other research groups [5,6] define chronic CRF/post-cancer fatigue as clinically-relevant CRF that continues for ≥3 months after the completion of cancer treatment.

CRF is assessed as a patient-reported outcome, and there are psychometric tools used for both routine screening [7], and the assessment of CRF severity [8]. Evidence-based guidelines for the screening, assessment and treatment of CRF have been developed, including pan-Canadian practice guidelines for clinical use [7,9]. In addition, there are consensus-developed diagnostic criteria for CRF, as proposed for the International Statistical Classification of Diseases and Related Health Problems, 10th revision (ICD-10) in 1998 [10]. The proposed ICD-10 criteria specify that six or more symptoms must have been present every day or nearly every day during the same two-week period over the past month, and at least one symptom must be "significant fatigue." Previous qualitative data have been interpreted as consistent with various descriptors in the ICD-10 case definition of CRF, including an excessive fatigue state, protracted in course, in relation to relatively limited physical activity [11]. Alongside items such as "perceived need to struggle to overcome inactivity" and "difficulty completing daily tasks attributed to feeling fatigued", "post-exertional malaise (PEM) lasting several hours" is also included in the ICD-10 criteria [10]. However, PEM has not been defined in the context of CRF, and is only vaguely addressed in the accompanying diagnostic interview guide, where the related question is: "Did you find yourself feeling sick or unwell for several hours after you had done something that took some effort?" [10].

PEM is a cardinal symptom of a separate and serious long-term illness called myalgic encephalitis and/or chronic fatigue syndrome (ME/CFS) [12]. In ME/CFS, PEM is described

by the Centers for Disease Control and Prevention as a worsening of symptoms after physical or mental activity that would not have caused a problem before illness [12]. PEM in ME/CFS has also been described as a constellation of extensively disabling signs and symptoms in response to exertion [13]. Historically, there has been difficulty in defining and measuring PEM (see [14] for more information), but for the purposes of this study, we will use PEM to denote concepts related to post-exertional symptom exacerbation [14].

The chronic nature of CRF in some cancer survivors has led to a comparison with ME/CFS [15]. Following a reanalysis of an influential study [16], and concerns among clinicians that exercise exacerbates or causes PEM, graded exercise (progressive aerobic activity) is no longer a recommended therapy in ME/CFS [13]. However, in people living with and beyond cancer, exercise (including moderate-vigorous exercise) is considered a safe and effective intervention to counteract the adverse physical and psychological effects of cancer and its treatment [17,18], including CRF (e.g. [19]). Considering a failure to adequately report adverse reactions to exercise in previous studies of ME/CFS [20], and the lack of data on PEM in people with chronic CRF, caution may be warranted. PEM is inadequately documented in the literature on chronic CRF and requires further investigation considering the potential implications for exercise prescription. The purpose of this study was to identify and describe self-reported incidences of PEM in a group of people with chronic CRF. Despite the inclusion of "PEM lasting several hours" in the proposed ICD-10 criteria for CRF [6], to our knowledge, this is the first attempt to measure PEM in people with chronic CRF. Based on our experience with an ongoing exercise trial for chronic CRF [21], we hypothesized that a minority of people with chronic CRF would experience PEM.

Methods

Participants

Participants were part of an ongoing prospective randomized controlled trial (ClinicalTrials.gov, NCT03049384 and [21]). Participants were recruited via the Alberta Cancer Registry (Alberta Health Services, Canada), and from clinicians and/or advertising at cancer centres local to the University of Calgary, Alberta, Canada. Participants were eligible if they were aged 18-75 years and had completed curative-intent cancer treatment (≥3 months and ≤5 years from enrolment), self-identified as feeling fatigued/tired, and self-identified that the fatigue/tiredness developed at any point during the course of the disease (e.g. during the treatment phase) and had not resolved. Time since treatment was calculated as the time between

the most recent (i.e. last) surgery, chemotherapy treatment, radiation treatment or stem cell

transplant (as relevant) and the date of the lab visit. Furthermore, participants were eligible if

they scored ≤34 [22] on the Functional Assessment of Chronic Illness Therapy Fatigue Scale

(FACIT-F) [23]. The FACIT-F is widely recommended for the assessment of CRF severity [8],

has a cut-off point (≤34) that correctly identifies over 90% of 'ICD-10 positive' cases, and has

been recommended for the diagnosis of CRF [22]. Based on these eligibility criteria, all

participants in the larger study were considered to have chronic CRF. Participants were

excluded if they had a contraindication to experimental procedures and/or exercise. Additional

inclusion and exclusion criteria have been reported previously [21]. Written informed consent

for all study procedures was obtained. This study was approved by the Health Research Ethics

Board of Alberta Cancer Committee (HREBA.CC-16-10-10) and was performed according to

the Declaration of Helsinki.

Laboratory Visit

Full details of the wider project have been reported elsewhere [21]. The data for this sub-study

were collected during and after an initial visit to the laboratory, previously described as Lab

Visit #1 [21]. The lab visit commenced between 8:00 – 9:30 am and lasted 2 – 3 h. Participants

were advised to consume breakfast 1.5 h prior to arrival at the laboratory. Participants were

instructed to arrive at the laboratory hydrated and to refrain from alcohol, caffeine and

strenuous activity for the preceding 24 h. Participants completed a Physical Activity Readiness

Questionnaire for Everyone (PAR-Q+) and were also screened for arrhythmia and

hypertension, determined during resting electrocardiography and blood pressure

measurements, respectively. Continuation with the study was conditional on the screening

process, and physician approval was sought at this stage where required. Otherwise,

participants were cleared for participation by a Certified Exercise Physiologist (Canadian

Society for Exercise Physiology). Following the measurement of stature (cm) and body mass

(kg), a maximal exercise test was conducted on a custom-built recumbent ergometer, whch

uses an electromagnetically-braked Velotron system (RacerMate Inc, Seattle, WA). Seat

position was adjusted for each participant, and self-selected cadence was determined during

the first minute of cycling. Participants were instrumented for the measurement of heart rate

and breath-by-breath pulmonary gas exchange and ventilation (Quark CPET, COSMED,

Rome, Italy). The starting power output (25-50 W) and increment (8-20 W) was adjusted on

an individual basis as that which was estimated to result in a test of 8-12 min duration. The

power was increased at 1-min intervals until volitional exhaustion. Rating of perceived exertion

(RPE) (Borg's RPE scale [24], administered according to published instructions [25]) and

dyspnea (Borg CR-10 scale [26]) was recorded every minute. For all cycling tests, verbal

encouragement was provided by the same experimenters every 20-60 s [27]. A fingertip blood

sample was collected at exercise cessation for the analysis of blood lactate (Lactate Scout$^+$,

EFK Diagnostics, Cardiff, UK), and participants completed a supervised cool down. Data

collected during the test and used for the main study (e.g. peak oxygen uptake, heart rate, RPE

and blood lactate) is reported for the purpose of characterizing the test and describing the sub-

study participants.

Measurement of Post-Exertional Malaise

Before the exercise test, participants completed a brief questionnaire to assess PEM over a 6-

month time-frame (the DePaul Symptom Questionnaire – Post-Exertional Malaise; DSQ-PEM)

[28,29]. The first five items (Table 2) were recommended by the National Institutes of

Health/Centers for Disease Control and Prevention Common Data Elements (CDE) PEM

working group [30] and five supplementary items (including symptom duration, see Table 2)

were designed to operationalize the CDE recommendations [23] further. In the pre-registration

for this study (https://osf.io/nygq5/), we planned to use the Edmonton Symptom Assessment

System (ESAS-r) [31,32] as a quantitative assessment of symptom exacerbation after the

exercise test. However, we modified the instruction "Please circle the number that best

describes how you feel NOW" to "Please circle the number that best describes your average

symptom over the past 24 hours". This was confusing for participants who were using the

ESAS-r (with the momentary instruction) as part of the wider project. Therefore, we removed

this outcome. To gain a comprehensive understanding of the participant experience, an open-

ended questionnaire was provided (hard copy or via email, participant preference), and

participants were instructed to complete this 96 h after the exercise test. This accounted for the

potential for prolonged adverse responses as reported in people with ME/CFS [33], while also

limiting the requirement for recall. The open-ended questionnaire was designed based on a

review of the literature on PEM in ME/CFS (e.g. [11,14,33,34]) and is available online

(https://osf.io/nygq5/).

Data Analysis

For the DSQ-PEM, items 1-5 were scored according to Cotler *et al.* [28], i.e., a frequency score

of ≥2 (i.e., "about half the time" or more frequently) in combination with a severity score of

≥2 (i.e. "moderate" severity or more severe) was considered relevant. Items 6-10 are considered

supplementary, but were also scored according to Cotler *et al.* [28]. For clarity, this
questionnaire was not administered to support a diagnosis of ME/CFS in our participants;
rather, these scores were used as a tool to identify a self-reported experience of PEM.

For the open-ended questionnaires, the intended qualitative data analysis described in the pre-
registration was modified based on the detail of the responses. The open-ended questionnaires
were analyzed with qualitative content analysis [35]. The written responses were de-identified,
and two authors (RT and SY) independently read all responses several times in order to become
familiarized with the data. Responses were assigned codes according to a classification scheme
that was developed using (i) the questions on the open-ended questionnaire/knowledge about
PEM (e.g. experience of worsening of fatigue) and (ii) recurrent responses that arose from the
data (e.g. the need to nap). Responses were then coded independently, and an agreement was
reached on all items by RT and SY. Coded responses were then counted and summarized
descriptively.
192
Results

Of 18 participants (14 female), two did not complete the DSQ-PEM (missing data), and another
completed the exercise test but did not complete the open-ended questionnaire (due to food
poisoning). Participant characteristics and parameters related to the maximal exercise test are
presented in Table 1.

Table 1. Participant Characteristics and the Maximal Exercise Test

n = 18	
Age, mean ± SD	51 ± 15
Sex	
Male, *n* (%)	4 (22.2)
Female, *n* (%)	14 (77.8)
Height (cm), mean ± SD	169 ± 10
Body Mass (kg), mean ± SD	80.0 ± 17.1
Fatigue Score (FACIT-F), mean ± SD	23 ± 8
Months Since Treatment*, mean ± SD	22 ± 14
Cancer Type	
Breast, *n* (%)	9 (50)
Hodgkin's Lymphoma, *n* (%)	2 (11.1)
Diffuse Large B Cell Lymphoma, *n* (%)	2 (11.1)
Non-Hodgkin's Lymphoma, *n* (%)	1 (5.6)
Prostate, *n* (%)	1 (5.6)
Myelodysplastic Syndrome (Leukaemia), *n* (%)	1 (5.6)
Thyroid, *n* (%)	1 (5.6)
Oligodendroglioma, n (%)	1 (5.6)
Treatment Type	
Surgery, n (%)	13 (72.2)
Chemotherapy, *n* (%)	11 (61.1)
Radiation Therapy, *n* (%)	11 (61.1)
Hormonal Therapy (completed), *n* (%)	2 (22.2)
Hormonal Therapy (ongoing), *n* (%)	6 (33.3)
Stem Cell Transplant, *n* (%)	3 (11.2)
Maximal Exercise Test	
Peak Oxygen Uptake (L·min^{-1})	1.78 ± 0.43
Peak Oxygen Uptake (mL·kg^{-1}·min^{-1})	22.6 ± 4.8
End-Exercise Rating of Perceived Exertion, median [IQR]	19 [17-19]
Maximal Heart Rate (bpm)	157 ± 18
Percentage of Age Predicted Maximal Heart Rate (%)	92 ± 8
End-Exercise Blood Lactate (mmol/L)	7.2 ± 2.4
Exercise Time (min)	9.9 ± 1.6
Peak Power Output (W)	119 ± 33

 DSQ-PEM

In this sample of people with chronic CRF, we found that 50% of respondents indicated that they experienced at least one of the first 5 items (see Table 2) at least half time at a moderate or greater severity. A sub-set of participants (4/16) indicated that they feel worse after activities for >24 h, and 5/16 indicated that they do not exercise because exercise makes their symptoms worse. In order to apply objective criteria to help identify participants for whom PEM could be an issue, we used all scoring steps recommended by Cotler *et al* [28] to identify people who scored ≥2 for frequency and severity for any item from 1-5, plus answered "Yes" to item 7 or 8, plus gave a response of >14 h for item 9. Three people in this sample (19%) met these criteria (and we note that all three answered ≥2 for frequency and severity on all items from 1-5, answered yes for both item 7 and 8, and gave a response of >24 h for item 9).

Table 2. DSQ-PEM responses.

DSQ-PEM Item	Response	*n* (%)
1. Dead, heavy feeling after starting to exercise	Frequency ≥2 *and* Severity ≥2*	5 (31)
2. Next day soreness or fatigue after non-strenuous, everyday activities	Frequency ≥2 *and* Severity ≥2*	7 (39)
3. Mentally tired after the slightest effort	Frequency ≥2 *and* Severity ≥2*	6 (33)
4. Minimum exercise makes you physically tired	Frequency ≥2 *and* Severity ≥2*	7 (39)
5. Physically drained or sick after mild activity	Frequency ≥2 *and* Severity ≥2*	5 (31)
Frequency ≥2 *and* Severity ≥2 for *at least 1* of the above items		9 (50)
Frequency ≥2 *and* Severity ≥2 for *all* of the above items		4 (22)
6. If you were to become exhausted after actively participating in extracurricular activities, sports, or outings with friends, would you recover within an hour or two after the activity ended?	No	8 (44)
7. Do you experience a worsening of your fatigue/energy related illness after engaging in minimal physical effort?	Yes	7 (39)
8. Do you experience a worsening of your fatigue/energy related illness after engaging in mental effort?	Yes	12 (67)
9. If you feel worse after activities, how long does this last?	>24 h	4 (22)
10. If you do not exercise, is it because exercise makes your symptoms worse?	Yes	5 (31)

*Frequency = "how often you have had this symptom" and a score of ≥2 (2-4) ranges from "about half the time" to "all of the time." Severity = "how much has this symptom bothered you," and a score ≥2 (2-4) ranges from "moderate" to "very severe."

Open-Ended Questionnaire

There were 12 codes used for the content analysis: (1) worsening of fatigue; (2) duration/recovery time; (3) change in routine; (4) reduced daily activities; (5) muscle soreness; (6) weakness (7) joint pain; (8) mood; (9) memory/concentration; (10) need to nap or lie down; (11) flu-like symptoms; (12) positive comments on exercise/the test. Of 17 participants, 13 reported a worsening of fatigue after the exercise test. Responses ranged from mild, e.g. "*Felt a little more tired the day after the lab visit"* to more severe, e.g. "*I was thoroughly exhausted."*

Of the 13 participants who experienced worsening of fatigue, seven participants had to reduce their daily activities, and six had to change their routines or plans as a result of how they felt after the exercise test. This ranged from relatively mild: "*The only change was lying down in the morning, which is unusual for me. Typically, I get through the morning and rest in some way after lunch.*" To more severe "*The next day I stayed in bed most of the day and evening. Unable to walk the dog for two days. On the third day, I cancelled a 'coffee date' with a friend.*" In an item that arose from the data, six people noted that they need to nap or lie down later on the same day of the test and/or in the following days, e.g. "*I lay down all 4 afternoons for a rest. I don't usually do that*". Interestingly, in response to a question on the need to change plans or reduce activities, one participant answered, "*No. I anticipated the fatigue*", indicating that they expected increased fatigue (specifically, the day after the exercise test) and had planned accordingly. Also of note, is one participant who did not need to change their routine because their routine is already reduced/disrupted by the fluctuation and unpredictable nature of their fatigue: "*I don't usually plan anything because I don't know how I'm going to feel especially next day, [I] only plan day to da*y."

It was difficult to obtain clear information on how long it took for participants fatigue to return to 'usual' or 'normal,' partly because some people were unsure about their 'usual' or 'normal,' e.g. in the case of someone who only felt more tired later on the day of the test, they also commented: "*I don't know what is abnormal for me, as things are variable.*" Another confounder was that three participants were unsure whether to attribute increased fatigue to the test, or to other activities that were perceived as strenuous that occurred within the data collection period, specifically, e.g. "*more activities with family,*" and "*more stairs than usual.*" Nevertheless, of the 12 people who indicated worse fatigue, two participants only experienced this on the day of the test, seven returned to typical levels within 1-3 days, e.g. "*Back to normal after the two days had passed.*" and four were unsure if they had recovered by 96 h (when the questionnaire was completed) e.g. "*Each evening, I seemed to be so tired that I couldn't do my*

normal activities (research on family history). I didn't have the energy or ability to concentrate.
I don't know if I have recovered that yet because I have very little energy left by evening."

Of 17 participants, 11 experienced muscle soreness after the test, localized to the legs and
mainly attributed to cycling in all but one participant, e.g. "*The pain was just mild muscle pain*
that a person feels after working muscles more than usual." Four people experienced joint pain,
and five people referred to weakness. In one person, this was prolonged: "*The weakness was*
extreme and lasted in its worst state for days 3 and 4." Seven people experienced changes in
mood, and seven people had worse memory/concentration, e.g. "*First day, I had significant*
problems with my short-term memory and concentration." It is also noteworthy that one of the
participants who did not report worsening of memory and concentration compared to 'normal'
stated that they "*never feel normal*," referring to ongoing issues with memory and
concentration since cancer treatment. Five people experienced flu-like symptoms (e.g. "*chills,*
cold, body wet, and very sore" and "*I felt generalized aching throughout my body, like the*
flu."). Finally, in the additional comments, four participants self-reported feeling better after
exercise in general, e.g. "*Usually feel better after exercise*" or after the test e.g. "*I think in the*
afternoon I felt invigorated and then the next two days quite a bit more tired," and 2 people
commented on the test itself e.g. *"I felt very supported and safe throughout the test."*

Overall, using content analysis of the open-ended questionnaires delivered after the exercise
test, we identified five participants (29% of this sample) for whom PEM could be an issue.
Specifically, these participants indicated that they experienced worsening of fatigue, plus had
to change routine or reduce daily activities, plus had to nap or lie down, plus experienced ≥3
other symptoms (including muscle soreness, weakness, joint pain, mood disturbance,
memory/concentration or flu-like symptoms), plus had a delayed recovery of >2 days.

Discussion
The purpose of this study was to investigate self-reported incidences of PEM in people with
chronic CRF. In line with our hypothesis, our results provide preliminary evidence that a
minority (up to 29% in this sample, a possible 6/18 participants) of people with chronic CRF
experience symptoms of PEM. This has potential implications for exercise professionals
working with people experiencing chronic CRF. We recommend that CRF is monitored
throughout an exercise program (including asking people about their symptoms between
individual exercise sessions, and after increases in, e.g. exercise intensity) and that trials

involving people with chronic CRF report this patient-level data to facilitate the understanding
of PEM in this context, and therefore help mitigate the potential for harm.

Content analysis of open-ended questionnaires delivered 96 h after a maximal exercise test
provided evidence of a constellation of symptoms, change in routine/plans, and duration of
recovery in people experiencing chronic CRF. Although many participants experienced a
worsening of fatigue, based on each participant's profile of responses, and with particular
consideration of the severity of symptoms and duration of their recovery, we identified five
people whose descriptions were consistent with PEM. Despite qualitative comparisons
between chronic CRF and ME/CFS [15] and the inclusion of "PEM lasting several hours" in
the proposed ICD-10 criteria for CRF [10], to our knowledge, this is the first post-exercise
symptom characterization reported in people experiencing chronic CRF. A number of previous
studies have used maximal exercise tests to investigate the effects of physical exertion on
ME/CFS symptoms (e.g. [33]). Prolonged adverse responses to maximal exercise tests are
common in women with CFS (96% took >2 days to recover) and largely absent in a sedentary
control group (where 87% indicated full recovery within 24 h and 100% by 2 days) [33].
Although a maximal exercise test may be more strenuous than a typical exercise session
performed in line with the updated international exercise guidelines for cancer survivors [18],
the DSQ-PEM asks questions relating to minimal physical effort and non-strenuous, everyday
activities. Thus, we do not know what level of exertion provokes PEM. The current study was
conceived due to the need to reduce exercise intensity and carefully monitor symptom
exacerbation in a minority of people with chronic CRF participating in an ongoing exercise
trial [21]. In that trial, three participants described overwhelming exhaustion between exercise
sessions; therefore, little to no progression in exercise intensity or duration was made. These
anecdotal descriptions are in line with recent qualitative data from Penner *et al.* [36], where
fatigue was described not only as a chronic experience of imbalance but as involving "energy
crashes" that forced participants to "radically reshape their day-to-day lives."

In this sample, up to 29% of participants were identified as potentially experiencing PEM (i.e.
met the objective criteria for the DSQ-PEM and/or described PEM in the open-ended
questionnaire, see also Supplementary Table 1). Two of the five participants who described
PEM in their responses to the open-ended questionnaire also met the DSQ-PEM criteria. Of
the remaining three participants, one did not complete the DSQ-PEM (missing data); a second
met all DSQ-PEM criteria except they answered "No" to items 7 and 8 which refer to minimal

effort; and the third indicated "next day soreness or fatigue after non-strenuous, everyday activities" of a moderate severity about half the time and "minimum exercise makes you physically tired" at 2 ("severe" but only "a little of the time"), plus "yes" to items 7 and 10 (experience a worsening of fatigue after engaging in minimal physical effort, and do not exercise because it makes their symptoms worse). Although these latter two participants did not meet all scoring criteria for the DSQ-PEM, PEM does seem to be an issue above a certain intensity (i.e., above minimal), possibly explaining why the open-ended questionnaire identified these participants when the DSQ-PEM did not. Finally, the DSQ-PEM identified one participant who was not identified from the open-ended questionnaire. This participant did report increased fatigue for 2-3 days after the exercise test but also noted, "*I like exercise. I like feeling strong. Like I am working toward health*". However, considering our experience of their later participation in the RCT, it is our clinical judgement that PEM was an issue for them (which also prevented returning to work).

Although this brief commentary about individuals is anecdotal, we believe it is important to communicate about individual profiles to minimize the potential for harm. Clearly, nuance is required in the interpretation of these two PEM measurements. Our goal is increased awareness of PEM as a potential issue within the wider context of "exercise is medicine" in oncology [37]. If exercise is medicine in chronic CRF, side-effects at a given exercise dose must be carefully detailed. This will enable exercise professionals to provide the best exercise prescription in people with PEM, one that can be achieved within a pacing strategy that focusses on balancing rest and activity to achieve daily tasks, maintain physical function, and avoid symptom exacerbation. The new exercise guidelines for cancer survivors [18] acknowledge that some proportion of cancer survivors may not be able to tolerate the recommendations (particularly because dose reductions may have been poorly reported in previous trials). The results of this study reinforce the need to monitor tolerance and symptom exacerbation, especially in those with chronic CRF. Without appropriate safeguards (e.g. monitoring of fatigue throughout the exercise intervention), we do not know if people with PEM continue with little symptom exacerbation (e.g. due to low effort, or individualization and regular communication with an exercise professional), continue despite symptom exacerbation (and experience "energy crashes" [36] that may significantly impact daily activities), or drop out. It has been recently highlighted that the reporting of exercise interventions and specifically the "precise reporting of exercise treatment adherence (tolerability)" [38] must be improved. We add that these reports must include worsening of

symptoms, including fatigue. However, we emphasize that, for the majority of people with chronic CRF, exercise is likely to be beneficial for multiple and interacting components of physical and mental health. With support, most participants can reach physical activity guidelines (e.g. 30 minutes of aerobic activity at RPE 12, three times per week [18]) with no adverse consequences. Higher intensities/longer durations can be appropriate for people with chronic CRF who do not experience any adverse responses, particularly if the goal is to maximize physiological adaptations.

Study Limitations and Future Directions

We did not include a control group and do not have data on the responses of healthy/sedentary adults. However, the main purpose was to provide preliminary, rather than comprehensive evidence that a minority of people with chronic CRF experience symptoms of PEM. We note that in a previous study, 100% of controls recovered within 2 days of a maximal exercise test, 87% experienced an increased sense of wellbeing relative to the test, and none experienced issues such as weakness, impaired memory/concentration or mood disturbance [33]. The DSQ-PEM is not validated for use in people with CRF. However, it has been used in people with multiple sclerosis and post-polio syndrome (other clinical conditions where chronic fatigue can be an issue) [28]. As stated previously, we did not use the DSQ-PEM as a method to support a diagnosis of ME/CFS study, especially considering the diagnostic complexity of ME/CFS [39]). Here, it is used as a tool to help identify people experiencing symptoms of PEM, and any further speculation on whether participants would meet any given diagnostic criteria for ME/CFS if assessed by an expert clinician is out of our scope of practice. Because we were unable to obtain a copy of the open-ended questionnaire used previously [33], we designed a questionnaire (https://osf.io/349ky/) based on a review of the literature on PEM in ME/CFS. Another limitation is that maximal incremental exercise is not a typical exercise intensity prescribed within an intervention, and provides no information about the intensity of physical effort (below maximal) that could result in PEM. Furthermore, although the FACIT-F has been recommended to support a diagnosis of CRF [40], there is no consensus on what "diagnosis" means in this context. The proposed ICD-10 criteria were not accepted, and few studies have implemented them [41], and current guidelines have differing recommendations for screening and assessment [7,42], and we recognize a lack of consensus for the identification of someone with clinically-relevant CRF.

In this preliminary study, up to 29% of people appeared to experience PEM, but all participants
self-selected themselves into an RCT that explicitly involved a 12-week exercise intervention.
It is reasonable to suggest that the sample in the RCT, and therefore this sub-study, may not
represent the prevalence of PEM in people with chronic CRF, because people who experience
PEM may be less likely to enrol in an exercise trial. Future studies should estimate the
prevalence of PEM in people with chronic CRF. In addition, future studies should include semi-
structured qualitative interviews to elucidate the cancer-specific symptoms and impact of PEM.
In this preliminary study, we investigated people with chronic CRF, identified using a cancer-
related symptom onset, and presence of CRF at least 3 months and within 5 years (22 ± 14
months) after curative-intent cancer treatment. Because CRF is more prevalent during cancer
treatment [43], future work must address the potential of PEM after exercise in interventions
delivered during treatment. One study pre-emptively reduced the exercise prescription to
encourage attendance when treatment side effects (including fatigue) were anticipated to peak
during a chemotherapy cycle [44], but to our knowledge, there has been no post-exertional
symptom characterization during cancer treatment.
403

Conclusion
We found preliminary evidence that a minority of people with chronic CRF experience PEM.
We recommend that exercise professionals not only monitor fatigue throughout an exercise
program (i.e. asking participants how they felt in between individual sessions) but also report
this data if the exercise program is part of a clinical trial. In conclusion, the field of exercise
oncology must be aware that adverse responses to exercise may be an issue for a minority of
people with chronic CRF, and exercise interventions should be designed and adapted to
mitigate symptom exacerbation and limit the potential for harm.

REFERENCES

[1]   Berger AM, Mooney K, Alvarez-Perez A, et al. Cancer-related fatigue, version 2.2015. *J Natl Compr Canc Netw*. 2015;13(8):1012-1039.

[2]   Jones JM, Olson K, Catton P, et al. Cancer-related fatigue and associated disability in post-treatment cancer survivors. *J Cancer Surviv*. 2016;10(1):51-61.

[3]   Goedendorp MM, Gielissen MFM, Verhagen CAHHVM, et al. development of fatigue in cancer survivors: a prospective follow-up study from diagnosis into the year after treatment. *J Pain Symptom Manage*. 2013;45(2):213-222.

[4]   Abrahams HJG, Gielissen MFM, Schmits IC, et al. Risk factors, prevalence, and course of severe fatigue after breast cancer treatment: a meta-analysis involving 12 327 breast cancer survivors. *Ann Oncol*. 2016;27(6):965-974.

[5]   Sandler CX, Goldstein D, Horsfield S, et al. Randomized evaluation of cognitive-behavioral therapy and graded exercise therapy for post-cancer fatigue. *J Pain Symptom Manage*. 2017;54(1):74-84.

[6]   Bruggeman-Everts FZ, Wolvers MDJ, van de Shoot R, et al. Effectiveness of two web-based interventions for chronic cancer-related fatigue compared to an active control condition: results of the "fitter na kanker" randomized controlled trial. *J Med Internet Res*. 2017;19(10):e336.

[7]   Howell D, Keller-Olaman S, Oliver TK, et al. A pan-Canadian practice guideline and algorithm: screening, assessment, and supportive care of adults with cancer-related fatigue. *Curr Oncol*. 2013;20(3):233.

[8]   Minton O, Stone P. A systematic review of the scales used for the measurement of cancer-related fatigue (CRF). *Ann Oncol*. 2008;20(1):17-25.

[9]   Pearson EJM, Morris ME, McKinstry CE. Cancer-related fatigue: appraising evidence-based guidelines for screening, assessment and management. *Support Care Cancer*. 2016;24(9):3935-3942.

[10]   Cella D, Peterman A, Passik S, et al. Progress toward guidelines for the management of fatigue. *Oncol*. 1998;12(11A):369-377.

[11]   Bennett B, Goldstein D, Friedlander M, et al. The experience of cancer-related fatigue and chronic fatigue syndrome: a qualitative and comparative study. *J Pain Symptom Manage*. 2007;34(2):126-135.

[12]   Centers for Disease Control and Prevention. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) | Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Retrieved from: https://www.cdc.gov/me-cfs/symptoms-diagnosis/symptoms.html [Accessed November 19].

[13]   Davenport TE, Stevens SR, Vanness JM, et al. Checking our blind spots: current status of research evidence summaries in ME/CFS. *Br J Sports Med*. 2019:53:1198.

[14]   Chu L, Valencia IJ, Garvert DW, et al. Deconstructing post-exertional malaise in myalgic encephalomyelitis/ chronic fatigue syndrome: a patient-centered, cross-sectional survey. *PloS one*. 2018;13(6):e0197811.

[15]   Bennett B, Goldstein D, Friedlander M, et al. The experience of cancer-related fatigue and chronic fatigue syndrome: a qualitative and comparative study. *J Pain Symptom Manage*. 2007;34(2):126-135.

[16]   Wilshire CE, Kindlon T, Courtney R, et al. Rethinking the treatment of chronic fatigue syndrome—a reanalysis and evaluation of findings from a recent major trial of graded exercise and CBT. *BMC Psychol*. 2018;6(1):6.

[17]   Cormie P, Atkinson M, Bucci L, et al. Clinical Oncology Society of Australia position statement on exercise in cancer care. *Med J Australia*. 2018;209(4):184-187.

[18]   Campbell KL, Winters-Stone KM, Wiskemann J, et al. Exercise guidelines for cancer
survivors: consensus statement from international multidisciplinary roundtable. *Med*
Sci Sports Exerc. 2019;51(11):2375-2390.
[19]   Oberoi S, Robinson PD, Cataudella D, et al. Physical activity reduces fatigue in patients
with cancer and hematopoietic stem cell transplant recipients: a systematic review and
meta-analysis of randomized trials. *Crit Rev Oncol Hemat*. 2018;122:52-59.
[20]   Vink M, Vink-Niese A. Graded exercise therapy for myalgic encephalomyelitis/chronic
fatigue syndrome is not effective and unsafe. Re-analysis of a Cochrane review. *Health*
Psychol Open. 2018;5(2):205510291880518.
[21]   Twomey R, Martin T, Temesi J, et al. Tailored exercise interventions to reduce fatigue
in cancer survivors: study protocol of a randomized controlled trial. *BMC Cancer*.
2018;18(1):757.
[22]   van Belle S, Paridaens R, Evers G, et al. Comparison of proposed diagnostic criteria
with FACT-F and VAS for cancer-related fatigue: proposal for use as a screening tool.
Support Care Cancer. 2005;13(4):246-254.
[23]   Yellen SB, Cella DF, Webster K, et al. Measuring fatigue and other anemia-related
symptoms with the Functional Assessment of Cancer Therapy (FACT) measurement
system. *J Pain Symptom Manage*. 1997;13(2):63-74.
[24]   Borg G. An Introduction to Borg's RPE-Scale. Ithaca, NY: Mouvement Publications;
1985.
[25]   Borg G. Borg's Perceived Exertion and Pain Scales. Leeds: Human Kinetics; 1998.
[26]   Mador MJ, Rodis A, Magalang UJ. Reproducibility of Borg scale measurements of
dyspnea during exercise in patients with COPD. *Chest*. 1995;107(6):1590-1597.
[27]   Andreacci JL, LeMura LM, Cohen SL, et al. The effects of frequency of encouragement
on performance during maximal exercise testing. *J Sports Sciences*. 2002;20(4):345-
352.
[28]   Cotler J, Holtzman C, Dudun C, et al. A brief questionnaire to assess post-exertional
malaise. *Diagnostics*. 2018;8(3):E66.
[29]   Jason LA, Evans MA, Porter N, et al. The development of a revised Canadian myalgic
encephalomyelitis chronic fatigue syndrome case definition. *Am J Biochem*
Biotechnol. 2010;6:120–35
[30]   NINDS Common Data Elements (CDE) Group Post-Exertional Malaise Subgroup
Summary. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Retrieved from:
https://www.commondataelements.ninds.nih.gov/sites/nindscde/files/Doc/MECFS/PE
M_Subgroup_Summary.pdf [Accessed October 19].
[31]   Watanabe SM, Nekolaichuk C, Beaumont C, et al. A multicenter study comparing two
numerical versions of the Edmonton Symptom Assessment System in palliative care
patients. *J Pain Symptom Manage*. 2011;41(2):456-468.
[32]   Bruera E, Kuehn N, Miller MJ, Selmser P, Macmillan K. The Edmonton Symptom
Assessment System (ESAS): a simple method for the assessment of palliative care
patients. *J Palliative Care*. 1991;7(2):6-9.
[33]   VanNess JM, Stevens SR, Bateman L, et al. Postexertional malaise in women with
chronic fatigue syndrome. *J Women's Health*. 2010;19(2):239-244.
[34]   Lindheimer JB, Meyer JD, Stegner AJ, et al. Symptom variability following acute
exercise in myalgic encephalomyelitis/chronic fatigue syndrome: a perspective on
measuring post-exertion malaise. *Fatigue: Biomed, Health Behav*. 2017;5(2):69-88.
[35]   Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. *Qual Health*
Res. 2005;15(9):1277-1288.

[36] Penner CK, Zimmerman C, Kaptchuck T, et al. Towards a holistic framework of cancer
related fatigue: qualitative data on the experience of embodiment in breast cancer
survivors. PsyArXiv; 2019. doi:10.31234/osf.io/eg7ur
[37] Schmitz KH, Campbell AM, Stuiver MM, et al. Exercise is medicine in oncology:
engaging clinicians to help patients move through cancer. *CA Cancer J Clin.*
2019;69:468–484
[38] Nilsen TS, Scott JM, Michalski M, et al. Novel methods for reporting of exercise dose
and adherence: an exploratory analysis. *Med Sci Sports Exerc*. 2018;50(6):1134-1141.
[39] Brurberg KG, Fønhus MS, Larun L, et al. Case definitions for chronic fatigue
syndrome/myalgic encephalomyelitis (CFS/ME): a systematic review. *BMJ Open*.
2014;4(2).
[40] Van Belle S, Paridaens R, Evers G, et al. Comparison of proposed diagnostic criteria
with FACT-F and VAS for cancer-related fatigue: proposal for use as a screening tool.
Support Care Cancer. 2005;13(4):246-254.
[41] Donovan KA, McGinty HL, Jacobsen PB. A systematic review of research using the
diagnostic criteria for cancer-related fatigue. *Psycho-Oncology*. 2013;22(4):737-744.
[42] Bower JE, Bak K, Berger A, et al. Screening, assessment, and management of fatigue
in adult survivors of cancer: an American Society of Clinical Oncology clinical practice
guideline adaptation. *J Clin Oncol*. 2014;32(17):1840-1850.
[43] Henry DH, Viswanathan HN, Elkin EP, et al. Symptoms and treatment burden
associated with cancer treatment: results from a cross-sectional national survey in the
U.S. *Support Care Cancer*. 2008;16(7):791-801.
[44] Kirkham AA; Bland KA; Zucker DS. "Chemotherapy-periodized" exercise to
accommodate for cyclical variation in fatigue. *Med Sci Sports Exerc*. 2019; doi:
10.1249/MSS.0000000000002151