Title

Long title

Minimal transmission in an influenza A (H3N2) human challenge-

transmission model within a controlled exposure environment

Short title

Human influenza A transmission model in a controlled environment

Authors

Jonathan S. Nguyen-Van-Tam[1], Ben Killingley[1*], Joanne Enstone[1], Michael Hewitt[1],

Jovan Pantelic[2,#a], Michael L. Grantham[2,#b], P. Jacob Bueno de Mesquita[2], Robert

Lambkin-Williams[3], Anthony Gilbert[3], Alexander Mann[3], John Forni[3,#c], Catherine J.

Noakes[4], Min Z. Levine[5], LaShondra Berman[5], Stephen Lindstrom[5], Simon

Cauchemez[6,#d], Werner Bischoff[7], Raymond Tellier[8], and Donald K. Milton[2], for the

EMIT Consortium[^].

Affiliations

[1] Health Protection and Influenza Research Group, Division of Epidemiology and
Public Heath, University of Nottingham School of Medicine, Nottingham, UK

26 [2] University of Maryland School of Public Health, Maryland Institute for Applied
27 Environmental Health, College Park, MD, USA

28
29 [3] hVIVO London, UK

30
31 [4] University of Leeds School of Civil Engineering, Leeds, UK

32
33 [5] Centers for Disease Control and Prevention, Influenza Division, Atlanta, GA, USA

34
35 [6] Imperial College London, MRC Centre for Outbreak Analysis and Modelling,
36 Department of Infectious Disease Epidemiology, London, UK

37
38 [7] Wake Forest School of Medicine, Winston-Salem, NC, USA

39
40 [8] University of Calgary, Cumming School of Medicine, Department of Pathology and
41 Laboratory Medicine, Calgary, Alberta, CA

42
43 [#a] Current Address: University of California, Center for the Built Environment,
44 Berkeley, CA, USA

45
46 [#b] Current Address: Missouri Western State University, St. Joseph, MO, USA

47
48 [#c] Current Address: Department of Acute and Specialty Care, MSD, London, UK

49
[#d] Current Address: Mathematical Modelling of Infectious Diseases Unit, CNRS
UMR2000: Génomique évolutive, modélisation et santé (GEMS), Center of
Bioinformatics, Biostatistics and Integrative Biology, Institut Pasteur, Paris, FR

[*] Corresponding author

E-mail: ben.killingley@nhs.net (BK)

[^] A complete list of the EMIT Consortium can be found in the Supporting

Information.

Author Contributions

Conceptualization: JSN-V-T, BK, JE; Methodology: all authors; Investigation: all

authors except RT (face shield tests: WB); Formal analysis: JSN-V-T, BK, JE, AM,

DKM, MG, MH, PJB, CJN; Data curation: all authors; Original draft preparation:

JSN-V-T, BK, JE, DKM, AM, RLW, PJB; Review, editing and approval: all authors.

Abstract

Uncertainty about the importance of influenza transmission by airborne

droplet nuclei generates controversy for infection control. Human challenge-

transmission studies have been supported as the most promising approach to fill this

knowledge gap. Healthy, seronegative volunteer 'Donors' (n=52) were randomly

selected for intranasal challenge with influenza A/Wisconsin/67/2005 (H3N2).

'Recipients' randomized to Intervention (IR, n=40) or Control (CR, n=35) groups

were exposed to Donors for four days. IRs wore face shields and hand sanitized

frequently to limit large droplet and contact transmission. One transmitted infection

was confirmed by serology in a CR, yielding a secondary attack rate of 2.9% among

CR, 0% in IR (p=0.47 for group difference), and 1.3% overall, significantly less than

16% (p<0.001) expected based on a proof-of-concept study secondary attack rate and

considering that there were twice as many Donors and days of exposure. The main

difference between these studies was mechanical building ventilation in the follow-on

study, suggesting a possible role for aerosols.

Author summary

Understanding the relative importance of influenza modes of transmission informs strategic use of preventive measures to reduce influenza risk in high-risk settings such as hospitals and is important for pandemic preparedness. Given the increasing evidence from epidemiological modelling, exhaled viral aerosol, and aerobiological survival studies supporting a role for airborne transmission and the potential benefit of respirators (and other precautions designed to prevent inhalation of aerosols) versus surgical masks (mainly effective for reducing exposure to large droplets) to protect healthcare workers, more studies are needed to evaluate the extent of risk posed airborne versus contact and large droplet spray transmission modes. New human challenge-transmission studies should be carefully designed to overcome limitations encountered in the current study. The low secondary attack rate reported herein also suggests that the current challenge-transmission model may no longer be a more promising approach to resolving questions about transmission modes than community-based studies employing environmental monitoring and newer, state-of-the-art deep sequencing-based molecular epidemiological methods.

Introduction

Influenza virus is a pathogen of global health significance, but human-to-human transmission remains poorly understood. In particular, the relative importance of the different modes of transmission (direct and indirect contact, large droplet, and aerosols (airborne droplet nuclei)) remains uncertain during symptomatic and asymptomatic infection (1–4).

The evidence base for influenza transmission is derived from studies that have assessed: virus deposition and survival in the environment; the epidemiology of disease; pharmaceutical and non-pharmaceutical interventions; animal models; and mathematical models of transmission. Those approaches have yet to produce conclusive data quantifying the relative importance of human-human transmission modes (1,2).

Infection control guidance for pandemic and seasonal influenza assumes that most transmission occurs during symptomatic infection, predominantly via large droplet spread at short range (1-2m) (1). Thus, social distancing measures are often proposed to mitigate the spread and impact of a pandemic; and hand washing and respiratory etiquette are promoted to reduce transmission. Evidence to support the possibility of aerosol transmission has grown over recent years (5–7). and leads to controversies about when and if filtering facepiece respirators (and other precautions designed to prevent inhalation of aerosols) versus surgical masks (mainly capable of reducing large droplets and some fine particles) should be used to protect healthcare workers, particularly during a severe pandemic (1,3,4,8–10).

An expert panel, after in-depth review of the challenges facing community- and workplace-based intervention studies and their failure thus far to provide definitive evidence regarding the relative contribution of the various modes,

concluded that a human challenge-transmission study would be a more promising

direction for future research (11). Influenza challenge studies in humans have been

conducted to investigate disease pathogenesis and the efficacy of antivirals and

vaccines. Challenge studies assessing human-to-human transmission had not been

performed (11). In 2009, we demonstrated proof-of-concept that healthy seronegative

volunteers inoculated intranasally with influenza A/Wisconsin/67/2005 (A/WI), an

H3N2 virus, would develop symptoms of influenza-like illness (ILI) and, under two

days of household-like conditions without environmental controls, transmit infection

to other seronegative volunteers. This suggested that larger scale human challenge-

transmission models might be useful to evaluate transmission modes. A subsequent

international workshop discussed the potential that human challenge-transmission

studies, with appropriate interventions, monitoring of aerosol shedding, and

environmental controls, could provide definitive results (12). Here, we report a large

follow-on study, including design factors (Fig 1) aimed at assessing the importance of

aerosol transmission in human-to-human transmission of influenza virus. Although

the study did not achieve the intended level of transmission required for a more

conclusive interpretation, it has revealed important lessons about potential airborne

transmission and study design that represent critical knowledge to support effective

large-scale, costly studies in this area.

Results

Participation and safety

Between January and June 2013, 496 seronegative (HAI≤10 to the challenge virus antigen) volunteers underwent study-specific screening and 166 entered the quarantine unit, of whom 127 proved suitable for final study entry (Fig 2 – trial profile; S5 Appendix – volunteer baseline characteristics). Thirty-nine subjects were discharged before inoculation or exposure per protocol as described in Methods. Three separate quarantine EEs (exposure events) took place in March, April, and June 2013 involving Q1: 41 (20 Donors; 11 CR; 10 IR), Q2: 31 (12 Donors; 9 CR; 10 IR) and Q3: 55 (20 Donors; 15 CR; 20 IR) subjects respectively, with 4 Donors and 4 to 7 Recipients per exposure group. No serious adverse events were recorded in volunteers who commenced the study.

Environmental control

In Q1 relative humidity averaged 40% (Standard Deviation 9%), room temperature averaged 20.2 ^{o}C (0.4 ^{o}C) and CO_2 concentration averaged 1430ppm (110ppm). For Q2 and Q3, respectively, the corresponding values were 44% (4%), 21.4^{o}C (0.3 ^{o}C), 1810ppm (160ppm), and 57% (4%), 21.4^{o}C (0.3 ^{o}C), 1810ppm (160ppm). Outdoor CO_2 concentration proxies, taken from the average of CO_2 measurements during 2:00am-3:00am were 418, 435, and 422 ppm, for Q1, Q2, and Q3, respectively.

Donor status

Donor status is summarised by quarantine study in Table 1. Over all quarantines combined, intranasal inoculation produced an infection rate of 81% (42/52) among inoculated volunteers. Of the 42 lab-confirmed infected Donors 25

176   (60%) had ILI and 10 (24%) were classified as asymptomatic (4 in Q1, 4 in Q2, and 2

177   in Q3).

179   **Table 1. Infected Donor Status**

Q[*]	Infected/Inoculated n/n (%)	Clinical Illness n (% of Infected)			Laboratory-confirmed Infection n (% of Infected)		
		Symptomatic	Febrile	ILI	PCR-confirmed	PCR-confirmed & Seroconversion	Seroconversion by HAI : MN : Either
1	15/20 (75)	11 (73)	4 (27)	8 (53)	12 (80)	11 (73)	12 : 14 : 14
2	11/12 (92)	7 (64)	0 (0)	5 (45)	10 (91)	8 (73)	9 : 7 : 9
3	16/20 (80)	14 (88)	2 (13)	12 (75)	14 (88)	12 (75)	14 : 11 : 14
All	42/52 (81)	32 (76)	6 (14)	25 (60)	36 (86)	31 (74)	35 : 32 : 37

[*]Quarantine number; Ten Donors had greater than anticipated pre-challenge immunity: 4 in Q1, 2 in Q2, 4 in Q3.

Ten Donors had greater immunity on admission, as identified by samples

collected on day -2 (HAI>10 or MN≥80), than at their earlier screening. Four of the

10 seroconverted (i.e. had a 4-fold rise in HAI or MN titres) between admission to

quarantine and follow-up. Five of the 10 met laboratory case definition by qRT-PCR

including all four who seroconverted. The one additional qRT-PCR positive Donor

had positive swabs on study days 2 and 3 in Q2.

Virus shedding by donors

Overall, 36 Donors had nasopharyngeal swab (NPS) that tested positive by

PCR for A/WI. Of these 36: 53% (n=19) were positive on day 1 post-challenge; 94%

(34) on day 2; 97% (35) on day 3; 86% (31) on day 4; 92% (33) on day 5; and 67%

(24) on day 6 (Fig 3).

Aerosol shedding was determined for 25 Donors on day 1, 31 on day 2, 30 on

day 3, and 24 on day 4, and for a total of 36 person-days in Q1, 34 person-days in Q2,

and 40 person-days in Q3. Aerosol shedding from infected Donors, detected in

Gesundheit-II samples, is summarised in Table 2. Six (7%) of the coarse and 14

(16%) of the fine aerosol samples had detectable viral RNA. We observed aerosol

shedding from 11 (26%) of the 42 successfully infected Donors. The geometric mean

(GM) and geometric standard deviation (GSD) for coarse and fine aerosol viral RNA

copy numbers per 30-min sample were 3.1E+3 (3.3) and 5.3E+3 (4.6), respectively.

The maximum levels of shedding into coarse and fine aerosols were 2.79E+4 and

8.02E+4 RNA copies, respectively (Fig 3).

Table 2. Exhaled Breath Viral RNA Detection and Copy Number Among
Infected Donors by Quarantine Event and Aerosol Fraction

Q[*]	n Subjects	n Samples	Coarse Aerosol (>5μm)			Fine Aerosol (≤5μm)		
			Positive Subjects (%)	Positive Samples (%)	Positive Sample Mean[†]	Positive Subjects (%)	Positive Samples (%)	Positive Sample Mean[†]
1	15	27	1 (7)	1 (4)	2.79e+04	3 (20)	5 (19)	3.32e+04
2	11	30	3 (27)	3 (10)	2.16e+03	3 (27)	4 (13)	1.80e+04
3	16	32	2 (13)	2 (6)	1.73e+03	5 (31)	5 (16)	1.70e+03
All	42	89	6 (14)	6 (7)	6.31e+03	11 (26)	14 (16)	1.76e+04

[*]Quarantine number;

[†]The arithmetic mean RNA copy number used positive samples only; The geometric means (GM) and geometric

standard deviations (GSD) over all positive samples were 3.14E+3 (3.33) and 5.31E+3 (4.59) for coarse and fine

aerosol samples, respectively.

Recipient status

Recipient status is shown in Table 3. There were similar rates of symptoms in

both IR and CR groups, although more in the CR group met the study definition of

ILI, the rates were not significantly different (p = 0.23); no Recipient developed fever.

One infection was confirmed by serology (HAI increased from ≤10 to 40 and MN

increased from 10 to 320) in a CR subject who was symptomatic and whose

symptoms met the definition of ILI, but whose qRT-PCR evaluations were

persistently negative. Two other CR were transiently qRT-PCR positive but neither

met laboratory positivity criteria (Table S2). Both were asymptomatic and had no

change from baseline serology. Thus, there was only one, confirmed transmission

event. The CR and IR group SARs (2.9% and 0%) were not significantly different

(p=0.47).

Table 3. Recipient Status

Q[*]	Recipient[†]	Infected/Exposed n/n (%)	Clinical Illness n (% of Exposed)			Laboratory-confirmed Infection n (% of Exposed)		
			Symptomatic	Febrile	ILI	PCR-confirmed	PCR-confirmed & Seroconversion	Seroconversion by HAI : MN : Either
1	CR	0/11 (0)	4 (36)	0 (0)	3 (27)	0 (0)	0 (0)	0 : 0 : 0
	IR	0/10 (0)	2 (20)	0 (0)	1 (10)	0 (0)	0 (0)	0 : 0 : 0
2	CR	1/9 (11)	2 (22)	0 (0)	2 (22)	0 (0)	0 (0)	1 : 1 : 1
	IR	0/10 (0)	3 (30)	0 (0)	2 (20)	0 (0)	0 (0)	0 : 0 : 0
3	CR	0/15 (0)	6 (40)	0 (0)	4 (27)	0 (0)	0 (0)	0 : 0 : 0
	IR	0/20 (0)	6 (30)	0 (0)	2 (10)	0 (0)	0 (0)	0 : 0 : 0
All	**CR**	**1/35 (3)**	**12 (34)**	**0 (0)**	**9 (26)**	**0 (0)**	**0 (0)**	**1 : 1 : 1**
	IR	**0/40 (0)**	**11 (28)**	**0 (0)**	**5 (13)**	**0 (0)**	**0 (0)**	**0 : 0 : 0**

[*]Quarantine number; Eleven recipients had greater than anticipated pre-challenge immunity: 3 CR and 3 IR for Q1;
no Q2 Recipients; 3 CR and 2 IR for Q3. None seroconverted.
[†]CR: Control Recipient; IR: Intervention Recipient.

To compare these results with the SAR from the proof-of-concept study, we

recomputed the latter results using the current, more stringent outcome criteria. The

adjusted proof-of-concept SAR was 8.3% giving an expected SAR of 16%. The

observed SAR for the current study was not significantly different than that of the

adjusted SAR from the proof-of-concept study, but was significantly lower than the

expected doubling of the SAR (1.3% overall, p<0.0001; and 2.9% for CR, p=0.035).

Comparisons of observed and expected SAR using the proof-of-concept study

outcome definitions were also statistically significant (p<0.0001, S6 Appendix).

Data availability

238   **Data availability**

239   Data required for reproduction of analyses is available upon request. Scripts

240   and other documentation to reproduce analyses are available at Digital Repositories at

241   the University of Maryland (13) and

242   https://gitlab.com/jacobbueno/emit_quarantine_main.

Discussion

To our knowledge, this is the largest human influenza challenge-transmission study undertaken to date. We applied measures to control and standardise environmental conditions and ventilation rates within and between exposure events, to emulate as far as possible indoor winter conditions when respiratory virus spread is maximal. We particularly sought to maintain low humidity conditions which have been associated with enhanced transmission (14) and increased virus viability (15), together with a low ventilation rate to maximize recipient exposure to airborne virus. The near absence of transmission to control Recipients suggests contact and large droplet spray did not contribute substantially to transmission under the conditions used in these EEs. The significantly lower than expected SAR in this study compared with the proof-of-concept study, which had much lower ventilation rates, suggests aerosols as an important mode of influenza virus transmission in this model. The overall low SAR suggests that Donors in this model were minimally contagious and prevents definitive assessment of the modes of transmission.

Having reported an SAR of 25% (3/12) in our earlier proof-of-concept study, we expected to observe an SAR of >25%, having doubled both the duration of the exposure and the number of Donors in each quarantine (16). Indeed, the study was designed to examine an SAR of 40% in CR versus 20% in IR which would have required 125 Recipient volunteers; this was not met (n=75) and the study was underpowered.

However, the outcome criteria used in the proof-of-concept study, which included as positive a single NPS positive by qRT-PCR without seroconversion, were made more stringent in the present study by requiring two or more NPS positive by qRT-PCR in the absence of seroconversion (S6 Appendix). Applying the proof-of-

concept criteria to the current study gives an SAR of 4% (3/75) overall, while

applying the stricter criteria used in this study to the proof-of-concept study gives an

SAR of 8.3% (1/12) rather than 25%. Given the lower than planned enrolment and the

stricter outcome criteria, this study was doubly underpowered.

These observations raise two questions: 1) Why were SARs, using stringent

criteria, low in both studies and what are the implications for future human challenge-

transmission studies? 2) Why was the SAR significantly lower in the present study

compared with the expected doubling of the rate observed for the proof-of-concept?

The low SAR in these studies suggests that, unless a much greater SAR can be

achieved, type II error associated with underpowering will be a major obstacle to

successful use of human challenge-transmission studies. Potential areas to consider

addressing in order to raise the SARs in future studies include the virus used, the route

of inoculation, susceptibility of the human volunteers, the rate of viral shedding into

NPS and aerosols, and reducing ventilation of exposure rooms.

In the proof-of-concept (2009) and follow-on (2013) studies we used a GMP

A/WI challenge virus manufactured by Baxter BioScience (Vienna, Austria). Both

studies produced similar clinical and serological infection rates (typically 60-70% and

70-75%, respectively) after inoculation via nasal instillation, and similar spectrums of

clinical illness severity in Donors. These rates were higher than reported by a previous

study using lower doses of the same viral preparation (17) and consistent with rates

reported in a review of 56 challenge studies (18). The illnesses we observed were

similar to the range seen in healthy adults in the community, from asymptomatic to

febrile symptomatic infection (19). Thus, skewed illness severity does not seem to

explain the low SAR.

The virus preparation has been used in other human challenge studies with

similar rates of infection via nasal instillation (20). Using deep sequencing, Sobel

Leonard and colleagues showed that a sample of the Baxter stock "was at least

partially adapted" to the egg and/or tissue environments in which it was produced

(20). They also found that nasal instillation of the stock into human volunteers

resulted in rapid purification selection, although a fixed variant in the HA gene

remained. We have performed a BLAST search and identified the fixed variant

(G70A/D8N) in deposited sequences of wild-type H3N2 viruses. This suggests that,

on its own, the fixed HA variant is unlikely to have been a key alteration. These

results suggest that it is unlikely that the virus stock was the primary cause of the low

SARs. But, the impact of positive selection of the challenge virus for growth in the

production environment, rather than for human transmissibility, remains a potential

contributing factor to consider in choosing challenge viruses for future transmission

studies.

The route of infection with influenza virus is known to matter in the setting of

experimental infection, with aerosolized virus infectious at lower doses and more

likely to result in 'typical influenza-like disease' (fever plus cough) than intranasal

inoculation (21,22). This anisotropic property (23) of influenza virus is not unique

among respiratory viruses; e.g. it is exploited by the live, unattenuated adenovirus

vaccine (24). The implication for human challenge-transmission studies, however,

may be that increased rates of lower respiratory tract infection via aerosol inoculation

might be required to achieve sufficiently high rates of donors with fever, cough, and

contagiousness to achieve a useful SAR.

In the current quarantine-based human challenge-transmission model,

consistent with historical precedent, screening for susceptibility was undertaken

primarily by HAI antibody screening, although it is recognised that screening by MN

titre or other assays (25,26) could be an alternative or adjunctive approach. The exact

correlates of immunity and severity using novel immunological assays have not been

validated and selecting subjects based on these assays would have added substantial

complexity and costs. Six Donors and five Recipients in the present study were

discovered, in retrospect, to have seroconverted during the 3 to 56-day interval prior

to entering the quarantine facility, despite having as short a delay as possible between

final screening for HAI and quarantine entry. However, the majority of Donors and

Recipients were susceptible according to the results of microneutralization tests. Prior

immunity, as measured by the HAI and MN assays, does not therefore, appear to have

been a major limitation nor account for failure to transmit from readily infected

Donors to identically screened Recipients. Regarding future studies, as novel immune

correlates of influenza protection and severity become established, additional

approaches beyond HAI and MN assays could be employed for volunteer selection.

This might enable selection of those likely to become infected, febrile, and have

greater symptomatology including more frequent and greater levels of cough and

runny nose. Unfortunately, such screening might also dramatically reduce the yield of

suitable volunteers and substantially increase overall study costs.

Results from serial nasopharyngeal swabs in Donors indicate that over 80%

were positive by qRT-PCR testing on one or more post-challenge days. Viral load

detected by swabs was substantial with qRT-PCR Ct values in the mid 20s on days

two and three (Fig 3). Thus, failure to shed virus into nasal secretions cannot explain

the low SARs. Minimal transmission despite days of prolonged exposure to Donors

shedding virus into nasal secretions provides strong evidence that contact and large

droplet transmission are not important in this model.

The results from breath sampling with the Gesundheit-II device indicate that

26% (11/42) of infected Donors had virus detectable in exhaled air during the same

period. By comparison, virus shedding into exhaled breath was detected in 84% (119

of 142) of influenza cases selected on the basis of having fever or a positive rapid test

and sampled on one to three days post onset of symptoms, mostly recruited from

young adults on a college campus (7). When compared on a per-sample basis,

infected Donors shed detectable virus less frequently than naturally infected college

campus cases (7) in both coarse (7% and 40%, respectively) and fine aerosols (15%

and 76%); all assays for both groups were performed in the same laboratory using the

same methods. However, when the comparison was limited to positive aerosol

samples from each study population, the average quantities of virus detected were

similar (within 1 log), for the Donors as for the college community cases (GM coarse

3.1E+3 and 1.2E+4, GM fine 5.3E+3 and 3.8E+4, respectively). The maximum

exhaled breath viral aerosol from the 11 Donors was two to four logs lower than from

the college campus cases selected for having fever or a high viral load in the NPS

(maximum coarse 2.8E+4 and 4.3E+8, and maximum fine 8.0E+4 and 4.4E+7,

respectively) (7). While this difference may merely represent the low probability of

sampling from the tail of a log-normal distribution with only 11, as compared with

119 cases, it may be relevant to the low SAR in the challenge-transmission model if

aerosols disseminated by more symptomatic individuals, such as the selected

community cases, and rare supershedders account for most transmission. If aerosols

are largely derived from the lower respiratory tract, as has been suggested by analysis

of the college community cases, this would also suggest that future challenge-

transmission studies should employ methods designed to increase the frequency of

lower respiratory tract infection.

The proof-of-concept study was conducted in a hotel room with closed

windows and thermal control provided only by a recirculating air conditioning unit.

While the ventilation rate was not measured, it was likely to have been extremely low

for the number of occupants, with only small, intermittent, bathroom extract and

natural building infiltration providing fresh air. The ventilation rate of 4 l/s/person

during the main study was low compared to 10 l/s/person recommended in UK design

standards (27) but was likely substantially greater than in the proof-of-concept study.

This would have produced significantly higher viral aerosol concentrations during the

proof-of-concept EEs, assuming similar generation rates from Donors in both

experiments. Given that the Donors in the two studies were similar in other respects,

differences in shedding rates seem unlikely. Therefore, the difference in SAR between

this study and the expected SAR based on design changes and prior results are

possibly due to differing ventilation conditions. The implication for future challenge-

transmission studies, given that the ventilation rate in the current study was as low as

possible with a single pass ventilation system, is that recirculating air conditioning

systems similar to that in the proof-of-concept study should be employed to limit

dilution ventilation and maximize exposure to aerosols. This will be especially

important if Donors in future studies continue to represent the lower end of the

aerosol shedding spectrum seen in naturally infected cases.

Achieving temperature and humidity to simulate winter conditions was

challenging, particularly in Q3, conducted in June 2013 when the average external

conditions were 16^{o}C and 64% relative humidity. It was necessary to strike a balance

between volunteer comfort and conditions favourable to transmission, both of which

were constrained by the capability of the mechanical systems in the building.

However, the relative humidity in the current study overlapped with that during the

proof-of-concept study, which ranged between 38 and 53%, and thus, eliminated

humidity as a potential explanation for the difference in transmission rates.

Despite this study not having produced the planned SAR, it yields important

findings. First, although fewer viral challenged subjects had virus-laden aerosols than

seen in people with natural infections presenting with influenza-like symptoms, those

volunteers who did produce viral aerosols did so at a rate similar to the average

symptomatic naturally infected case. Second, given that a subset of the infected

volunteers had moderate viral aerosol shedding in this model, observation of

transmission via aerosols in quarantine studies may be strongly dependent on the

dilution ventilation rate. Third, low risk of transmission to Control Recipients

suggests that contact and large droplet spray transmission were not important modes

of transmission in this model. The overall low SAR compared to that observed in the

proof-of-concept study suggests that, given the main difference between the studies

was the indoor air ventilation rate, aerosol transmission may be an important mode of

influenza virus transmission between adults. Finally, sensitivity of transmission to

details of the Donors selected, environment, and activity during exposure events,

suggest that if a successful transmission model can be developed, carefully designed

studies may be useful for investigating specific, targeted intervention strategies for

prevention of specific transmission modalities. However, sensitivity to experimental

conditions also demonstrates that it will be challenging to generalize the results of the

quarantine-based transmission model to broad conclusions about the relative

importance of aerosol, droplet spray, and contact modes of transmission. These

complexities of the challenge-transmission model suggest that community-based

transmission studies employing deep-sequencing based molecular epidemiologic

methods in natural experiments, e.g. comparing high and low ventilation dormitories

or barracks, may be more attractive alternatives than previously thought.

Unfortunately, although an important role for aerosols in transmission of influenza, at

least between adults, is hinted at when comparing the proof-of-concept and current

studies, this challenge model cannot provide a definitive answer to the importance of

this mode for influenza virus transmission between humans.

Materials and methods

Ethics statement

The randomized challenge-transmission trial took place from March to June

2013 in a closed quarantine facility, with written informed consent from healthy

volunteers in accordance with the principles of the Declaration of Helsinki, in

compliance with UK regulatory and ethical (IRB) requirements (under auspices of the

UK Health Research Authority (HRA) National Research Ethics Service (NRES)

Committee London – City & East; reference number 12/LO/1277), and registered

with ClinicalTrials.gov (number NCT01710111).

Overview

Volunteers, screened for serologic susceptibility, were randomly selected for

intranasal challenge with A/WI (16) – becoming 'Donors'. After allowing for a short

incubation period, Donors were introduced to other sero-susceptible volunteers –

'Recipients' – under controlled household-like conditions for four days. Recipients

were randomised as Intervention Recipients (IR) or Control Recipients (CR). IRs

wore face shields evaluated to interrupt large droplet transmission but to be

permissive to aerosols (S3 Appendix); in addition, IRs hand sanitised (using alcohol-

based Deb® InstantFOAM, 72% ethyl alcohol) once every 15 minutes to minimise

the possibility of contact transmission. IRs were only allowed to touch face via single-

use wooden spatulas. Thus, IRs would be exposed to influenza only via aerosols. CRs

did not wear face shields or use hand sanitiser and were allowed to touch face freely;

therefore, CRs would have been exposed via all routes of transmission consistent with

close proximity human-human contact. Fig 1 shows an overview of the study design.

Influenza Virus

Influenza A/WI manufactured and processed under current good

manufacturing practices (cGMP) was obtained from Baxter BioScience, (Vienna,

Austria). Stocks of this virus preparation have been sequenced and its evolution in the

upper respiratory track of inoculated volunteers extensively analysed (20).

Screening

Volunteers were screened from 3-56 days in advance of the experiment to

determine humoral immunity to A/WI before undergoing further screening against

inclusion and exclusion criteria (S4 Appendix). Volunteers needed to be healthy,

between the ages of 18 and 45 years, not living with anyone deemed at high risk of

influenza complications on discharge, and not to have had a seasonal influenza

vaccine in the last 3 years. Blood samples from volunteers were collected

immediately before quarantine entry for repeat serology, although results were not

available until after the study. An initial screening haemagglutination inhibition assay

(HAI) titre of ≤10 was considered evidence of susceptibility to infection.

Power calculation

We calculated, based on the reported SAR of the proof-of-concept study (16)

(S4 Appendix), that over a range of scenarios the statistical power of the whole study

would be 63%-84%, typically 80% in the most realistic scenario, if a sample size of

125 Recipients was achieved (70 IR, 55 CR). To increase the SAR from that observed

in the proof-of-concept study, we opted to inoculate 4 (rather than 2) Donors per 5

Recipients and to conduct exposure events on days 1-4 (rather than 2-3) post

inoculation.

Pre-exposure

Screened, eligible, volunteers entered a closed, quarantine unit on Day -2 and

were randomised to Donor or Recipient (IR and CR) groups; thereafter Donors and

Recipients were segregated. Donors and Recipients were immediately screened for a

panel of 7 imported 'contaminant' respiratory viruses (influenza A, B; adenovirus;

respiratory syncytial virus; parainfluenza 1, 2, 3) by a direct fluorescence antibody

assay (DFA) (LIGHT DIAGNOSTICS™ SimulFluor1 Respiratory Screen, Merck

Millipore) and any with a positive test were immediately discharged. On day 0,

Donors were inoculated intranasally, via pipette in a supine position, with 0.5ml per

nostril of a suspension containing 5.5 $\log_{10}TCID_{50}$/ml of influenza A/WI (17,28).

Exposure Events

The study was conducted in three, separate, identically-designed quarantine

events (Q1, Q2, and Q3). From Day 1 to Day 4 of each quarantine event, all

volunteers took part in an Exposure Event (EE). Individual Donors and Recipients

were each allocated to a single exposure room per day where they interacted at close

distances for approximately 15 hours/day, for four consecutive days. In-room staff

supervised activities such as playing board games, pool, and table football, and

watching films, whilst ensuring that volunteers mixed freely, and that IRs complied

with face shield use, hand sanitisation, and no-touch-face rules. Donor, IR and CR

groups were moved into different corners of the rooms for meal breaks, and Donor

and Recipient groups were housed separately at night, including further separation

and withdrawal of any Recipients with symptoms to prevent any contamination of the

results by Recipient-Recipient secondary transmission. Five exposure rooms were

used ranging from 17-30m^2 floor area and 50-87m^3 volume. Four Donors were non-

randomly allocated to each exposure group to ensure even distribution of subjects

actively shedding virus. This was achieved by assessing symptom scores and the

results of influenza rapid tests (Quidel Sofia®) performed on Days 1 and 2. From Day

2 onwards, Donors remained in their allocated group and were not redistributed

further. Once assigned to an exposure group, Recipients remained in the same group

until the end of the EE or until they developed ILI and were withdrawn to a separate

isolation area. On each day of EE, each exposure group rotated to a different exposure

room.

Environmental controls

Each exposure room was assessed pre-quarantine by building and ventilation

engineers and modified to achieve a ventilation rate of approximately

4L/second/person (based on planned occupancy during EEs), temperature 18-22^{o}C,

and relative humidity 45-65%, to produce conditions favourable to influenza

transmission (14), balanced against tolerability for occupants, and the capability of

building systems to provide controlled environments comparable across all three

quarantine studies. During each EE, rooms were monitored at 5-minute intervals for

CO_2 concentration (as a proxy for ventilation rate), temperature and humidity;

heating, cooling and humidity were remotely adjusted to maintain optimal conditions.

Clinical assessment

All subjects underwent thrice daily monitoring of respiratory and systemic

symptoms; each symptom was reported as grade 0 (not present) to 3 (severe). Paired

venous blood specimens for serology were taken on Day -2 and Day 28. NPS were

taken daily from all subjects. Respiratory specimens were analysed by quantitative

reverse-transcriptase PCR (qRT-PCR) and serological specimens by HAI and

microneutralisation (MN) assays. The qRT-PCR and HAI were performed in

duplicate at the MRC University of Glasgow Centre for Virus Research (with Fast

Track Diagnostics qRT-PCR kit) and the U.S. Centers for Disease Control and

Prevention (CDC), Atlanta; MN assays were performed by CDC as described

previously (29).

Exhaled breath sampling

Donors were assigned to provide exhaled breath samples on two, randomly

selected days within the exposure event, collected using a Gesundheit-II cone

collection apparatus allowing for fractionation of particle sizes into 'fine' <5μm and

'coarse' ≥5μm aerosol samples (8,30). Each sample collection session lasted for 30

minutes. Breath samples were concentrated, extracted, stored at -80°C, and evaluated

by qRT-PCR using protocols and materials specified by the CDC with RNA copy

number estimated as previously described (7).

Discharge

After completion of the EE, Donors were discharged on active treatment with

oseltamivir (75mg b.i.d. 5 days), whereas Recipients were observed for 7 days, then

discharged with oseltamivir on day 8. All volunteers attended for 28-day (+/-3 days)

post virus exposure outpatient follow-up and study dismissal.

Outcome Definitions

Respiratory symptoms were defined as self-reported grade ≥1 of runny nose,

stuffy nose, sneeze, sore throat, cough, or shortness of breath 'lasting ≥24 hours' (S4

Appendix). Fever was defined as temperature >37.9 °C. Symptomatic was defined as

evidence of any respiratory symptom lasting ≥24 hours during study days 1-6.

Influenza-like Illness (ILI) was defined as an illness >24 hours duration with: either

fever and at least one respiratory symptom; or two or more symptoms of grade ≥1,

one of which must have been respiratory; eligible non-respiratory symptoms were

headache, muscle/joint ache, and malaise; lasting ≥24 hours means report of a

respiratory symptom during 3/3 observations within a single day, or at least once per

day over two consecutive days. Laboratory confirmed infection was defined as: a 4-

fold or greater rise in HAI or MN titres between Day -2 (baseline) and Day 28; or two

or more positive NPS test results by qRT-PCR. These differed from the proof-of-

concept study, which used seroconversion or a single positive nasal wash (S6

Appendix).

Statistics

Comparisons were made between groups using Fisher's exact tests for

binomial proportions and between observed and expected outcomes using binomial

tests. Tests were two-tailed. All data to reproduce analyses is available from the

EMIT Consortium upon request.

Acknowledgements

The authors thank Blanca Beato Arribas and William Booth of BSRIA Limited, Chang Chen, Sheryl Ehrman, Neil Ferguson, Lisa Grohskopf, F. Liaini Gross, Andrew Hayward, Ashley Kang, Jacqueline Katz, John Oxford, Massimo Palmarini, Walt Adamson, Jennifer Wang, and the hVivo clinical team for their advice and contributions to the study. The authors are grateful to other Scientific Advisory Board members: Allan Bennett, Ben Cowling, and Arnold Monto.

References

1. Brankston G, Gitterman L, Hirji Z, Lemieux C, Gardam M. Transmission of influenza A in
human beings. Lancet Infect Dis. 2007 Apr;7(4):257–65.

2. Killingley B, Nguyen-Van-Tam J. Routes of influenza transmission. Influenza and Other
Respiratory Viruses. 2013 Sep;7:42–51.

3. Tellier R. Review of aerosol transmission of influenza A virus. Emerging Infect Dis. 2006
Nov;12(11):1657–62.

4. Tellier R. Aerosol transmission of influenza A virus: a review of new studies. J R Soc
Interface. 2009 Dec 6;6 Suppl 6:S783-790.

5. Cowling BJ, Ip DKM, Fang VJ, Suntarattiwong P, Olsen SJ, Levy J, et al. Aerosol
transmission is an important mode of influenza A virus spread. Nat Commun.
2013;4(1):1935.

6. Kormuth KA, Lin K, Prussin AJ, Vejerano EP, Tiwari AJ, Cox SS, et al. Influenza Virus
Infectivity Is Retained in Aerosols and Droplets Independent of Relative Humidity. J
Infect Dis. 2018 Jul 24;218(5):739–47.

7. Yan J, Grantham M, Pantelic J, Bueno de Mesquita PJ, Albert B, Liu F, et al. Infectious
virus in exhaled breath of symptomatic seasonal influenza cases from a college
community. Proc Natl Acad Sci USA. 2018 Jan 30;115(5):1081–6.

8. Milton DK, Fabian MP, Cowling BJ, Grantham ML, McDevitt JJ. Influenza Virus
Aerosols in Human Exhaled Breath: Particle Size, Culturability, and Effect of Surgical
Masks. Fouchier RAM, editor. PLoS Pathogens. 2013 Mar 7;9(3):e1003205.

9. Makison Booth C, Clayton M, Crook B, Gawn JM. Effectiveness of surgical masks against
influenza bioaerosols. Journal of Hospital Infection. 2013 May 1;84(1):22–6.

10. Bischoff WE, Swett K, Leng I, Peters TR. Exposure to Influenza Virus Aerosols During
Routine Patient Care. J Infect Dis. 2013 Apr 1;207(7):1037–46.

11. Killingley B, Enstone J, Booy R, Hayward A, Oxford J, Ferguson N, et al. Potential role
of human challenge studies for investigation of influenza transmission. Lancet Infect
Dis. 2011 Nov;11(11):879–86.

12. Snider D, Bridges C, Weissman D. Meeting summary of the workshop 'Approaches to
better understand human influenza transmission''.' In Tom Harkin Global
Communications Center, Centers for Disease Control and Prevention, Atlanta, Georgia:
Centers for Disease Control and Prevention; 2010. Available from:
https://www.cdc.gov/influenzatransmissionworkshop2010/pdf/Influenza_Transmission_
Workshop_Summary_508.pdf

13. Bueno de Mesquita PJ, Jacob P. Code to reproduce analysis of manuscript: Minimal
transmission in an influenza A (H3N2) human challenge-transmission model with
exposure events in a controlled environment. 2019 Dec 10 [cited 2019 Dec 10];
Available from: http://drum.lib.umd.edu/handle/1903/25315

14. Lowen AC, Mubareka S, Steel J, Palese P. Influenza virus transmission is dependent on
relative humidity and temperature. PLoS Pathog. 2007 Oct 19;3(10):1470–6.

15. Yang W, Elankumaran S, Marr LC. Relationship between Humidity and Influenza A
Viability in Droplets and Implications for Influenza's Seasonality. PLOS ONE. 2012
Oct 3;7(10):e46789.

16. Killingley B, Enstone JE, Greatorex J, Gilbert AS, Lambkin-Williams R, Cauchemez S,
et al. Use of a human influenza challenge model to assess person-to-person
transmission: proof-of-concept study. J Infect Dis. 2012 Jan 1;205(1):35–43.

17. Zaas AK, Chen M, Varkey J, Veldman T, Hero AO, Lucas J, et al. Gene expression
signatures diagnose influenza and other symptomatic respiratory viral infections in
humans. Cell Host Microbe. 2009 Sep 17;6(3):207–17.

18. Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, Leach S, et al. Time lines
of infection and disease in human influenza: a review of volunteer challenge studies.
Am J Epidemiol. 2008 Apr 1;167(7):775–85.

19. Hayward AC, Fragaszy EB, Bermingham A, Wang L, Copas A, Edmunds WJ, et al.
Comparative community burden and severity of seasonal and pandemic influenza:
results of the Flu Watch cohort study. Lancet Respir Med. 2014 Jun;2(6):445–54.

20. Sobel Leonard A, McClain MT, Smith GJD, Wentworth DE, Halpin RA, Lin X, et al.
Deep Sequencing of Influenza A Virus from a Human Challenge Study Reveals a
Selective Bottleneck and Only Limited Intrahost Genetic Diversification. J Virol. 2016
Dec 15;90(24):11247–58.

21. Alford RH, Kasel JA, Gerone PJ, Knight V. Human influenza resulting from aerosol
inhalation. Proc Soc Exp Biol Med. 1966 Jul;122(3):800–4.

22. Henle W, Henle G, Stokes J, Maris EP. Experimental Exposure of Human Subjects to
Viruses of Influenza. J Immunol. 1946;52(2):145–65.

23. Milton DK. What was the primary mode of smallpox transmission? Implications for
biodefense. Front Cell Infect Microbiol. 2012;2:150.

24. Couch RB, Chanock RM, Cate TR, Lang DJ, Knight V, Huebner RJ. Immunization with
types 4 and 7 adenovirus by selective infection of the intestinal tract. Am Rev Respir
Dis. 1963 Sep;88:SUPPL 394-403.

25. Park J-K, Han A, Czajkowski L, Reed S, Athota R, Bristol T, et al. Evaluation of
Preexisting Anti-Hemagglutinin Stalk Antibody as a Correlate of Protection in a
Healthy Volunteer Challenge with Influenza A/H1N1pdm Virus. MBio. 2018 Jan
23;9(1):e02284-17.

26. Tang JW. Pre-existing immunity in human challenge studies of influenza transmission.
The Lancet Infectious Diseases. 2012 Oct 1;12(10):744.

27. Chartered Institution of Building Services Engineers. GVA/15 CIBSE Guide A:
Environmental Design 2015 [Internet]. London: CIBSE; 2015 [cited 2018 Oct 12]. 402
p. Available from: https://www.cibse.org/knowledge/knowledge-
items/detail?id=a0q20000008I79JAAS

28. Zaas AK, Burke T, Chen M, McClain M, Nicholson B, Veldman T, et al. A host-based
RT-PCR gene expression signature to identify acute respiratory viral infection. Sci
Transl Med. 2013 Sep 18;5(203):203ra126.

29. World Health Organization, editor. Manual for the laboratory diagnosis and virological
surveillance of influenza. Geneva: World Health Organization; 2011. 139 p.

30. McDevitt JJ, Koutrakis P, Ferguson ST, Wolfson JM, Fabian MP, Martins M, et al.
Development and Performance Evaluation of an Exhaled-Breath Bioaerosol Collector
for Influenza Virus. Aerosol Sci Technol. 2013 Jan 1;47(4):444–51.

Figure legends

Fig 1. Schematic of study design showing timelines, environmental controls and monitoring, physical segregation arrangements, exposure intervention, and volunteer movements during quarantine study. DFA: direct fluorescence assay; RH: relative humidity; NPS: nasopharyngeal swab.

Fig 2. Trial profile. Intervention Recipients: wore face shields, used hand sanitizer every 15 min and only allowed to touch face with single-use wooden spatula; Control Recipients: did not use face shields or the specified hand hygiene protocol.

Fig 3. Viral detection in Donors by day of exposure event. A) Columns show the proportion of all infected donors (n=42) who were qRT-PCR positive for viral shedding for coarse (>5µm) and fine (≤5µm) aerosols, and nasopharyngeal swabs. B) Mean and standard deviation error bars for qRT-PCR cycle threshold values from the positive nasopharyngeal swabs (n=19 on day 1; n=34 on day 2; n=35 on day 3; n=31 on day 4). C) Virus quantified from detectable exhaled coarse (n=6) and fine (n=14) breath aerosols by qRT-PCR; the boxes show the inner-quartile range (IQR) with a band to indicate the median, and whiskers extending to highest and lowest data points within 1.5 IQR.

Supporting information captions

S1 Appendix. EMIT Consortium Team Members

S2 Appendix. Role of Funding Source

S3 Appendix. Efficacy of a Face Shield to Reduce Transmission of Influenza

Virus in Large Droplets. Contains Figs. S1-S5.

S4 Appendix. Full Inclusion and Exclusion Criteria, Power Calculation, and

Sample Handling

S5 Appendix. Baseline Characteristics. Contains Table S1.

S6 Appendix. Comparing the Current Study and Proof-of-Concept Study Using

Current Outcome Criteria and Applying Infection Criteria from The Proof of

Concept Study. Contains Tables S2-S4.