Comparison of first trimester dating methods for gestational age estimation and

their implication on preterm birth classification in a North Indian cohort

Ramya Vijayram[1,2]*, Nikhita Damaraju[1,2]*, Ashley Xavier[1,2]*, Bapu Koundinya Desiraju[3,4],

Ramachandran Thiruvengadam[3,4], Sumit Misra[3,4], Shilpa Chopra[3,4], Ashok Khurana[5], Nitya

Wadhwa[3,4], GARBH-Ini Study Group[4], Raghunathan Rengaswamy[2,6,7], Himanshu Sinha[1,2,7]†,

Shinjini Bhatnagar[3,4]†

[1] Department of Biotechnology, Bhupat and Jyoti Mehta School of Biosciences, Indian

Institute of Technology Madras, Chennai, India

[2] Initiative for Biological Systems Engineering, Indian Institute of Technology Madras,

Chennai, India

[3] Maternal and Child Health Program, Translational Health Science and Technology Institute,

Faridabad, India

[4] Interdisciplinary Group for Advanced Research on Birth Outcomes - DBT India Initiative,

Translational Health Science and Technology Institute, Faridabad, India

[5] The Ultrasound Lab, Defence Colony, New Delhi, India

[6] Department of Chemical Engineering, Indian Institute of Technology Madras, Chennai,

India

[7] Robert Bosch Centre for Data Science and Artificial Intelligence, Indian Institute of

Technology Madras, Chennai, India

* Ramya Vijayram, Nikhita Damaraju, and Ashley Xavier are joint first authors.

† Correspondence: Shinjini Bhatnagar, Maternal and Child Health Program, Translational

Health Science and Technology Institute, Faridabad, India shinjini.bhatnagar@thsti.res.in;

Himanshu Sinha, Department of Biotechnology, Bhupat and Jyoti Mehta School of

Biosciences, Indian Institute of Technology Madras, Chennai, India sinha@iitm.ac.in

Abstract

Background: Different formulae have been developed globally to estimate gestational age

(GA) by ultrasonography in the first trimester of pregnancy. In this study, we develop an

Indian population-specific dating formula and compare its performance with published

formulae. Finally, we evaluate the implications of the choice of dating method on preterm

birth (PTB) rate. This study's data was from GARBH-Ini, an ongoing pregnancy cohort of

North Indian women to study PTB.

Methods: Comparisons between ultrasonography-Hadlock and last menstrual period (LMP)

based dating methods were made by studying the distribution of their differences by Bland-

Altman analysis. Using data-driven approaches, we removed data outliers more efficiently

than by applying clinical parameters. We applied advanced machine learning algorithms to

identify relevant features for GA estimation and developed an Indian population-specific

formula (Garbhini-GA1) for the first trimester. PTB rates of Garbhini-GA1 and other

formulae were compared by estimating sensitivity and accuracy.

Results: Performance of Garbhini-GA1 formula, a non-linear function of crown-rump length

(CRL), was equivalent to published formulae for estimation of first trimester GA (LoA, -

0.46,0.96 weeks). We found that CRL was the most crucial parameter in estimating GA and

no other clinical or socioeconomic covariates contributed to GA estimation. The estimated

PTB rate across all the formulae including LMP ranged 11.27 – 16.50% with Garbhini-GA1

estimating the least rate with highest sensitivity and accuracy. While the LMP-based

method overestimated GA by three days compared to USG-Hadlock formula; at an

individual level, these methods had less than 50% agreement in the classification of PTB.

Conclusions: An accurate estimation of GA is crucial for the management of PTB. Garbhini-

GA1, the first such formula developed in an Indian setting, estimates PTB rates with higher

accuracy, especially when compared to commonly used Hadlock formula. Our results

reinforce the need to develop population-specific gestational age formulae.

Keywords

55   Gestational age; Crown-rump length; CRL; Preterm birth; Last menstrual period; GARBH-Ini;

56   Machine learning

Background

Preterm birth (PTB) is conventionally defined as a birth that occurs before 37 completed weeks of gestation [1,2]. Globally, complications arising from preterm birth were the leading cause of child (less than 5 years of age) mortality in 2016, accounting for 35% of neonatal deaths [3]. PTB is a unique disease in the way it is defined by the duration of gestation and not by a pathological process. The duration of gestation is the period between the date of conception and date of delivery. While the date of delivery can be documented with fair accuracy, ascertaining the date of conception is challenging. The estimation of gestational age (GA) during the antenatal period also called as the dating of pregnancy has been conventionally done using the first day of the recall-based last menstrual period (LMP) or measurement of foetal biometry by ultrasonography (USG) [4,5]. Each of these methods poses a unique set of challenges. The accuracy of dating by LMP method is dependent on accurate recall, and regularity of menstrual cycle [4,6] which, is affected by numerous physiological and pathological conditions such as obesity [7], polycystic ovarian syndrome [8], breastfeeding [9] and use of contraceptive methods [10].

The USG method is based on foetal biometry using crown-rump length (CRL) in the first trimester. Several formulae exist to estimate GA using CRL, including Hadlock formula [11], based on a US population-based study widely used in India [12]. However, the choice of dating formula might influence dating accuracy, as these formulae have been developed from studies that differed both in the study population and study design [13]. The error and bias due to the choice of dating formula need to be quantitatively studied to estimate the rate of PTB in a specific population [14]. In addition to its public health importance, accurate

dating is essential for clinical decision making during the antenatal period, such as

scheduling monitoring visits and recommending appropriate antenatal care [4].

This study first quantified the discrepancy between LMP and USG-based (Hadlock) dating

methods during the first trimester in an Indian population. We characterised how each

method could contribute to the discrepancy in calculating the GA. We then built a

population-specific model from the GARBH-Ini cohort (Interdisciplinary Group for Advanced

Research on BirtH outcomes - DBT India Initiative), Garbhini-GA1, and compared its

performance with the published 'high quality' formulae for the first-trimester dating [13] –

McLennan and Schluter [15], Robinson and Fleming [16], Sahota [17] and Verburg [18],

INTERGROWTH-21st [19], and Hadlock's formula [11] (Table S1). Finally, we quantified the

implications of the choice of dating methods on PTB rates in our study population.

Methods

Methods

Study design

GARBH-Ini is a collaborative program, initiated by Translational Health Science and

Technology Institute, Faridabad with partners from Regional Centre of Biotechnology,

Faridabad; National Institute of Biomedical Genomics, Kalyani; Civil Hospital, Gurugram;

Safdarjung hospital, New Delhi. The GARBH-Ini cohort is a prospective observational cohort

of pregnant women initiated in May 2015 at the District Civil Hospital that serves a mostly

rural and semi-urban population in the Gurugram district, Haryana, India. The cohort study's

objective is to develop an effective risk stratification that facilitates timely referral for

women at high risk of PTB, particularly in low- and middle-income countries. Women in the

GARBH-Ini cohort are enrolled within 20 weeks of gestation and are followed three times

during pregnancy till delivery and once postpartum [20]. After a verbal consent to be

interviewed, informed consent for screening is obtained for women at < 20-weeks of

gestational age (GA) calculated by the last menstrual period. A dating ultrasound is

performed within the week to confirm a viable intrauterine pregnancy with

< 20-weeks GA using standard foetal biometric parameters. A time-series data on a large set

of clinical and socioeconomic variables are collected across pregnancy to help stratify

women into defined risk groups for PTB. The dating ultrasound is performed by a qualified

radiologist specifically trained in the study protocol. The clinical and demographic

information is collected by trained, dedicated research staff under medically qualified

research officers' supervision. The data acquisition protocols and quality control measures

are detailed elsewhere [20].

Sampling strategy and participant datasets derived for the study

This analysis's samples were derived from the first 3499 participants enrolled in the GARBH-

Ini study (between May 2015 to November 2017). We included 1721 participants (N_p =

1721) who had information on the LMP, CRL and had singleton pregnancy which advanced

beyond 20 weeks of gestation, i.e. the pregnancy did not end in a spontaneous abortion. If

more than one scan was performed < 14 weeks, data from both the scans were included as

unique observations (N_o). Therefore, 1721 participants contributed a total of 2562

observations (N_o = 2562) that was used for further analyses, and this dataset of

observations was termed as the TRAINING DATASET (Figure 1). This dataset was used to

develop a population-based dating model named Garbhini-GA1, for the first trimester.

It is essential to independently evaluate models on data that was not used for building the

model in order to eliminate any biases that may have been incorporated due to the iterative

learning process of the model building dataset and estimate the expected performance

when applying the model on new data in the real world. We used an unseen TEST DATASET

created from 999 participants enrolled after the initial set of 3499 participants in this cohort

(Figure 1). The TEST DATASET was obtained by applying identical processing steps as

described for the TRAINING DATASET (N_o = 808 from N_p = 559; Figure 1).

Assessment of LMP and CRL

The date of LMP was ascertained from the participant's recall of the first day of the last

menstrual period. CRL from an ultrasound image (GE Voluson E8 Expert, General Electric

Healthcare, Chicago, USA) was captured in the midline sagittal section of the whole foetus

by placing the callipers on the outer margin skin borders of the foetal crown and rump. The

CRL measurement was done thrice on three different ultrasound images, and the average of

the three measurements was considered for estimation of CRL-based GA. Under the

supervision of medically qualified researchers, study nurses documented the clinical and

sociodemographic characteristics [20].

Development and validation of the population-specific gestational dating model

The gold standard or ground truth for development of first-trimester dating model was

derived from a subset of participants with the most reliable GA based on last menstrual

period. We used two approaches to create subsets from the TRAINING DATASET for

developing the first-trimester population-based dating formula. The first approach excluded

participants with potentially unreliable LMP or high risk of foetal growth restriction, giving

us the CLINICALLY-FILTERED DATASET (N_o = 980 from N_p = 650; Figure 1, Table S2).

The second approach used Density-Based Spatial Clustering of Applications with Noise

(DBSCAN) method to remove outliers based on noise in the data points. DBSCAN identifies

noise by classifying points into clusters if there are a sufficient number of neighbours that lie

within a specified Euclidean distance or if the point is adjacent to another data point

meeting the criteria [21]. DBSCAN was used to identify and remove outliers in the TRAINING

DATASET using the parameters for distance cut-off (*epsilon, eps*) 0.5 and the minimum

number of neighbours (*minpoints*) 20. A range of values for eps and *minpoints* did not

markedly change the clustering result (Table S3). The resulting dataset that retained reliable

data points for the analysis was termed as the DBSCAN DATASET (N_o = 2156 from N_p = 1476;

Figure 1).

The use of CRL for dating of pregnancy is restricted to the first trimester of pregnancy in

clinical practice. This is because of the technical difficulties in obtaining accurate CRL

measurements beyond this period. The same was practised in the GARBH-Ini cohort as it is

an observational study. When an ultrasonographic examination was performed during early

pregnancy, the radiologist refrained from measuring CRL if she/he was not assured of its

accuracy. Instead, the radiologist measured the other foetal biometry (biparietal diameters,

abdominal and head circumference and femur length to ascertain the gestational age). This

resulted in a dataset with GA by CRL truncated at 14 weeks of gestation. When used for

training models, such a truncated dataset may lead to inaccuracies in the model fitting

particularly at the margins of the distribution around 14 weeks [22]. We considered multiple

approaches used in the literature [22] and overcame this by supplementing our dataset with

simulated observations from the Hadlock dataset, which measured the relationship

between CRL and GA in the range of 15 – 18 weeks [11]. This supplemented dataset was

used to build fractional polynomial models of GA as a function of CRL (see Figure S1, Table

S7).

Development of a first trimester dating formula was done by fitting fractional polynomial

regression models of GA (weeks) as a function of CRL (cm) on CLINICALLY-FILTERED and

DBSCAN datasets. The performance of the chosen formula was validated in the TEST

DATASET.

In addition to CRL as a primary indicator, a list of 282 candidate variables was explored by

feature selection methods on the DBSCAN DATASET to identify other variables which may

be predictive of GA during the first trimester. These methods helped to find uncorrelated,

non-redundant features that might improve GA prediction accuracy (Table S4). First, the

feature selection was done using Boruta [23], a random forest classifier, which identified six

features and second, by implementing Generalised Linear Modelling (GLM) that identified

two candidate predictors of GA. A union of these features (Table S5), gave a list of six

candidate predictors. Equations were generated using all combinations of these predictors

in the form of linear, logarithmic, polynomial and fractional power equations. The best fit

model was termed Garbhini-GA1 formula and was validated for its performance in the TEST

DATASET.

Comparison of LMP- and USG-based dating methods during the first-trimester

We calculated the difference between LMP- and USG-based GA for each participant and

studied the distribution of the differences by Bland-Altman (BA) analysis [24]. Additionally,

we estimated the effect of factors that could contribute to the discrepancy between GA by

LMP and ultrasound. This may be due to an unreliable LMP or foetal growth restriction. We

repeated the comparative analysis in our population's subsets with accurate LMP and no

risk factors for foetal growth restriction (see Additional File 1).

The mean difference between the methods and the limits of agreement (LoA) for 95% CI

were reported. The PTB rates with LMP- and USG-based methods were reported per 100

live births with 95% CI. We compared different USG-based formulae using correlation

analysis.

The data analyses were carried out in R versions 3.6.1 and 3.5.0. DBSCAN was implemented

using the package *dbscan*, and the random forests feature selection was performed using

the *Boruta* package [23]. Statistical analysis for comparing PTB rate as estimated using

different dating formulae was carried out using standard t-test with or without Bonferroni

multiple testing correction or using Fisher's Exact test wherever appropriate.

Results

Description of participants included in the study

The median age of the participants enrolled in the cohort was 23.0 years (IQR 21.0 – 26.0),

with the median weight and height as 47.0 kg (IQR 42.5 – 53.3) and 153.0 cm (IQR 149.2 –

156.8), respectively and with 59.93% of the participants having a normal first trimester BMI

(median 20.09, IQR 18.27 – 22.59). Almost half of them were primigravida. Most of the

participants (98.20%) were from middle or lower socioeconomic strata [25]. The

participants selected for this analysis had a median GA of 11.71 weeks (IQR 9.29 – 13.0). The

other baseline characteristics are given in Table 1.

Comparison of USG-Hadlock and LMP-based methods for estimation of GA in the first

trimester

The mean difference between USG-Hadlock and LMP-based dating at the time of enrolment

was found to be -0.44 ± 2.02 weeks (Figure 2a) indicating that the LMP-based method

overestimated GA by nearly three days. The LoA determined by BA analysis was -4.39, 3.51

weeks, with 8.82% of participants falling beyond these limits (Figure 2b) suggesting a high

imprecision in both the methods. The LoA between USG-Hadlock and LMP-based dating

marginally narrowed when tested on participants with reliable LMP (LoA -4.22, 3.28) or

those with low-risk of foetal growth restriction (LoA -4.13, 3.21). The wide LoA that

persisted despite ensuring reliable LMP and standardised CRL measurements represent the

residual imprecision due to unknown factors in GA's estimation.

Development of Garbhini-GA1 formula for first-trimester dating

To remove noise from the TRAINING DATASET for building population-specific first-trimester

dating models, two methods were used – clinical criteria-based filtering and DBSCAN (Figure

1). When clinical criteria (Figure 1) were used, more than two-third observations (68.46%)

were excluded (Figure 3a). However, when DBSCAN was implemented, less than one-sixth

observations (15.85%) were removed (Figure 3b). Models for first-trimester dating using

CLINICALLY-FILTERED and DBSCAN datasets with CRL as the only predictor was done using

fractional polynomial regression to identify the best predictive model (Figure S2). The

DBSCAN approach provided a more accurate dataset (i.e. no artefacts as observed in the

CLINICALLY-FILTERED DATASET) with lesser outliers. We, therefore, used DBSCAN DATASET

for building dating models. Comparison among various dating models showed that the best

regression coefficient (R^2) was for quadratic regression (R^2 = 0.86, Table S6). This provided

235   the basis for using the following quadratic formula as the final model for estimating GA in

236   the first trimester and was termed as Garbhini-GA1 formula:

237                    $GA = -0.02294(CRL)^2 + 1.15018(CRL) + 6.73526$

where GA is in weeks, and CRL is in cm.

A multivariate dating model including CRL and the six additional predictors identified by

data-driven approaches (GLM and Random forests): resident state, weight, BMI, abdominal

girth, age, and maternal education, did not improve the performance of the CRL-based

dating model (Figure S3, Table S6).

Comparison of published formulae and Garbhini-GA1 formula for estimation of GA

The actual test of the validity of a formula is to estimate GA reliably in an unseen sample

population. We tested the published formulae's performance (Table S1) and Garbhini-GA1

formula independently on the TEST DATASET (Figure S4). It was observed that Garbhini-GA1

had an R^2 value of 0.58 (Table S8). All other formulae performed identically to Garbhini-GA1

on the TEST DATASET (Table S8). Furthermore, all possible pairwise BA analysis of these

formulae (including Garbhini-GA1) showed that the mean difference of estimated GA varied

from -0.17 to 0.50 weeks (Table 2). This result shows that Garbhini-GA1 performs equally

well as other formulae.

Impact of the choice of USG dating formula on the estimation of the rate of PTB

The PTB rates estimated using different methods ranged between 11.27 and 16.5% with

Garbhini-GA1 estimating the least (11.27%; CI 9.70, 13.00), followed by LMP (13.99%; CI

12.25, 15.86), Hadlock (14.53%; CI 12.77, 16.43), and Robinson-Fleming formula being the

highest (16.50%; CI 14.64, 18.49). Among all pairwise comparisons performed, the

differences in PTB rates estimated by Garbhini-GA1 compared with Robinson-Fleming or

McLennan-Schluter were statistically significant (Fisher's Exact test with Bonferroni

correction for *p* < 0.05, Table S9). Furthermore, Garbhini-GA1 formula had the highest

sensitivity and balanced accuracy (Table S10).

When these methods were used to determine PTB at an individual level, the Jaccard

similarity coefficient (a statistic used for gauging the similarity and diversity of sample sets)

ranged between 0.49 – 0.98 (Table 3). Interestingly, even though the two most used

methods of dating, LMP and USG-Hadlock had similar PTB rates (13.99 and 14.53%,

respectively) at the population-level, the Jaccard similarity coefficient was only 0.49

suggesting a poor agreement between the methods at an individual-level (Figure 2C, Table

3).

Discussion

Principal findings

This study's primary objectives were to compare different methods and formulae used for

GA estimation during the first trimester, develop a population-specific dating model for the

first trimester, and study the differences in PTB rate estimation using these formulae. Our

findings show that the LMP-based method overestimates GA by three days compared to the

USG (Hadlock) method. While this bias does not impact at the population level with similar

overall PTB rates determined by both methods, interestingly, there is less than 50%

agreement between these methods on who are classified as preterm at an individual level.

This is consistent with the pattern observed in a recent study from a Zambian cohort [26].

The Hadlock formula for USG-based estimation of GA was developed on a Caucasian

population and has been used for several decades globally [12]. We developed and tested

population-specific dating formula to estimate GA in an Indian setting. The CRL-based

Garbhini-GA1 formula performed the best and addition of other clinical and

sociodemographic predictors identified from machine learning tools did not improve the

performance of CRL-based Garbhini-GA1 formula. While most of the dating formulae

estimated similar PTB rates, Garbhini-GA1 formula estimated the lowest PTB rate and had

the best sensitivity to determine preterm birth.

Strengths of the study

The Garbhini-GA1 formula developed from Indian population overcomes the low

representativeness of existing dating formulae. Using advanced data-driven approaches, we

evaluated multiple combinations of various clinical and sociodemographic parameters to

estimate gestational age. We conclusively show that CRL is the sufficient parameter for first-

trimester dating of pregnancy and the addition of other clinical or social parameters do not

improve the performance of the dating model. Further, to build Garbhini-GA1 formula, we

used a data-driven approach to remove outliers that retained more observations for

building the model than would have been possible if the clinical criteria-based method had

been used to develop the reference standard. Another important strength of our study is

the standardised measurement of CRL. This reduces the imprecision to the minimum and

makes USG-based estimation of gestational age accurate.

Limitations of the data

For the development of Garbhini-GA1 model, it would have been ideal to have used

documented LMP collected pre-conceptionally. Since our GARBH-Ini cohort enrols

participants in the first trimester of pregnancy, clinical criteria based on data collected using

a questionnaire was used to derive a subset of participants with reliable LMP. This was

relatively incomplete as we had residual imprecision, which was not accounted for by the

clinical criteria. We tried to overcome this limitation by using data-driven approaches to

improve precision.

To address the truncation problem [22], we supplemented observations simulated from

Hadlock distribution. This is the best possible solution because CRL is not measured as a

current standard clinical practice beyond 14 weeks of gestation.

Interpretation

The LMP-based dating is prone to errors from recall and irregularity of menstrual cycles due

to physiological causes and pathological conditions. The overestimation of GA by the LMP-

based method seen in our cohort has been reported in other populations from Africa and

North America [26,27]. However, the magnitude of overestimation varies, as seen in studies

done earlier [26-28]. These differences could be attributed to the precision and accuracy

with which these cohorts' participants recalled their LMP. In our study, the bias in LMP-

based dating was not reflected in the population-level PTB rates; however, at an individual

level, LMP and USG-Hadlock had less than 50% agreement in the classification of PTB. Such

considerable discordance is concerning as the clinical decisions during the early neonatal

period largely depend on GA at birth. Further, any clinical and epidemiological research

studying the risk factors and complications of PTB will be influenced by choice of dating

method.

Garbhini-GA1 formula based on first-trimester CRL of our study population can be

interchangeably used with Hadlock, INTERGROWTH-21st, Verburg and Sahota but not with

McLennan-Schluter and Robinson-Fleming formulae. The higher sensitivity of Garbhini-GA1

formula to detect PTB in our study population is encouraging but should be externally

validated in other populations within the country before it can be recommended for

application. The comparable performance of Garbhini-GA1 formula with most of those

formulae developed globally may be explained by the negligible differences in the early

foetal growth as measured by CRL across populations. It would be useful to evaluate the

performance of population-specific formulae for second and third trimesters of gestation as

ethnic differences in foetal growth might manifest during this period.

Conclusions

LMP overestimates GA by three days compared to USG-Hadlock method, and only half of

the preterm birth were classified correctly by both these methods. CRL-based USG method

is the best for GA estimation in the first trimester, and the addition of clinical and

demographic features does not improve its accuracy. Garbhini-GA1 formula is an Indian-

population based formula for estimating GA in the first trimester based on CRL as the prime

parameter. It has better sensitivity than the more commonly used Hadlock formula in

estimating the PTB rate. Our results reinforce the need to develop population-specific GA

formulae. These results need to be further validated in subsequent multi-ethnic cohorts

before being applied for broader use.

List of Abbreviations

346 LMP = Last Mensural Period, GA = Gestational Age, CRL = Crown Rump Length, PTB =

347 Preterm Birth Rate, USG = Ultrasonography, CI = Confidence Interval, GLM = Generalised

348 Linear Model, LoA = Limits of Agreement, BA = Bland-Altman, IQR = Inter-Quartile Range,

349 BMI = Body Mass Index

Declarations

Ethics approval and consent to participate

Ethics approvals were obtained from the Institutional Ethics Committees of Translational

Health Science and Technology Institute; District Civil Hospital, Gurugram; Safdarjung

Hospital, New Delhi (ETHICS/GHG/2014/1.43); and Indian Institute of Technology Madras

(IEC/2019-03/HS/01/07). Written informed consent was obtained from all study participants

enrolled in the GARBH-Ini cohort. For an illiterate woman, details of the study were

explained in the presence of a literate family member or a neighbour who acted as the

witness; a verbal consent and a thumb impression were taken from her along with the

signature of the witness. All the methods were performed in accordance with the relevant

guidelines and regulations.

Consent for publication

Not applicable

Availability of data and materials

The datasets used or analysed during the current study are available from the

corresponding author on reasonable request. All the codes used for this paper are available

at https://github.com/HimanshuLab/GARBH-Ini_GA1

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by an alumni endowment from Prakash Arunachalam to the Initiative

for Biological Systems Engineering, IIT Madras (BIO/18-19/304/ALUM/KARH). GARBH-Ini

cohort study is funded by Department of Biotechnology, Government of India

(BT/PR9983/MED/97/194/2013) and for some components of the biospecimen and

ultrasound repository by the Grand Challenges India-All Children Thriving Program

(supported by the Programme Management Unit), Biotechnology Industry Research

Assistance Council, Department of Biotechnology, Government of India

(BIRAC/GCI/0114/03/14-ACT). The data analysis exercise was supported by the Grand

Challenges India- ki' Data Challenge for Maternal and Child Health grant (supported by the

Programme Management Unit), Biotechnology Industry Research Assistance Council,

Department of Biotechnology, Government of India (BT/kiData0394/06/18).

Authors' contributions

R.T., H.S. and S.B. conceived this study, R.V., N.D. and A.X., performed data and statistical

analyses, K.D. and R.T. performed data exports and contributed to data analysis, A.K., N.W.,

S.M. S.C. and GARBH-Ini Study Group developed and implemented the clinical data

collection methods and data management in GARBH-Ini cohort, R.R. provided critical

feedback on data analysis, K.D., R.T., H.S. and S.B. interpreted the results, R.V., N.D., A.X.,

R.T. and H.S. wrote the first draft of the manuscript and all listed authors critically revised

and edited subsequent manuscript drafts. All authors approved the final draft of the

manuscript.

Acknowledgements

We thank all the participants of GARBH-Ini study. We thank Karthik Raman and Nirav Bhatt

from Department of Biotechnology, Bhupat and Jyoti Mehta School of Biosciences and

Initiative of Biological Systems Engineering, IIT Madras, and Gagandeep Kang from Christian

Medical College, Vellore, for their valuable suggestions.

Members of GARBH-Ini Study Group: Translational Health Science and Technology Institute,

NCR Biotech Cluster, Faridabad, India-Coordinating Institute (Vineeta Bal, Shinjini Bhatnagar

(PI), Bhabatosh Das, Mahadev Dash, Bapu Koundinya Desiraju, Pallavi Kshetrapal, Sumit

Misra, Uma Chandra Mouli Natchu, Satyajit Rath, Kanika Sachdeva, Dharmendra Sharma,

Amanpreet Singh, Shailaja Sopory, Ramachandran Thiruvengadam, Nitya Wadhwa); National

Institute of Biomedical Genomics, Kalyani, West Bengal, India (Arindam Maitra, Partha P

Majumder (Co-PI) Souvik Mukherjee); Regional Centre for Biotechnology, NCR Biotech

Cluster, Faridabad, India (Tushar K Maiti); Clinical Development Services Agency,

Translational Health Science and Technology Institute, NCR- Biotech Cluster, Faridabad,

India (Monika Bahl, Shubra Bansal); Gurugram Civil Hospital, Haryana, India (Umesh Mehta,

Sunita Sharma, Brahmdeep Sindhu); Safdarjung Hospital, New Delhi, India (Sugandha Arya,

Rekha Bharti, Harish Chellani, Pratima Mittal); Maulana Azad Medical College, New Delhi,

India (Anju Garg, Siddharth Ramji), The Ultrasound Lab, Defence Colony, New Delhi, India

(Ashok Khurana); Hamdard Institute of Medical Sciences and Research, Jamia Hamdard

University, New Delhi, India (Reva Tripathi); All India Institute of Medical Sciences, New

Delhi, India (Alpesh Goyal, Yashdeep Gupta, Smriti Hari, Nikhil Tandon); Government of

Haryana, India (Rakesh Gupta); International Centre For Genetic Engineering and

Biotechnology, New Delhi, India (Dinakar M Salunke Co-PI); G Balakrish Nair (Rajiv Gandhi

Centre for Biotechnology, Trivandrum); Gagandeep Kang (Christian Medical College,

Vellore).

References

1. WHO: recommended definitions, terminology and format for statistical tables
related to the perinatal period and use of a new certificate for cause of perinatal
deaths. Acta Obstet Gynecol Scand. 1977;56:247–53.

2. Preterm birth. World Health Organisation. 2018. Available from:
https://www.who.int/news-room/fact-sheets/detail/preterm-birth Accessed 21 Oct
2020.

3. You D, Sharrow D, Hug L. Levels and trends in child mortality: Report 2017. World
Bank. 2017.
http://documents.worldbank.org/curated/en/358381508420391876/Levels-and-
trends-in-child-mortality-report-2017 Accessed 21 Oct 2020.

4. Committee Opinion No 700: Methods for estimating the due date. Obstet Gynecol.
2017;129:e150–4.

5. Butt K, Lim KI. Guideline No. 388-Determination of gestational age by ultrasound. J
Obstet Gynaecol Can. 2019;41:1497–507.

6. Nakling J, Buhaug H, Backe B. The biologic error in gestational length related to the
use of the first day of last menstrual period as a proxy for the start of pregnancy.
Early Hum Dev. 2005;81:833–9.

7. Wei S, Schmidt MD, Dwyer T, Norman RJ, Venn AJ. Obesity and menstrual
irregularity: associations with SHBG, testosterone, and insulin. Obesity.
2012;17:1070–6.

8. Lobo RA. What are the key features of importance in polycystic ovary syndrome?
Fertil Steril. 2003;80:259–61.

9. Chowdhury R, Sinha B, Sankar MJ, Taneja S, Bhandari N, Rollins N, et al.
Breastfeeding and maternal health outcomes: a systematic review and meta-
analysis. Acta Paediatr. 2015;104:96–113.

10. Creinin MD, Keverline S, Meyn LA. How regular is regular? An analysis of menstrual
cycle regularity. Contraception. 2004;70:289–92.

11. Hadlock FP, Shah YP, Kanon DJ, Lindsey JV. Fetal crown-rump length: reevaluation of
relation to menstrual age (5-18 weeks) with high-resolution real-time US. Radiology.
1992;182:501–5.

12. Aggarwal. Fetal ultrasound parameters: Reference values for a local perspective.
Indian J Radiol Imaging. 2020;30:149.

13. Napolitano R, Dhami J, Ohuma EO, Ioannou C, Conde-Agudelo A, Kennedy SH, et al.
Pregnancy dating by fetal crown–rump length: a systematic review of charts. Br J
Obstet Gynaecol. 2014;121:556–65.

14. Chawanpaiboon S, Vogel JP, Moller AB, Lumbiganon P, Petzold M, Hogan D, et al.
Global, regional, and national estimates of levels of preterm birth in 2014: a
systematic review and modelling analysis. Lancet Glob Health. 2019;7:e37–e46.

15. McLennan AC, Schluter PJ. Construction of modern Australian first trimester
ultrasound dating and growth charts. J Med Imaging Radiat Oncol. 2008;52:471–9.

16. Robinson HP, Fleming JEE. A critical evaluation of sonar "crown-rump length"
measurements. Br J Obstet Gynaecol. 1975;82:702–10.

17. Sahota DS, Leung TY, Leung TN, Chan OK, Lau TK. Fetal crown-rump length and

estimation of gestational age in an ethnic Chinese population. Ultrasound Obstet

Gynecol. 2009;33:157–60.

18. Verburg BO, Steegers EA, De Ridder M, Snijders RJ, Smith E, Hofman A, et al. New

charts for ultrasound dating of pregnancy and assessment of fetal growth:

longitudinal data from a population-based cohort study. Ultrasound Obstet Gynecol.

2008;31:388–96.

19. Villar J, Cheikh Ismail L, Victora CG, Ohuma EO, Bertino E, Altman DG, et al.

International standards for newborn weight, length, and head circumference by

gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-

21st Project. Lancet. 2014;384:857–68.

20. Bhatnagar S, Majumder PP, Salunke DM, Interdisciplinary Group for Advanced

Research on Birth Outcomes—DBT India Initiative (GARBH-Ini). A pregnancy cohort

to study multidimensional correlates of preterm birth in India: study design,

implementation, and baseline characteristics of the participants. Am J Epidemiol.

2019;188:621–31.

21. Ester M, Kriegel H-P, Sander J, Xu X. A density-based algorithm for discovering

clusters in large spatial databases with noise. In: Simoudis E, Han J, Fayyad U, editors.

KDD Proceedings of the Second International Conference on Knowledge Discovery

and Data Mining (KDD-96). Menlo Park CA, AAAI Press. 1996. P 226–31.

22. Ohuma EO, Papageorghiou AT, Villar J, Altman DG. Estimation of gestational age in

early pregnancy from crown-rump length when gestational age range is truncated:

the case study of the INTERGROWTH-21 st Project. BMC Med Res Methodol.

2013;13:1–14.

23. Kursa MB, Rudnicki WR. Feature selection with the Boruta package. J Stat Softw.

2010;036.

24. Martin Bland J, Altman D. Statistical methods for assessing agreement between two

methods of clinical measurement. Lancet. 1986;327:307–10.

25. Wani R. Socioeconomic status scales-modified Kuppuswamy and Udai Pareekh's

scale updated for 2019. J Family Med Prim Care. 2019;8:1846.

26. Price JT, Winston J, Vwalika B, Cole SR, Stoner MCD, Lubeya MK, et al. Quantifying

bias between reported last menstrual period and ultrasonography estimates of

gestational age in Lusaka, Zambia. Int J Gynecol Obstet. 2018;144:9–15.

27. Savitz DA, Terry JW Jr, Dole N, Thorp JM Jr, Siega-Riz AM, Herring AH. Comparison of

pregnancy dating by last menstrual period, ultrasound scanning, and their

combination. Am J Obstet Gynecol. 2002;187:1660–6.

28. Hoffman CS, Messer LC, Mendola P, Savitz DA, Herring AH, Hartmann KE.

Comparison of gestational age at birth based on last menstrual period and

ultrasound during the first trimester. Paediatr Perinat Epidemiol. 2008;22:587–96.

Figures

Figure 1: Outline of the data selection process for different datasets – (A) TRAINING

DATASET and (B) TEST DATASET. Coloured boxes indicate the datasets used in the analysis.

The names of each of the dataset are indicated below the box. Exclusion criteria for each

step are indicated. N_p indicates the number of participants included or excluded by that

particular criterion and N_o indicates the number of unique observations derived from the

participants in a dataset. Biologically implausible CRL values (either less than 0 or more than

10 cm) for the first trimester were excluded, b Biologically implausible GA values (either less

than 0 and more than 45 weeks) were excluded.

Figure 2: (A) Distribution of the difference between USG- and LMP-based GA. The x-axis is

the difference between USG and LMP-based GA in weeks, and the y-axis is the number of

observations. (B) BA analysis to evaluate the bias between USG and LMP-based GA. The x-

axis is mean of Hadlock and LMP-based GA in weeks, and the y-axis is the difference

between Hadlock and LMP-based GA in weeks. Regression line with 95% CI is shown. (C)

Comparison of individual-level classification of preterm birth by Hadlock- and LMP-based

methods. Green (term birth for both), red (preterm birth for both), blue (term birth for LMP

but preterm birth for Hadlock) and purple (term for Hadlock but preterm for LMP).

Figure 3: Comparison of data chosen to be reference data for the development of dating

formula by (A) clinical and (B) data-driven (DBSCAN) approaches. The x-axis is CRL in cm, and

the y-axis is GA in weeks (LMP-based are datapoints, Garbhini-GA1 is regression line). After

filtering, the data points selected (TRUE) are coloured black and points not selected (FALSE)

are white.

Tables

Table 1: Baseline characteristics of the participants included in the TRAINING DATASET (N_o = 2562) to compare different methods of dating.

Sociodemographic characteristics	Median (IQR) or N (%) or Mean ± SD
Age (year)	23 (21 – 26)
GA at enrolment by LMP (weeks)	11.31 ± 2.67
GA at enrolment by USG-Hadlock (weeks)	10.87 ± 2.28
BMI at enrolment into the cohort [a]	
Underweight	27.20%
Normal weight	59.93%
Obese	9.09%
Overweight	1.66%
Haemoglobin (g/dL)	8.8 (8.2 – 9.2)
Height (cm)	153.0 (149.2 – 156.8)
Socioeconomic status [b]	
Upper class	0.66%
Upper middle class	15.40%
Lower middle class	33.98%
Upper lower class	48.96%
Lower class	0.43%
Undetermined	0.57%
Parity (number)	
0	49.53%
1	33.55%
2	12.60%
3	3.34%
4	0.74%
5	0.14%
Level of education	

	Illiterate	21.58%
	Literate or primary school	8.63%
	Middle school	15.09%
	High school	18.61%
	Post high school diploma	20.89%
	Graduate	12.23%
	Post-graduate	2.94%
Occupation		
	Unemployed	93.48%
	Unskilled worker	3.34%
	Semi-skilled worker	0.97%
	Skilled worker	1.40%
	Clerk, shop, farm owner	0.17%
	Semi-professional	0.26%
	Professional	0.34%
Religion		
	Hindu	92.14%
	Muslim	6.60%
	Sikh	0.40%
	Christian	0.74%
	Buddhist	0.00%
	More than one religion	0.09%
Fuel used for cooking [c]		
	Biomass fuel	7.86%
	Clean fuel [d]	92.14%
Source of drinking water		
	Safe water [e]	49.80%
	Unsafe water	50.20%
Second-hand tobacco smoke		
	Exposed	19.23%
	Unexposed	80.57%
	Undetermined	0.20%

History of any chronic illnesses [f]	
Absent	99.03%
Present	0.97%
History of hypertensive disease of pregnancy	
Absent	99.57%
Present	0.43%
History of contraceptive at the time of conception	
Absent	90.79%
Present	7.30%

[a] Pre-pregnancy BMI was calculated as weight (kg)/height^2 (m) from participants' weight and

height measured at enrolment. BMI categories were defined as underweight (< 18.5);

normal (18.5 – 24.9); overweight (25.0 – 29.9); obese (≥ 30.0).

[b] Socioeconomic status was assessed using Modified Kuppuswamy's socioeconomic scale

[25], calculated using education and occupation of the head of the family and monthly

family income.

[c] Indoor air pollution: use of biomass fuel for cooking or presence of a smoker in the

residential compound, as reported by the participant.

[d] Clean fuel includes liquefied petroleum gas and electricity.

[e] Safe water includes bottled water or piped water into the residence.

[f] Chronic illnesses include a history of hypertension, diabetes, cardiac disease and thyroid

disorders.

Table 2: Pairwise comparison of mean difference (LoA) between different first-trimester dating formulae (Difference: Column formula - Row

formula). Values shown in white are for the TRAINING DATASET (N_o = 2562) and values shown in grey are for the TEST DATASET (N_o = 808) (see

Methods for details).

Formula	Hadlock	McLennan-Schluter	Robinson-Fleming	Sahota	Verburg	INTERGROWTH-21st	Garbhini-GA1
Hadlock		-0.16 (-0.40, 0.079)	-0.17 (-0.36, 0.016)	0.034 (-0.22, 0.29)	0.037 (-0.41, 0.48)	0.079 (-0.54, 0.70)	0.38 (-0.11, 0.87)
McLennan-Schluter	0.14 (-0.032, 0.31)		-0.015 (-0.16, 0.13)	0.19 (0.05, 0.34)	0.20 (-0.10, 0.50)	0.24 (-0.36, 0.83)	0.54 (-0.02, 1.10)
Robinson-Fleming	0.17 (-0.019, 0.35)	0.024 (-0.095, 0.14)		0.21 (0.082, 0.33)	0.21 (-0.097, 0.52)	0.25 (-0.35, 0.85)	0.56 (0.02, 1.09)
Sahota	-0.052 (-0.30, 0.19)	-0.19 (-0.33, -0.057)	-0.22 (-0.35, -0.088)		0.002 (-0.20, 0.20)	0.044 (-0.46, 0.55)	0.35 (-0.15, 0.85)
Verburg	-0.065 (-0.51, 0.39)	-0.21 (-0.52, 0.11)	-0.23 (-0.54, 0.08)	-0.013 (-0.22, 0.19)		0.042 (-0.45, 0.53)	0.35 (-0.26, 0.95)

INTERGROWTH-21st	-0.12 (-0.79, 0.55)	-0.26 (-0.90, 0.38)	-0.28 (-0.94, 0.38)	-0.066 (-0.62, 0.49)	-0.053 (-0.59, 0.49)		0.30 (-0.03, 0.64)
Garbhini-GA1	-0.40 (-0.93, 0.13)	-0.54 (-1.12, 0.04)	-0.57 (-1.14, 0.01)	-0.35 (-0.87, 0.18)	-0.34 (-0.96, 0.29)	-0.28 (-0.61, 0.05)	

546 **Table 3:** The Jaccard similarity coefficient of PTB prediction between each pair of the method.

Formula	LMP	Hadlock	McLennan-Schluter	Robinson-Fleming	Sahota	Verburg	INTERGROWTH-21st	Garbhini-GA1
LMP	1.00	0.49	0.50	0.50	0.52	0.53	0.53	0.50
Hadlock		1.00	0.90	0.88	0.88	0.81	0.80	0.77
McLennan-Schluter			1.00	0.98	0.83	0.82	0.80	0.69
Robinson-Fleming				1.00	0.82	0.81	0.79	0.68
Sahota					1.00	0.92	0.89	0.83
Verburg						1.00	0.87	0.83
INTERGROWTH-21st							1.00	0.87
Garbhini-GA1								1.00

Additional files

550   Additional File 1: Supplementary information (PDF)

551   Additional File 2: Supplementary information tables (XLS)

553   **Figure 1**

556   **Figure 2**

558   **Figure 3**