Title Page

Interprofessional Learning in Immediate Life Support training improves simulated patient related outcomes.

Jeremy Charles Morse, University of Aberdeen, Clinical Skills Centre, Suttie Centre for Teaching and Learning in Healthcare, University of Aberdeen, Foresterhill, Aberdeen, AB25 2XG
Craig William Brown, NHS Grampian, Emergency Department, Aberdeen Royal Infirmary, Foresterhill, Aberdeen, AB25 2XG

Corresponding Author
Jerry Morse

Clinical Skills Centre
Suttie Centre for Teaching and Learning in Healthcare
University of Aberdeen
Foresterhill
Aberdeen
AB25 2XG

Email: jerry.morse@abdn.ac.uk

Telephone: +44 1224437733

Word count of abstract: 303

Word count of article: 2674 (excluding abstract)

Abstract

Aim of the study
To assess team performance in implementing time critical key interventions during a simulated resuscitation after participating in either an interprofessional-learning (IPL) or uniprofessional-learning (UPL) Immediate Life Support training course (ILS).

Introduction
Much of the published work on team-based simulation training has measured the lower levels of Kirkpatrick's hierarchy of evidence and effectiveness. This study aimed to ascertain if interprofessional team training could improve a higher level of outcome such as behaviour and patient outcomes.

Methods
A retrospective quantitative analysis of time critical points in a simulated cardiac arrest resuscitation, from a previous randomised study on the effects of Interprofessional Learning in ILS. The video recordings from the original study consisting of medical (n=48) and nursing (n=48) were analysed to mark when either the IPL or UPL team performed a time critical intervention.

Results
Five time-critical points for interventions were identified; confirmation of cardiac arrest, commencement of initial CPR, rhythm check, time to 1^{st} shock and delay in restarting CPR. Parametric testing of each of these time-based criteria were subjected to an independent sample *t-test* with statistically significant findings in three of the five criteria in favour of those who had undertaken the interprofessional learning.

Conclusion
Our results demonstrate that through an IPL approach in ILS there was a statistically different improvement in mean times to performing time-critical interventions, which if transferred to the clinical environment could improve and impact on both change of behaviour and patient outcomes in Kirkpatrick's higher levels of evidence and effectiveness.

Though this study shows that team behaviour and performing time-critical interventions can improve in the short-term, we acknowledge that further longitudinal studies are required to ascertain the effect of such improvement over time. So that both as researchers and educators we can make healthcare teams work safer and more efficiently to improve patient outcomes.

What this paper adds

What is already known:
Like simulation, intuitively we know that Interprofessional Learning and Team training should make a difference to clinical practice for those involved. The majority of research reported, is measured at the lower levels of Kirkpatrick's hierarchy. This retrospective video analysis of a previous study aimed to look at the functioning between interprofessionally and uni-professionally trained teams in performing time-critical interventions in a simulated cardiac arrest.

What this study adds:
Our study suggests that the use of Interprofessional Learning in resuscitation training does have an effect on performance during the management of a simulated cardiac arrest which could improve and impact on both change of behaviour and patient outcomes in Kirkpatrick's higher levels of evidence and effectiveness.

Introduction

Resuscitation training continues to be a core element of many undergraduate healthcare curricula. Standardised resuscitation courses, such as the Resuscitation Council UK's Immediate Life Support (ILS) course[1], are often structured to provide healthcare learners with an algorithmic approach to managing the critically unwell patient. Whilst these courses are delivered to learners from a number of disciplines at postgraduate level, often at an undergraduate healthcare level they are provided in a uniprofessional context. However, this is despite in practice almost always occurring as a multi-professional team-based response. Though studies of simulated cardiac arrest have been reported, the investigation of team-based factors in real life cardiac arrest scenarios remains challenging.[2] Therefore, simulation-based studies such as this one, can offer educational researchers surrogate with markers in respect to measuring complex team-based indicators in a cardiac-arrest resuscitation.[3,4]

From the resuscitation literature it is known that in regards to patient outcome, for each minute of delay to defibrillation, when the patient is in a shockable cardiac rhythm, the probability of survival to discharge reduces by 10-12%.[5] Therefore, the effective management of a cardiac arrest resuscitation requires a team-based approach both in training and in practice[4]. With previous studies showing that teams who were trained using an interprofessional ILS method, had improved overall team-working scores than those who had been trained uni-profesionally.[6,7,8]

Much of the published work on team-based simulation training has measured outcomes on the lower levels of Kirkpatrick's hierarchy of evidence of effectiveness, i.e. learner outcomes such as self-reported views, perceptions or attitudes, rather than the process elements of quality care[9,10].

This study, in line with the 2015 report from the Institute of Medicine, hopes to provide another step towards building more solid evidence in linking interprofessional learning to

patient outcomes.[11] It was therefore hypothesised that if interprofessional learning can improve team performance, then this could also be seen in time critical points throughout the simulated resuscitation, thus leading to potential better patient survival and provide potential evidence at Kirkpatrick levels 3 & 4 ('behaviour' and 'patient outcomes').

Ethical Approval:

Ethical approval for the original study was obtained from both the School of Nursing Ethical Review Panel, Robert Gordon University (RGU) and the College Ethics Review Board, College of Life Sciences and Medicine, University of Aberdeen (UoA) and included approval for limited disclosure to minimise any perceptual bias regarding Inter-Professional Learning (IPL). In this way all participants were blinded as to the study objectives regarding the effect on the team performance/interaction during a simulated resuscitation. The participants were however made aware that it was a study to evaluate the teaching of ILS to final year medical and nursing students.

Methods:

Design: A retrospective quantitative analysis of time critical points in a simulated cardiac arrest resuscitation, from a previous randomised study on the effects of Interprofessional Learning in Immediate Life Support (ILS).[8]

From the original data previously reported, this subsequent study would retrospectively analyse, the video recordings taken of the final year healthcare students (nursing n=48 and medical n=48) recruited and who made up the final teams for the simulated resuscitation. These formed sixteen teams each of six participants who were randomly assigned into equal numbers of medical and nursing students based on their original ILS course. Namely those who had undertaken either the uniprofessional (UPL) or interprofessional (IPL) resuscitation teaching.

Having consented to participation and video recording of the session, each of the teams were invited to participate in a cardiac arrest simulation scenario which was recorded using the SMOTS ™ fixed recording system (Scotia UK plc, Edinburgh UK) installed in the simulation laboratory allowing for retrospective detailed analysis of the team's performances. To ensure consistency the same resuscitation scenario was delivered to all groups, which was a patient who presented in cardiac arrest and in a shockable rhythm (ventricular fibrillation). Each of the groups simulated resuscitations were recorded, then downloaded and saved for analysis by the blinded reviewer in the original study.

Following advice from the department of medical statistics at the University of Aberdeen it was decided to employ independent sample t-test to compare the time taken to complete key time-based interventions between the two cohorts, with a starting point from when the team entered the room. These key time-based criteria, measured in seconds were; confirming cardiac arrest, commencement of chest compressions, time to delivery of first shock and delay in restarting CPR post shock. The resultant qualitative data, from the retrospective analysis, were then imputed into the SPSS 10 statistical software and parametric testing applied.

Results:

Video recorded data for eight 'IPL' and eight 'UPL' trained teams, comprising of a total of ninety-six participants were available for the follow up cardiac arrest simulation scenario and each time-based criterion was subjected to an independent sample t-test are summarised and displayed in *table 1*

Table 1 – Mean Times in seconds from entering the room

Mean Times (seconds) to Intervention and Statistical Significance						
		Mean (Seconds)	SD	*t*	df	*p*
Confirm Cardiac	IPL	24.50	11.15	-0.271	14	0.790

Arrest	UPL	25.75	6.78			
Commence Initial CPR	IPL	42.88	14.74	-1.338	14	.202
	UPL	52.50	14.03			
Check Rhythm	IPL	88.38	12.55	-3.57	14	0.003
	UPL	116.25	18.15			
Time to 1st Shock	IPL	110.25	10.93	-3.370	14	.005
	UPL	159.50	39.86			
Delay in restarting CPR	IPL	22.00	5.86	-2.625	14	.020
	UPL	43.25	22.13			

An independent-samples t-test was conducted to compare the time taken in seconds, to confirm cardiac arrest by the IPL and UPL teams which showed that there was no statistical significance in the time taken for UPL teams (M = 25.75, SD = 6.78) and IPL teams (M = 24.5, SD =11.15; t (-.271), p =0.790). (*Figure 1*). Similarly, in regards time to commencement of CPR an independent-samples t-test was conducted on the number of seconds taken by the IPL and UPL teams which again showed despite the IPL cohort commencing compressions in a mean time of 10 seconds faster than the UPL teams, there was no statistical significance in time taken between the UPL teams (M = 52.5, SD = 14.03) and IPL teams (M = 42.88, SD =14.74; t (-1.34.), p =0.202). (*Figure 2*).

When analysing the differences between the time taken between the groups to perform the first 'rhythm check' (a process involving connecting the mannequin to the defibrillator whilst CPR was ongoing) an independent-samples t-test was conducted that showed statistical significance in time (seconds) taken between the UPL teams (M = 116.25, SD = 18.15) and IPL teams (M = 88.38, SD =12.55; t (-3.57.), p =0.003). (*Figure 3*). In regards the time critical point of delivering the 'first shock', the independent-samples t-test conducted also showed there was statistical significance in the time in seconds taken between the UPL teams (M = 159.50, SD = 39.86) and IPL teams (M = 110.25, SD =10.93; t (-3.37.), p =0.010). (*Figure 4*). The final key-

time based criteria analysed was the recommencement of CPR after the first shock. The independent-samples t-test was conducted and once again showed that there was statistical significance in time taken between the UPL teams (M = 43.25, SD = 22.13) and IPL teams (M = 22.00, SD =5.86; t (-2.625.), p =0.030).

Discussion:

It is recognised that interprofessional learning is essential in the training and education of healthcare professionals in order to equip them with the prerequisite skills, knowledge and attitudes to be able to work together in meeting the needs of patients.[12] However, despite this there remains no clear path on how to assess the effective preparation of healthcare professionals to work collaboratively. It is therefore hoped that this study will assist in not only highlighting the positive impact that collaborative working can achieve, but also the impact on patient outcomes in a simulated environment, which can then be transferred to the clinical setting. [13,14] Within clinical practice most in-hospital cardiac arrests are managed by interprofessional teams, as opposed to uniprofessional groups or isolated responders, with an emphasis on the importance of good team work and leadership. Currently within the UK at undergraduate level, many of the mandatory life support courses continue to be delivered in a uniprofessional manner despite increased regulatory emphasis on team-based approaches to healthcare[15, 16].

In the first component of this programme of research, it was identified that undergraduate students who were trained in an interprofessional format with ILS performed significantly better on TEAM scoring in a simulated cardiac arrest scenario[8]. Although the use of the TEAM score[17] allowed the for the assessment of IPL versus UPL resuscitation team performance. This retrospective analysis has shown that IPL trained teams not only can achieve improved team performance but more importantly are able to perform better time-based outcomes in checking initial rhythm and most importantly a delivery of the 'first shock' in a simulated cardiac arrest than those trained uni-professionally. Whilst it can be argued that enhancing performance

through interprofessional learning is a pre-curser to improving patient outcomes, it could also be suggested that this would impact on process related outcomes ultimately reflected in patient survival.

It has been well documented that in cardiac arrest situations, shorter times taken to complete time-critical outcomes were improved through standardised delivery of resuscitation training programmes such as the UK resuscitation Council ILS course.[1,12] Indeed, it is the algorithmic teaching of such courses that encourage learners to promptly diagnose cardiac arrest, commence effective CPR and perform early defibrillation when indicated, that can maximise the chance of survival both in the pre-hospital and in-hospital situation. Furthermore, that in clinical practice, any delay in any of these key components or prolonged pauses in the algorithmic process are associated with poorer patient outcomes.[18,19]

From the analysis of this study's data there are statistically significant findings that suggest interprofessional trained teams achieved the key time critical points of performing rhythm check and delivering the first shock faster than those who were trained uniprofessionally. Although it is acknowledged this remains a simulation-based resuscitation, these statistically significant results assist in progressing the related simulation-based education (SBE) literature on Kirkpatrick's hierarchy of effectiveness at least to level 3 'behavioural change' and potentially level 4 'impact on patient outcomes'.

To achieve this then practising as a team and receiving feedback with the aim of improving overall performance has to be an integral component of interprofessional learning environment. This was the case in the delivery the initial ILS courses for both cohorts', where all the participants would have received feedback and learning conversations upon their performance. Whereas with any feedback the aim is to identify any potential performance gaps and allow participants to reflect and reframe their thought processes and actions for future situations. Therefore, whilst those in the UPL cohort would have been given ways to improve on performance, it can be suggested that those in the IPL cohort benefited further from the

feedback and learning conversations from a multidisciplinary team perspective and sharing the experience to work collaboratively in a resuscitation scenario. As a result, it was this feedback to the IPL teams, combined with the focused and deliberate and shared practise on the ILS algorithmic process that led to the statistically significant results identified in this study.

It has been reported that such deliberate practise in healthcare procedures leads to improved chronometry i.e. time taken to complete procedures.[20] Whilst there was no data reported from either of the cohort's initial resuscitation courses, the retrospective study data showed that those trained interprofessionally had significantly improved chronometry. Especially in relation to the delivery of first shock and recommencement of CPR, two of the key factors in the chain of survival and improved patient outcomes of patients in cardiac arrest.

Interprofessional learning as a pedagogical approach to training continues to rise as educators across the many professions embrace the concept that ultimately healthcare on the 'shop floor' occurs in teams. Further, that for IPL to be truly effective and have an effect on outcomes then all members should train together and be aware of each other's roles, capabilities and ability to perform as part of a multidisciplinary team. It was strikingly evident during the video analysis that there was a notable difference in the way that the IPL trained cohort were aware of and respected each other whilst working together as a more cohesive team.

Although the impact of IPL on patient outcomes remains unclear, it is hoped that the evidence provided through this study will assist in furthering the knowledge base on which further research can continue to expand and promote the benefits of interprofessional learning.[21]

As SBE continues to be widely adopted and the drive for patient safety continues, interprofessional learning and deliberate practice such as in the case of resuscitation skills, allows for students to understand the equality of roles within the team. Indeed, the student participants in this study interacted and responded more positively following the combined profession training than those completing the uniprofessional training. Yet despite the growing

evidence and the willingness from many educators to further utilise interprofessional learning, this is often still limited to just two professions rather than all those involved in a collaborative practice. A situation which, while challenging, may make such team-based practice less relevant.[22 CAIPE] However, as has been demonstrated by the results of this study, the use of IPL between medical and nursing students significantly improved the overall team's performance. Furthermore, that the use of IPL in resuscitation training allowed the team members to see the equality of role expectations, rather than the professional boundaries often assumed at an undergraduate level.

Limitations to the study

It is acknowledged that analysing higher level Kirkpatrick related outcomes in randomized interprofessional simulation studies remain difficult, in part due to the total number of participants involved being divided into teams, in this case ninety-six participants forming sixteen teams. Also, that if there was a delay at the initial recognition of cardiac arrest then this would have a 'knock on effect' to the other key markers within the study parameters. However, the same could also occur in clinical practice and the statistically significant results of the study indicate that IPL has mitigated some of the argument.

Conclusion

The initial study methodology of incorporating the TEAM tool to quantitively measure the performance of IPL and UPL trained medical and nursing students, indicated a statistically significant difference in team performance for those from the IPL cohort. This second and retrospective study of the data has further demonstrated that as well as increased team performance, those who underwent interprofessional learning completed the time critical key tasks of initial rhythm check and first defibrillation in a cardiac arrest faster than those who had the uni-professional training. Moreover, this study has identified and evaluated that as a direct result of interprofessional learning patient outcomes can be improved and adds to the evidence

base that training in a multi-professional educational experience can have an impact not only on IPL but on collaborative care on patient outcomes and health both in the UK and worldwide.

Supporting recent recommendations,[23] this study indicates that in undergraduate healthcare curricula and for which either basic or intermediate life support is required, such training should be done so wherever possible interprofessionally. Furthermore, that despite the low number of studies measuring the outcomes of behavioural change and benefits to patient outcomes, this research adds to the evidence that the use of IPL in resuscitation training can make a difference. To this end, further longitudinal studies are required to ascertain the effect of such improvement over time, so that both as researchers and educators we can make healthcare teams work safer and more efficiently to improve patient outcomes.

Contributors: JM: participated in study design, data analysis and interpretation, drafting and revising the manuscript and approved the final version. CB: participated in study design, data interpretation, drafting and revising the manuscript and approved the final version.

Conflicts of Interests:

The authors report no conflict of interest. The authors alone are responsible for the writing and content of this paper.

Acknowledgments:

The Department of Medical Statistics for their continued support and advice

Funding Statement:

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

References

[1]. Course Information, Resuscitation Council (UK), London, Resuscitation Council (UK), 2017. (Accessed 19 October 2017, athttps://www.resus.org.uk/information-on-courses/)

[2]. Mundell WC, Kennedy CC, Szostek JH, Cook DA. Simulation technology for resuscitation training: a systematic review and meta-analysis. Resuscitation 2013;84(9):1174-1183.

[3]. Norris EM, Lockey AS. Human factors in resuscitation teaching. Resuscitation 2012;83(4):423-427.

[4]. Connell CJ, Endacott R, Jackman JA, Kiprillis NR, Sparkes LM, Cooper SJ. The effectiveness of education in the recognition and management of deteriorating patients: A systematic review. Nurse Educ Today 2016; 44:133-145.

[5]. Brock D, Abu-Rish E, Chiu C, Hammer D, Wilson S, Vorvick L, et al. Interprofessional education in team communication: working together to improve patient safety. BMJ Qual Saf 2013;22(5):414-423.

[6]. Castelao EF, Russo SG, Riethmüller M, Boos M. Effects of team coordination during cardiopulmonary resuscitation: A systematic review of the literature. J Crit Care 2013;28(4):504-521.

[7]. Reeves S, Fletcher S, Barr H, Birch I, Boet S, Davies N, et al. A BEME systematic review of the effects of interprofessional education: BEME Guide No. 39. Med Teach 2016;38(7):656-668.

[8]. Morse JC, Brown CW, Morrison I, Wood C. Interprofessional learning in immediate life support training does effect TEAM performance during simulated resuscitation. BMJ Simulation and Technology Enhanced Learning 2018: bmjstel-2018-000394.

[9]. Garling P. Commissioner. Final report of the Special Commission of Inquiry into Acute Care Services in NSW Public Hospitals. Sydney: State of NSW, 2008.

[10]. Lapkin S, Levett-Jones T, Gilligan C. A systematic review of the effectiveness of interprofessional education in health professional programs. Nurse Education Today 2013;33(2):90-102.

[11]. Institute of Medicine (IOM), Committee on Measuring the Impact of Interprofessional Education on Collaborative Practice and Patient Outcomes. The National Acadamies Press, 2015.

[12]. Schultz AM, McCoy MA, Graham R. Strategies to improve cardiac arrest survival: a time to act: National Academies Press; 2015.

[13]. World Health Organization. Framework for action on interprofessional education and collaborative practice 2010.

[14]. Reid A, Fielden SA, Holt J, MacLean J, Quinton ND. Learning from interprofessional education: a cautionary tale. Nurse Educ Today 2018; 69:128-133.

[15]. Hunziker S, Tschan F, Semmer NK, Zobrist R, Spychiger M, Breuer M, et al. Hands-on time during cardiopulmonary resuscitation is affected by the process of teambuilding: a prospective randomised simulator-based trial. BMC emergency medicine 2009;9(1):3.

[16]. Hunziker S, Johansson AC, Tschan F, Semmer NK, Rock L, Howell MD, et al. Teamwork and leadership in cardiopulmonary resuscitation. J Am Coll Cardiol 2011;57(24):2381-2388.

[17]. Cooper S, Cant R, Porter J, Sellick K, Somers G, Kinsman L, et al. Rating medical emergency teamwork performance: development of the Team Emergency Assessment Measure (TEAM). Resuscitation 2010;81(4):446-452.

[18]. Schneider T, Mauer D, Diehl P, Eberle B, Dick W. Does standardized mega-code training improve the quality of pre-hospital advanced cardiac life support (ACLS)? Resuscitation 1995;29(2):129-134.

[19]. Birchak J, Abdul-Kafi O, Pham T, Viner M, Nehmer M, Rao B, et al. Prolonged pauses in cardiopulmonary resuscitation are associated with poor survival during in-hospital cardiac arrest. J Am Coll Cardiol 2018;71(11 Supplement): A459.

[20]. Pusic MV, Brydges R, Kessler D, Szyld D, Nachbar M, Kalet A. What's your best time? Chronometry in the learning of medical procedures. Med Educ 2014;48(5):479-488.

[21]. Cox M, Cuff P, Brandt B, Reeves S, Zierler B. Measuring the impact of interprofessional education on collaborative practice and patient outcomes 2016.

[22]. Barr H, Ford J, Gray R, Helme M, Hutchings M, Low H, et al. Interprofessional education guidelines. London: CAIPE 2017.

[23]. McNaughton S. The long-term impact of undergraduate interprofessional education on graduate interprofessional practice: A scoping review. Journal of interprofessional care 2018;32(4):426-435.

TIME LINE OF CRITICAL EVENTS

Figure 1 – Time to confirming cardiac arrest

TIME LINE OF CRITICAL EVENTS

Figure 1 – time to commencement of CPR

TIME LINE OF CRITICAL EVENTS

◆ Confirm CA   ■ Commence CPR   ▲ Check Rhythm

Figure 3 – time to check cardiac rhythm

TIME LINE OF CRITICAL EVENTS

Figure 1 – time to delivering first shock