Title:

Clinical presentation and differential splicing of *SRSF2*, *U2AF1* and *SF3B1* mutations in

patients with Acute Myeloid Leukaemia

Running Head:

Characterization of SF mutations in AML

Stefanos A. Bamopoulos[1,2], Aarif M. N. Batcha[3,4], Vindi Jurinovic[1,3], Maja Rothenberg-Thur-

ley[1], Hanna Janke[1], Bianka Ksienzyk[1], Julia Philippou-Massier[5], Alexander Graf[5], Stefan

Krebs[5], Helmut Blum[5], Stephanie Schneider[1,6], Nikola Konstandin[1], Maria Cristina Sauer-

land[7], Dennis Görlich[7], Wolfgang E. Berdel[8], Bernhard J. Woermann[9], Stefan K. Bohlander[10],

Stefan Canzar[11], Ulrich Mansmann[3,4,12,13], Wolfgang Hiddemann[1,12,13], Jan Braess[14], Karsten

Spiekermann[1,12,13], Klaus H. Metzeler[1,12,13] and Tobias Herold[1,12,15]

1 Laboratory for Leukemia Diagnostics, Department of Medicine III, University Hospital, LMU

Munich, Munich, Germany

2 Department of Hematology and Oncology (CBF), Charité University Medicine, Berlin, Ger-

many

3 Institute for Medical Information Processing, Biometry and Epidemiology, LMU Munich, Mu-

nich, Germany

4 DIFUTURE (Data integration for Future Medicine (DiFuture, www.difuture.de), LMU Munich,

Munich, Germany

5 Laboratory for Functional Genome Analysis (LAFUGA), Gene Center, LMU Munich, Munich,

Germany

6 Institute of Human Genetics, University Hospital, LMU Munich, Munich, Germany

7 Institute of Biostatistics and Clinical Research, University of Münster, Münster, Germany

8 Department of Medicine, Hematology and Oncology, University of Münster, Münster, Ger-

many

9 German Society of Hematology and Oncology, Berlin, Germany

10 Leukaemia and Blood Cancer Research Unit, Department of Molecular Medicine and Pa-

thology, University of Auckland, Auckland, New Zealand

11 Gene Center, LMU Munich, Munich, Germany

12 German Cancer Consortium (DKTK), Partner Site Munich, Munich, Germany

13 German Cancer Research Center (DKFZ), Heidelberg, Germany

14 Department of Oncology and Hematology, Hospital Barmherzige Brüder, Regensburg,

Germany

15 Research Unit Apoptosis in Hematopoietic Stem Cells, Helmholtz Zentrum München, Ger-

man Center for Environmental Health (HMGU), Munich, Germany

Corresponding Authors:

Stefanos A. Bamopoulos; stefanos.bamopoulos@protonmail.com

and

Tobias Herold, MD; Marchioninistr. 15; 81377 Munich; Germany; Phone: +49 89 4400-0; FAX:

+49 89 4400-74242; Email: tobias.herold@med.uni-muenchen.de

Abstract

Previous studies have demonstrated that splicing factor mutations are recurrent events in hematopoietic malignancies. Their clinical characteristics and aberrant splicing patterns have been explored in myelodysplasia, however, their functional consequences in acute myeloid leukaemia are largely unknown. The aim of this study was the comprehensive clinical and functional analysis of mutations in the three most commonly afflicted splicing factor genes: *SRSF2*, *U2AF1* and *SF3B1*. To this end, we examined the prognostic role of splicing factor mutations in two large independent cohorts, encompassing a total of 2678 acute myeloid leukaemia patients treated with intensive chemotherapy. The clinical analysis was complemented by RNA-sequencing of 246 patients to identify targets of splicing dysregulation. Results were validated in an additional RNA-sequencing dataset of 177 patients. Patients with splicing factor mutations show inferior relapse-free and overall survival, however, these mutations do not represent independent prognostic markers. Differential isoform expression analysis revealed a characteristic expression profile for each splicing factor mutation with a strong dysregulation of several isoforms. Furthermore, by establishing a custom differential splice junction usage pipeline we accurately detected aberrant splicing in splicing factor mutated samples. Mutated samples were characterized by predominantly decreased splice junction utilization of a large number of genes. A large proportion of differentially used spliced junctions were novel. Targets of splicing dysregulation included several genes with a known role in acute myeloid leukaemia. In *SRSF2*(P95H) mutants we further explored the possibility of a cascading effect through the dysregulation of the splicing pathway.

Taken together, our findings suggest that splicing factor mutations does not represent independent prognostic markers. However, they do have genome-wide consequences on gene splicing leading to dysregulated isoform expression of several genes.

Introduction

The discovery of recurring somatic mutations within splicing factor genes in a large spectrum of human malignancies has brought attention to the critical role of splicing and its complex participation in carcinogenesis [1–3]. The spliceosome is a molecular machine assembled from small nuclear RNA (snRNA) and proteins and is responsible for intron removal (splicing) in pre-messenger RNA. In acute myeloid leukaemia (AML), splicing factor mutations occur most frequently in *SRSF2, U2AF1* and *SF3B1*. The splicing factors encoded by these genes are all involved in the recognition of the 3'-splice site during pre-mRNA processing.[4] Splicing factor (SF) mutations are especially common in haematopoietic malignancies, where they occur early on and remain stable throughout the disease evolution of myelodysplastic syndromes (MDS) [1,5–9]. SF mutations are also prevalent in acute myeloid leukaemia (AML), which is often the result of myeloid dysplasia progression, with reported frequencies of 6-10%, 4-8% and 3% for *SRSF2, U2AF1* and *SF3B1* mutations respectively [2,4,10,11].

SF mutations rarely co-occur within the same patient, implying the lack of a synergistic effect or synthetic lethality [1,2,6]. They are typically heterozygous point mutations, frequently coincide with other recurrent mutations in haematopoietic malignancies and are associated with aberrant splicing in genes recurrently mutated in AML [2,4,8]. Notably, the aberrant splicing patterns are distinct for each SF mutation, suggesting that SF mutations do not share the same mechanism of action and should be recognized as individual alterations [4,9,12–17].

The clinical characteristics and outcome of patients with SF mutations are well defined in MDS [1,3,8,9]. Meanwhile, attempts at determining the role of SF mutations as independent prognostic markers in AML have often been limited to specific subgroups and it remains unclear whether the inferior survival associated with SF mutations is confounded by their association with older age or accompanying mutations [10,18]. Additionally, while evidence of aberrant

splicing due to SF mutations has emerged for many genes relevant in AML, it is yet uncertain

whether and how these changes directly influence disease initiation or evolution.

The aim of this study was a comprehensive analysis of the prognostic implications of SF

mutations in two well-characterized and intensively treated adult AML patient cohorts

amounting to a total of 2678 patients. In addition, the core functional consequences of SF

mutations were explored using targeted amplicon sequencing in conjunction with RNA-

sequencing on two large datasets.

Patients and Methods

Patients

Our primary cohort included a total of 1138 AML patients treated with intensive chemotherapy in two randomized multicenter phase 3 trials of the German AML Cooperative Group (AMLCG). Treatment regimens and inclusion criteria are described elsewhere [2]. A cohort of 1540 AML patients participating in multicenter clinical trials of the German-Austrian AML Study Group (AMLSG), was used for validation [19]. Cohort composition and filtering criteria are outlined in the supplementary.

Molecular Workup

All participants of the AMLCG cohort received cytogenetic analysis, as well as targeted DNA-sequencing as described previously [2]. The subset of the AMLSG cohort included in this study received a corresponding molecular workup, described elsewhere [19].

RNA-Sequencing and data processing

Using the Sense mRNA Seq Library Prep Kit V2 (Lexogen; Vienna, Austria) 246 samples underwent poly(A)-selected, strand-specific, paired-end sequencing on a HiSeq 1500 instrument (Illumina; San Diego, CA, USA). A subset of the Beat AML cohort (n=177) was used for validation [20]. The same bioinformatics analysis was used for both datasets and is described in the supplementary. The samples were aligned to the reference genome (Ensembl GRCh37 release 87) using the STAR [21] aligner with default parameters. Splice junctions from all samples were pooled, filtered and used to create a new genomic index. Multi-sample 2-pass alignments to the re-generated genome index followed, using the STAR recommended parameters for gene-fusion detection. Read counts of transcripts and genes were measured with salmon [22]. Read counts of splice junctions were extracted from the STAR output.

Differential expression analysis and differential splice junction usage (DSJU)

A minimum expression filter was applied prior to each differential analysis. Differentially expressed isoforms were identified with the limma [23] package after TMM-normalization [24] with edgeR [25] and weighting with voom [26,27]. A surrogate variable analysis step using the sva [28] package was included to reduce unwanted technical noise. DSJU was quantified similarly using the diffSplice function of the limma package. Both analyses are described in detail in the supplementary.

Nanopore cDNA sequencing and analysis

Total RNA was transcribed into cDNA using the TeloPrime Full-Length cDNA Amplification Kit (Lexogen) which is highly selective for polyadenylated full-length RNA molecules with 5'-cap structures. Two barcoded samples for multiplexed analysis were sequenced on the Oxford Nanopore Technologies MinION platform. Alternative isoform analysis was performed with FLAIR [29].

Statistics

Statistical analysis was performed using the R-3.4.1 [30] software package. Correlations between variables were performed using the Mann-Whitney U test and the Pearson's chi-squared test. In case of multiple testing, p-value adjustment was performed as described in the supplementary. Survival analysis was performed and visualized using the Kaplan-Meier method and the log-rank test was utilized to capture differences in relapse free survival (RFS) and overall survival (OS). Patients receiving an allogeneic stem cell transplant were censored at the day of the transplant, for both RFS and OS. Additionally, Cox regressions were performed for all available clinical parameters and recurrent aberrations. Cox multiple regression models were then built separately for RFS and OS, using all variables with an unadjusted p-value < 0.1 in the single Cox regression models.

Results

Clinical features of AML patients with SF mutations

We characterized SF mutations in two independent patient cohorts (the AMLCG and AMLSG cohorts). Our primary cohort (AMLCG) consisted of 1119 AML patients (Figure S1), 236 (21.1%) of which presented with SF mutations. The three most commonly affected SF genes, *SRSF2*, *U2AF1* and *SF3B1* were mutated in 12.1% (n=136), 3.4% (n=38) and 4.1% (n=46) of the patients (Figure 1A). In agreement with previous findings [19], SF mutations were in their majority mutually exclusive, heterozygous hotspot mutations (Figure 1B). The four most common point mutations were *SRSF2*(P95H) (n=69), *SRSF2*(P95L) (n=27), *U2AF1*(S34F) (n=18), and *SF3B1*(K700E) (n=18) mutations (Figure 1C-E). The clinical characteristics of patients harboring SF mutations are summarized in Tables 1 and S1 (AMLSG cohort), along with a statistical assessment between cohorts (Table S2). We observed a high overall degree of similarity regarding clinical features of SF mutated patients between the AMLCG and AMLSG cohorts, despite their large median age difference. Mutations in *SRSF2*, *U2AF1* and *SF3B1* occurred more frequently in secondary AML (44.7% compared to 18.2% in *de novo* AML) and were all associated with older age. As reported previously [1], *SRSF2* and *U2AF1* mutated patients were predominantly male (76.7% and 76.3%, respectively). Furthermore, patients harboring *SRSF2* mutations presented with a lower white blood cell count (WBC; median 13.3 10^9/L vs. 22.4 10^9/L) while *U2AF1* mutated patients presented with a reduced blast percentage in their bone marrow when compared to SF wildtype patients (median 60% vs 80%).

Associations of SF mutations and other recurrent alterations in AML

In a second step, we investigated associations between SF mutations and recurrent cytogenetic abnormalities and gene mutations in AML (Figure 2). Notably, SF mutations were not found in inv(16)/t(16;16) patients (n=124), with the exception of one inv(16)/t(16;16) patient harboring a

U2AF1(R35Q) mutation. The same held true for t(8;21) patients (n=98), where only one patient

had a rare deletion in *SRSF2*. Additionally, all patients in the AMLCG cohort presenting with an

isolated trisomy 13 (n=9) also harbored an *SRSF2* mutation ($p<0.001$), as described previously

[31].

Mutations in all SF genes correlated positively with mutations in *BCOR* and *RUNX1* and

negatively with mutations in *NPM1*. Expectedly, *SRSF2*(P95H) and *SRSF2*(P95L) mutations

shared a similar pattern of co-expression including significant pairwise associations with

mutations in *ASXL1*, *IDH2*, *RUNX1* (both $p<0.001$) and *STAG2* ($p<0.001$ and $p=0.002$,

respectively). However, apart from *IDH2* mutations where co-occurrence was comparable (OR:

3.4 vs 5.1), mutations in *ASXL1*, *RUNX1* and *STAG2* coincided more frequently with

SRSF2(P95H) mutations. Despite this, *SRSF2*(P95L) mutations showed a slightly increased

co-occurrence with other recurrent AML mutations (median 5 vs 4 mutations, $p=0.046$).

Prognostic relevance of SF mutations for relapse-free survival and overall survival

The prognostic impact of *SRSF2*, *U2AF1* and *SF3B1* mutations was initially assessed using

Kaplan-Meier graphs and log-rank testing. All SF mutations presented with both inferior relapse-

free survival (RFS) and overall survival (OS) compared to SF wildtype patients (Figures S3.1A-

C and S3.2). The effect was most pronounced in *U2AF1* mutated patients with an one-year

survival rate of only 29.1%, followed by *SF3B1* (40.6%) and *SRSF2* mutated patients (49.2%).

Different point mutations inside the same SF gene did not differ significantly in their effect on

OS.

To confirm the observed prognostic impact of SF mutations we performed single Cox

regressions on all available clinical and genetic parameters. In agreement with the Kaplan-

Meier estimates, patients harboring *SRSF2*(P95H), *SRSF2*(P95L), *U2AF1*(S34F) and

SF3B1(K700E) mutations had significantly reduced RFS and OS (Figures S3.1D and S4.1). To

test whether any SF mutation was an independent prognostic marker, multiple Cox regression models (Figure 3) were built by integrating all parameters significantly associated (p < 0.1) with RFS and OS in the single Cox regression models. Along with several known predictors, only *U2AF1*(S34F) mutations presented with prognostic relevance for both RFS (Hazard Ratio=2.81, p=0.012) and OS (HR=1.90, p=0.034) in the AMLCG cohort, but not in the AMLSG cohort. However, when aggregating mutations at the gene level, mutations in *SRSF2* and *SF3B1* presented with prognostic relevance for RFS in the AMLSG cohort (HR=1.77, p=0.008; HR=2.15, p=0.014; respectively), while not reaching significance in the AMLCG cohort (Table S4.2). When looking only at de novo AML patients, the prognostic impact of *U2AF1*(S34F) mutations diminished, yet the prognostic impact observed for *SRSF2* and *SF3B1* remained significant in the AMLSG cohort (HR=1.84, p=0.009; HR=2.43, p=0.015; respectively) (Tables S4.1-S4.2).

Differential isoform expression in SF mutated patients

We next assessed the impact of SF mutations on mRNA expression. To this end, whole-transcriptome RNA-sequencing was performed on 246 AML patients, 29 of which harbored a mutation in the SF genes of interest, while 199 SF wildtype patients were used as a control (Figure S2). The remaining patients either presented with a different SF mutation (n=17) or exhibited more than one SF mutation (n=1) and were excluded. In addition, a subset of the Beat AML cohort (n=177) with matched DNA- and RNA-sequencing data was used for validation [20].

After low-coverage filtering we performed a differential isoform expression analysis for ~90 000 isoforms. Differential expression was restricted to a small fraction of all expressed isoforms (<0.5%; Figure 4A and Table S6.1). Little overlap of differentially expressed (DE) isoforms was found when different SF mutation groups were compared to the control, consistent with previous

observations [33]. However, 10 isoforms were reported as DE in both *SRSF2*(P95H) and

SRSF2(P95L) mutated samples, all with the same fold-change direction (Figure 4B). Out of

those, the isoforms in *GTF2I*, *H1F0*, *INHBC*, *LAMC1* and one of the isoforms of *METTL22*

(*ENST00000562151*) were also significant in the validation cohort for both *SRSF2*(P95H) and

SRSF2(P95L). Additionally, the isoform of *H1F0* was also reported as DE for *U2AF1*(S34F)

mutants in both cohorts. For *SRSF2*(P95H) mutants 107 of all DE isoforms also reached

significance in the validation cohort (40.1%), while for the other SF mutation subgroups

validation rates ranged from 15.1 to 27.3% increasing with larger mutant sample sizes. Notably,

mutated and wildtype samples showed large differences in the expression levels of several

isoforms (Figure 4C and Figure S5). The top two overexpressed isoforms in *SRSF2*(P95H) both

corresponded to *INTS3*, which was recently reported as dysregulated in *SRSF2*(P95H) mutants

co-expressing *IDH2* mutations [32]. Several DE isoforms identified in SF mutated patients

correspond to cancer-related genes, many of which have a known role in AML. Specifically,

genes with DE isoforms included, but were not limited to *BRD4* [34], *EWSR1* [35] and *YBX1*

[36] in *SRSF2*(P95H) mutated samples, *CUX1* [37], *DEK* [15,38] and *EZH1* [39] in

U2AF1(S34F) mutated samples, as well as *PTK2* [40] in *SF3B1*(K700E) mutated patients

(Tables S7.1-S7.2).

Hierarchical clustering using DE isoforms was performed for all samples to assess the

expression homogeneity of SF mutations. A tight clustering of samples harboring identical SF

point mutations was observed, indicating an isoform expression profile highly characteristic for

each individual SF mutation (Figures S6.1-S6.3). When using DE isoforms resulting from the

comparison of all *SRSF2* mutated samples against SF wildtype samples, the samples did not

cluster as well. This stands in agreement with the limited overlap of differentially expressed

isoforms found between the two *SRSF2* point mutations examined and suggests at least some

heterogeneity among them. The same also held true for *U2AF1* mutated samples, however all *SF3B1* mutated samples still clustered together when compared as a single group to the control.

Differential splicing in SF mutants

Previous studies have reported differential splicing as causal for isoform dysregulation in SF mutants [41,42]. To detect aberrant splicing in our dataset, we quantified the usage of all unique splice junctions by pooling information from all samples (Figure S7). After filtering out junctions with low expression, 235 730 junctions were assigned to genes. Only junctions within an annotated gene were considered, leading to the exclusion of 11 617 (4.9%) junctions. Junctions present in genes with low-expression or genes with a single junction were excluded, leaving 221 249 unique junctions (19.3% novel) across 15 526 genes (Table S6.1). Applying the same workflow to the Beat AML cohort yielded 194 158 junctions (8.3% novel). Notably, of the 172 518 junctions shared across both datasets, 10 029 (5.8%) were novel. The novel junctions passing our filtering criteria were supported by a high amount of reads and samples with a distribution comparable to that of annotated junctions (Figure 5A). Neither the number of novel junctions nor the number of reads supporting them correlated with the presence of SF mutations, suggesting that novel splicing events are not increased in SF mutants.

In consideration of the high proportion of novel junctions in both datasets, we employed a customized pipeline that can quantify the differential splice junction usage (DSJU) of each individual junction, by harnessing usage information from all junctions inside one gene. Of the several hundred junctions reported as differentially used in our primary cohort ($p<0.05$, $\log_2$(fold change) > 1), 20.2-45.9% constituted novel junctions (Tables S7.1-S7.3 and Tables S9.1-S9.2) and were classified according to their relationship with known acceptor and donor sites as described previously (Figure 5B) [15]. Unsurprisingly, validation rates increased with larger

mutant sample sizes, ranging from 9.3% (*SF3B1*(K700E); n = 3) to 74.0% (all *SRSF2* mutants;

n = 26). Furthermore, validation rates were higher for novel junctions (mean 39.3% vs. 21.5%

known junctions), likely due to the stricter initial filtering criteria applied. By performing nanopore

sequencing of one *SRSF2*(P95H) mutant and one SF wildtype sample we were able to confirm

the usage of several novel junctions and detect resulting novel isoforms as exemplified for

IDH3G in Figure 6A-D. A tendency towards decreased junction usage was observed for all SF

point mutations and was most evident in *SF3B1*(K700E) mutants (1 423/1 927; 73.9% of

differentially used junctions). The total number of splicing events, however, was not reduced in

SF mutants (mean 9,275,359 events vs. 9 192 697 in wildtype patients), suggesting that

decreased splicing is limited to selected junctions rather than being a global effect.

We systematically compared the genes with at least one DE isoform and those reported as

differentially spliced in all SF mutation subgroups (Tables S9.3-S9.4). For *SRSF2* mutants,

genes significant in both analyses included *EWSR1*, *H1F0*, *INTS3* and *YBX1*. In general, out

of the genes examined in both analyses only 9.8-23.3% (depending on the SF mutation) of

genes reported as having a DE isoform were also reported as being differentially spliced.

Conversely, 3.3-28.5% of differentially spliced genes were also reported as having a DE

isoform. These findings suggest that differential gene splicing does not always lead to altered

isoform expression while at the same time differential isoform expression cannot always be

attributed to an explicit splicing alteration. Considering the complementary nature of the

analyses, we performed gene ontology (GO) analysis by combining the genes with evidence of

differential isoform usage or differential splicing (Tables S10.1-S10.7). Interestingly, GO terms

enriched for both *SRSF2* mutants included "mRNA splicing, via spliceosome" (p<0.001 and

p=0.046, respectively) and "mRNA splice site selection" (p=0.022 and p=0.019, respectively)

(Figure 5C).

In an additional step, the splice junction counts reported by *Okeyo-Owuor et al.* were used to

detect DSJU between CD34+ cells with U2AF1(S34) mutations (n=3) and SF wildtype (n=3)

via the same pipeline applied to the AMLCG and Beat AML cohorts. While no identical junctions

were differentially used in all three datasets, 16 genes were reported as differentially spliced in

all, including leukemia or cancer-associated genes (*ABI1*, *DEK*, *HP1BP3*, *MCM3* and *SET*), as

well as *HNRNPK* (a major pre-mRNA binding protein), thereby further refining our list of genes

with strong evidence of differential splicing between *U2AF1*(S34F) mutants and SF wildtype

samples (Table S11).

The effect of SF mutations on the splicing pathway

It is well established that SF mutations dysregulate splicing by changing RNA binding affinities

or altering 3' splice site recognition of the corresponding splicing factors. However, SRSF2 also

plays a splicing-independent role in transcriptional pausing by translocating the positive

transcription elongation factor complex (P-TEFb) from the 7SK complex to RNA polymerase II

[43]. A recent study reported that mutant SRSF2(P95H) enhances R-loop formation due to

impaired transcriptional pause release, thus providing evidence that altered splicing does not

account for the entirety of the *SRSF2*(P95H) mutant phenotype [33]. In another study no

difference was found in the total mRNA of 12 Serine/arginine rich splicing factors and 14 out of

16 major heterogeneous nuclear ribonucleoprotein (hnRNP) splicing factors between

SRSF2(P95H) mutant and WT CRISPR clones [44].

We cross-referenced our differential expression and differential splicing analysis results with a

list of all genes involved in splicing (GO:0000398; mRNA splicing, via spliceosome). Of the 347

splicing-related genes (317 of which were expressed in our dataset) 101 were dysregulated in

at least one SF mutant group. On average 30.5 (range 6-52) splicing-related genes were

dysregulated per SF point mutation. Of note, both *SRSF2* point mutations associated with

differential splicing of *HNRNPA1* and *HNRNPUL1*, as well as *PCF11* and *TRA2A*. A query of

the STRING database suggests protein-protein interactions between SRSF2 and the proteins

of the above genes, which are also interconnected (Figure 6E). Interestingly, one of the

differential splicing events reported in both *SRSF2* mutants involves the under-usage of the

same novel splice junction in *TRA2A* (Figure 6F). *TRA2A* has previously been shown to be

differentially spliced in mouse embryo fibroblasts upon *SRSF2* knockout [45]. Furthermore, it

has been shown that both HNRNPA1 and SRSF2 interact with the loop 3 region of 7SK RNA

and by favoring the dissociation of SRSF2, HNRNPA1 may lead to the release of active P-TEFb

[46]. Taken together, our results indicate a strong dysregulation of the splicing pathway in SF

mutants including several genes whose gene products closely interact with SRSF2.

324
325

Discussion

The clinical relevance of SF mutations and their aberrant splicing patterns have been explored in myelodysplasia, while comparable data for AML is lacking. In this study we examined two AML patient cohorts, encompassing a total of 2678 patients from randomized prospective trials, to characterize SF mutations clinically. This analysis was complemented by RNA-sequencing analysis of two large datasets to reveal targets of aberrant splicing in AML.

We show that SF mutations are frequent alterations in AML, identified in 21.4% of our primary patient cohort, especially in elderly patients and in secondary AML. SF mutations are associated with other recurrent mutations in AML, such as *BCOR* and *RUNX1* mutations, however *SRSF2*(P95L) mutations co-occur less often with those mutations when compared to *SRSF2*(P95H) mutations, albeit showing a slightly increased mutational load. This suggests a more diverse co-expression profile of *SRSF2*(P95L).

Previous studies have demonstrated the predictive value of SF mutations in clonal haematopoiesis of indeterminate potential (CHIP) [47], MDS [6,8,48–50] and AML [10,18,19,51]. However, survival analyses in AML were, in their majority, hampered by small sample sizes and limited availability of further risk factors. Therefore, we examined whether SF mutations impact survival while accounting for recently proposed risk parameters included in the ELN 2017 classification [52]. In our analysis, *SRSF2* and *SF3B1* mutations were no independent prognostic markers for OS in AML. *U2AF1*(S34F) mutations displayed poor OS in the AMLCG cohort, which we were unable to validate in the AMLSG cohort. The discrepancy in survival of SF mutated patients between the two cohorts lied most likely in the large age difference of the participants (median age difference of 8 years), which also led to a higher percentage of patients receiving allogeneic transplants in the AMLSG cohort (56.5% vs. 30.6% in the AMLCG cohort). In summary, SF mutations are early evolutionary events and define

prognosis and transformation risk in CHIP and MDS patients, yet there is no clear independent

prognostic value of SF mutations in AML.

Two large RNA-sequencing studies have been performed previously, to detect aberrantly

spliced genes in SF mutants, both of which focused on MDS patients [41,42]. In this study we

described a distinct differential isoform expression profile for each SF point mutation.

Furthermore, we evaluated differential splicing for the four most common SF point mutations

via a customized pipeline to determine differential usage of both known and novel splice

junctions. Our pipeline enables the differential quantification of individual splice junctions

without restricting the analysis to annotated alternative splicing events. We argue that the

strength of our analysis lies in the accurate detection of single dysregulated junctions

(especially in cases where splice sites are shared by multiple junctions) in an annotation-

independent manner achieving validation rates up to 74.0% in our largest mutant sample group

(*SRSF2*, n=19). Limitations of the analysis include the restriction to junctions with both splice

sites within the same gene (a restriction shared by most differential splicing algorithms) and

genes with at least two junctions. However, the reduced requirements of our analysis could

prove valuable in the study of differential splicing in organisms with lacking annotation.

All SF point mutations shared a tendency towards decreased splice junction usage, which did

not affect the global number of splicing events in SF mutants. Surprisingly, we observed a

limited overlap between genes with differentially expressed isoforms and differentially spliced

genes. In addition, a recent study by *Liang et al.* reported that the majority of differential binding

events in *SRSF2*(P95H) mutants do not translate to alternative splicing [53]. Taken together,

these findings indicate a "selection" or possibly a compensation of deregulatory events from

differential binding through differential splicing to finally differential isoform expression.

Furthermore, the enrichment of aberrant splicing in splicing-related genes opens the possibility

of a cascading effect on transcription via the differential alternative splicing of transcriptional

components. A congruent hypothesis was stated by *Liang et al.*, where an enrichment of

SRSF2(P95H) targets in RNA processing and splicing was shown, further supporting the notion

of an indirect effect of mutant SRSF2 facilitated through additional splicing components. Further

investigations may provide a mechanistic link between the differential splicing of selected genes

and the impairment of transcription and specifically transcriptional pausing observed in SF

mutant cells, which contributes to the MDS phenotype [33].

To the best of our knowledge our study represents the most comprehensive analysis of SF

mutations in AML to date, both in terms of clinical and functional characterization. This enabled

us to study *SRSF2*(P95H) and *SRSF2*(P95L) separately, thereby not only outlining their

differences but also identifying common and likely core targets of differential splicing in *SRSF2*

mutants. We conclude that SF mutated patients represent a distinct subgroup of AML patients

with poor prognosis that is not attributable solely to the presence of SF mutations. SF mutations

induce aberrant splicing throughout the genome including the dysregulation of several genes

associated with AML pathogenesis, as well as a number of genes with immediate, functional

implications on splicing and transcription. Further studies are required to identify which splicing

events are critical in leukaemogenesis and whether they are accessible to new treatments

options, such as splicing inhibitors [54] and immunotherapeutic approaches.

Acknowledgments

The authors thank all participants and recruiting centers of the AMLCG, BEAT and AMLSG trials.

Funding

This work is supported by a grant of the Wilhelm-Sander-Stiftung (no. 2013.086.2) and the Physician Scientists Grant (G-509200-004) from the Helmholtz Zentrum München to T.H. and the German Cancer Consortium (Deutsches Konsortium für Translationale Krebsforschung, Heidelberg, Germany). K.H.M., K.S. and T.H. are supported by a grant from Deutsche Forschungsgemeinschaft (DFG SFB 1243, TP A06 and TP A07). S.K.B. is supported by Leukaemia & Blood Cancer New Zealand and the family of Marijanna Kumerich. A.M.N.B. is supported by the BMBF grant 01ZZ1804B (DIFUTURE).

Author Contributions

S.A.B., A.M.N.B. and T.H. conceived and designed the analysis. S.A.B., A.M.N.B., V.J., M.R.-T., H.J., A.G., S.C., N.K., K.S., K.H.M and T.H. provided and analyzed data. A.M.N.B., V.J. and U.M. provided bioinformatics support. J.P.-M., S.K. and H.B. managed the Genome Analyzer IIx platform and the RNA-sequencing of the AMLCG samples. M.R.-T., H.J., B.K., S.S., N.K., S.K.B., K.H.M. and K.S. characterized patient samples; M.C.S., D.G., W.B., B.W., J.B. and W.H. coordinated the AMLCG clinical trials. S.A.B. and T.H. wrote the manuscript. All authors approved the final manuscript.

Additional Information

The authors declare no conflicts of interest.

References

1. Papaemmanuil E, Gerstung M, Malcovati L, Tauro S, Gundem G, Van Loo P, et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. Blood. 2013;122(22):3616–27.

2. Metzeler KH, Herold T, Rothenberg-Thurley M, Amler S, Sauerland MC, Görlich D, et al. Spectrum and prognostic relevance of driver gene mutations in acute myeloid leukemia. Blood [Internet]. 2016 Aug 4 [cited 2018 May 23];128(5):686–98. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27288520

3. Makishima H, Visconte V, Sakaguchi H, Jankowska AM, Kar SA, Jerez A, et al. Mutations in the spliceosome machinery, a novel and ubiquitous pathway in leukemogenesis. Blood [Internet]. 2012;119(14):3203–10. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3321850/

4. Larsson CA, Cote G, Quintás-Cardama A. The changing mutational landscape of acute myeloid leukemia and myelodysplastic syndrome. Mol Cancer Res [Internet]. 2013 Aug [cited 2016 Oct 25];11(8):815–27. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23645565

5. Yoshida K, Sanada M, Shiraishi Y, Nowak D, Nagata Y, Yamamoto R, et al. Frequent pathway mutations of splicing machinery in myelodysplasia. Nature [Internet]. 2011 Sep 11 [cited 2016 Oct 23];478(7367):64–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21909114

6. Thol F, Kade S, Schlarmann C, Löffeld P, Morgan M, Krauter J, et al. Frequency and prognostic impact of mutations in SRSF2, U2AF1, and ZRSR2 in patients with myelodysplastic syndromes. Blood. 2012;119(15):3578–84.

7.   Dolatshad H, Pellagatti A, Fernandez-Mercado M, Yip BH, Malcovati L, Attwood M, et

al. Disruption of SF3B1 results in deregulated expression and splicing of key genes and

pathways in myelodysplastic syndrome hematopoietic stem and progenitor cells.

Leukemia [Internet]. 2015 May [cited 2018 May 3];29(5):1092–103. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25428262

8.   Wu S, Kuo Y, Hou H, Li L, Tseng M, Huang C, et al. The clinical implication of SRSF2

mutation in patients with myelodysplastic syndrome and its stability during disease

evolution. Blood [Internet]. 2014;120(15):3106–12. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/22932795

9.   Graubert TA, Shen D, Ding L, Okeyo-owuor T, Cara L, Shao J, et al. Recurrent

Mutations in the U2AF1 Splicing Factor In Myelodysplastic Syndromes. Nat Genet

[Internet]. 2012 Dec 11 [cited 2016 Oct 23];44(1):53–7. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/22158538

10.   Hou H-A, Liu C-Y, Kuo Y-Y, Chou W-C, Tsai C-H, Lin C-C, et al. Splicing factor

mutations predict poor prognosis in patients with de novo acute myeloid leukemia.

Oncotarget [Internet]. 2016 [cited 2017 Jun 19];7(8). Available from:

www.impactjournals.com/oncotarget

11.   Cho Y-U, Jang S, Seo E-J, Park C-J, Chi H-S, Kim D-Y, et al. Preferential occurrence of

spliceosome mutations in acute myeloid leukemia with a preceding myelodysplastic

syndrome and/or myelodysplasia morphology. Leuk Lymphoma [Internet]. 2014 Aug 3

[cited 2016 Oct 25];8194(November):1–25. Available from:

http://www.tandfonline.com/doi/full/10.3109/10428194.2014.995648

12.   Moon H, Cho S, Loh TJ, Jang HN, Liu Y, Choi N, et al. SRSF2 directly inhibits intron

splicing to suppresses cassette exon inclusion. BMB Rep [Internet]. 2017 Aug [cited

2018 Jan 7];50(8):423–8. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28712387

13.   Kim E, Ilagan JO, Liang Y, Daubner GM, Lee SCW, Ramakrishnan A, et al. SRSF2

Mutations Contribute to Myelodysplasia by Mutant-Specific Effects on Exon

Recognition. Cancer Cell [Internet]. 2015 May 11 [cited 2016 Oct 25];27(5):617–30.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/25965569

14.   Alsafadi S, Houy A, Battistella A, Popova T, Wassef M, Henry E, et al. Cancer-

associated SF3B1 mutations affect alternative splicing by promoting alternative

branchpoint usage. Nat Commun [Internet]. 2016 Feb 4 [cited 2016 Oct 19];7:10615.

Available from: http://www.nature.com/doifinder/10.1038/ncomms10615

15.   Okeyo-Owuor T, White BS, Chatrikhi R, Mohan DR, Kim S, Griffith M, et al. U2AF1

mutations alter sequence specificity of pre-mRNA binding and splicing. Leukemia

[Internet]. 2015 Apr [cited 2018 Jan 28];29(4):909–17. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25311244

16.   Przychodzen B, Jerez A, Guinta K, Sekeres MA, Padgett R, Maciejewski JP, et al.

Patterns of missplicing due to somatic U2AF1 mutations in myeloid neoplasms. Blood

[Internet]. 2013 Aug 8 [cited 2018 May 23];122(6):999–1006. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23775717

17.   Shirai CL, Ley JN, White BS, Kim S, Tibbitts J, Shao J, et al. Mutant U2AF1 Expression

Alters Hematopoiesis and Pre-mRNA Splicing In Vivo. Cancer Cell [Internet]. 2015 May

11 [cited 2018 May 23];27(5):631–43. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/25965570

18.   Yang J, Yao D, Ma J, Yang L, Guo H, Wen X, et al. The prognostic implication of SRSF2

486   mutations in Chinese patients with acute myeloid leukemia. Tumor Biol [Internet]. 2016

487   Aug 28 [cited 2018 May 23];37(8):10107–14. Available from:

488   http://www.ncbi.nlm.nih.gov/pubmed/26820131

489   19.   Papaemmanuil E, Gerstung M, Bullinger L, Gaidzik VI, Paschka P, Roberts ND, et al.

490   Genomic Classification and Prognosis in Acute Myeloid Leukemia. N Engl J Med

491   [Internet]. 2016 Jun 9 [cited 2019 Jan 18];374(23):2209–21. Available from:

492   http://www.ncbi.nlm.nih.gov/pubmed/27276561

493   20.   Tyner JW, Tognon CE, Bottomly D, Wilmot B, Kurtz SE, Savage SL, et al. Functional

494   genomic landscape of acute myeloid leukaemia. Nature [Internet]. 2018 Oct 17 [cited

495   2019 Sep 8];562(7728):526–31. Available from: http://www.nature.com/articles/s41586-

496   018-0623-z

497   21.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

498   universal RNA-seq aligner. Bioinformatics [Internet]. 2013 Jan 1 [cited 2017 May

499   8];29(1):15–21. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23104886

500   22.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

501   aware quantification of transcript expression. Nat Methods [Internet]. 2017 Apr [cited

502   2018 Jan 29];14(4):417–9. Available from:

503   http://www.ncbi.nlm.nih.gov/pubmed/28263959

504   23.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential

505   expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res

506   [Internet]. 2015 Apr 20 [cited 2017 May 8];43(7):e47–e47. Available from:

507   http://www.ncbi.nlm.nih.gov/pubmed/25605792

24. Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. Genome Biol [Internet]. 2010 [cited 2017 May 8];11(3):R25.

Available from: http://download.springer.com/static/pdf/905/art%253A10.1186%252Fgb-

2010-11-3-

r25.pdf?originUrl=http%3A%2F%2Fgenomebiology.biomedcentral.com%2Farticle%2F1

0.1186%2Fgb-2010-11-3-

r25&token2=exp=1494254026~acl=%2Fstatic%2Fpdf%2F905%2Fart%25253A10.1186

%25252Fgb

25. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics [Internet]. 2009 Jan

1 [cited 2018 May 23];26(1):139–40. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19910308

26. Liu R, Holik AZ, Su S, Jansz N, Chen K, Leong HS an, et al. Why weight? Modelling

sample and observational level variability improves power in RNA-seq analyses.

Nucleic Acids Res [Internet]. 2015 [cited 2017 Apr 10];43(15):e97. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4551905/pdf/gkv412.pdf

27. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model

analysis tools for RNA-seq read counts. Genome Biol [Internet]. 2014 [cited 2017 Feb

12];15(2):R29. Available from:

http://genomebiology.biomedcentral.com/articles/10.1186/gb-2014-15-2-r29

28. Leek JT. Svaseq: Removing batch effects and other unwanted noise from sequencing

data. Nucleic Acids Res [Internet]. 2014 Dec 1 [cited 2018 Jul 3];42(21):e161. Available

from: https://academic.oup.com/nar/article/42/21/e161/2903156

29.   Tang AD, Soulette CM, Baren MJ van, Hart K, Hrabeta-Robinson E, Wu CJ, et al. Full-
532   length transcript characterization of SF3B1 mutation in chronic lymphocytic leukemia
533   reveals downregulation of retained introns. bioRxiv [Internet]. 2018 Sep 6 [cited 2019
534   Dec 9];410183. Available from: https://www.biorxiv.org/content/10.1101/410183v1

535   30.   R Core Team. R: A Language and Environment for Statistical Computing [Internet].
536   Vienna, Austria: R Foundation for Statistical Computing; Available from: https://www.r-
537   project.org/

538   31.   Herold T, Metzeler KH, Vosberg S, Hartmann L, Ollig C, Olzel FS, et al. Isolated trisomy
539   13 defines a homogeneous AML subgroup with high frequency of mutations in
540   spliceosome genes and poor prognosis. 2014;124(8):1304–11.

541   32.   Yoshimi A, Lin K-T, Wiseman DH, Rahman MA, Pastore A, Wang B, et al. Coordinated
542   alterations in RNA splicing and epigenetic regulation drive leukaemogenesis. Nature
543   [Internet]. 2019 Oct 2 [cited 2019 Dec 10];574(7777):273–7. Available from:
544   http://www.nature.com/articles/s41586-019-1618-0

545   33.   Chen L, Chen J-Y, Huang Y-J, Gu Y, Qiu J, Qian H, et al. The Augmented R-Loop Is a
546   Unifying Mechanism for Myelodysplastic Syndromes Induced by High-Risk Splicing
547   Factor Mutations. Mol Cell [Internet]. 2018 Feb 1 [cited 2018 Feb 1];69(3):412-425.e6.
548   Available from: http://linkinghub.elsevier.com/retrieve/pii/S1097276517310080

549   34.   Roe J-S, Vakoc CR. The Essential Transcriptional Function of BRD4 in Acute Myeloid
550   Leukemia. Cold Spring Harb Symp Quant Biol [Internet]. 2016 [cited 2019 Sep
551   8];81:61–6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28174254

552   35.   Endo A, Tomizawa D, Aoki Y, Morio T, Mizutani S, Takagi M. EWSR1/ELF5 induces
553   acute myeloid leukemia by inhibiting p53/p21 pathway. Cancer Sci [Internet]. 2016 Dec

[cited 2019 Sep 8];107(12):1745–54. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27627705

36.   Perner F, Jayavelu AK, Schnoeder TM, Mashamba N, Mohr J, Hartmann M, et al. The

Cold-Shock Protein Ybx1 Is Required for Development and Maintenance of Acute

Myeloid Leukemia (AML) in Vitro and In Vivo. Blood [Internet]. 2017 [cited 2019 Sep

8];130(Suppl 1). Available from:

http://www.bloodjournal.org/content/130/Suppl_1/792?sso-checked=true

37.   McNerney ME, Brown CD, Wang X, Bartom ET, Karmakar S, Bandlamudi C, et al.

CUX1 is a haploinsufficient tumor suppressor gene on chromosome 7 frequently

inactivated in acute myeloid leukemia. Blood [Internet]. 2013 Feb 7 [cited 2019 Sep

8];121(6):975–83. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23212519

38.   McGarvey T, Rosonina E, McCracken S, Li Q, Arnaout R, Mientjes E, et al. The acute

myeloid leukemia-associated protein, DEK, forms a splicing-dependent interaction with

exon-product complexes. J Cell Biol [Internet]. 2000 Jul 24 [cited 2019 Sep

8];150(2):309–20. Available from: http://www.ncbi.nlm.nih.gov/pubmed/10908574

39.   Fujita S, Honma D, Adachi N, Araki K, Takamatsu E, Katsumoto T, et al. Dual inhibition

of EZH1/2 breaks the quiescence of leukemia stem cells in acute myeloid leukemia.

Leukemia [Internet]. 2018 Apr 27 [cited 2019 Sep 8];32(4):855–64. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28951561

40.   Pallarès V, Hoyos M, Chillón MC, Barragán E, Prieto Conde MI, Llop M, et al. Focal

Adhesion Genes Refine the Intermediate-Risk Cytogenetic Classification of Acute

Myeloid Leukemia. Cancers (Basel) [Internet]. 2018 Nov 13 [cited 2019 Sep 8];10(11).

Available from: http://www.ncbi.nlm.nih.gov/pubmed/30428571

41.   Shiozawa Y, Malcovati L, Gallì A, Sato-Otsubo A, Kataoka K, Sato Y, et al. Aberrant

splicing and defective mRNA production induced by somatic spliceosome mutations in

myelodysplasia. Nat Commun [Internet]. 2018 [cited 2019 Jan 21];9(1):3649. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/30194306

42.   Pellagatti A, Armstrong RN, Steeples V, Sharma E, Repapi E, Singh S, et al. Impact of

spliceosome mutations on RNA splicing in myelodysplasia: Dysregulated

genes/pathways and clinical associations. Blood [Internet]. 2018 Sep 20 [cited 2019

Feb 6];132(12):1225–40. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29930011

43.   Ji X, Zhou Y, Pandit S, Huang J, Li H, Lin CY, et al. SR Proteins Collaborate with 7SK

and Promoter-Associated Nascent RNA to Release Paused Polymerase. Cell [Internet].

2013 May 9 [cited 2019 Nov 12];153(4):855–68. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23663783

44.   Zhang J, Lieu YK, Ali AM, Penson A, Reggio KS, Rabadan R, et al. Disease-associated

mutation in SRSF2 misregulates splicing by altering RNA-binding affinities. Proc Natl

Acad Sci [Internet]. 2015 Aug 25 [cited 2017 May 31];112(34):E4726–34. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/26261309

45.   Skrdlant L, Stark JM, Lin R-J. Myelodysplasia-associated mutations in serine/arginine-

rich splicing factor SRSF2 lead to alternative splicing of CDC25C. BMC Mol Biol

[Internet]. 2016 [cited 2019 Nov 12];17(1):18. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27552991

46.   Lemieux B, Blanchette M, Monette A, Mouland AJ, Wellinger RJ, Chabot B. A Function

for the hnRNP A1/A2 Proteins in Transcription Elongation. Caputi M, editor. PLoS One

[Internet]. 2015 May 26 [cited 2019 Nov 12];10(5):e0126654. Available from:

https://dx.plos.org/10.1371/journal.pone.0126654

47.   Abelson S, Collord G, Ng SWK, Weissbrod O, Mendelson Cohen N, Niemeyer E, et al.

Prediction of acute myeloid leukaemia risk in healthy individuals. Nature [Internet]. 2018

Jul 9 [cited 2019 Feb 14];559(7714):400–4. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/29988082

48.   Malcovati L, Papaemmanuil E, Bowen DT, Boultwood J, Della Porta MG, Pascutto C, et

al. Clinical significance of SF3B1 mutations in myelodysplastic syndromes and

myelodysplastic/myeloproliferative neoplasms. Blood [Internet]. 2011 Dec 8 [cited 2019

Feb 24];118(24):6239–46. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21998214

49.   Papaemmanuil E, Cazzola M, Boultwood J, Malcovati L, Vyas P, Bowen D, et al.

Somatic SF3B1 mutation in myelodysplasia with ring sideroblasts. N Engl J Med

[Internet]. 2011 Oct 13 [cited 2019 Feb 24];365(15):1384–95. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21995386

50.   Wu L, Song L, Xu L, Chang C, Xu F, Wu D, et al. Genetic landscape of recurrent

ASXL1, U2AF1, SF3B1, SRSF2, and EZH2 mutations in 304 Chinese patients with

myelodysplastic syndromes. Tumor Biol [Internet]. 2016 Apr 28 [cited 2019 Feb

24];37(4):4633–40. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26508027

51.   Zhang S-J, Rampal R, Manshouri T, Patel J, Mensah N, Kayserian A, et al. Genetic

analysis of patients with leukemic transformation of myeloproliferative neoplasms

shows recurrent SRSF2 mutations that are associated with adverse outcome. Blood

[Internet]. 2012 May 10 [cited 2019 Jan 31];119(19):4480–5. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/22431577

52.   Döhner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Büchner T, et al.

Diagnosis and management of AML in adults: 2017 ELN recommendations from an

international expert panel [Internet]. Vol. 129, Blood. 2017 [cited 2019 Feb 14]. p. 424–

47. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27895058

53.   Liang Y, Tebaldi T, Rejeski K, Joshi P, Stefani G, Taylor A, et al. SRSF2 mutations drive

oncogenesis by activating a global program of aberrant alternative splicing in

hematopoietic cells. Leukemia. 2018;

54.   Lee SCW, Abdel-Wahab O. Therapeutic targeting of splicing in cancer [Internet]. Vol.

22, Nature Medicine. NIH Public Access; 2016 [cited 2019 Jan 31]. p. 976–86. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/27603132

Table 1: Clinical characteristics of SF mutations in the AMLCG cohort.

Variables	SF wildtype	*SRSF2*	*P*	*U2AF1*	*P*	*SF3B1*	*P*
No. of patients	903	133	-	38	-	45	-
Age, years, median (range)	55 (18-86)	65 (25-80)	**<0.001**	64 (23-74)	**0.007**	65 (31-78)	**0.001**
Female sex, no. (%)	490 (54.3)	31 (23.3)	**<0.001**	9 (23.7)	**0.003**	19 (42.2)	0.387
Hemoglobin, g/dL, median (range)	9 (3.5-16)	8.9 (3.8-14.7)	0.466	9.2 (6-13.6)	1.000	8.9 (6.8-13.4)	1.000
WBC count, 10^9/L, median (range)	22.4 (0.1-798.2)	13.3 (0.5-406)	**0.008**	7 (0.7-666)	0.079	22.4 (0.9-269.5)	1.000
Platelets, 10^9/L, median (range)	55 (0-1760)	49.5 (0-643)	0.736	47 (11-132)	0.466	67 (5-585)	0.744
LDH, U/L, median (range)	448 (76-19624)	362 (150-14332)	0.118	346 (128-3085)	0.313	472 (142-7434)	0.950
BM Blasts, %, median (range)	80 (6-100)	76 (15-100)	0.455	60 (10-95)	**0.002**	70 (13-95)	0.206
Performance Status (ECOG) > 1, no. (%)	157 (25.9)	18 (26.1)	1.000	8 (27.6)	1.000	4 (16)	0.941
primary AML, no. (%)	786 (87)	100 (75.2)	**0.016**	24 (63.2)	**0.003**	29 (64.4)	**0.002**
secondary AML, no (%)	69 (7.6)	30 (22.6)	**<0.001**	13 (34.2)	**<0.001**	11 (24.4)	**0.021**
therapy-related AML, no (%)	48 (5.3)	3 (2.3)	0.507	1 (2.6)	1.000	5 (11.1)	0.380
Allogeneic transplant, no. (%)	296 (32.8)	27 (20.3)	**0.022**	7 (18.4)	0.273	12 (26.7)	0.797
Complete Remission, no. (%)	641 (71)	71 (53.4)	**<0.001**	18 (47.4)	**0.015**	19 (42.2)	**0.001**
Relapse, no (%)	366 (63.5)	46 (82.1)	**0.034**	11 (78.6)	0.713	14 (87.5)	0.266
Deceased, no (%)	575 (63.7)	112 (84.2)	**<0.001**	34 (89.5)	**0.010**	41 (91.1)	**0.002**

637 WBC: white blood cells, LDH: lactate dehydrogenase, BM: bone marrow, ECOG: Eastern Co-

638 operative Oncology Group performance score

 Figure Legends

Figure 1: Frequency and location of SF mutations. (A) Distribution of SF mutations in the

AMLCG cohort. (B) Variant allele frequency of SF mutations in both study cohorts. (C) Mutation

plots showing the protein location of all SF mutations in the genes *SRSF2*, *U2AF1*, and *SF3B1*

for all patients of the AMLCG cohort. Number of patients harboring SF mutations are

additionally provided for both cohorts.

Figure 2: Correlations between SF mutations and recurrent abnormalities. Correlation

matrix depicting the co-occurrence of SF mutations and recurrent mutations in AML (A), as well

as cytogenetic groups, as defined in the ELN 2017 classification (B). Only variables with a

frequency >1% in the AMLCG cohort are shown. FLT3-ITD: *FLT3* internal tandem duplication

mutation; FLT3-TKD: *FLT3* tyrosine kinase domain mutation; CN-AML: cytogenetically normal

AML

Figure 3: Multiple Cox regression models for overall survival. Multiple Cox regression

models (for OS) were performed separately for patients in the AMLCG cohort (on the left) and

AMLSG cohort (on the right). The models include all variables with $p < 0.1$ in the single Cox

regression models of the primary cohort (AMLCG cohort). Significant p-values (<0.05) are

marked with a star. CN-AML: cytogenetically normal AML; LDH: lactate dehydrogenase; pAML:

primary AML; sAML: secondary AML; tAML: therapy-related AML; WBC: white blood cells

Figure 4: Differential isoform expression analysis in the AMLCG cohort. (A) Number of

differentially expressed isoforms for each SF point mutation. (B) Isoforms reported as

differentially expressed in the same direction among patients with different SF mutations vs. SF

wildtype patients. HGNC symbols of the genes in which the common isoforms are located are

shown. The corresponding Ensembl isoform identifiers are: 1. ENST00000340857, 2.

ENST00000481621, 3. ENST00000309668, 4. ENST00000258341, 5. ENST00000426532, 6.

ENST00000562151, 7. ENST00000564133, 8. ENST00000368817, 9. ENST00000566524,

10. ENST00000530211. (C) Volcano plots showing the magnitude of differential isoform

expression for *SRSF2*(P95H) and *SRSF2*(P95L). The x-axis corresponds to the $\log_2$(fold

change) of each isoform between mutated and wildtype samples, while the y-axis corresponds

to $-\log 10$(FDR), where FDR represents the adjusted p-value for each isoform (False Discovery

Rate).

Figure 5: Differential splice junction usage. (A) Scatterplot displaying the number of samples

as well as the total number of reads supporting each splice junction, separately for known and

novel splice junctions in both RNA-Seq datasets. To preserve visibility 20000 random junctions

are shown for each group. (B) Barchart showing the annotation status of splice junctions

reported as differentially used. Novel splice junctions where classified into 5 groups based on

their annotation status as described previously[15] ("DA": annotated junctions, "NDA": unknown

combination of known donor and acceptor sites, "D": known donor, but novel acceptor site, "A":

novel donor, but known acceptor site, "N": previously unknown donor and acceptor site). (C)

Venn diagram showing the overlap of GO terms between *SRSF2*(P95H) and *SRSF2*(P95L)

mutants for the "biogical process" domain.

Figure 6: Splicing dysregulation in *SRSF2* mutants. (A) DSJU of all splice junctions inside

IDH3G (ENSG00000067829) are shown for *SRSF2*(P95H) mutants compared to SF wildtype

patients in the AMLCG and Beat AML cohort. To determine significance the log fold-change

(logFC) of each splice junction (SJ) is compared to the logFC of all other junctions inside the

same gene. The x-axis denotes individual splice junctions defined by their chromosomal

coordinates (note the high number of splice sites shared by multiple splice junctions). (B)-(D)

Nanopore sequencing results of one *SRSF2*(P95H) mutated sample and one SF wildtype

sample of the AMLCG cohort. The yellow boxes highlight examples of exon skipping (same

exons highlighted in (C) and (D). (B) FLAIR distribution of transcripts. Only one known isoform

is expressed in the samples. Additionally, two novel isoforms were detected which are virtually

mutually exclusive in the *SRSF2*(P95H) mutant and the SF wildtype sample. (C) Exon

composition of known and novel isoforms detected (D). Sashimi plots showing the exon

sequence coverage as well as the splice junction usage of the *SRSF2*(P95H) mutated and SF

wildtype samples. The black arrow indicates the novel splice junction that is differentially used

in *SRSF2*(P95H) mutated samples compared to SF wildtype samples in both RNA-Seq

datasets shown in (A). (E) STRING plot depicting an interaction between SRSF2 and several

other splicing-related proteins including *TRA2A*. (F) Differential splice junction usage of all

splice junctions inside *TRA2A* (ENSG00000164548) are shown for *SRSF2*(P95H) and

SRSF2(P95L) mutants compared to SF wildtype patients. Annotation same as (A).

Figure 1

A

B

C

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6