A prospective evaluation of inter-rater agreement of routine medical records

audits at a large general hospital in São Paulo, Brazil.

Ana Carolina Cintra Nunes Mafra*[,1,2], João Luiz Miraglia[1], Fernando Antonio Basile

Colugnati[3], Gilberto Soares Lourenço Padilha[4], Renata Rafaella Santos Tadeucci[1,2],

Ederson Almeida[1], Mario Maia Bracco[1,2,4]

[1]Hospital Israelita Albert Einstein, São Paulo, Brasil

[2]Hospital Municipal Dr. Moysés Deutsch – M´Boi Mirim, São Paulo, Brasil

[3]Universidade Federal de Juiz de Fora, School of Medicine, Juiz de Fora, Brasil

[4]Centro de Estudos e Pesquisas Dr. João Amorim – CEJAM, São Paulo, Brasil

*Corresponding author

Avenida Brigadeiro Faria Lima, 1188, Jardim Paulistano, São Paulo – SP, Brasil CEP

01451-001; e-mail: ana.mafra@einstein.br; telephone number: 55 11 2151-0906.

Abstract

Background: The quality of the patient's medical records is strictly related to patient

safety. Besides, its data are widely used in observational studies. However, the

reliability of the information extracted from them is a matter of concern in audit

processes to ensure inter-rater agreement (IRA). Thus, the objective of this study is to

evaluate the IRA among members of the Patient's Health Record Review Board

(PHRRB), in routine auditing of medical records, and the impact of periodic discussions

of results with raters.

Methods: Prospective longitudinal study conducted between July of 2015 and April of 2016 at Hospital Municipal Dr. Moysés Deutsch, a large public hospital in São Paulo. The PHRRB was composed of 12 physicians, 9 nurses and 3 physiotherapists, who audited medical records, monthly, with the number of raters changing throughout the study. It was carried out PHRRB meetings to reach a consensus on criteria that the members have to rate in the auditing process. It was created a review chart that raters should verify the registry of patient's secondary diagnosis, chief complaint, history of presenting complaint, past medical history, medication history, physical exam and diagnostic testing. It was obtained the IRA every three months. The Gwet´s AC1 coefficient and Proportion of Agreement (PA) were calculated to evaluate the IRA for each item over time.

Results: The study included 1884 items from 239 records with an overall full agreement among raters of 71.2%. A significant IRA increase by 16.5% (OR=1.17; 95% CI=1.03—1.32; p=0.014) was found in the routine PHRRB auditing, with no significant differences between the PA and the Gwet´s AC1, that showed a similar evolution over time. The PA decreased by 27.1% when at least one of the raters was absent from the review meeting (OR=0.73; 95% CI=0.53—1.00; p=0.048).

Conclusions: Medical record quality has been associated with the quality of care and could be optimized and improved by targeted interventions. The PA and the Gwet´s AC1 are suitable agreement coefficients that are feasible to be incorporated in the routine of PHRRB evaluation process.

Keywords: Inter-rater agreement; Longitudinal agreement; medical quality register; audit; Gwet´s AC1.

Background

Adequate medical recordkeeping is an essential part of good health professional practice that makes it possible to evaluate and improve the quality of health care. Besides, medical records should also be used as a learning tool, far beyond the medical management of patients, but also improving the coordination and continuity of care and supporting decision making, avoiding adverse events that can compromise patient safety, mainly in hospitals.[1,2]

Auditing patient's medical records is a practice that aims to ensure the quality of patient care throughout reliable information registered by health professionals during patient visits or admissions in healthcare units. The Brazilian Medical Council establishes that patients' records review commissions is mandatory for health services, since 2002.[3] However, the reliability of the auditing processes is a matter closely related to the inter-rater agreement (IRA) when different raters assign the same precise value for each item being observed[4,5]. Some review studies assessing adverse events have been shown to suffer from poor to moderate inter-rater reliability (IRR)[6,7]. Also, IRR is rarely described or discussed in research papers based on data abstracted from medical records and there are no standard methods for assessing IRR[8]. Moreover, time constraints and work overload are frequent situations faced by health staff to perform tasks involving data management resulting in low data quality that can affect managerial decision-making. Therefore, the evaluation of suitable methods for data abstraction from this source is essential.[9]

When such studies employ multiple raters it is important to have a strategy to document adequate levels of agreement between them, and the Cohen's Kappa coefficient (κ) is a well-known measure.[10] However, it is affected by the skewed distributions of categories

(the prevalence paradox) and by the degree to which raters disagree (the bias

problem).[11,12]

Kilem Li Gwet proposed a new agreement coefficient to fix those limitations and which

can be used with any number of raters requiring a simple categorical rating system.[13,14]

The objective of this study was to evaluate the IRA of routine audits of medical records

and the impact of periodic discussions among raters, to refine auditing criteria, in a large

general hospital as part of an intervention to improve the quality of the medical records

related to essential content. Besides, the study also aimed to compare the estimates of

the percent agreement (PA) to the Gwet´s agreement coefficient (AC1) and to identify

possible factors associated with the PA.

Methods

Population and setting

This was a prospective longitudinal study conducted between July of 2015 and April of

2016 at the Hospital Municipal Dr. Moysés Deutsch (HMMD), a large public general

hospital (300 beds) located in the southern zone in the city of São Paulo, Brazil, an

impoverished region encompassing approximately 600,000 inhabitants. The present

study was part of a larger intervention aimed at improving the quality of patient care

throughout a tailored integration strategies among health facilities in its Regional Health

Care Network, that used a Lean Six Sigma methodology to get improvements of data

quality registered in the medical records, with potential benefits to the patients, to

decision-making actions and processes, and to obtain scientific quality over these data

for research purposes, as published elsewhere.[15]

Audit of medical records and review meetings

The HMMD maintains a routine auditing process that includes 13% of all medical

records of patients discharged in the previous month, carried out by the Patient's

Medical Record Review Board (PMRRB), that was composed by 24 nominated health

professionals from several HMMD staff, medical, nurses and multi-professional staff

coordinators, or delegated by them, from all clinical departments, 12 physicians, 9

nurses and 3 physiotherapists. It is a time-costly procedure because competes with the

patient-care tasks among these professionals. As a consequence, is common that the

audits have happened isolated by each PMRRB member without any criteria alignment,

to rate the items in the audit chart, which compromise the whole quality of the auditing

process. However, the patient's medical charts were selected in a non-aleatory way,

lacking representativeness over the results achieved, compromising the accuracy and

generalizability of these data.

The planned intervention has used the Lean-6-Sigma methodology, that is largely

utilized to aggregate values in several HMMD quality improvement processes, that is

part of the environment culture among the professionals.[15]

The proposed actions included at least one team-leader from each HMMD clinical

department, preferably its coordinator, which increased the PMRRB components,

reducing the total medical charts to be reviewed by each member. It was refined the

audit chart by all members through discussions about the relevance of the information

that should be registered by their health teams, answering the question: 'Which

information cannot be missed in the patient's medical record?" The chosen items were

then discussed, to define the criteria that should be rated as adequate, inadequate,

incomplete, or not applicable (Table 1). Finally, the patient's medical records have

become selected randomly, weighed by the discharge proportion of each department.

Table 1: Audited items.

Audited items	Options to rate
Secondary diagnosis	Adequate or inadequate
Chief complaint	Not applicable, adequate or inadequate
History of presenting complaint	Not applicable, adequate or inadequate
Past medical history	Not applicable, adequate or inadequate
Medication history	Not applicable, adequate or inadequate
Complete medical history	Not applicable, adequate, inadequate or incomplete
Physical exam	Adequate or inadequate
Diagnostic testing	Not applicable, adequate or inadequate

The number of raters varying throughout the study as shown in Table 2.

Every three months during the study period, and in addition to the routine audits, five to

six medical records were randomly allocated to the same two or three independent raters

of the same professional category to evaluate the IRA. The study also included review

meetings conducted every three months to align assessment criteria based on the results

of the IRA evaluation and the auditing processes.

Table 2: Number of audited medical records and raters over time.

Audit period	Number of medical records for IRA	Number of raters
1. 2015 July	54	18
2. 2015 October	45	19
3. 2016 January	84	21
4. 2016 April	56	16

Statistical analysis

The Gwet's AC1 and PA were calculated to evaluate the IRA for each item over time

and were compared through line graphs including 95% confidence intervals (CIs). The

Gwet's AC1 95% CIs[14] were calculated, while the PA were modelled by generalized

estimating equations (GEE), without an intercept[16,17]. The agreement measures were

interpreted following the categories proposed by Altman.[18]

Logistic GEE was used to model the PA of all raters along the time, using the value of 1

to full agreement and of 0 to some disagreement, and combining all items for each item

individually. The analyses employed an exchangeable working correlation matrix and

items in a single medical record were considered to be correlated. The model included

as independent variables: professional category, review meeting attendance, and time

(audits 1 to 4). A forward stepwise approach was used to variable selection employing a

p-value lesser than or equal to 0.200 in the unadjusted model, and lesser than or equal to

0.050 in the multiple-variable model.

 The analyses were performed with the R software version 3.2.2[19] with geepack[20].

Results

The study included 1884 items from 239 records with an overall full agreement among

raters of 71.2%. The estimated mean PA was found to be larger than the Gwet´s AC1

for all audited items (Figure 1), however, these differences were not statistically

significant and the evolution of the two agreement coefficients was similar throughout

the study period. Although a positive trend was found in the agreement of almost all

items, their CIs did not indicate any statistically significant changing over time. Also,

the coefficients measurements got closer as the agreement got greater. During the study

period, the greatest agreement was "chief complaint", while the lowest one was "secondary diagnosis".

Figure 1 here

The logistic GEE model that included all items (Table 3) found a statistically significant increase of 17% over time for the PA, but when at least one of the raters was absent from the review meeting, the PA decreased by 27%. Physiotherapists and physicians showed higher PA when compared to nurses.

In the analysis by item, there was a non-significant positive trend to higher PA for History of presenting complaint while physicians presented a significantly higher PA over time when compared to nurses for "secondary diagnosis", "medication history", and "diagnostic testing". Physiotherapists presented a significantly higher PA over time when compared to nurses for "medication history". Finally, when at least one of the raters was absent from the review meeting the PA decreased by 60.5% for "diagnostic testing" (Table 4).

Table 3: Estimated odds ratios (OR) for percent agreement. N=1884 items from 239 records.

	OR[a] (95%CI)	p value	OR[b] (95% CI)	p value
Time (audits 1 to 4)	1.20 (1.06–1.35)	0.004	1.17 (1.03–1.32)	0.014
Absent from meeting (Yes)	0.82 (0.60–1.11)	0.195	0.73 (0.53–1.00)	0.048
Professional category (physiotherapists)	1.49 (0.99–2.26)	0.058	1.66 (1.10–2.51)	0.016
Professional category	1.45 (1.08–1.93)	0.013	1.44 (1.07–1.93)	0.015

(physicians)

182   Table 4 here.

Discussion

A significant increase in the IRA among PHRRB members was found along the time in

routine medical record auditing processes when periodic evaluations of the agreement

were performed and discussed by them. On the other hand, but supporting this finding,

the absence of a member in a review meeting had a negative impact in the PA. Besides,

the PA and the Gwet´s AC1 were comparable and presented a similar evolution over

time. Complete medical history was a composite of chief complaint, history of

complaint, past medical history, and medication history. It was considered complete

whether all of them were complete, too. Thus, it showed a positive evolution in both PA

and Gwet AC1 over time from moderate to substantial according to Altman's

categories[18]. Only the IRA of secondary diagnosis has remained moderate. These

findings can indicate raters learning curve regarding the positive evolution of some

variables across agreement ranges. Nevertheless, the degree of agreement is arbitrary

making it impossible to define an acceptable level[5]. Thus, the interpretation of these

IRA values is following the main study objective, i.e., the rater's concordance in a

particular category.

The greater IRA among physicians and physiotherapists, when compared to nurses, can

reflect some inconsistency over the evaluations that can be attributed by rater's

selection, training, and accountability[5], that could be influenced by a misunderstanding

about rating the Complete History item.

The strategy applied for the IRA was feasible to be carried out in this real-world

scenario, aggregating value to the auditing process, providing more accurate

information that can be used by health leadership. The use of PA and Gwet´s AC1 for

that purpose was successful because they demand a relatively small sample of PMRs to

be audited by each rater, and can provide two data consistency measures.[5,21] Both of the

used indices have reached acceptable levels of agreement[18,22], according to study

purposes.

Following and evaluating the progress of the agreement among raters of PMRs allows

setting up goals and identifying associated factors to improve the audit processes, but

previously proposed models worked with continuous variables[23] or with the Kappa

coefficient[24], so the use of PA and Gwet AC1 made it possible to model the agreement

of more than two raters over time.

The increased IRA highlights the need for more careful planning and evaluation of

medical record audits since this activity is closely related to health care quality and

patient safety improvements efforts.[8,9]

Since the present study was conducted under real-world conditions and included

different health providers as raters, this intervention has the potential to be applicable in

other similar settings, taking into consideration that it was carried out in only one

hospital that has a culture of evidence-based improvement interventions, during a short-

term follow-up. Furthermore, the literature on the quality of medical records keeping

and IRA or IRR is scarce what is reflected by the fact that no reviews on the subject

could be identified making the results of this study relevant to improve the body of

knowledge, in the era of data-driven institutions and big data from patient's health

records.

Finally, this study did not include an evaluation of the impact in the quality of medical

records what should be the final goal of any routine audit, and therefore should be

evaluated in future studies.

Abbreviations

AC1: agreement coefficient

CI: confidence interval

GEE: generalized estimating equations

HMMD: Hospital Municipal Dr. Moysés Deutsch

IRA: inter-rater agreement

IRR: inter-rater reliability

OR: odds ratio

PA: proportion of agreement

PHRRB: patient's health record review board

Declarations

Ethics approval

245 The study was approved by the research ethics committee of the São Paulo Municipal

246 Health Department, and the partners' institutions (26981514.3.0000.0086).[25]

Consent for publication

249 Not applicable.

Availability of data and materials

The dataset supporting the conclusions of this article is available to researchers who want to explore the data. To request, please send an email to ana.mafra@einstein.br.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the Brazilian Ministry of Health and São Paulo State Research Foundation through Research Program for the Unified Health System-PPSUS grant 2012/51228-9.

Author's contribution

ACCNM conceptualized the study design, drafted the initial manuscript, carried out the random sampling, the statistical analysis, and revised the manuscript. RRST and EA elaborated and operationalized the intervention, contributed to the study design and reviewed the manuscript. FABC contributed to the study design and reviewed the manuscript. JLM and GP contributed to the interpretation of data for the work and revised the manuscript critically for important intellectual content. MMB elaborated the study design, operationalized the intervention, drafted and revised the manuscript. All authors approved the final manuscript as submitted.

Acknowledgements

We thank all the MRRC members who conducted the audit records and support the process, the members of the archiving sector as well as the hospital leadership.

References

1. Pirkle CM, Dumont A, Zunzunegui M-V. Medical recordkeeping, essential but overlooked aspect of quality of care in resource-limited settings. *International Journal for Quality in Health Care.* 2012;**24**(6):564–7. doi: 10.1093/intqhc/mzs034.

2. Zegers M, de Bruijne MC, Spreeuwenberg P, Wagner C, Groenewegen PP, van der Wal G. Quality of patient record keeping: an indicator of the quality of care? *BMJ Quality & Safety.* 2011;**20**(4):314–8. doi: 10.1136/bmjqs.2009.038976.

3. Conselho Federal de Medicina. Resolução nº 1638. Diário Oficial União nº 153, seção 1, 09/08/2002, p. 184-5. Available: https://sistemas.cfm.org.br/normas/visualizar/resolucoes/BR/2002/1638 [Accessed 30 Dec 2019].

4. Gisev N, Bell JS, Chen TF. Interrater agreement ant interrater reliability: Key concepts, approaches, and applications. *Research in Social and Administrative Pharmacy.* 2013; **9**:330-338. doi: 10.1016/j.sapharm.2012.04.004.

5. Bajpai S, Bajpai R, Chaturvedi HK. Evaluation of Inter-Rater Agreement and Inter-Rater Reliability for Observational Data: An Overview of Concepts and Methods. *Journal of the Indian Academy of Applied Psychology*. 2015, **41** (3): 20-27.

6. Lilford R, Edwards A, Girling A, Hofer T, Di Tanna GL, Petty J, *et al.* Inter-rater reliability of case-note audit: a systematic review. *J Health Serv Res Policy.* 2007;**12**(3):173–80. doi: 10.1258/135581907781543012

7.   Thomas EJ, Lipsitz SR, Studdert DM, Brennan TA. The reliability of medical

record review for estimating adverse event rates. *Ann Intern Med.*

2002;**136**(11):812–6. doi: 10.7326/0003-4819-136-11-200206040-00009

8.   Yawn BP, Wollan P. Interrater Reliability: Completing the Methods Description in

Medical Records Review Studies. *Am J Epidemiol.* 2005;**161**(10):974–7. doi:

10.1093/aje/kwi122

9. Liddy C, Wiens M, Hogg W. Methods to achieve high interrater reliability in data

collection from primary care medical records. *Ann Fam Med* 2011;**9**:57-62.doi:

10.1370/afm.1195

10. Cohen J. A Coefficient of Agreement for Nominal Scales. *Educational and*

Psychological Measurement. 1960;**20**(1):37–46. doi:

10.1177/001316446002000104

11. Zec S, Soriani N, Comoretto R, Baldi I. High Agreement and High Prevalence:

The Paradox of Cohen's Kappa. *The Open Nursing Journal.* 2017, **11**, (Suppl-1,

M5) 211-218. doi: 10.2174/1874434601711010211

12. Eugenio BD, Glass M. The Kappa Statistic: A Second Look. *Computational*

Linguistics. 2004;**30**(1):95–101. doi: 10.1162/089120104773633402

13. Wongpakaran N, Wongpakaran T, Wedding D, Gwet KL. A comparison of

Cohen's Kappa and Gwet's AC1 when calculating inter-rater reliability

coefficients: a study conducted with personality disorder samples. *BMC Medical*

Research Methodology. 2013; **13**:61. doi:10.1186/1471-2288-13-61.

14.   Gwet KL. Handbook of inter-rater reliability: the definitive guide to measuring the

324   extent of agreement among raters. 4. ed. Gaithersburg, MD: Advanced Analytics,

325   LLC; 2014.

326   15.   Bracco MM, Mafra ACCN, Abdo AH, Colugnati FAB, Dalla MDB, Demarzo

327   MMP, et al. Implementation of integration strategies between primary care units

328   and a regional general hospital in Brazil to update and connect health care

329   professionals: a quasi-experimental study protocol. *BMC Health Serv Res* 2016;

330   **16**:380. doi: 10.1186/s12913-016-1626-9

331   16.   Prentice RL, Zhao LP. Estimating equations for parameters in means and

332   covariances of multivariate discrete and continuous responses. *Biometrics*

333   1991;**47**(3):825–39. doi: 10.2307/2532642

334   17.   Liang K-Y, Zeger SL. Longitudinal data analysis using generalized linear models.

335   *Biometrika* 1986;**73**(1):13–22. doi: 10.1093/biomet/73.1.13

336   18.   Altman DG. Practical statistics for medical research. 1st ed. London: Chapman and

337   Hall; 1991

338   19.   R Core Team (2019). R: A Language and Environment for Statistical Computing.

339   R Foundation for Statistical Computing. Vienna. Austria. Available:

340   http://www.R-project.org/. [Accessed 30 Dec 2019].

341   20. Højsgaard S, Halekoh U, Yan J. The R Package geepack for Generalized Estimating

342   Equations. *Journal of Statistical Software* 2005; **15**:2. doi: 10.18637/jss.v015.i02.

343   21.   Walter SD, Eliasziw M, Donner A. Sample size and optimal designs for reliability

344   studies. *Stat Med.* 1998;**17**(1):101–10. doi:10.1002/(SICI)1097-

345   0258(19980115)17:1<101::AID-SIM727>3.0.CO;2-E.

22. Stemler, SE. A comparison of consensus, consistency, and measurement approaches

to estimating interrater reliability. *Practical Assessment, Research & Evaluation*.

2004, **9**:4.

23. Hill EG, Slate EH. A semi-parametric Bayesian model of inter- and intra-examiner

agreement for periodontal probing depth. *Ann Appl Stat*.2014;**8**(1):331–51. doi:

0.1214/13-AOAS688

24. Williamson JM, Lipsitz SR, Manatunga AK. Modeling kappa for measuring

dependent categorical agreement data. *Biostatistics* 2000;**1**(2):191–202. doi:

10.1093/biostatistics/1.2.191.

25. PlataformaBrasil. Availabe: http://aplicacao.saude.gov.br/plataformabrasil/login.jsf.

[Accessed 15 Apr 2019].

357
358

360   **Figure 1:** Estimated percent agreement (PA) and Gwet´s AC1 by audited item.

361   CI: Confidence interval. AC: Agreement coefficient.

363   **Table 4:** Estimated odds ratios (OR) of percent agreement by item. N= 239 records.

	OR (95% CI)	p value
Secondary diagnosis		
Time (audits 1 to 4)	1.02 (0.80–1.30)	0.878
Absent from meeting (Yes)	0.65 (0.35–1.23)	0.187
Prof. category (physiotherapists)	1.74 (0.72–4.22)	0.221
Prof. category (physicians)	1.82 (1.00–3.29)	0.048
Chief complaint		
Time (audits 1 to 4)	1.20 (0.80–1.80)	0.370

	OR (95% CI)	p value
Absent from meeting (Yes)	0.71 (0.26–1.95)	0.507
Prof. category (physiotherapists)	1.22 (0.29–5.16)	0.786
Prof. category (physicians)	0.74 (0.30–1.83)	0.519
History of presenting complaint		
Time (audits 1 to 4)	1.31 (0.99–1.74)	0.056
Absent from meeting (Yes)	0.58 (0.28–1.19)	0.138
Prof. category (physiotherapists)	1.78 (0.68–4.61)	0.238
Prof. category (physicians)	1.56 (0.81–3.01)	0.182
Past medical history		
Time (audits 1 to 4)	1.14 (0.86–1.51)	0.375
Absent from meeting (Yes)	0.70 (0.34–1.44)	0.334
Prof. category (physiotherapists)	1.12 (0.43–2.91)	0.817
Prof. category (physicians)	1.35 (0.69–2.63)	0.378
Medication history		
Time (audits 1 to 4)	1.25 (0.96–1.62)	0.099
Absent from meeting (Yes)	0.83 (0.44–1.57)	0.569
Prof. category (physiotherapists)	4.25 (1.53–11.77)	0.005
Prof. category (physicians)	1.87 (1.02–3.41)	0.041
Complete medical history		
Time (audits 1 to 4)	1.22 (0.95–1.57)	0.118
Absent from meeting (Yes)	0.74 (0.38–1.42)	0.359
Prof. category (physiotherapists)	1.15 (0.47–2.82)	0.753
Prof. category (physicians)	0.96 (0.52–1.76)	0.893
Physical exam		
Time (audits 1 to 4)	1.07 (0.82–1.39)	0.616
Absent from meeting (Yes)	1.32 (0.68–2.58)	0.410

	OR (95% CI)	p value
Prof. category (physiotherapists)	2.59 (0.70–9.57)	0.154
Prof. category (physicians)	1.02 (0.54–1.91)	0.950
Diagnostic testing		
Time (audits 1 to 4)	1.23 (0.91–1.66)	0.175
Absent from meeting (Yes)	0.39 (0.18–0.89)	0.024
Prof. category (physiotherapists)	1.52 (0.59–3.92)	0.387
Prof. category (physicians)	3.11 (1.53–6.30)	0.002

364 Prof.: professional.