Global DNA hypomethylation in epithelial ovarian cancer: passive demethylation and association with genomic instability

Wa Zhang[1], David Klinkebiel[2], Carter J Barger[1], Sanjit Pandey[3], Chittibabu Guda[3], Austin Miller[4], Stacey N. Akers[5] , Kunle Odunsi[5, 6, 7], and Adam R. Karpf[1,*]

[1]Eppley Institute, Departments of [2]Biochemistry and Molecular Biology, [3]Genetics, Cell Biology, and Anatomy, University of Nebraska Medical Center, Omaha, NE, 68198. Departments of [4]Biostatistics, [5]Gynecologic Oncology, [6]Immunology, and [7]Center for Immunotherapy, Roswell Park Cancer Institute, Buffalo, NY, 14263.

* Corresponding Author: Eppley Institute for Research in Cancer, Fred & Pamela Buffett Cancer Center, University of Nebraska Medical Center, Omaha, NE, 68198. Phone: 402-559-6115. Fax: 402-599-4651. Email: adam.karpf@unmc.edu.

Running Title: Characterization of DNA hypomethylation in ovarian cancer

Keywords: DNA hypomethylation, epithelial ovarian cancer, gene expression, repetitive elements, genomic instability, DNMTs

Non-standard abbreviations:

CG gene, cancer germline or cancer testis gene

CIN, chromosomal instability

CNA, copy number alteration

DAC, decitabine, 5-aza-2'-deoxycytidine

DEG, differentially expressed gene

DE-CG gene, differentially expressed CG gene

DMC, differentially methylated CpG site

DMR, differentially methylated region

DNMT, cytosine DNA methyltransferase

EOC, epithelial ovarian cancer

FDR, false discovery rate

FTE, fallopian tube epithelia

GDHO, global DNA hypomethylation

GSEA, gene set enrichment analysis

HGSOC, high grade serous ovarian cancer

IPA, Ingenuity Pathway Analysis

LAD, lamina-associated domain

NE, normal epithelia (OSE + FTE)

NO, bulk normal ovary

OSE, ovarian surface epithelia

PCA, principal component analysis

RE, repetitive DNA elements

RMA, robust multichip average

TET, Ten-eleven translocation methylcytosine dioxygenase

TF, transcription factor

Abstract

A hallmark of human cancer is global DNA hypomethylation (GDHO), but the mechanisms accounting for this defect and its pathological consequences are ill-defined. In human epithelial ovarian cancer (EOC), GDHO was associated with advanced disease and reduced overall and disease-free survival. GDHO(+) EOC was enriched for a proliferative gene expression signature, including FOXM1 activation. DNA hypomethylation preferentially occurred within genomic blocks (hypo blocks) overlapping late-replicating, lamina-associated domains, PRC2 binding, and H3K27me3. Increased proliferation coupled with hypo block formation at late replicating regions suggested passive hypomethylation, which was further supported by the observation that cytosine DNA methyltransferases (*DNMTs)* and *UHRF1* showed significantly reduced expression in GDHO(+) EOC, after normalization to proliferation markers. Importantly, GDHO(+) EOC showed elevated chromosomal instability (CIN), and copy number alterations (CNA) were enriched at hypo blocks. Together, these findings implicate a passive demethylation mechanism for GDHO that promotes genomic instability and poor prognosis in EOC.

Altered DNA methylation, a fundamental characteristic of human cancer, includes both gains and losses of methylation[1,2]. DNA hypermethylation leads to tumor suppressor gene silencing and occurs frequently at genomic regions occupied by polycomb group proteins in ES cells[1,3]. In contrast, DNA hypomethylation is referred to as "global," as 5-methyl-deoxycytidine (5mdC) levels are often reduced in cancer[4-6]. In agreement, DNA hypomethylation occurs at repetitive elements (RE), including the interspersed retrotransposon *LINE-1*[7], which accounts for ~17% of the genome. Global DNA hypomethylation (GDHO) is also associated with hypomethylation and activation of cancer-testis or cancer-germline (CG) genes[8-10]. More recently, epigenomic approaches have revealed that global DNA hypomethylation may not be random or driven by changes at RE, but rather is localized at large genomic regions called hypo blocks[11-13]. Hypo blocks overlap lamina-associated domains (LADs) and, interestingly, appear to contain epigenetically silenced genes, as well as genomic regions showing high gene expression variability[12,14,15].

GDHO, which is commonly measured using RE methylation markers such as *LINE-1*, is associated with poor clinical outcome, but the reasons for this association have not been established[16]. A number of plausible mechanisms could account for this link. First, GDHO may promote chromosomal instability (CIN), as genetically induced DNA hypomethylation in mouse tumor models and human cancer cell lines leads to aneuploidy, chromosomal translocations, and copy number alterations (CNA)[17-19]. Supporting this model, DNA hypomethylation and genomic alterations have been linked in some human cancers[20-23]. Second, aberrant gene expression, including oncogene activation, RE expression, or CG antigen gene activation may promote oncogenic phenotypes and/or disease progression[2,24-26]. Third, GDHO and the associated hypo block formation could promote gene expression variability and provide a selective growth advantage, e.g. chemotherapy resistance or accelerated disease progression[27].

In addition to identifying the targets and consequences of GDHO, it is of high interest to understand its origin.   At least two general types of mechanisms may underlie GDHO. First, "active" hypomethylation, caused by a specific molecular alteration that disrupts DNA methylation or enhances DNA demethylation, could be involved. Potential related mechanisms could include, for example, mutations in *DNMTs* or *Ten-eleven translocation methylcytosine dioxygenase* (*TET*) genes.   Second, "passive" hypomethylation is predicated on the fact that DNA methylation is a post-replicative DNA modification that could become unlinked from DNA replication. In the passive model, hypomethylation is proposed to indirectly result from cellular transformation and the accompanied increase in cell proliferation[28].

RNA sequencing (RNA-seq) is commonly used to study promoter variation and splice variants, which are difficult to measure using microarrays[29]. It is less appreciated that RNA-seq can also measure RE-derived transcripts, which have been excluded from microarray designs and are traditionally omitted from studies of the cancer transcriptome. Notably, total RNAseq revealed frequent and widespread expression of RE in pancreatic cancer and other tumor types[30]. The mechanisms accounting for RE expression in cancer are unknown, but could potentially include epigenetic activation by DNA hypomethylation.

Epithelial ovarian cancer (EOC), and specifically the high-grade serous subtype, HGSOC, is the most lethal gynecologic malignancy, due to a lack of diagnostic approaches coupled with poor long-term survival for patients with late stage disease[31]. EOC is characterized by widespread CNA, *TP53* mutations, defects in homologous recombination (including BRCA genes), Rb pathway dysregulation, and FOXM1 activation[32]. In addition to genetic changes, EOC shows altered DNA methylation, including both hyper- and hypo-methylation[33]. More specifically, GDHO is a common phenotype observed in EOC[9,34,35]. Here, we studied GDHO in EOC, including its clinicopathological context, molecular underpinnings, and relationship to CIN.

Results

***LINE-1* hypomethylation is associated with disease progression and reduced survival in EOC.** Previously, we reported that *LINE-1* methylation is an appropriate biomarker for global DNA methylation in EOC, and that *LINE-1* is hypomethylated in EOC as compared to bulk normal ovary (NO), ovarian surface epithelia (OSE), or fallopian tube epithelia (FTE) tissues[9,36]. Here, we investigated the relationship between *LINE-1* methylation and EOC clinicopathology. Notably, *LINE-1* hypomethylation was associated with increased clinical stage and histopathological grade, and correlated with reduced overall and disease-free survival (Fig. 1a-d). Furthermore, the association with disease free survival remained significant in a proportional hazards model after adjustment for age (HR P=0.03).

Global gene expression patterns are distinct in GDHO(+) EOC. We used gene expression microarrays to profile: i) EOC samples showing significant *LINE-1* hypomethylation [GDHO(+) EOC; N=20], ii) EOC samples with *LINE-1* methylation levels similar to NO [GDHO(−) EOC; N=20], and iii) NO; N=3 (Fig. 2a). For expression studies, we used NO as a control because of difficulty in obtaining high quality RNA from primary OSE and FTE tissues. Hierarchical clustering of differentially expressed genes (DEG) revealed distinct patterns of gene expression in GDHO(+) and GDHO(−) EOC (Fig. 2b). Using a cutoff of P<0.01, 1696 Affymetrix probe sets (genes) were differentially expressed in the two EOC groups, with 958 (56%) of these upregulated in GDHO(+) EOC (Fig. 2b; Supplementary Table 1). Clustering of DEG using a false discovery rate (FDR) cutoff of 0.1 resulted in a similar separation of GDHO(+) from GDHO(−) samples (Supplementary Fig. 1a). Because GDHO is associated with disease progression (Fig. 1), we conducted sub-group analysis using disease-matched tumors, in which we compared age-matched, stage 3/4, grade 3, serous EOC (i.e. HGSOC) samples from each group [GDHO(+) N=13; GDHO(−) N=9]. Importantly, disease matched GDHO(+) and GDHO(−) EOC gene expression patterns remained distinct (and provided improved group separation) in

this comparison (Fig. 2c).   At P<0.01, 752 Affymetrix probe sets (genes) were differentially expressed, with 357 (47%) upregulated in GDHO(+) EOC (Fig. 2C; Supplementary Table 1). Clustering of DEG using a false discovery rate (FDR) cutoff of 0.1 resulted in similar separation of disease-matched GDHO(+) vs. GDHO(−) EOC (Supplementary Fig. 1b).   A list of all DEG identified in both comparisons [GDHO(+) vs. GDHO(−): overall and disease-matched samples] is provided in Supplementary Table 2, and the microarray raw intensity files have been deposited in GEO (see *Methods*).

Proliferative expression signature and FOXM1 activation in GDHO(+) EOC. We used Ingenuity Pathway Analysis (IPA) of microarray expression data to identify cellular pathways involved in GDHO. We observed significant alterations in both cancer and reproductive disease pathways (Fig. 2d; Supplementary Table 3). In addition, functional pathways related to cell growth, proliferation, and division were altered; together these data suggested that increased proliferation is a key characteristic of GDHO(+) EOC (Fig 2d; Supplementary Table 3). To test this, we measured the expression of the canonical cell proliferation marker *MKI67*[37] in GDHO(+) and GDHO(−) EOC. Both microarray and RT-qPCR analysis demonstrated significantly elevated *MKI67* expression in GDHO(+) EOC, as well as a significant negative association with *LINE-1* methylation (Supplementary Fig. 2).   In addition, two key drivers of EOC proliferation, *CCNE1* and *FOXM1*, were markedly upregulated in GDHO(+) EOC (Fig. 3a,b). Based on the emerging role of the FOXM1 pathway in EOC and a report that FOXM1 may be linked to changes in DNA methylation[32,38], we tested the association between *FOXM1* expression and *LINE-1* methylation using an expanded set of EOC samples. Notably, the data revealed a strong inverse association (Fig. 3c). In addition to the mRNA, expression of the FOXM1 protein and also FOXM1 target genes involved in cell cycle progression, including *PLK1*, *AURKB*, *BIRC5*, and *CCNB1*[32], were significantly upregulated in GDHO(+) EOC, consistent with functional activation of FOXM1 (Fig 3d; Supplementary Fig. 3).   Furthermore, gene set

enrichment analysis (GSEA) indicated a significant enrichment of the FOXM1 transcription factor network (p <0.01, FDR <0.25) and G2/M checkpoint genes (p < 0.05, FDR <0.25) in GDHO(+) EOC. Together, these data implicate increased FOXM1 expression as a prominent feature of GDHO in EOC. Activation of *CCNE1* and *FOXM1* in GDHO(+) EOC could potentially result from direct hypomethylation of their promoter regions, resulting from a passenger effect in GDHO(+) tumors. However, bisulfite sequencing analyses indicated that the *CCNE1* and *FOXM1* promoters were constitutively hypomethylated in NO, GDHO(−) and GDHO(+) EOC (Supplementary Fig. 4), suggesting that increased proliferation may promote GDHO, rather than *vice versa*.

Widespread activation of cancer germline (CG) genes and altered expression of epigenetic regulators and histone genes in GDHO(+) EOC. Expression of CG genes in association with GDHO in cancer has been reported, but previous studies have only studied one or a limited number of genes[8-10]. Our current data set allowed for a comprehensive and global examination of this association. Almost one-fifth of annotated CG genes were differentially expressed in GDHO(+) vs. GDHO(−) EOC, with each showing upregulation (Fig. 4; Supplementary Table 4). In agreement, GSEA analysis of CG genes showed significant enrichment in GDHO(+) EOC (p<0.01, FDR<0.25).   Over half of the CG genes activated in GDHO(+) EOC were also induced in the disease-matched comparison, including both X-chromosome and autosomal genes (Fig. 4; Supplementary Table 4). The differentially expressed CG genes (DE-CG genes) included cancer vaccine targets and genes with oncogenic function, including *MAGEA* genes, *NY-ESO-1*, *XAGE-1*, and *PRAME* (Supplementary Table 4). To test whether these genes are directly regulated by DNA methylation, we used Affymetrix microarrays to measure their expression in EOC cell models before and after decitabine (DAC) treatment.   Notably, the majority of the DE CG genes were upregulated following decitabine treatment (Fig 4; Supplementary Table 4). This observation,

coupled with extensive prior data regarding CG gene hypomethylation in cancer[10], strongly

suggests that many of the DE CG genes are directly regulated by DNA methylation.

Two additional gene groups of note had altered expression in GDHO(+) EOC. First,

genes with known roles in epigenetic regulation were altered. These genes included

EHMT2/*G9a, ATAD2,* and *HDAC1*, which have reported oncogenic activity in EOC[39-43]

(Supplementary Table 5). Increased expression of *ATAD2* in GDHO(+) tumors could be due to

promoter hypomethylation[44]. Interestingly, there was significantly increased expression both of

epigenetic regulators involved in gene activation (e.g. *JMJD2a*, *ATAD2*, *ASF1b*), as well as

gene repression (e.g. *G9a*, *HDAC1*, *LSD1*) in GDHO(+) EOC (Supplementary Table 5). These

changes may reinforce the epigenetic impact of GDHO, or alternatively could become activated

as a feedback inhibitory response to GDHO. In addition to direct epigenetic regulators, we

noted that approximately half of all histone genes were upregulated in GDHO(+) EOC as

compared to GDHO(−) EOC (Supplementary Table 6). In part, this could reflect the increased

proliferation signature observed in these tumors; in addition, direct hypomethylation of histone

genes in EOC has recently been reported[45]. Bisulfite sequencing analysis of select histone

genes indicated that some are constitutively hypomethylated in NO and EOC, while others are

hypomethylated in GDHO(+) EOC, as compared to GDHO(−) EOC (Supplementary Fig. 5).

DNA Methylome analysis of GDHO(+) EOC. For initial analysis of the DNA methylation

landscape of GDHO(+) EOC, we used Illumina Infinium 450K bead arrays (450K)[46]. In addition

to profiling both groups of EOC, we analyzed normal epithelia (NE; OSE + FTE average) as a

control. Both unsupervised hierarchical clustering and principal component analyses (PCA) of

450K data revealed that GDHO(+) and GDHO(−) EOC have distinct DNA methylomes, and that

each are distinct from NE (Fig. 5a,b). As a complementary approach, and to define the

methylome in greater depth, we performed methylome sequencing (Methyl-seq) using a solution

hybridization selection method[47] (Agilent SureSelect). 450K and Methyl-seq data were highly

concordant (Pearson r = 0.97; N=347,357 CpG sites), and hierarchical clustering and PCA analyses of Methyl-seq confirmed that GDHO(+) EOC, GDHO(–) EOC, and NE have distinct methylomes (Fig. 5c,d). As expected, DNA methylation overall was reduced in GDHO(+) EOC, as determined by either DNA methylome method (Fig. 5 e,f).

Both 450K and Methyl-seq indicated that a large number of promoters were significantly differentially methylated in the two EOC groups, with a vast majority, regardless of CpG island context, showing hypomethylation in GDHO(+) EOC (Supplementary Fig. 6).   In contrast, promoter hypermethylation and hypomethylation was almost evenly split when comparing EOC to NE, and, strikingly, CpG island promoters were mostly hypermethylated while non-CpG island promoters were mostly hypomethylated in EOC (Supplementary Fig. 6).   We also defined differentially methylated regions (DMR) globally and found that the vast majority of DMR were hypomethylated in GDHO(+) as compared to GDHO(−) EOC (Supplementary Fig. 7).   The majority of DMR were also hypomethylated in EOC as compared to the NE control (Supplementary Fig. 7). As methylation at different genomic locations has distinct consequences, we calculated differentially methylated CpGs (DMC) independently for genes, CpG islands, CpG shores, CpG shelves, and CpG open seas, as described previously[46]. Most DMC were hypomethylated in GDHO(+) as compared to GDHO(−) EOC, regardless of genomic context (Supplementary Fig. 7 c,d), while hypermethylation and hypomethylation was more evenly split for EOC vs. NE (Supplementary Fig. 7 e,f).   In the latter comparison, CpG rich regions of the genome favored hypermethylation in EOC, while CpG poor regions (i.e. open seas) favored hypomethylation in EOC.

Hypomethylated genomic blocks (hypo blocks) in GDHO(+) EOC. To investigate whether hypo blocks[11,12] are present in EOC, and to determine their relationship to GDHO, we first visually inspected Methyl-seq data using the *UCSC Genome Browser*.   Examples of the resulting data are shown for chromosome 11 (Fig. 6a,b). These observational data suggested

that hypo blocks were present and enriched in GDHO(+) EOC.   To formally test this, we used a

quantitative approach to determine the number and size of hypo blocks in the different EOC

groups, using NE as the control (see *Methods*).   While both 450K and Methyl-seq were capable

of detecting and quantifying hypo blocks, we only present Methyl-seq data here, due to its

significantly greater genomic coverage.   We observed a large enrichment in both the number

and size of hypo blocks in GDHO(+) EOC as compared with GDHO(–) EOC, with approximately

14% of the genome residing in hypo blocks in the former (Supplementary Table 7).   These data

indicate that hypo block formation is heterogeneous in EOC, and may be closely linked to more

common measures of GDHO, such as *LINE-1* hypomethylation.   Based on previous findings in

other cancer types, we next quantified the overlap between EOC hypo blocks, lamina-

associated domains (LADs), and late-replicating genomic regions.   We observed significant

enrichment of hypo blocks at LADs, and a similar enrichment at late-replicating regions (Fig 6b;

Table 1).   Genes and CpG islands were also enriched in hypo blocks, but this likely reflects the

bias of our Methyl-seq approach to coverage of these regions (Table 1; Supplementary Table

8). We also analyzed hypo blocks for overlap with specific transcription factor (TF) binding

sites, histone modifications, and RE, using ENCODE and the UCSC genome browser database.

Among TF binding sites showing strong enrichment in EOC hypo blocks were the polycomb

repressor complex 2 (PRC2) components EZH2 and SUZ12, and the genomic insulator CTCF

(Table 1; Supplementary Table 9). Hypo blocks were also highly enriched for H3K27me3, the

modification recognized by the PRC2 complex (Table 1; Supplementary Table 10). These data

suggest an association between DNA hypomethylation and repressive chromatin formation in

EOC, as was reported previously in a breast cancer cell line[14].   Notably, hypo blocks have been

proposed to facilitate variable gene expression, which may provide a selective growth

advantage during tumorigenesis[13].   In agreement, we found that genes with high expression

variability were enriched in the EOC hypo blocks (Fig 6c).

Maintenance methylation components have reduced expression relative to proliferation markers in GDHO(+) EOC. The proliferative gene expression signature found in GDHO(+) EOC, coupled with the presence of hypo blocks that overlap late-replicating regions, suggests a passive DNA hypomethylation mechanism.   Paradoxically, however, elevated *DNMT* expression (relative to normal tissues) has been commonly observed in cancer.   To address this discrepancy, we first analyzed *DNMT* expression in our samples using standard Robust Multichip Average (RMA) Affymetrix normalization. As expected, *DNMT1* and *DNMT3B* were upregulated in EOC compared to NO, while *DNMT3A* and *DNMT3L* were expressed at similar levels (Supplementary Fig. 8). However, it is important to note that maintenance DNA methylation is restricted to S phase, where *DNMT* expression is also increased[48]. Therefore, we hypothesized that *DNMT* expression normalized to cell proliferation status is a more relevant measure of maintenance methylation capacity in tumors. Remarkably, after normalization to the canonical cancer cell proliferation marker *MKI67*[37], all four DNMTs showed significantly reduced expression in EOC as compared to NO (Supplementary Fig. 9 and 10), and, more importantly, in GDHO(+) as compared to GDHO(−) EOC (Fig. 7a). In agreement, *DNMT1* and *DNMT3A* expression, when normalized to *MKI67*, inversely correlated with *LINE-1* methylation in an expanded set of EOC samples (Fig. 7b-d).   In addition to DNMTs, UHRF1 is a critical component of maintenance DNA methylation[49].   Similar to *DNMTs,* after normalization to *MKI67*, *UHRF1* was downregulated in GDHO(+) as compared to GDHO(−) EOC (Fig. 7a). *UHRF1*, after *MKI67* normalization, also showed lower expression in EOC vs. NO (Supplementary Fig. 9e). Importantly, normalization of *DNMTs* and *UHRF1* to other established cancer proliferation markers, including *PLK1* and *BUB1*[37], provided similar results as seen using *MKI67* normalization (Supplementary Figures 11 and 12). Together, these data strongly suggest that GDHO(+) EOC has a reduced capacity for maintenance DNA methylation.

Repetitive element (RE) expression and hypomethylation in GDHO(+) EOC. As described

above, GDHO is most commonly defined using RE methylation as a biomarker, e.g. *LINE-1*

hypomethylation. As widespread RE expression has recently been reported to occur in

cancer[30], we wanted to investigate the potential relationship between GDHO and RE

expression. For this task, we used total RNA-seq and Methyl-seq, methods that allow precise

genomic mapping of repeat sequences. Importantly, RE expression was generally elevated in

GDHO(+) EOC (Fig. 8a, b). However, this effect was not uniform, but rather was class-specific

(Fig. 8a; Supplementary Table 11). Most of the upregulated RE in GDHO(+) EOC were

hypomethylated at their corresponding genomic loci (Fig. 8 a,b). We additionally examined the

relationship between RE methylation and hypo blocks. RE showed specific patterns of

enrichment or depletion in hypo blocks, including *LINE*, which, in most instances, was enriched

in hypo blocks. In contrast, *Alu/SINE* and satellite sequences were depleted from hypo blocks

(Table 1; Supplementary Table 12). Importantly, Methyl-seq data indicated that, as was

reported previously in colon cancer[12], hypomethylation in EOC was mostly a consequence of

hypo block formation, rather than RE or *LINE-1* hypomethylation (Fig. 8 c,d).

Elevated chromosomal instability (CIN) and copy number alterations (CNA) in GDHO(+)

EOC. To further understand the oncogenic consequences of GDHO, we focused our attention

on CIN, a hallmark of EOC[32], using two complementary approaches. We first used a previously

reported 25-gene expression signature of CIN to interrogate our gene expression data[50]. This

signature (CIN25) was highly elevated in GDHO(+) as compared to GDHO(−) EOC, both when

analyzing all samples as well as in diseased-matched samples (Fig. 9a). Second, we directly

measured CNA in an additional set of 40 disease-matched GDHO(+) and GDHO(−) EOC (all

HGSOC), using copy number/SNP arrays. The data revealed that CNA was significantly

elevated in GDHO(+) EOC (Fig. 9b). Moreover, CNA was uniformly high in GDHO(+) EOC

(with one exception), while it was highly variable in GDHO(−) EOC (Fig. 9b). These data

suggest that other important contributors to CIN may be variably present in GDHO(−) EOC.

Finally, we observed significant enrichment of CNA in hypo blocks (Fig. 9c), suggesting that

hypo block formation could promote the acquisition of CIN in EOC.

Discussion

We report here an evaluation of the molecular pathology associated with GDHO in EOC. We used *LINE-1* hypomethylation as a biomarker for GDHO based on our earlier study[9], and observed that GDHO was associated with disease progression (advanced stage and grade) and reduced survival (disease-free and overall). By analyzing the mRNA transcriptome, RE expression, the DNA methylome, and copy number variation in parallel, we uncovered the major molecular features of GDHO. These include: i) distinct patterns of gene expression, including activation of key drivers of EOC cell proliferation including *CCNE1* and *FOXM1*, ii) widespread activation of CG antigen genes; iii) deregulated expression of epigenetic regulators and histone genes, iv) increased expression of RE, often in conjunction with hypomethylation at the associated loci, and v) increased CNA and CIN. Each of these features has oncogenic potential that could contribute to the disease progression and poor prognosis observed. In addition to the characteristics described above, DNA methylome analyses revealed the formation of hypo blocks, which were not ubiquitous in EOC, but rather occurred most dramatically in GDHO(+) EOC. Importantly, all known genes involved in maintenance DNA methylation displayed significantly reduced expression, relative to cell proliferation markers, in GDHO(+) EOC. Together, increased proliferation, hypo block formation (which predominated at late-replicating regions), and decreased *DNMT* and *UHRF1* expression strongly suggests that passive demethylation is a key contributor to GDHO in EOC.

DNA hypomethylation was the first epigenetic alteration discovered in cancer[2]. Early seminal studies showed that DNA hypomethylation can target oncogenes, coincides with overall loss of 5mdC, and is enriched in metastatic tumors[4-6,51]. Despite its early recognition, the mechanisms underlying GDHO are still not well understood. The recent discovery of TET-assisted oxidative DNA demethylation, involving intermediates including 5-hydroxymethylcytosine (5hmC), suggests TET activation as a potential mechanism for GDHO. However, this appears to be unlikely, as TETs are often mutated or downregulated in cancer,

and cancer tissues show dramatically reduced levels of 5hmC[52]. Other possible mechanisms

underlying GDHO have been proposed. For example, we earlier reported that an increased

(*BORIS*) *CTCFL/CTCF* ratio correlates with *LINE-1* hypomethylation in EOC[9]. In contrast, we

found that BORIS overexpression was insufficient to promote DNA hypomethylation[53]. Other

recent studies have suggested that UHRF1 overexpression (in hepatocellular carcinoma), and

PIWI protein repression (in testicular cancer) may contribute to GDHO[54,55]. However, in the

current study, we found no evidence for these mechanisms in EOC. It is possible that the

mechanisms involved in driving GDHO in cancer are tissue or context dependent.

Our data suggest that GDHO in EOC results, at least in part, from a proliferation-

dependent process. The most statistically significant functional pathways upregulated in GDHO

involved cell growth and proliferation, and, this was true for both unselected and disease-

matched samples, suggesting a fundamental link to GDHO.   We hypothesize that increased

proliferation may overwhelm the capacity for EOC cells to perform faithful maintenance DNA

methylation. Consistent with this model, a study of replication timing and DNA methylation

found that late-replicating regions were generally hypomethylated and that this accumulated

over multiple cell divisions[56].   Also supportive of this model is the recent observation of a short

delay between DNA replication and the completion of maintenance methylation in cancer cells[57].

A passive model for GDHO predicts that DNA hypomethylation would be concentrated at the

genomic regions most difficult to methylate, e.g. heterochromatic and late replicating regions,

including LADs, which is precisely what we and others have observed[11-14]. The association of

GDHO with LADs should be viewed with caution, however, as LADs have not been

simultaneously mapped with DNA methylation in parallel. In this context, a recent study found a

link between DNA hypomethylation and late replication timing, and also showed that the

association of each of these elements with LADs was secondary[58].

Because early data indicated that DNMT expression is increased in cancer, it has long

been assumed that DNMTs show a gain of function in cancer, which could in turn promote the

epigenetic silencing of tumor suppressor genes. However, this concept is paradoxical to the global reduction of methylation frequently observed in cancer[2]. Moreover, tumor suppressor functions for DNMTs have now been demonstrated[59]. Because DNMTs are cell cycle regulated[48], and cancers, especially at late stages, often have an elevated proliferation index, we hypothesized that it is appropriate to normalize DNMT expression to cell proliferation, in order to more accurately gauge the functional capacity of maintenance methylation. Remarkably, after normalization to canonical cancer proliferation markers such as *MKI67*, we observed significantly reduced expression of all *DNMTs* and *UHFR1* in GDHO(+) EOC. In addition to *DNMT1*, reduced expression of *DNMT3A/3B/3L* would be expected to impair maintenance methylation, as these enzymes contribute to maintenance methylation of heterochromatic genomic regions, presumably including LOCKs (large organized chromatin K9 modifications) and LADs[60,61]. Although we did not directly measure DNMT protein expression in the current study, it was recently reported that DNMT protein expression is downregulated in high malignant potential vs. low malignant potential EOC, and, moreover, that reduced DNMT protein expression correlates with poorer survival in EOC[62]. These critical observations affirm our mRNA expression data and are also consistent with our data showing that GDHO correlates with reduced survival.

The link observed between *FOXM1* and GDHO is of particular interest. While FOXM1's association with GDHO may be indirect due to its role in promoting cell cycle progression, this pathway may be particularly relevant in HGSOC[32]. Moreover, a recent study reported that FOXM1 overexpression leads to DNA methylation changes in oral keratinocytes, including a global hypomethylation phenotype similar to that observed in a relevant squamous cell carcinoma cell line[38]. Coupled with our current data, an investigation of the potential mechanistic connection between FOXM1 and DNA hypomethylation in EOC is warranted.

A seminal early study demonstrated that expression of the prototype CG antigen gene *MAGEA1* was linked to global DNA hypomethylation[8]. Cadieux et al. later reported a link

between GDHO, *MAGEA1* expression, and cell proliferation in glioblastoma[63]. In the current

study, we measured expression of all CG genes in EOC using microarrays and found that

approximately one-fifth were activated in conjunction with GDHO. The lack of ubiquitous

expression of all CG genes in GDHO(+) EOC suggests additional regulatory mechanisms for

CG antigen expression, e.g. oncogenic transcription factors[10].   More generally, dramatically

enhanced expression of CG antigens in association with GDHO suggests that patients

harboring such tumors could be an optimal group for CG antigen-directed immunotherapy.

Notably, immune checkpoint modulators, such as α-CTLA4 or α-PD1/PD-L1 antibodies, require

tumor antigens for their clinical activity[64]. These agents, as well as CG antigen-directed

vaccines, may be most effective in patients with GDHO(+) tumors[65].   It will be relevant to test

this concept in prospective clinical studies, in which the diagnostic tumor is used for GDHO

assessment.

Pivotal work using murine models and human cell lines established the initial link

between DNA hypomethylation and genomic instability[17,18]. This link has also been observed in

studies using primary human tumor samples, but to date these studies have focused on a

limited number of CpG methylation sites and chromosomal alterations. In the current study, we

comprehensively determined CNAs in HGSOC, which among human cancers studied to date,

shows the most extreme level of CIN[66]. GDHO(+) tumors showed significantly and consistently

higher levels of CIN, and most notably, CNAs were enriched in hypo blocks. These data point

towards a previously unknown molecular connection between DNA hypomethylation and CIN.

Consistent with our data, a recent study reported that chromosomal breakpoint regions in breast

cancer co-localize with DMRs, which were found to be typically hypomethylated in cancer[67].

Determining the mechanism by which the acquisition of hypomethylation at specific genomic

regions promotes CIN is an important area for future study.

Methods

Human tissue samples. NO, OSE, FTE, and EOC samples were obtained from patients undergoing surgical resection at Roswell Park Cancer Institute (RPCI), using Institutional Review Board-approved protocols. These samples have been described previously[9,36], and clinicopathological information is provided in Supplementary Table 13. NE controls (OSE and FTE) were obtained from patients without malignancy, and EOC samples contained >90% neoplastic cells. Processing of frozen tissue samples was described previously[34]. Supplementary Table 14 lists all sample groups used in this study and the genomic and epigenomic analyses conducted on each.

DNA, RNA, and protein extractions. Genomic DNA was isolated using the Puregene Tissue Kit (Qiagen).   Total RNA was purified using TRIzol (Invitrogen).   Total cellular protein was extracted using RIPA buffer. Extractions were performed as described previously[9,34].

Cell lines and drug treatments. IOSE-121 and OVCAR3 cells were described previously[68]. Cells at ~50% confluence were treated with 1μM decitabine (DAC) (day 0), passaged at day 2, re-treated with 1μM DAC at day 3, and harvested for RNA and DNA extractions at day 5.   PBS was used as the vehicle control. The data describing DAC treatment in Figure 4 is compiled from treatment of these two cell lines.

Gene expression microarrays. Affymetrix HG 1.0ST array analysis was performed at the University at Buffalo Center of Excellence in Bioinformatics and Life Sciences.   Microarray probe cell intensity data (.cel) were normalized using the Affymetrix Expression Console (version 1.3.0.187) software running the RMA workflow. We used a regularized t test analysis of control versus treatment comparisons using a Bayesian approach to estimate the within-treatment variation among replicates, using Cyber-T software. Expression heat maps were

created using the TM4 microarray software suite Multi Experiment Viewer (MeV) software

hierarchical clustering routine based on a Pearson correlation metric and average linkage.

Functional identification of gene networks was performed using IPA (Ingenuity Systems).   Gene

set enrichment analysis (GSEA, http://www.broadinstitute.org/gsea/index.jsp) was used to

determine the enrichment of the FOXM1 transcription factor network

(http://www.broadinstitute.org/gsea/msigdb/cards/PID_FOXM1PATHWAY.html), genes involved

in the G2/M checkpoint

(http://www.broadinstitute.org/gsea/msigdb/cards/HALLMARK_G2M_CHECKPOINT.html), and

CG genes (http://cancerimmunity.org/resources/ct-gene-database) in GDHO(+) EOC vs.

GDHO(−) EOC.

Reverse Transcriptase-Quantitative PCR (RT-qPCR). RT-qPCR was performed as

described previously[9].   Briefly, DNase-treated RNA was converted to cDNA, and samples were

analyzed in triplicate using a BioRad CFX Connect system and the SYBR green method.

Expression data were normalized to *18s rRNA*. All primer sequences are provided in

Supplementary Table 15.

Western blot analysis. Western blotting was performed as described previously[9], and used

the rabbit polyclonal primary anti-FOXM1 antibody (K19, Santa Cruz Biotechnology, sc-500,

1:500 dilution), and the goat anti-rabbit IgG-HRP secondary antibody (Santa Cruz

Biotechnology, sc-2004, 1:5000 dilution).   Ponceau S (Acros) staining was used as an

additional loading control.

Bisulfite clonal sequencing and pyrosequencing. Genomic DNA was converted using the

EZ DNA Methylation Kit (Zymo Research).   Bisulfite sequencing was accomplished as

described previously[69] and DNA sequences were analyzed using Lasergene (DNASTAR).

LINE-1 pyrosequencing was performed as described previously[34], using the PSQ HS96 System (Qiagen). All primer sequences are provided in Supplementary Table 15.

DNA methylome analyses. Illumina Infinium 450K BeadChip analysis was performed at the RPCI Genomics Core and the University of Utah Genomics Core Facilities. Agilent SureSelect Methylome sequencing, a targeted solution hybridization bisulfite sequencing method (SHBS-seq)[47], was performed at the UNMC Epigenomics Core Facility. This method encompasses ~15% of genomic CpG sites (3.5×10^6 CpGs). High-throughput sequencing of library tags was performed at the UNMC Sequencing Core Facility, using an Illumina HiSeq 2000 Genome Analyzer.   Sequence tags were aligned to the human genome (hg19) using the methylated sequence aligner Bismark (http://www.bioinformatics.babraham.ac.uk/projects/bismark/). The coverage of different genomic regions by Methyl-seq is shown in Supplementary Table 8. Differentially methylated regions (DMR) corresponded to genomic regions of any length containing ≥ 3 CpG sites, with ≥ 1 CpG site showing a mean methylation change of >20% at p ≤ 0.05. RnBeads was used to analyze both 450K and Methyl-seq data, in order to define methylation changes including DMR, promoters, CpG sites, and other genomic elements (http://rnbeads.mpi-inf.mpg.de/).

Determination of EOC hypo blocks. We defined hypo blocks essentially as described previously[13].   Briefly, we used RnBeads to determine hypo blocks using Methyl-seq data for NE control [(FTE + OSE average (N=2)] versus GDHO(+) EOC (N=2), using a ≤ 0.05 false discovery rate (FDR) of 5 kb tiled regions, with a ≥ 35% average methylation decrease. We combined regions that were ≤ 250kb distance apart, and selected only final blocks containing ≥ 5 CpGs.

Correlation of EOC hypo blocks with genomic features. Spatial correlations were calculated between the reference hypo block genomic intervals determined from GDHO(+) EOC Methyl-Seq data and the query genomic intervals of specific genomic features, using the R package GenometriCorr (http://genometricorr.sourceforge.net/).   CTCF, EZH2 and SUZ12 transcription binding sites were acquired from the UCSC Genome Browser hg19 table browser wgEncodeRegTfbsClusteredV3 table. *LINE-1*, *SINE-Alu,* and satellite repeat genomic locations were acquired from the UCSC Genome Browser hg19 table browser RepeatMasker. LAD genomic locations were acquired from the UCSC Genome Browser hg19 table browser NKI LADs (Tig3) database. Early and late replicating genomic regions were acquired from the UCSC Genome Browser hg19 table browser Replication Timing by Repli-chip from ENCODE/FSU table IMR90 1, wgEncodeFsuRepliChipH1hescWaveSignalRep1. Histone modifications were acquired from the ENCODE ChIP-seq of mammary epithelial cells, using H3K9me3 (accession ENCFF001SWV), H3K4me3 (accession ENCFF001SXB), H3K27ac (accession ENCFF001SWW), H3K36me3 (accession ENCFF001SWY), and H3K27me3 (accession ENCFF001SWX).

Total RNA sequencing (RNA-seq). RNA-seq was performed at the UNMC Sequencing Core Facility using the TruSeq Stranded Total RNA kit (Illumina) and an Illumina HiSeq 2000 Genome Analyzer. The starting material was 1.0 µg total RNA/sample. The resulting sequence tags were aligned to the UCSC Genome Browser reference human genome (hg19) mRNAs and REs using the software TopHat (v2.0.8)[70]. Cufflinks (v2.1.1) and Cuffdiff (v2.1.1) were used to estimate the expression values and determine differential expression of REs[70].

Genomic copy number analysis. Genomic copy number analysis was performed at the UNMC Sequencing Core Facility, using Affymetrix Cytoscan HD microarrays. The total number and size (segments) of CNA per sample were determined using the Affymetrix Chromosome

Analysis Suite (ChAS) Software. A CNA index was calculated for each sample based on the percent of the genome that resulted in either a copy number loss or gain. Base overlap (CNA inside hypo blocks) and non-overlap (CNA outside hypo blocks) between each EOC CNA segment and hypo blocks was determined using the Bedtools intersect routine (http://bedtools.readthedocs.org/en/latest/content/tools/intersect.html).

Correlation of *LINE-1* methylation with EOC patient survival. Overall survival was defined as the number of months between the diagnosis date and death, and patients still alive were censored at their date of last follow-up. For disease-free survival, patients who were alive and disease-free were censored at the date of the last visit. *LINE-1* methylation, as determined by bisulfite pyrosequencing, was segregated into three groups as shown in Fig. 1c,d, and survival was compared using Kaplan Meier analyses. The null hypothesis of no difference in the survival distributions was assessed using the Logrank test.   The age-independent association of *LINE-1* methylation with survival was tested using a proportional hazards model.

Genomic data deposit and public access. Upon publication of this paper, all data from each genomic parameter platform will be deposited into the NCBI Gene Expression Omnibus (GEO). Illumina Infinium Human Methylation 450K BeadChip array data will be deposited as *.idat files. High throughput sequencing analysis Total RNA-seq and Agilent's SureSelect Target Enrichment Bisulfite Sequencing data will be deposited as *.fastq files. Affymetrix GeneChip Human Gene 1.0 ST and Cytoscan HD microarray data will be deposited as *.cel files.

Acknowledgements

We thank Drs. Jennifer Black and Michael Green of UNMC for critical review of the manuscript. We thank the following core facilities for support: University of Buffalo Center of Excellence Genomics and Bioinformatics, RPCI Bioinformatics and Genomics, UNMC DNA Sequencing and Microarray and Bioinformatics and Systems Biology (both received support from NIGMS 8P20GM103427 and P20GM103471), and Epigenomics, and University of Utah Genomics. A.R.K. was supported by NIH RO1CA116674, DOD OCRP W81XWH-12-1-0456, Otis Glebe Medical Research Foundation, Betty J. and Charles D. McKinsey Ovarian Cancer Research Fund, Fred & Pamela Buffett Cancer Center (NCI Cancer Center Support Grant P30 CA036727), and RPCI Alliance Foundation. C.J.B. was supported by NIH T32CA009476. S.N.A. was supported by NIH T32CA108456. KO was supported by NCI Cancer Center Support Grant P30 CA016056, NIH 1R01CA158318, and RPCI-UPCI Ovarian Cancer SPORE P50CA159981, and the Roswell Park Alliance Foundation.

Author Contributions

W.Z. and A.R.K. designed the research; W.Z., D.K., and C.J.B. performed the research; S.N.A., and K.O. contributed reagents; W.Z., D.K., S.P., C.G., C.J.B., A.M., and A.R.K. analyzed the data; and W.Z, D.K., and A.R.K. wrote the paper.

Competing Financial Interests

The authors declare no competing financial interests.

Figure Legends

Figure 1. *LINE-1* **hypomethylation is associated with EOC disease progression and reduced survival.** *LINE-1* methylation was determined by sodium bisulfite pyrosequencing. **(a)** *LINE-1* methylation vs. clinical stage. (**b**) *LINE-1* methylation vs. pathological grade. For **a** and **b**, mean +/- SD is plotted, and Mann Whitney p-values are indicated. **(c,d)** Kaplan-Meier survival analyses and log rank test p-values of EOC patients separated based on tumor *LINE-1* methylation values. **(c)** *LINE-1* methylation vs. overall survival. Patients were separated into three groups based on *LINE-1* methylation values: low (<51.0%), middle (51.0%-66.5%), and high (>66.5%). A key for the three groups, and the number of patients in each, is shown. **(d)** *LINE-1* methylation vs. disease-free survival. Patients were separated into three groups based on *LINE-1* methylation values: low (<51.0%), middle (51.0%-67.0%), and high (>67.0%). A key for the three groups, and the number of patients in each, is shown.

Fig 2. Distinct patterns of gene expression in GDHO(+) EOC. (a) *LINE-1* methylation in sample groups used for Affymetrix gene expression analyses: bulk normal ovary (NO), GDHO(+) EOC (*LINE-1* hypomethylated), GDHO(–) EOC (*LINE-1* hypermethylated). **(b)** Hierarchical clustering heat map of genes differentially expressed (p<0.01) between GDHO(+) vs. GDHO(–) EOC (all samples). **(c)** Hierarchical clustering heat map of genes differentially expressed between disease-matched (age-matched, stage 3/4, grade 3, serous EOC) GDHO(+) vs. GDHO(–) EOC. The number of genes showing significant differential expression in each comparison is indicated. Sample identities are indicated at top (see key). **(d)** Selected IPA data for the GDHO(+) vs. GDHO(–) EOC microarray comparison.

Fig 3. *CCNE1* and FOXM1 overexpression in GDHO(+) EOC. (a) *CCNE1* expression in NO, GDHO(+) EOC, and GDHO(–) EOC, as determined by Affymetrix microarray. **(b)** *FOXM1* expression in NO, GDHO(+) EOC, and GDHO(–) EOC, as determined by Affymetrix

microarray. Means +/− SD are plotted, and Mann Whitney p-values are indicated. **(c)** *FOXM1* mRNA expression vs. *LINE-1* methylation in EOC. *FOXM1* expression was measured by RT-qPCR while *LINE-1* methylation was measured by pyrosequencing. Spearman test results and p-value are shown. **(d)** Western blot analysis of FOXM1 protein expression in NO, GDHO(+) EOC, and GDHO(−) EOC.   The two blots are comprised of different sets of samples. Ponceau S staining is shown as a loading control.

Fig 4. Differential expression of CG antigen genes in GDHO(+) EOC and regulation by DNA methylation. CG antigen gene list was obtained from the CT gene database on March 19, 2015: http://www.cta.lncc.br/.   **Left:** the proportion of CG genes (i.e., the corresponding Affymetrix probes) differentially expressed between GDHO(+) and GDHO(−) EOC (p<0.01) as determined by Affymetrix microarray. **Right:** Direction of expression change for the DEGs in all samples or disease-matched samples, their chromosomal localization, and the effect of decitabine (DAC) treatment on their expression in EOC cell lines, as determined by Affymetrix microarray.

Fig 5. DNA methylome analysis of GDHO(+) EOC. (a) Dendogram showing relatedness of OSE, FTE, GDHO(+) EOC, and GDHO(−) EOC methylomes, using Illumina 450K beadarray analysis. Data for the 1000 most variable CpG sites is shown. **(b)** Principal component analysis of sample groups using all 450K data.   **(c)** Dendogram showing relatedness of OSE, FTE, GDHO(+) EOC, and GDHO(−) EOC methylomes, using Agilent Sure Select Methyl-seq. The 1000 most variable CpG sites is shown. **(d)** Principal component analysis of sample groups using all Methyl-seq data. **(e)** Total genomic methylation of OSE, FTE and EOC sample groups as determined by 450K. Median and interquartile ranges are plotted, and the unpaired two-tailed t-test p-values are shown. **(f)** Total genomic methylation levels of OSE, FTE and EOC sample groups as determined by Methyl-seq.

Fig 6. Hypomethylated genomic blocks (hypo blocks) in GDHO(+) EOC. (a) Circos plot (http://circos.ca/) of Methyl-seq data of chromosome 11 for OSE, FTE, GDHO(+) EOC, and GDHO(−) EOC. Methyl-seq data for each sample group, hypo blocks, late replicating regions, and LADs are indicated (see panel **b** for key). The enclosed region of the circos plot is enlarged in panel **b**. **(b)** Methyl-seq data for a selected region of chromosome 11 for OSE, FTE, GDHO(+) EOC, and GDHO(−) EOC. Calculated hypo block regions are indicated at bottom, along with UCSC genome browser view of lamina-associated domains (LADs), late-replicating (Late rep) regions and gene positions. The Methyl-seq data is plotted on a scale from 0-100% methylation. **(c)** The relationship between hypo blocks and gene expression variability in GDHO(+) EOC. Gene expression variability at hypo blocks and non-block regions was determined from Affymetrix microarray data, by calculating the median average deviation (MAD) for individual gene expression values amongst the 20 GDHO(+) EOC samples. Hypo blocks were calculated from Methyl-seq data by comparing GDHO(+) EOC to normal epithelia (NE; FTE + OSE average value). The 2-tailed paired t-test p-value is shown. The results indicate enrichment of hypervariable gene expression inside of hypo blocks.

Fig 7. *DNMTs* and *UHRF1* expression in GDHO(+) EOC. (a) Affymetrix (microarray) gene expression data for *DNMT1*, *DNMT3A*, *DNMT3B*, *DNMT3L*, and *UHRF1*, normalized to *MKI67*, in GDHO(+) and GHDO(−) EOC. Median with interquartile range is plotted, and Mann-Whitney test p-values are shown. **(b-d)** Gene expression of **(b)** *DNMT1*, **(c)** *DNMT3A*, and **(d)** *DNMT3B*, normalized to *MKI67*, in EOC, as compared to *LINE-1* methylation in matched samples. Gene expression data were obtained by RT-qPCR, and *LINE-1* methylation was determined by bisulfite pyrosequencing. Spearman correlation analysis test results are shown.

Fig 8. Expression and methylation of repetitive elements (RE) in GDHO(+) EOC. (a)
Comparison of the expression and methylation of major sub-classes of REs in GDHO(+) vs.
GDHO(–) EOC. Expression data were obtained from RNA-seq and methylation data from
Methyl-seq. EOC83 and EOC91 are matched for stage and grade (stage 3C, grade 3). **(b)** RE
expression vs. methylation in GDHO(+) vs. GDHO (–) EOC, using the same samples as in
panel **a**. RE expression and methylation were determined by RNA-seq and Methyl-seq,
respectively. Data points correspond to the average % methylation and log2 expression change
across RE families. The results indicate that most RE families show increased expression and
decreased methylation in GDHO(+) EOC. **(c)** DNA hypomethylation at hypo blocks and/or all
RE, in GDHO(+) as compared to NE, as determined by Methyl-seq. **(d)** DNA hypomethylation at
hypo blocks and/or all *LINE-1* elements, in GDHO(+) as compared to NE, as determined by
Methyl-seq. In **c-d**, the median + interquartile range values are indicated. The results indicate
that most hypomethylation is due to hypo block formation rather than RE or *LINE-1*
hypomethylation.

**Fig 9. Chromosomal instability (CIN) and copy number alterations (CNA), are significantly
increased in GDHO(+) EOC, and CNA are enriched at hypo blocks. (a)** Expression of
CIN25 signature genes in the indicated sample comparisons.   Gene expression was
determined by Affymetrix microarrays, and log2 fold change values are shown. + p<0.05, ++
p<0.01. **(b)** CNA index in GDHO(+) EOC vs. GDHO(–) EOC. EOC samples are disease-
matched and represent HGSOC. CNA were determined using Affymetrix Cytoscan HD chips as
described in *Methods*. Median with interquartile range is plotted, and the Mann Whitney p-value
is shown. **(c)** CNA are enriched in hypo blocks. The total base pair overlap between CNA and
hypo blocks, or CNA outside hypo blocks (i.e. non-hypo block) regions, in GDHO(+) EOC.
Hypo blocks were determined by Methyl-seq analysis of GDHO(+) EOC vs. normal epithelia
(NE; OSE + FTE average), and CNA were determined for each tumor sample, as described in

panel **b**. Hypo block and non-hypo block regions were normalized according to the percent of

genomic coverage for each. The two-way ANOVA p-value is shown. The asterisk indicates that

the values are too low to be visible on the scale.

References

Jones, P. A. & Baylin, S. B. The epigenomics of cancer. *Cell* **128**, 683-692, doi:10.1016/j.cell.2007.01.029 (2007).

Feinberg, A. P. & Tycko, B. The history of cancer epigenetics. *Nature reviews. Cancer* **4**, 143-153, doi:10.1038/nrc1279 (2004).

Widschwendter, M. *et al.* Epigenetic stem cell signature in cancer. *Nature genetics* **39**, 157-158, doi:10.1038/ng1941 (2007).

Feinberg, A. P., Gehrke, C. W., Kuo, K. C. & Ehrlich, M. Reduced genomic 5-methylcytosine content in human colonic neoplasia. *Cancer research* **48**, 1159-1161 (1988).

Bedford, M. T. & van Helden, P. D. Hypomethylation of DNA in pathological conditions of the human prostate. *Cancer research* **47**, 5274-5276 (1987).

Gama-Sosa, M. A. *et al.* The 5-methylcytosine content of DNA from human tumors. *Nucleic acids research* **11**, 6883-6894 (1983).

Beck, C. R., Garcia-Perez, J. L., Badge, R. M. & Moran, J. V. LINE-1 elements in structural variation and disease. *Annual review of genomics and human genetics* **12**, 187-215, doi:10.1146/annurev-genom-082509-141802 (2011).

De Smet, C. *et al.* The activation of human gene MAGE-1 in tumor cells is correlated with genome-wide demethylation. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 7149-7153 (1996).

Woloszynska-Read, A. *et al.* Coordinated cancer germline antigen promoter and global DNA hypomethylation in ovarian cancer: association with the BORIS/CTCF expression ratio and advanced stage. *Clinical cancer research : an official journal of the American Association for Cancer Research* **17**, 2170-2180, doi:10.1158/1078-0432.CCR-10-2315 (2011).

Akers, S. N., Odunsi, K. & Karpf, A. R. Regulation of cancer germline antigen gene expression: implications for cancer immunotherapy. *Future oncology* **6**, 717-732, doi:10.2217/fon.10.36 (2010).

Berman, B. P. *et al.* Regions of focal DNA hypermethylation and long-range hypomethylation in colorectal cancer coincide with nuclear lamina-associated domains. *Nature genetics* **44**, 40-46, doi:10.1038/ng.969 (2012).

Hansen, K. D. *et al.* Increased methylation variation in epigenetic domains across cancer types. *Nature genetics* **43**, 768-775, doi:10.1038/ng.865 (2011).

Timp, W. *et al.* Large hypomethylated blocks as a universal defining epigenetic alteration in human solid tumors. *Genome medicine* **6**, 61, doi:10.1186/s13073-014-0061-y (2014).

Hon, G. C. *et al.* Global DNA hypomethylation coupled to repressive chromatin domain formation and gene silencing in breast cancer. *Genome research* **22**, 246-258, doi:10.1101/gr.125872.111 (2012).

Hinoue, T. *et al.* Genome-scale analysis of aberrant DNA methylation in colorectal cancer. *Genome research* **22**, 271-282, doi:10.1101/gr.117523.110 (2012).

Schulz, W. A., Steinhoff, C. & Florl, A. R. Methylation of endogenous human retroelements in health and disease. *Current topics in microbiology and immunology* **310**, 211-250 (2006).

Karpf, A. R. & Matsui, S. Genetic disruption of cytosine DNA methyltransferase enzymes induces chromosomal instability in human cancer cells. *Cancer research* **65**, 8635-8639, doi:10.1158/0008-5472.CAN-05-1961 (2005).

Eden, A., Gaudet, F., Waghmare, A. & Jaenisch, R. Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* **300**, 455, doi:10.1126/science.1083557 (2003).

Gaudet, F. *et al.* Induction of tumors in mice by genomic hypomethylation. *Science* **300**, 489-492, doi:10.1126/science.1083558 (2003).

Rodriguez, J. *et al.* Chromosomal instability correlates with genome-wide DNA demethylation in human primary colorectal cancers. *Cancer research* **66**, 8462-9468, doi:10.1158/0008-5472.CAN-06-0293 (2006).

Suzuki, K. *et al.* Global DNA demethylation in gastrointestinal cancer is age dependent and precedes genomic damage. *Cancer cell* **9**, 199-207, doi:10.1016/j.ccr.2006.02.016 (2006).

Daskalos, A. *et al.* Hypomethylation of retrotransposable elements correlates with genomic instability in non-small cell lung cancer. *International journal of cancer. Journal international du cancer* **124**, 81-87, doi:10.1002/ijc.23849 (2009).

Richards, K. L. *et al.* Genome-wide hypomethylation in head and neck cancer is more pronounced in HPV-negative tumors and is associated with genomic instability. *PloS one* **4**, e4941, doi:10.1371/journal.pone.0004941 (2009).

Epping, M. T. *et al.* The human tumor antigen PRAME is a dominant repressor of retinoic acid receptor signaling. *Cell* **122**, 835-847, doi:10.1016/j.cell.2005.07.003 (2005).

Karpf, A. R., Bai, S., James, S. R., Mohler, J. L. & Wilson, E. M. Increased expression of androgen receptor coregulator MAGE-11 in prostate cancer by DNA hypomethylation and cyclic AMP. *Molecular cancer research : MCR* **7**, 523-535, doi:10.1158/1541-7786.MCR-08-0400 (2009).

Oricchio, E. *et al.* Distinct roles for LINE-1 and HERV-K retroelements in cell proliferation, differentiation and tumor progression. *Oncogene* **26**, 4226-4233, doi:10.1038/sj.onc.1210214 (2007).

Timp, W. & Feinberg, A. P. Cancer as a dysregulated epigenome allowing cellular growth advantage at the expense of the host. *Nature reviews. Cancer* **13**, 497-510, doi:10.1038/nrc3486 (2013).

Wild, L. & Flanagan, J. M. Genome-wide hypomethylation in cancer may be a passive consequence of transformation. *Biochimica et biophysica acta* **1806**, 50-57, doi:10.1016/j.bbcan.2010.03.003 (2010).

Eswaran, J. *et al.* RNA sequencing of cancer reveals novel splicing alterations. *Scientific reports* **3**, 1689, doi:10.1038/srep01689 (2013).

Ting, D. T. *et al.* Aberrant overexpression of satellite repeats in pancreatic and other epithelial cancers. *Science* **331**, 593-596, doi:10.1126/science.1200801 (2011).

Vaughan, S. *et al.* Rethinking ovarian cancer: recommendations for improving outcomes. *Nature reviews. Cancer* **11**, 719-725, doi:10.1038/nrc3144 (2011).

Integrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609-615, doi:10.1038/nature10166 (2011).

Nephew, K. P., Balch, C., Zhang, S. & Huang, T. H. Epigenetics and ovarian cancer. *Cancer treatment and research* **149**, 131-146, doi:10.1007/978-0-387-98094-2_6 (2009).

Woloszynska-Read, A., Mhawech-Fauceglia, P., Yu, J., Odunsi, K. & Karpf, A. R. Intertumor and intratumor NY-ESO-1 expression heterogeneity is associated with promoter-specific and global DNA methylation status in ovarian cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* **14**, 3283-3290, doi:10.1158/1078-0432.CCR-07-5279 (2008).

Widschwendter, M. *et al.* DNA hypomethylation and ovarian cancer biology. *Cancer research* **64**, 4472-4480, doi:10.1158/0008-5472.CAN-04-0238 (2004).

Akers, S. N. *et al.* LINE1 and Alu repetitive element DNA methylation in tumors and white blood cells from epithelial ovarian cancer patients. *Gynecologic oncology* **132**, 462-467, doi:10.1016/j.ygyno.2013.12.024 (2014).

Whitfield, M. L., George, L. K., Grant, G. D. & Perou, C. M. Common markers of proliferation. *Nature reviews. Cancer* **6**, 99-106, doi:10.1038/nrc1802 (2006).

Teh, M. T. *et al.* FOXM1 induces a global methylation signature that mimics the cancer epigenome in head and neck squamous cell carcinoma. *PloS one* **7**, e34329, doi:10.1371/journal.pone.0034329 (2012).

Hua, K. T. *et al.* The H3K9 methyltransferase G9a is a marker of aggressive ovarian cancer that promotes peritoneal metastasis. *Molecular cancer* **13**, 189, doi:10.1186/1476-4598-13-189 (2014).

Wan, W. N. *et al.* ATAD2 is highly expressed in ovarian carcinomas and indicates poor prognosis. *Asian Pacific journal of cancer prevention : APJCP* **15**, 2777-2783 (2014).

Hayashi, A. *et al.* Type-specific roles of histone deacetylase (HDAC) overexpression in ovarian carcinoma: HDAC1 enhances cell proliferation and HDAC3 stimulates cell migration with downregulation of E-cadherin. *International journal of cancer. Journal international du cancer* **127**, 1332-1346, doi:10.1002/ijc.25151 (2010).

Yang, G. *et al.* Aurora kinase A promotes ovarian tumorigenesis through dysregulation of the cell cycle and suppression of BRCA2. *Clinical cancer research : an official journal of the American Association for Cancer Research* **16**, 3171-3181, doi:10.1158/1078-0432.CCR-09-3171 (2010).

Jimeno, A. *et al.* Phase I study of ON 01910.Na, a novel modulator of the Polo-like kinase 1 pathway, in adult patients with solid tumors. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* **26**, 5504-5510, doi:10.1200/JCO.2008.17.9788 (2008).

Wrzeszczynski, K. O. *et al.* Identification of tumor suppressors and oncogenes from genomic and epigenetic features in ovarian cancer. *PloS one* **6**, e28503, doi:10.1371/journal.pone.0028503 (2011).

Liao, Y. P. *et al.* Hypomethylation signature of tumor-initiating cells predicts poor prognosis of ovarian cancer patients. *Human molecular genetics* **23**, 1894-1906, doi:10.1093/hmg/ddt583 (2014).

Sandoval, J. *et al.* Validation of a DNA methylation microarray for 450,000 CpG sites in the human genome. *Epigenetics : official journal of the DNA Methylation Society* **6**, 692-702 (2011).

Lee, E. J. *et al.* Targeted bisulfite sequencing by solution hybrid selection and massively parallel sequencing. *Nucleic acids research* **39**, e127, doi:10.1093/nar/gkr598 (2011).

Robertson, K. D., Keyomarsi, K., Gonzales, F. A., Velicescu, M. & Jones, P. A. Differential mRNA expression of the human DNA methyltransferases (DNMTs) 1, 3a and 3b during the G(0)/G(1) to S phase transition in normal and tumor cells. *Nucleic acids research* **28**, 2108-2113 (2000).

Bostick, M. *et al.* UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science* **317**, 1760-1764, doi:10.1126/science.1147939 (2007).

Carter, S. L., Eklund, A. C., Kohane, I. S., Harris, L. N. & Szallasi, Z. A signature of chromosomal instability inferred from gene expression profiles predicts clinical outcome in multiple human cancers. *Nature genetics* **38**, 1043-1048, doi:10.1038/ng1861 (2006).

Feinberg, A. P. & Vogelstein, B. Hypomethylation distinguishes genes of some human cancers from their normal counterparts. *Nature* **301**, 89-92 (1983).

Pfeifer, G. P., Xiong, W., Hahn, M. A. & Jin, S. G. The role of 5-hydroxymethylcytosine in human cancer. *Cell and tissue research* **356**, 631-641, doi:10.1007/s00441-014-1896-7 (2014).

Woloszynska-Read, A. *et al.* BORIS/CTCFL expression is insufficient for cancer-germline antigen gene expression and DNA hypomethylation in ovarian cell lines. *Cancer immunity* **10**, 6 (2010).

Ferreira, H. J. *et al.* Epigenetic loss of the PIWI/piRNA machinery in human testicular tumorigenesis. *Epigenetics : official journal of the DNA Methylation Society* **9** (2013).

Mudbhary, R. *et al.* UHRF1 Overexpression Drives DNA Hypomethylation and Hepatocellular Carcinoma. *Cancer cell* **25**, 196-209, doi:10.1016/j.ccr.2014.01.003 (2014).

Aran, D., Toperoff, G., Rosenberg, M. & Hellman, A. Replication timing-related and gene body-specific methylation of active human genes. *Human molecular genetics* **20**, 670-680, doi:10.1093/hmg/ddq513 (2011).

Desjobert, C. *et al.* Combined analysis of DNA methylation and cell cycle in cancer cells. *Epigenetics : official journal of the DNA Methylation Society*, 0, doi:10.1080/15592294.2014.995542 (2014).

Shipony, Z. *et al.* Dynamic and static maintenance of epigenetic memory in pluripotent and somatic cells. *Nature*, doi:10.1038/nature13458 (2014).

Peters, S. L. *et al.* Tumor suppressor functions of Dnmt3a and Dnmt3b in the prevention of malignant mouse lymphopoiesis. *Leukemia* **28**, 1138-1142, doi:10.1038/leu.2013.364 (2014).

Jeong, S. *et al.* Selective anchoring of DNA methyltransferases 3A and 3B to nucleosomes containing methylated DNA. *Molecular and cellular biology* **29**, 5366-5376, doi:10.1128/MCB.00484-09 (2009).

Wen, B., Wu, H., Shinkai, Y., Irizarry, R. A. & Feinberg, A. P. Large histone H3 lysine 9 dimethylated chromatin blocks distinguish differentiated from embryonic stem cells. *Nature genetics* **41**, 246-250, doi:10.1038/ng.297 (2009).

Bai, X. *et al.* Clinicopathological significance and prognostic value of DNA methyltransferase 1, 3a, and 3b expressions in sporadic epithelial ovarian cancer. *PloS one* **7**, e40024, doi:10.1371/journal.pone.0040024 (2012).

Cadieux, B., Ching, T. T., VandenBerg, S. R. & Costello, J. F. Genome-wide hypomethylation in human glioblastomas associated with specific copy number alteration, methylenetetrahydrofolate reductase allele status, and increased proliferation. *Cancer research* **66**, 8469-8476, doi:10.1158/0008-5472.CAN-06-1547 (2006).

Coulie, P. G., Van den Eynde, B. J., van der Bruggen, P. & Boon, T. Tumour antigens recognized by T lymphocytes: at the core of cancer immunotherapy. *Nature reviews. Cancer* **14**, 135-146, doi:10.1038/nrc3670 (2014).

Odunsi, K. *et al.* Epigenetic potentiation of NY-ESO-1 vaccine therapy in human ovarian cancer. *Cancer immunology research* **2**, 37-49, doi:10.1158/2326-6066.CIR-13-0126 (2014).

Ciriello, G. *et al.* Emerging landscape of oncogenic signatures across human cancers. *Nature genetics* **45**, 1127-1133, doi:10.1038/ng.2762 (2013).

Tang, M. H. *et al.* Major chromosomal breakpoint intervals in breast cancer co-localize with differentially methylated regions. *Frontiers in oncology* **2**, 197, doi:10.3389/fonc.2012.00197 (2012).

Woloszynska-Read, A. *et al.* DNA methylation-dependent regulation of BORIS/CTCFL expression in ovarian cancer. *Cancer immunity* **7**, 21 (2007).

James, S. R., Link, P. A. & Karpf, A. R. Epigenetic regulation of X-linked cancer/germline antigen genes by DNMT1 and DNMT3b. *Oncogene* **25**, 6975-6985, doi:10.1038/sj.onc.1209678 (2006).

Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature protocols* **7**, 562-578, doi:10.1038/nprot.2012.016 (2012).

Zhang et al., Fig.1.

Zhang et al., Fig.2.

a

d

Ingenuity Pathway Data	Score/P-value	# molecules
Networks		
Cell cycle, cell assembly, DNA replication	42	
Diseases		
Cancer	10^{-19}-10^{-4}	414
Reproductive system	10^{-17}-10^{-4}	154
Molecular and Cellular Functions		
Cell cycle	10^{-16}-10^{-4}	122
DNA replication, recombination, repair	10^{-15}-10^{-4}	47
Cell growth and proliferation	10^{-11}-10^{-4}	206

b

1696 DEGs, p<0.01

c

752 DEGs, p<0.01

GDHO(+) EOC   GDHO(–) EOC

Zhang et al., Fig.3.

Zhang et al., Fig.4.

Zhang et al., Fig.5.

Zhang et al., Fig.6.

Zhang et al., Fig. 7.

Zhang et al., Fig. 8.

Zhang et al., Fig. 9.

Table 1. Correlation of EOC hypomethylated blocks with genomic features[1,2]

Genomic Region	# of regions	Size of regions (bp)	% of genome	% regions in hypo blocks	Correlation	Jaccard Test P-value[3,4]
EOC Hypo Blocks	2,208	8.81E+08	29	100	N/A	N/A
LADs	1,231	1.13E+09	37	36	Direct	<0.01
Early Replicating	910	7.10E+08	23	25	Indirect	<0.01
Late Replicating	1,155	5.81E+08	19	36	Direct	<0.01
CpG islands	27,537	2.10E+07	1	32	Direct	<0.01
Genes	28,489	1.41E+09	46	32	Direct	<0.01

	# binding sites	# binding sites in hypo blocks	% binding sites in hypo blocks	Fold Enrichment	Correlation	Projection Test P-value
EZH2 Sites	14,818	6,103	41	1.41	Direct	<0.01
SUZ12 Sites	5,772	2,497	43	1.48	Direct	<0.01
CTCF Sites	162,209	52,216	32	1.10	Direct	<0.01

	# peaks	# peaks in hypo blocks	% peaks in hypo blocks	Fold Enrichment	Correlation	Jaccard Test P-value
H3K4me3	33,116	8,655	26	0.90	Indirect	<0.01
H3K9me3	49,328	18,264	37	1.28	Direct	<0.01
H3K27ac	67,989	17,935	26	0.91	Indirect	<0.01
H3K27me3	40,126	15,428	38	1.32	Direct	<0.01
H3K36me3	33,342	6,303	19	0.65	Indirect	<0.01

	# of repeats	# repeats in hypo blocks	% repeats in hypo blocks	Fold Change	Correlation	Jaccard Test P-value
LINE-1	951,780	264,352	28	0.96	Direct	<0.01
Alu/SINE	1,194,734	292,495	24	0.84	Indirect	<0.01
Satellites	6,775	1,052	16	0.64	Indirect	<0.01

1 Hypo blocks were determined from Methyl-seq comparison of NE (OSE + FTE average) to GDHO(+) EOC

2 Genomic features were retrieved from public databases, as described in *Methods*

3 Statistical tests were performed using GenometriCorr (Favorov et. al, *PLoS Computational Biology*, 2012), with hypo block coordinates used as the reference value.

4 All tests, other than CpG islands, gave the same result when using hypo blocks as query and the genomic features as the reference value. In contrast, the association with CpG islands was not significant.