

Associations between dietary patterns and gene expression pattern in peripheral blood mononuclear cells: a cross-sectional study

Jacob J. Christensen, PhD^{1,2}, Stine M. Ulven, PhD², Magne Thoresen, PhD³, Kenneth Westerman, PhD^{4,5}, Kirsten B. Holven, PhD^{1,2}, and Lene F. Andersen, PhD²

¹Norwegian National Advisory Unit on Familial Hypercholesterolemia, Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, Forskningsveien 2B, 0373 Oslo, Norway

²Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Sognsvannsveien 9, 0372 Oslo, Norway

³Department of Biostatistics, Institute of Basic Medical Sciences, University of Oslo, Sognsvannsveien 9, 0372 Oslo, Norway

⁴Clinical and Translation Epidemiology Unit, Department of Medicine, Massachusetts General Hospital, Boston, MA, USA

⁵Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA

Correspondence: Jacob J. Christensen, PhD, j.j.christensen@medisin.uio.no

Short running head: Dietary patterns and gene expression

Abbreviations: DP, dietary pattern; PBMC, peripheral blood mononuclear cell; WGCNA, weighted gene co-expression network analysis; GO, gene ontology; PPI, protein-protein interaction

1 **Abstract**

2 **Background:** Diet may alter gene expression in immune cells involved in cardio-metabolic
3 disease susceptibility. However, we still lack a robust understanding of the association between
4 diet and immune cell-related gene expression in humans.

5 **Objective:** Our objective was to examine the associations between dietary patterns (DPs) and
6 gene expression profiles in peripheral blood mononuclear cells (PBMCs) in a population of
7 healthy, Norwegian adults.

8 **Methods:** We used factor analysis to define *a posteriori* DPs from food frequency questionnaire-
9 based dietary assessment data. In addition, we derived interpretable features from microarray-
10 based gene expression data (13 967 transcripts) using two algorithms: CIBERSORT for estimation
11 of cell subtype proportions, and weighted gene co-expression network analysis (WGCNA) for
12 cluster discovery. Finally, we associated DPs with either CIBERSORT-predicted PBMC leukocyte
13 distribution or WGCNA gene clusters using linear regression models. All analyses were gender-
14 stratified (n = 130 women and 105 men).

15 **Results:** We detected three DPs that broadly reflected *Western*, *Vegetarian*, and *Low*
16 *carbohydrate* diets. CIBERSORT-predicted percentage of monocytes associated strongly and
17 negatively with the *Vegetarian* DP in both women and men. For women, the *Vegetarian* DP
18 associated most strongly with a large gene cluster consisting of 600 genes mainly involved in
19 regulation of DNA transcription. For men, the *Western* DP inversely associated most strongly
20 with a smaller cluster of 36 genes mainly involved in regulation of metabolic and inflammatory

21 processes. In subsequent protein-protein interaction network analysis, the most important
22 *driver genes* within these WGCNA gene clusters seemed to physically interact in biological
23 networks.

24 **Conclusions:** DPs may affect percentage monocytes and regulation of key biological processes
25 within the PBMC pool. Although the present findings are exploratory, our analysis pipeline
26 serves a useful framework for studying the association between diet and gene expression.

27 **Keywords:** dietary patterns, gene expression, PBMC, microarray, monocytes, CIBERSORT,
28 WGCNA

29 **Introduction**

30 Cardio-metabolic diseases are the main causes of death worldwide (1). They are mainly caused
31 by life-long exposure to classical risk factors such as obesity, hypertension, dyslipidemia and
32 dysglycemia (2). Diet affects these risk factors and thereby contributes to the rate of disease
33 progression (3). Diet can also influence gene expression in immune cells directly, and so
34 potentially affect cardio-metabolic disease susceptibility (4–6). However, we still lack a thorough
35 understanding of the association between diet and immune cell-related gene expression.

36 Free-living humans consume a variety of foods in combination. To capture this variation
37 meaningfully, we often define so-called dietary patterns (DPs). *A posteriori* DPs are data-driven;
38 they are defined based on the co-consumption of foods in the population under study (7).

39 Naturally, *a posteriori* DPs reflect local food culture and have high internal validity. As such, a
40 *a posteriori* DPs may constitute robust measures of global diet exposure, and could be used to
41 strengthen the reliability of associations between diet and biomarkers within a population. This
42 may be especially relevant in order to examine high-variance biomarkers, such as gene
43 expression profiles.

44 Peripheral blood mononuclear cells (PBMCs) are directly involved in the underlying
45 pathophysiology of cardio-metabolic disease (8). They represent a mixture of cells that are
46 transiently part of a specialized niche in the circulation, of which some move to sites of
47 inflammation. Affected by a number of input signals, PBMCs adapt to their environment; dietary
48 metabolites, interleukins and chemokines, classical risk factors, and a host of other factors all
49 influence the transcriptome of the PBMCs (9).

50 Many previous studies in humans that have associated diet with PBMC gene expression have
51 used a classical gene expression-wide association (gxWA) strategy (10,11). The underlying
52 correlation structure of the transcriptome, however, provides an opportunity to improve upon
53 gxWA methods. Biologically-relevant dimensionality reduction algorithms, such as CIBERSORT
54 and weighted gene co-expression network analysis (WGCNA), simplify whole-genome gene
55 expression matrices into interpretable features (12,13). These methods also increase the signal-
56 to-noise ratio and thereby robustness of the features, while they simultaneously reduce the
57 multiple testing burden (14).

58 The objective of the present study was to examine the associations between a posteriori-
59 defined DPs and derived gene expression features in PBMCs in a population of healthy,
60 Norwegian adults. We hypothesized that DPs would associate with PBMC gene expression, and
61 that the associations would point to specific biological mechanisms that potentially mediate the
62 effects of diet on cardio-metabolic diseases.

63 **Subjects and methods**

64 ***Study design and participants***

65 The present study is based on cross-sectional data from the screening visit of a randomized
66 controlled dietary intervention, presented in detail elsewhere (15). We included all participants
67 from whom we had both dietary assessment data and PBMC gene expression data, in addition
68 to standard clinical and biochemical measurements. After excluding four participants with self-
69 reported energy intake above 25 MJ/d, we included 235 participants in the analyses (n = 130
70 women, n = 105 men).

71 The subject characteristics are presented in **Table 1**. Briefly, the men were younger than the
72 women, but had a more unhealthy body composition and subsequent clinical sequelae. Both
73 genders had moderate hypercholesterolemia.

74 ***Data types***

75 We used a food-frequency questionnaire (FFQ) to assess habitual food intake from the
76 preceding year (16). From the originally 323 food items, we removed 41 items due to unclear
77 interpretation, and grouped the remaining 282 into 33 food groups, based on food category and
78 nutrient content (**Table 2**). Self-reported intake of foods and nutrients are presented in
79 **Supplementary Table 1, Supplementary Table 2** and **Supplementary Table 3**. Furthermore, we
80 collected PBMCs and extracted RNA according to standardized protocols, as previously
81 described (6).

82 See Supplemental Methods for an extended description of the data types.

83 ***Statistical and bioinformatics analyses***

84 Here we describe the statistical and bioinformatic analyses related to DPs, gene expression
85 clusters, and statistical modeling. All analyses were performed in R version 3.6.2 (17). We refer
86 to R packages and functions where appropriate, and using the following notation:
87 `package::function`. Important deviations from default function setting are written in
88 parentheses.

89 The flow of the analysis pipeline is outlined in **Supplementary Figure 1**. Women and men were
90 analyzed separately, as preliminary analyses suggested a strong gender-related signal in both
91 the DPs and gene expression dataset.

92 ***Dietary patterns***

93 We used a combination of principal component analysis (PCA) and factor analysis to determine
94 DPs. Factor analysis is a dimensionality reduction method similar to PCA, but it results in more
95 interpretable features. However, because factor analysis is informed by the same covariance
96 matrix as PCA, we used PCA-derived component variances (`stats::prcomp`) to determine a
97 meaningful number of factors to retain; the results are presented in **Supplementary Figure 2**.
98 For both genders, the eigenvalue-one criterion suggested around 12 principal components
99 (PCs), but there was little change between components from component 3-5 and onwards; the
100 scree test suggested around 3-5 components; the per component variance explained suggested
101 that about 7-17 % of the variance could be explained until about three components, and then
102 stabilized at 4-5 % at 4-6 components, with little change thereafter. We decided to extract three

103 components using factor analysis (`stats::factanal`). The extracted DPs made sense from a
104 dietary perspective (**Figure 1**), thus meeting the interpretability criterion.

105 *Gene expression features*

106 Two main mechanisms are central in studies of diet-related associations with cardio-metabolic
107 disease mechanisms in PBMCs: dietary effects on leukocyte subset distributions, and biological
108 modulation independent of leukocyte subset distribution. As a result, we performed analyses to
109 examine each of these aspects, as outlined in the upper right corner of Supplementary Figure 1.

110 Leukocyte subsets

111 We used CIBERSORT to perform *in silico* flow cytometry (13). This method uses support vector
112 regression to conduct robust deconvolution of a heterogenous cell population, and returns
113 predicted relative levels of various cell subsets. We used the raw, untransformed, whole-
114 genome gene expression data matrix as input. Although the algorithm provides 22 leukocyte
115 subsets, we filtered on the top most relevant cell types for the PBMC population, mainly
116 monocyte and lymphocytes subsets, and thereby retained 12 cell subsets (**Supplementary**
117 **Figure 3**). Note that although we had standard blood cell differential counts available,
118 CIBERSORT resulted in a richer set PBMC cell subsets unique to the gene expression profile of
119 each sample.

120 Gene expression clusters

121 We used WGCNA to identify highly correlated (“co-expressed”) clusters of genes (18). The
122 WGCNA package (CRAN, Bioconductor) provides a well-established and popular framework to
123 perform the WGCNA analysis (12). The details of the implementation can be found in (12); in

124 Supplemental Methods we give a brief outline of key steps in the WGCNA-based gene
125 expression cluster analysis pipeline.

126 To both avoid confounding by sex chromosomes and to aid compatibility with the dietary data,
127 we performed the analysis separate for women and men. We examined the stability and validity
128 of the resulting gene expression clusters between genders with module preservation statistics
129 (19). Also, to avoid confounding by cell types, we adjusted for measured percentage monocytes
130 and lymphocytes (standard differential counts), and extracted the residuals. Therefore, input for
131 this analysis was the *residuals for the complete gene expression matrix* ($p = 13967$ variables)),
132 after removing the main effect of monocytes and lymphocytes.

133 In order to highlight a few of the more important genes within interesting clusters, we
134 performed a *driver gene* analysis. First, we calculated *cluster membership*, which is defined as
135 the absolute correlation between gene expression and cluster eigengene, and can be
136 interpreted as the degree to which each gene contributes to that cluster's overall behavior, and
137 contributes to its variation. Secondly, we calculated *DP significance*, which is the absolute
138 correlation between gene expression and DP score. A positive correlation between cluster
139 membership and DP significance indicates that those genes that drive the variation in the
140 cluster eigengene are the same that drive the association with the specific DP (*driver genes*).
141 Finally, to rank driver genes, driver gene estimates were calculated as the sum of the cluster
142 membership and DP significance.

143 In order to describe relevant gene expression clusters biologically, we performed gene ontology
144 (GO) enrichment using the GO Consortium database (20,21). Also, to link statistical findings with

145 existing biological knowledge, we performed protein-protein interaction (PPI) network analyses
146 using The Protein Interaction Network Analysis (PINA) 2.0 database (22).

147 *Linear models*

148 We associated DPs with two types of outcomes: CIBERSORT-predicted cell counts, and the
149 eigengenes from the gene expression clusters using linear models (Supplementary Figure 1).

150 **Supplementary Figure 4** shows the directed acyclic graphs (DAGs) used in model development.

151 We used the open-access dagitty.net/dags web-resource to evaluate these relationships.

152 Minimal sufficient adjustment sets for estimating the *total effect* of dietary pattern on gene
153 expression were age and education (three levels: lower, middle, higher). For predicted cell
154 counts, we additionally adjusted for adiposity (total fat mass, measured by bioelectrical
155 impedance analysis) in sensitivity analyses. Also, in sensitivity analyses for the gene expression
156 clusters, we estimated the *direct effect* (see Supplemental Methods).

157 Note that for all models, technical covariates were considered in upstream batch correction
158 (Supplementary Figure 1). Percentage of total leukocyte count of monocytes and lymphocytes
159 (which make up the pool of PBMC subsets) were adjusted for in the gene expression pre-
160 processing pipeline (as shown in Supplementary Figure 1), prior to WGCNA only. Finally, to aid
161 interpretation of the results, we normalized (`base::scale`) both DP scores and cluster
162 eigengenes to a standard normal distribution (mean = 0, sd = 1) before modeling.

163 *Miscellaneous*

164 The dietary intervention study was powered to detect a significant change in LDL-C (15);
165 however, this does not apply to the present exploratory study. Because this study is exploratory,

166 we did not evaluate associations by standard significance level cut-offs. Instead, we evaluated
167 the strength and direction of associations, and their interrelations.

168 **Results**

169 ***Construction and description of study features***

170 *Dietary patterns*

171 First, we constructed gender-specific DPs from self-reported FFQ data, yielding three DPs
172 (Figure 1 and **Supplementary Table 4**). We considered the patterns to reflect typical *Western*
173 (DP1), *Vegetarian* (DP2), and *Low carbohydrate* (DP3) diets. These three DPs explained 14.1, 8.0
174 and 6.6 %, and 16.6, 9.3 and 7.0 % of the variance, for women and men, respectively. Although
175 there were some overlap, the *Vegetarian* and *Low carbohydrate* DPs were more unique to each
176 gender compared to the *Western* DP. This was also supported by the DP loading for various
177 foods (**Supplementary Figure 5**). For both genders, the *Western* DP associated with intake of
178 meat and eggs, fastfood, snacks, dairy, and fiber-poor carbohydrate sources. The *Vegetarian* DP
179 associated positively with a number of foods perceived as healthy, including plant foods, whole
180 grains, nuts and seeds, and tea. Additionally, the association with animal products, fast food,
181 dairy, and fiber-poor carbohydrate foods was low or negative. For women, the association with
182 high-fat dairy and snacks was slightly positive. The *Low carbohydrate* DP was generally a mixture
183 of the two former, reflected in positive associations for both plants and animal products. The
184 association with fastfoods, snacks and carbohydrate-rich foods, however, was negative. Wine
185 associated positively, whereas sweet beverages associated negatively with the *Low*
186 *carbohydrate* DP for women and men, respectively.

187 In addition to the direct link with food intake, the DP scores correlated with both macronutrient
188 intake (**Supplementary Figure 6**) and clinical variables (**Supplementary Figure 7**). The *Western*

189 DP correlated with energy intake and negatively with fiber intake in both genders. The
190 *Vegetarian* DP correlated positively with fiber and negatively with saturated fat intake in men.
191 In women, the *Vegetarian* DP correlated weakly, but positively, with energy, healthy fats, fiber
192 and sugar. The *Low carbohydrate* DP was negatively associated with carbohydrate and sugar
193 intake in both genders, and with higher protein and fat intake in men.

194 For the clinical variables, the negative association between *Western* DP and age was most
195 notable, which indicates that the younger part of the study sample adhere to a more unhealthy
196 diet. Additionally, The *Vegetarian* DP associated negatively with multiple obesity-related
197 markers, including immune cells and CRP. Again, the *Low carbohydrate* DP was a mixture of the
198 two, with positive correlations for age and lipids.

199 *Leukocyte subsets*

200 We used the CIBERSORT algorithm to computationally estimate the distribution of 12 leukocyte
201 subsets (13). As expected, predicted leukocyte cell proportions associated with multiple clinical
202 variables, although most notably for the differential count measures and obesity-related
203 measures (**Supplementary Figure 8**).

204 *Gene expression clusters*

205 Using the WGCNA algorithm, we detected 45 and 37 unique gene expression clusters for
206 women and men, respectively, which by default were named different colors (12). Although
207 there were large differences in cluster size (range = 67-307 and 85-438 genes for women and
208 men, respectively), most clusters explained a large proportion of the variance of the genes they
209 comprised (range = 32-39 and 33-40 % for women and men, respectively) (**Supplementary**
210 **Figure 9A and B, and Supplementary Table 5**). For men, explained variance inversely associated

211 with cluster size (**Supplementary Figure 9C**). In addition, genes in all clusters were generally
212 distributed over all chromosomes, with certain exceptions, such as chromosome 1 and 19
213 (**Supplementary Figure 9D**). The gene expression clusters displayed some correlation within
214 each gender, but they could largely be considered unique features (**Supplementary Figure 10**).
215 Between genders, the module preservation was acceptable for most medium- and large-sized
216 clusters, and poor for the smaller clusters (**Supplementary Figure 11, Supplementary Figure 12,**
217 **Supplementary Figure 13**).

218 Numerous gene expression clusters correlated with clinical phenotypes (**Supplementary Figure**
219 **14**). Most prominent was the global associations with body composition- and lipid-related
220 markers.

221 ***Associations of derived gene expression features with dietary patterns***

222 *Dietary patterns and leukocyte subsets*

223 Predicted percentage of monocytes associated negatively with the *Vegetarian* DP for both
224 women ($\beta = -0.21$, $P = 0.02$) and men ($\beta = -0.33$, $P = 0.0008$) (**Figure 2** and **Supplementary**
225 **Figure 15**), suggesting a link between this particular cell subset and diet. Interestingly, when
226 adjusting for adiposity, this association was only attenuated for women ($\beta = -0.15$, $P = 0.11$ for
227 women, and $\beta = -0.33$, $P = 0.001$ for men).

228 *Dietary patterns and gene expression clusters*

229 In general, relatively few associations were evident between DP scores and gene expression
230 cluster eigengenes (**Figure 3**). For women, the positive association between the *Vegetarian* DP
231 and the yellow cluster was strongest. The yellow cluster contained 600 genes involved in

232 regulation of transcription (**Supplementary Figure 16**). For men, the *Western* DP associated with
233 multiple clusters, of which the association with the darkmagenta cluster was strongest. This
234 cluster contained 36 genes related to metabolic and inflammatory processes, including
235 sterol/cholesterol transport (Supplementary Figure 16). Similarly, both the pink and greenyellow
236 clusters associated negatively with the *Western* DP, although not as strongly as darkmagenta.
237 The pink cluster consisted of 475 genes involved in regulation of viral processes,
238 endosome/vacuolar transport, UDP-GlcNAc metabolism, and monocyte and lymphocyte
239 stimulation. On the other hand, the greenyellow cluster consisted of 338 genes involved in
240 regulation of protein synthesis and degradation, and acyl carnitine transport. The top 20 most
241 enriched GO terms (for all three ontologies) for the top most significant cluster for each gender
242 are listed in **Supplementary Table 6**.

243 ***Identification of driver genes***

244 Next, we examined the most relevant gene expression clusters more in detail, using a driver
245 gene analysis to identify genes with both high DP significance and high cluster membership.
246 Interestingly, DP significance and cluster membership associated strongly (**Figure 4A and B**, and
247 **Supplementary Table 7**), which suggests that genes that associated with DPs were also among
248 the most important parts of the clusters that associated with that DP.

249 The five top driver genes for the association between the *Vegetarian* DP and the yellow cluster
250 in women were GIMAP7 (GTPase, IMAP family member 7), ZNF200 (zinc finger protein 200),
251 LCMT2 (leucine carboxyl methyltransferase 2), GPR18 (G protein-coupled receptor 18), ASTE1
252 (asteroid homolog 1) (Figure 4A). Proteins from these genes regulate aspects of biosynthetic

253 processes, including cell signaling, DNA transcription and repair, and protein synthesis (20,21).
254 For these genes, the correlation coefficients with DP2 score were in the range 0.19 to 0.26 ($P =$
255 0.03 - 0.003), and with the cluster eigengene in the range 0.83 to 0.90 ($P < 0.001$)
256 (Supplementary Table 7). This means that women who consumed a *Vegetarian* DP tended to
257 have *higher* expression of these genes in PBMCs.

258 The five top driver genes for the association between the *Western* DP and darkmagenta cluster
259 in men were DMWD (DM1 locus, WD repeat containing), SYTL3 (synaptotagmin like 3), ABCA2
260 (ATP binding cassette subfamily A member 2), TSEN54 (tRNA splicing endonuclease subunit 54)
261 and C9orf142/XLS (XRCC4-like small protein) (Figure 4B). Proteins from these genes are involved
262 in lysosomal transport, cholesterol homeostasis, mRNA processing and DNA repair (20,21). The
263 correlation coefficients with DP1 score were in the range -0.29 to -0.22 ($P = 0.03 - 0.003$), and
264 with the cluster eigengene in the range 0.75 - 0.83 ($P < 0.001$) (Supplementary Table 7). This
265 means that men who consumed a *Western* DP tended to have *lower* expression of these genes
266 in PBMCs.

267 ***Identification of hub proteins***

268 Finally, to examine if these driver genes were part of physically interacting biological networks,
269 we filtered them through the PINA database (22). For the strongest associations for each
270 gender, we then created protein-protein interaction (PPI) networks (**Figure 4C** and **D**, and
271 **Supplementary Table 8**). These proteins can be considered *hub proteins*; they likely exert a
272 higher degree of control over the protein network, as more proteins physically interact with this

- 273 hub in order to influence signaling pathways. For women and men, key hub proteins included
- 274 PPARGC1B (PPARG coactivator 1 beta) and UBC (ubiquitin C), respectively.

275 **Discussion**

276 In the present study of 235 Norwegian adults, we detected novel associations between DPs and
277 gene expression features in PBMCs. Our results suggest that diet affects a number of specific
278 cell types and pathways, of which the most pronounced are: predicted proportion of
279 monocytes, regulation of transcription, and regulation of metabolic and inflammatory
280 processes.

281 ***We detected three DPs commonly consumed in Norway***

282 Using data-driven analyses, we detected three DPs commonly consumed in Norway: *Western-*
283 *type, Vegetarian-type, and Low carbohydrate-type* DP (Figure 1). These DPs were neither
284 unexpected nor surprising: Norwegian adults follow trends, and this includes the vegetarian and
285 low carbohydrate trends. In previous studies, similar names have been used to characterize the
286 detected DPs. In a cohort of Norwegian postmenopausal women, Markussen and co-workers
287 found four DPs, including the *Western* and *Vegetarian* DPs (23). In addition, they found a *High-*
288 *protein* pattern that resembled our *Low carbohydrate* pattern. Their DPs, similar to ours, share
289 characteristics and therefore also names, with DPs throughout Europe and the US. This
290 emphasizes an important point: although the DPs retained in factor analyses are never exactly
291 equal, as opposed to *a priori* methods, our three DPs share characteristics with many other DPs
292 both in Norway and elsewhere (7,23–25).

293 The three DPs associated with food items, nutrient intake and clinical parameters to give a
294 consistent picture of the DPs: in general, the *Low carbohydrate* DP appeared neutral compared

295 to the *Western*-type and *Vegetarian*-type DPs, which associated with a number of unhealthy or
296 healthy behaviors, respectively.

297 ***Vegetarian DP associated with monocytes***

298 The *Vegetarian* DP associated with CIBERSORT-predicted levels of monocytes (Figure 2),
299 suggesting that gene expression related to monocyte differentiation and activity may be
300 affected by diet. These results are corroborated by previous reports by others and us (26–29).
301 Craddock and co-workers recently reviewed the evidence that vegetarian diets affect
302 inflammatory and immune biomarkers, concluding that vegetarian diets associate with lower
303 CRP, fibrinogen, and total leukocyte concentrations (26). Similarly, Eichelmann showed that
304 plant-based diets cause reductions in obesity-related inflammatory biomarkers such as CRP, IL6
305 and sICAM (27). Indeed, our observed association between diet and monocyte level might be
306 related to the degree of obesity in the population; however, we found only a slight attenuation
307 of the association for women when adjusting for adiposity. In previous work, we have shown
308 that both diet and risk factors may affect PBMC leukocyte distribution (28,29). We found that
309 plasma omega 6 fatty acid level, as a marker of dietary intake of omega 6 fatty acids, associated
310 with predicted leukocyte distribution (28). Vegetarian diets tend to have high content of
311 vegetable oils, which may have affected our present results also. Similarly, we recently showed
312 that children with familial hypercholesterolemia displayed an altered leukocyte distribution
313 (29).

314 Most studies that examine the association between diet and immune cells use a modest
315 number of established biomarkers, such as standard differential count or protein biomarkers. In

316 the current analysis, however, we used approximately 14 000 mRNA transcripts from PBMCs,
317 potentially making it a more sensitive test of associations with immune cell type distribution
318 specifically, and inflammation in general (13). Additionally, our finding is important since it adds
319 to the evidence that cell type distribution in cell mixtures can influence the association between
320 diet and gene expression. This must be taken into account when interpreting PBMC gene
321 expression results.

322 ***DPs associated with few gene expression clusters***

323 Few WGCNA-based gene expression clusters were associated with DPs, after correcting for
324 variation in monocytes and lymphocytes number (Figure 3). This indicates that most of the co-
325 variation between diet and gene expression in PBMCs relates to leukocyte cell type distribution.
326 Nevertheless, in women, the *Vegetarian* DP associated most strongly with a cluster of genes
327 involved in regulation of transcription, and in men, the *Western* DP associated most strongly
328 with a cluster of genes related to metabolic and inflammatory processes, including
329 sterol/cholesterol transport.

330 In previous reports, dietary intake of a healthy Nordic diet or omega-3 associated with
331 expression of genes related to mitochondrial function, cell cycle, endoplasmic reticulum stress,
332 apoptosis, and inflammatory processes (30–33). *Regulation of transcription* is another such
333 unspecific term. Although highly unspecific, regulation of transcription may be a process related
334 to age-related global or pathway-specific DNA methylation and gene expression (34–36).
335 *Sterol/cholesterol transport*, on the other hand, is a highly specific biological process that is
336 dramatically affected by diet and that affects disease risk (36,37). Plasma LDL-C is mainly

337 determined by cellular sterol status and the functionality and activity of the LDL receptor; this in
338 turn is a key determinant of disease risk (37). However, although cholesterol metabolism in liver
339 and monocytes are tightly coupled and similarly regulated, our observed association likely
340 results from molecular events occurring within the pool of PMBCs as they deal with cholesterol-
341 related metabolic challenges. Nevertheless, PBMC expression of genes related to
342 sterol/cholesterol transport could prove a robust marker of dietary variation (10).

343 Interestingly, the second and third most significant clusters that associated with the *Western DP*
344 in men contained genes related to other metabolic processes, such as UDP-GlcNAc and acyl
345 carnitine metabolism. While UDP-GlcNAc is involved in cellular glucose sensing, acyl carnitines
346 are involved in fatty acid transport into the mitochondria (38,39). Indeed, in previous work, we
347 found that plasma levels of acyl carnitines of specific chain lengths may be directly altered by
348 changes in fatty acid quality of the diet (6). Taken together, these may be processes particularly
349 sensitive to variation in dietary intake.

350 ***We identified top driver genes and hub proteins***

351 Finally, the WGCNA cluster analysis detected top driver genes that have been shown to
352 physically interact in protein-protein interaction networks (Figure 4) (22). This is an important
353 finding, as it provides further biological meaning to the statistical associations, and strengthens
354 our belief that the top driver genes may be more than just spurious associations (40). The
355 network analysis highlighted a few hub proteins that may act as central communicators within
356 each cluster, such as UBC and PPARGC1B. The UBC protein is a key cell signaling molecule,
357 especially related to ubiquitination, cytokine signaling, toll-like receptors, and nuclear factor

358 kappa B (NFkB); in mouse models, knock-down of the ubiquitin system shows protection from
359 diet-induced obesity (41). Furthermore, PPARGC1B is a transcriptional co-regulator involved in a
360 number of biological processes, including thermogenesis, bone turnover and regulation of
361 energy expenditure by fat and glucose oxidation. For example, Yin and co-workers recently
362 showed that PPARGC1B affects PPAR alpha to protect against cardiomyopathy (42).

363 ***Strengths and limitations***

364 To the best of our knowledge, nobody has used CIBERSORT and WGCNA to study molecular
365 associations with DPs. We believe these dimension reduction algorithms may be well suited to
366 examine diet-related effects on PBMC cell type distribution using sensitive gene expression
367 data. Although we have taken steps to minimize the probability of chance findings, we cannot
368 rule this out completely. Our study sample is also relatively small, compared to for example Lin
369 and co-workers (40). In addition to higher risk of false positive findings, low sample size also
370 increases the risk of false *negative* findings, for example for the associations between DPs and
371 WGCNA gene clusters. Furthermore, our study sample represents a highly selected part of the
372 Norwegian population, limiting the generalizability of our results. In line with this limitation, our
373 results should not be overinterpreted. Pathways related to metabolic regulation could
374 potentially be biomarkers of dietary intake, and also potentially predict future risk (40,43).
375 However, more prediction research is needed before this can be realized.

376 ***Conclusions***

377 In conclusion, we detected novel associations between DPs and gene expression features in
378 PBMCs. Our results suggest that DPs may affect monocytes proportions and regulation of

379 biological processes, such as regulation of transcription and metabolic and inflammatory
380 processes. Although the present findings are exploratory, our analysis pipeline serves as a useful
381 framework for future studies on the association between diet and gene expression. More
382 research is needed before our results can be translated into clinically meaningful biomarkers.

383 **Acknowledgements**

384 ***Funding***

385 The work in this study was funded by Oslo University Hospital and the University of Oslo.

386 ***Conflicts of interest***

387 Dr. Christensen has received research grants and/or personal fees from Mills DA, unrelated to
388 the content of this manuscript. Dr. Ulven has received research grants and/or personal fees
389 from Mills DA, Tine DA, and Olympic Seafood, none of which are related to the content of this
390 manuscript. Dr. Holven has received research grants and/or personal fees from Tine DA, Mills
391 DA, Olympic Seafood, Amgen, Sanofi, and Pronova, none of which are related to the content of
392 this manuscript. The other authors declare no conflicts of interest.

393 ***Authorship***

394 Conception and design: LFA, KBH, SMU. Data analysis: JJC, MT, KW. Data interpretation: JJC, MT,
395 KBH, KW, LFA. Wrote paper: JJC, SMU, KW. In addition, the microarray hybridization and gene
396 expression data pre-processing were performed at the Genomics Core Facility at NTNU,
397 Trondheim, Norway.

398

References

1. Global Burden of Disease Study 2017 (GBD 2017) Results. Seattle, United States: Global Burden of Disease Collaborative Network; 2018.
2. Khera AV, Kathiresan S. Genetics of coronary artery disease: Discovery, biology and clinical translation. *Nature Reviews Genetics*. 2017;18:331–44.
3. Müller H, Kirkhus B, Pedersen JI. Serum cholesterol predictive equations with special emphasis on trans and saturated fatty acids. An analysis from designed controlled studies. *Lipids*. 2001;36:783–91.
4. Chawla A, Repa JJ, Evans RM, Mangelsdorf DJ. Nuclear Receptors and Lipid Physiology: Opening the X-Files. *Science*. 2001;294:1866–70.
5. Øyri LKL, Narverud I, Bogsrud MP, Hansson P, Leder L, Byfuglien MG, Veierød MB, Thoresen M, Ulven SM, Holven KB. Postprandial changes in gene expression of cholesterol influx and efflux mediators after intake of SFA compared with n-6 PUFA in subjects with and without familial hypercholesterolaemia: Secondary outcomes of a randomised controlled trial. *Journal of Nutritional Science*. 2019;8.
6. Ulven SM, Christensen JJ, Nygård O, Svardal A, Leder L, Ottestad I, Lysne V, Laupsa-Borge J, Ueland PM, Midttun Ø, et al. Using metabolic profiling and gene expression analyses to explore molecular effects of replacing saturated fat with polyunsaturated fat—a randomized controlled dietary intervention study. *The American Journal of Clinical Nutrition*. 2019;109:1239–50.

7. Jannasch F, Riordan F, Andersen LF, Schulze MB. Exploratory dietary patterns: A systematic review of methods applied in pan-European studies and of validation studies. *British Journal of Nutrition*. 2018;120:601–11.
8. Libby P, Buring JE, Badimon L, Hansson GK, Deanfield J, Bittencourt MS, Tokgözoğlu L, Lewis EF. Atherosclerosis. *Nature Reviews Disease Primers*. 2019;5:1–18.
9. Martinez FO, Gordon S, Locati M, Mantovani A. Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: New molecules and patterns of gene expression. *Journal of Immunology (Baltimore, Md: 1950)*. 2006;177:7303–11.
10. Mello VDF de, Kolehmanien M, Schwab U, Pulkkinen L, Uusitupa M. Gene expression of peripheral blood mononuclear cells as a tool in dietary intervention studies: What do we know so far? *Molecular Nutrition & Food Research*. 2012;56:1160–72.
11. Olsen KS, Skeie G, Lund E. Whole-Blood Gene Expression Profiles in Large-Scale Epidemiological Studies: What Do They Tell? *Current Nutrition Reports*. 2015;4:377–86.
12. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network analysis. *BMC bioinformatics*. 2008;9:559.
13. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, Hoang CD, Diehn M, Alizadeh AA. Robust enumeration of cell subsets from tissue expression profiles. *Nature Methods*. 2015;12:453–7.

14. Nguyen LH, Holmes S. Ten quick tips for effective dimensionality reduction. *PLOS Computational Biology*. 2019;15:e1006907.
15. Ulven SM, Leder L, Elind E, Ottestad I, Christensen JJ, Telle-Hansen VH, Skjetne AJ, Raael E, Sheikh NA, Holck M, et al. Exchanging a Few Commercial, Regularly Consumed Food Items with Improved Fat Quality Reduces Total Cholesterol and LDL-Cholesterol: A Double-Blind, Randomised Controlled Trial. *The British journal of nutrition*. 2016;116:1383–93.
16. Carlsen MH, Lillegaard IT, Karlsen A, Blomhoff R, Drevon CA, Andersen LF. Evaluation of energy and dietary intake estimates from a food frequency questionnaire using independent energy expenditure measurement and weighed food records. *Nutrition Journal*. 2010;9:37.
17. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2019.
18. Zhang B, Horvath S. A general framework for weighted gene co-expression network analysis. *Statistical Applications in Genetics and Molecular Biology*. 2005;4:Article17.
19. Ghazalpour A, Doss S, Zhang B, Wang S, Plaisier C, Castellanos R, Brozell A, Schadt EE, Drake TA, Lusis AJ, et al. Integrating Genetic and Network Analysis to Characterize Genes Related to Mouse Weight. *PLOS Genetics*. 2006;2:e130.
20. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: Tool for the unification of biology. *Nature genetics*. 2000;25:25–9.

21. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic Acids Research*.

2019;47:D330–8.

22. Cowley MJ, Pinese M, Kassahn KS, Waddell N, Pearson JV, Grimmond SM, Biankin AV,

Hautaniemi S, Wu J. PINA v2.0: Mining interactome modules. *Nucleic Acids Research*.

2012;40:D862–865.

23. Markussen MS, Veierød MB, Sakhi AK, Ellingjord-Dale M, Blomhoff R, Ursin G, Andersen LF.

Evaluation of dietary patterns among Norwegian postmenopausal women using plasma

carotenoids as biomarkers. *British Journal of Nutrition*. 2015;113:672–82.

24. Chocano-Bedoya PO, O'Reilly EJ, Lucas M, Mirzaei F, Okereke OI, Fung TT, Hu FB, Ascherio A.

Prospective study on long-term dietary patterns and incident depression in middle-aged and

older women. *The American Journal of Clinical Nutrition*. 2013;98:813–20.

25. Weismayer C, Anderson JG, Wolk A. Changes in the stability of dietary patterns in a study of

middle-aged Swedish women. *The Journal of Nutrition*. 2006;136:1582–7.

26. Craddock JC, Neale EP, Peoples GE, Probst YC. Vegetarian-Based Dietary Patterns and their

Relation with Inflammatory and Immune Biomarkers: A Systematic Review and Meta-Analysis.

Advances in Nutrition (Bethesda, Md). 2019;10:433–51.

27. Eichelmann F, Schwingshackl L, Fedirko V, Aleksandrova K. Effect of plant-based diets on

obesity-related inflammatory profiles: A systematic review and meta-analysis of intervention

trials. *Obesity Reviews: An Official Journal of the International Association for the Study of*

Obesity. 2016;17:1067–79.

28. Christensen JJ, Bakke SS, Ulven SM, Retterstøl K, Henriksen T, Bollerslev J, Espevik T, Aukrust P, Halvorsen B, Holven KB. Serum Omega-6 Fatty Acids and Immunology-Related Gene Expression in Peripheral Blood Mononuclear Cells: A Cross-Sectional Analysis in Healthy Children. *Molecular Nutrition & Food Research*. 2019;1800990.
29. Christensen JJ, Osnes LT, Halvorsen B, Retterstøl K, Bogsrud MP, Wium C, Svilaas A, Narverud I, Ulven SM, Aukrust P, et al. Altered leukocyte distribution under hypercholesterolemia: A cross-sectional study in children with familial hypercholesterolemia. *Atherosclerosis*. 2017;256:67–74.
30. Myhrstad MCW, Ulven SM, Günther C-C, Ottestad I, Holden M, Ryeng E, Borge GI, Kohler A, Brønner KW, Thoresen M, et al. Fish oil supplementation induces expression of genes related to cell cycle, endoplasmic reticulum stress and apoptosis in peripheral blood mononuclear cells: A transcriptomic approach. *Journal of Internal Medicine*. 2014;276:498–511.
31. Myhrstad MCW, Mello VD de, Dahlman I, Kolehmainen M, Paananen J, Rundblad A, Carlberg C, Olstad OK, Pihlajamäki J, Holven KB, et al. Healthy Nordic Diet Modulates the Expression of Genes Related to Mitochondrial Function and Immune Response in Peripheral Blood Mononuclear Cells from Subjects with Metabolic Syndrome A SYSDIET Sub-Study. *Molecular Nutrition & Food Research*. 2019;63:1801405.
32. Kolehmainen M, Ulven SM, Paananen J, de Mello V, Schwab U, Carlberg C, Myhrstad M, Pihlajamäki J, Dungner E, Sjölin E, et al. Healthy Nordic diet downregulates the expression of genes involved in inflammation in subcutaneous adipose tissue in individuals with features of the metabolic syndrome. *The American Journal of Clinical Nutrition*. 2015;101:228–39.

33. Ulven SM, Holven KB, Rundblad A, Myhrstad MCW, Leder L, Dahlman I, Mello VD de, Schwab U, Carlberg C, Pihlajamäki J, et al. An Isocaloric Nordic Diet Modulates RELA and TNFRSF1A Gene Expression in Peripheral Blood Mononuclear Cells in Individuals with Metabolic SyndromeA SYSDIET Sub-Study. *Nutrients*. 2019;11:2932.
34. Field AE, Robertson NA, Wang T, Havas A, Ideker T, Adams PD. DNA Methylation Clocks in Aging: Categories, Causes, and Consequences. *Molecular Cell*. 2018;71:882–95.
35. Horvath S. DNA methylation age of human tissues and cell types. *Genome Biology*. 2013;14:3156.
36. Hahn O, Grönke S, Stubbs TM, Ficz G, Hendrich O, Krueger F, Andrews S, Zhang Q, Wakelam MJ, Beyer A, et al. Dietary restriction protects from age-associated DNA methylation and induces epigenetic reprogramming of lipid metabolism. *Genome Biology*. 2017;18:56.
37. Goldstein JL, Brown MS. A Century of Cholesterol and Coronaries: From Plaques to Genes to Statins. *Cell*. 2015;161:161–72.
38. Adeva-Andany MM, Carneiro-Freire N, Seco-Filgueira M, Fernández-Fernández C, Mouriño-Bayolo D. Mitochondrial β -oxidation of saturated fatty acids in humans. *Mitochondrion*. 2019;46:73–90.
39. Anthonisen EH, Berven L, Holm S, Nygård M, Nebb HI, Grønning-Wang LM. Nuclear receptor liver X receptor is O-GlcNAc-modified in response to glucose. *The Journal of Biological Chemistry*. 2010;285:1607–15.

40. Lin H, Rogers GT, Lunetta KL, Levy D, Miao X, Troy LM, Jacques PF, Murabito JM. Healthy diet is associated with gene expression in blood: The Framingham Heart Study. *The American Journal of Clinical Nutrition*. 2019;110:742–9.

41. Setsuie R, Suzuki M, Kabuta T, Fujita H, Miura S, Ichihara N, Yamada D, Wang Y-L, Ezaki O, Suzuki Y, et al. Ubiquitin C-terminal hydrolase-L3-knockout mice are resistant to diet-induced obesity and show increased activation of AMP-activated protein kinase in skeletal muscle. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology*. 2009;23:4148–57.

42. Yin Z, Zhao Y, He M, Li H, Fan J, Nie X, Yan M, Chen C, Wang DW. MiR-30c/PGC-1 β protects against diabetic cardiomyopathy via PPAR α . *Cardiovascular Diabetology*. 2019;18:7.

43. Wang X-b, Cui N-h, Liu X, Ming L. Identification of a blood-based 12-gene signature that predicts the severity of coronary artery stenosis: An integrative approach based on gene network construction, Support Vector Machine algorithm, and multi-cohort validation. *Atherosclerosis*. 2019;291:34–43.

Tables

Table 1. *Study sample: clinical characteristics.*

Variables	Women	Men
Age, years	58 (13)	50 (15)
BMI, kg/m ²	23.7 (4.2)	26.4 (4.1)
Fat mass, kg	34.2 (7.9)	23.4 (9.2)
Fat free mass, kg	65.8 (7.9)	76.7 (9.2)
Creatinine, mmol/L	69 (12)	85.5 (13)
ASAT, U/L	20 (9)	22 (11)
ALAT, U/L	22 (8)	28 (13)
TG, mmol/L	0.9 (0.6)	1.2 (0.9)
TC, mmol/L	6.6 (1.2)	6.2 (1.4)
HDL-C, mmol/L	1.8 (0.6)	1.4 (0.4)
LDL-C, mmol/L	3.9 (1)	3.9 (1.1)

Values are median (interquartile range).

Table 2. *Groupings of food items used as input in the dietary pattern analysis.*

Food group	Food items
Butter	Butter, butter-based margarines, Melange margarine
Margarine	Standard and low-fat margarine, standard and low-fat Vita, standard and low-fat Soft Flora
Cheese, high-fat	Regular Norwegian brown cheese, regular hard cheese, regular cheese spread, regular cream cheese
Cheese, low-fat	Low-fat Norwegian brown cheese, low-fat hard cheese, low-fat cheese spread, low-fat cream cheese
Dairy, high-fat	High-fat milk, flavoured milk, sour cream, ice cream, whipped cream, high-fat yogurt
Dairy, low-fat	Skimmed and semi-skimmed milk, cultured/probiotic low-fat milk, low-fat yogurt drink, low-fat yogurt
Dairy, semi low-fat	Cultura, light milk, natural yoghurt, fruit yoghurt, Go Morgen yoghurt, Biola, chocolate milk
Coffee	Boiled coffee, espresso, filter coffee, instant coffee
Tea	Black tea, green tea, herb tea
Sweet beverages	Artificially sweetened soft drinks, artificially sweetened ice tea, fruit juices with added sugar, squash with sugar, sugar-sweetened soft drinks, iced tea with sugar, orange juice, apple juice, Mana juice
Beer	Non-alcoholic beer, light beer, regular beer, alcopops
Wine	Wine, red wine, white wine
Chocolate and sweets	Chocolate, dark chocolate, extra dark chocolate, sweets/jelly sweets, sweet

	pastilles/candy
	Cookies, wheat bun, pastry, pastry bun, waffles, chocolate cake, cream cake,
Desserts and snacks	muffins, pudding, potatoe chips
Sweet spreads	Regular and low-sugar jam, honey, chocolate/nut spread, other sweet spreads
	Meat balls, minced meat sauce, taco, kebab, lasagne, grilled/wiener sausage, minced meat sausage,
	bacon, pork chops, regular and low-fat liver paste, regular and low-fat saveloy,
Processed meat	salami
Red meat	Roast of lamb/beef/pork, beef
Poultry	Chicken
Eggs	Eggs in dishes, whole eggs
	Salmon, herring, mackerel, mackerel in tomatoe (spread), achovy, caviar, cod,
Fish and shellfish	fish balls, fish cakes, tuna, shellfish
Legumes	Legumes
	Carrot, cabbage, swede, cauliflower, broccoli, Brussels sprout, onion, spinach, sweet pepper, avocado,
	tomato, maize, frozen vegetables, mixed salad (with lettuce, cucumber, tomato and sweet pepper),
Vegetables	vegetables as spread
	Apple, pear, banana, orange, clementine, peach/nectarine, kiwi, grapes, melon, pomegranate, fruit as spreads, fresh fruit salad, prune, raisins, other dried fruits, blackberry, blueberry, raspberry, strawberry, cherry, cloudberry, rose
Fruits and berries	hips, redcurrant, blackcurrant, cowberry
Rice	Rice

Potatoes	Potatoes, potato powder
Pasta	Spaghetti, nudles
Whole grains	Semi- and whole-grain bread, crisp bread (whole meal), oatmeal porridge and cereal, unsweetened muesli/breakfast cereal
Nuts and seeds	Almonds, hazelnuts, walnuts, peanuts, peanut butter, pine nuts, linseeds, sunflower seeds, nut mix, cashew nuts
Oil and oil products	Vegetable oils, oil-based dressings, dressing mix, mayonnaise-based salads, pesto
Mixed meals	Pizza, various mixed meals, including wraps and spring rolls
Potatoes, fat-rich	Pommes frites
Refined grains	White bread, crisp bread (wheat flour), hot dog bun, sweet muesli/breakfast cereal, pancake, rice porridge, waffle, sweet bun
Supplements, fish oil	Fish oil supplements, Tran

Figure legends

Figure 1

Figure 1. *DPs for women and men.* The figure shows factor loadings for all foods with a loading > 0.3. See Supplementary Table 4 for full table of all foods (and their uniqueness). Abbreviations: DP, dietary pattern.

Figure 2

Figure 2. *DPs associate with CIBERSORT-predicted cell types.* The figure displays heatmaps of linear regression β coefficients between DP scores (as the exposure variable, shown in rows), and CIBERSORT-predicted cell types (as the outcome variable, in columns), for both women and men. Asterix indicate significance level: ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$. See Supplementary Figure 1, Supplementary Figure 4 and Methods for a thorough explanation of the flow of analyses and adjustment levels. Abbreviations: DP, dietary pattern.

Figure 3

Figure 3. *DPs associate with gene expression clusters.* The figure displays heatmaps of linear regression β coefficients between DP scores (as the exposure variable, shown in rows), and gene expression cluster eigengenes (as the outcome variable, in columns), for both women and men. The clusters are sorted by size. Asterix indicate significance level: ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$. See Supplementary Figure 1, Supplementary Figure 4 and Methods for a thorough explanation of the flow of analyses and adjustment levels. Abbreviations: DP, dietary pattern.


Figure 4

Figure 4. *Identification of driver genes and hub proteins.* A) and B) display the association between *cluster membership* and *DP significance* for the strongest DP and gene expression cluster associations, for each gender. Cluster membership is defined as the absolute correlation between gene expression and cluster eigengene, and can be interpreted as the degree to which each gene belongs in that certain cluster, and contributes to its variation. DP significance is the absolute correlation between gene expression and DP score. A positive correlation between cluster membership and DP significance indicates that those genes that drive the variation in the cluster eigengene are the same that associate with the specific DP (*driver genes*). Finally, to rank driver genes, driver gene estimates were calculated by the sum of the cluster membership and DP significance. The darker the color, the higher the driver gene estimate; the top five genes driving this association are annotated. Note strong positive correlations for both comparisons, as is also evident from the linear regression line. C) and D) show networks of protein-protein interactions (PPI) for the same DP and gene expression cluster associations as above. Each network was created by the top 20 driver genes identified by the driver gene plot. The figures display *hub proteins* that are of particular interest to the gene regulatory network.


Abbreviations: DP, dietary pattern (see Supplementary Table 7 and Supplementary Table 8 for all abbreviations).

Women


DP1






DP2





DP3



Men




Coefficient estimate




A

Women




B

Men



C



D

