Association of the use of hearing aids with the conversion from mild cognitive impairment to dementia and progression of dementia: a longitudinal retrospective study

Magda Bucholc[a], Paula L. McClean[b], Sarah Bauermeister[c], Stephen Todd[d], Xuemei Ding[a,e], Qinyong Ye[f], Desheng Wang[g], Wei Huang[g], Liam P. Maguire[a]

[a] Cognitive Analytics Research Lab, School of Computing, Engineering & Intelligent Systems, Ulster University, BT48 7JL, United Kingdom

[b] Northern Ireland Centre for Stratified Medicine, Biomedical Sciences Research Institute, Ulster University, BT47 6SB, United Kingdom

[c] Department of Psychiatry, Warneford Hospital, University of Oxford, OX3 7JX, United Kingdom

[d] Altnagelvin Area Hospital, Western Health and Social Care Trust, BT47 6SB, United Kingdom

[e] Fujian Provincial Engineering Technology Research Centre for Public Service Big Data Mining and Application, College of Mathematics and Informatics, Fujian Normal University, Fuzhou, Fujian, 350108, China

[f] Department of Neurology, Fujian Medical University Union Hospital, Fujian, Fuzhou 350001, China.

[g] Department of Otolaryngology, Fujian Medical University Union Hospital, Fujian, Fuzhou 350001, China.

Corresponding author: Magda Bucholc; Cognitive Analytics Research Lab, School of Computing, Engineering & Intelligent Systems, Ulster University, BT48 7JL, United Kingdom. E-mail: m.bucholc@ulster.ac.uk

Abstract

INTRODUCTION: Hearing aid usage has been linked to improvements in cognition, communication, and socialization, but the extent to which it can affect the incidence and progression of dementia is unknown. Such research is vital given the high prevalence of dementia and hearing impairment in older adults, and the fact that both conditions often coexist. In this study, we examined for the first time the effect of the use of hearing aids on the conversion from mild cognitive impairment (MCI) to dementia and progression of dementia.

METHODS: We used a large referral-based cohort of 2114 hearing-impaired patients obtained from the National Alzheimer's Coordinating Center. Survival analyses using multivariable Cox proportional hazards regression model and weighted Cox regression model with censored data were performed to assess the effect of hearing aid use on the risk of conversion from MCI to dementia and risk of death in hearing-impaired participants. Disease progression was assessed with CDR® Dementia Staging Instrument Sum of Boxes (CDRSB) scores. Three types of sensitivity analyses were performed to validate the robustness of the results.

RESULTS: MCI participants that used hearing aids were at significantly lower risk of developing all-cause dementia compared to those not using hearing aids (hazard ratio [HR] 0.73, 95%CI, 0.61-0.89; false discovery rate [FDR] P=0.004). The mean annual rate of change (standard deviation) in CDRSB scores for hearing aid users with MCI was 1.3 (1.45) points and significantly lower than for individuals not wearing hearing aids with a 1.7 (1.95) point increase in CDRSB per year (P=0.02). No association between hearing aid use and risk of death was observed. Our findings were robust subject to sensitivity analyses.

DISCUSSION: Among hearing-impaired adults, hearing aid use was independently associated with reduced dementia risk. The causality between hearing aid use and incident dementia should be further tested.

Keywords: dementia; Alzheimer's disease; mild cognitive impairment; MCI; dementia onset; dementia incidence; hearing impairment; hearing loss; hearing aid; risk factor; disease progression; cognitive decline; survival analysis; National Alzheimer's Coordinating Center

Highlights

- High prevalence of dementia and hearing impairment in older adults

- Hearing aid (HA) use associated with a lower risk of incident dementia

- Slower cognitive decline in users than non-users of HA with mild cognitive impairment

- The relationship between hearing impairment and dementia should be further tested

1 Background

The escalating costs and devastating psychological and emotional impact of dementia on affected individuals, their families and caregivers makes the prevention, diagnosis, and treatment of dementia a national public health priority worldwide [1]. While the process of drug development to delay the onset of dementia has been slower than initially hoped, there is evidence that behavioural and lifestyle interventions might reduce dementia risk [2-4]. Numerous studies have investigated the effects of physical exercise, healthy diet, and management of medical conditions, such as diabetes and heart disease, on cognitive decline and risk of developing dementia [2,5]. However, there is a paucity of research on hearing impairment and dementia. Such research is vital, given the high prevalence of dementia and hearing impairment in older adults, and the fact that both conditions often coexist.

The prevalence of hearing loss is higher among older than younger individuals, with over 70% of adults aged 70 and older having hearing loss in at least 1 ear [6]. In the United States (U.S), mild hearing loss affects 23% of the population over the age of 12, with moderate hearing loss more prevalent in those over 80 years [6]. More importantly, approximately 23 million adults with hearing loss in the U.S. do not use hearing aids even though the negative impact of untreated hearing loss has been widely documented [7-9]. The low level of hearing aid adoption is associated with stigma and affordability of hearing aids [8, 15]. Adults with impaired hearing, who do not wear hearing aids demonstrate significantly higher rates of depression, anxiety, and other psychosocial disorders [9,10]. Hearing loss was also associated with increased risk of incident dementia [11-14].

Despite the observed relationship between hearing loss and cognition, surprisingly

few studies have investigated the association between the use of hearing aids,

cognitive decline, and risk for developing dementia [16-17]. One study found little

evidence that hearing aids promoted cognitive function but acknowledged that they

may be effective in reducing hearing handicap [18]. Other work showed that hearing

aid use was associated with better cognition while controlling for confounding by

age, sex, general health, and socioeconomic status [19]. A recent randomized pilot

study examined the changes in cognition due to treatment of hearing loss, with some

promising results for proximal outcomes (perceived hearing handicap, loneliness)

that may mediate a relationship between hearing and cognition [10]. Better

understanding of the relationship between use of hearing aids cognitive function, and

risk of dementia is of utmost importance since it has the potential to significantly

impact public health, as hearing aids represent a minimally invasive, cost effective

treatment to mitigate the impact of hearing loss on dementia. In fact, it is postulated

that up to 9% of dementia cases could be prevented with proper hearing loss

management [14]. Slower conversion from MCI to dementia and progression of

dementia following hearing aid use could potentially lead to the reduced incidence of

dementia and extended preservation of functional independence in people with

dementia.

In this study, we used data from a large referral-based cohort to examine for the first

time the effect of the use of hearing aids on the conversion from Mild Cognitive

Impairment (MCI) to dementia and risk of death. We tested if the use of hearing aids

is independently associated with a decreased risk of incident all-cause dementia

diagnosis for MCI patients and reduced risk of death in individuals with dementia.

We also examined if the rate of cognitive decline is slower for hearing aid users

when compared to those not using hearing aids in participants with both MCI and
dementia at baseline.

2   Methods

2.1   Participants

We conducted a retrospective analysis of the demographic and clinical data obtained
from the National Alzheimer's Coordinating Center (NACC) [20]. The NACC
database consists of data from Alzheimer's Disease Research Centers (ADRCs)
supported by the National Institute on Aging (NIA) (grant U01AG016976). Details
about the NACC consortium, data collection process, and design and
implementation of the NACC database have been reported previously [20,21]. The
data set used in our longitudinal investigation was the NACC Uniform Data Set
(UDS) [21].

The analytic sample for this study included 2114 participants (age >50) with hearing
impairment who had UDS data in the NACC database available between 2005 and
2018 (Fig. 1). All subjects were classified into two groups according to the disease
stage. Group 1 included 939 individuals that were diagnosed with MCI at baseline,
namely, 497 MCI converters (MCI-c) and 442 MCI non-converters (MCI-nc). Group 2
consisted of 1175 participants that were diagnosed with dementia at baseline: 349 of
those died during the follow-up. The 829 dementia participants who did not die
during the study follow-up were censored at their last clinical evaluation. Note that
only patients that clearly progressed from one stage to another were included in the
study. In addition, only active participants who continued to return for annual follow-
up visits were taken into account and hence, any subject that missed a scheduled
appointment (7%) was discarded from the analysis.

2.2 Clinical assessment and diagnosis

The incidence of MCI and all-cause dementia was determined based on the clinical

diagnosis made by a single clinician or a consensus panel. The clinical diagnosis

took into account patient's medical history, medication use, neuropsychological test

performance, and other modifying factors, such as educational and cultural

background, and behavioural assessments. In NACC-UDS Version 1 and 2, the

procedure of clinical diagnosis of all-cause dementia depended on the diagnostic

protocol of the ADRC, but each Center generally adhered to standardized clinical

criteria as outlined by the DSM-IV or NINDS-ADRDA guidelines [22,23]. In NACC-

UDS Version 3, the coding guidebook criteria for all-cause dementia were modified

from the McKhann all-cause dementia criteria (2011) [24]. Diagnoses of MCI were

made using the modified Petersen criteria [25].

In addition, we used the CDR® Dementia Staging Instrument Sum of Box (CDRSB)

scores to assess a decline due to cognitive changes in six functional domains,

namely, memory, orientation, judgment and problem solving, community affairs,

home and hobbies, and personal care [26]. CDRSB has a total score range from 0 to

18 points, with higher scores indicating greater cognitive impairment. CDRSB has

been commonly used as a reliable tool for assessing dementia severity [27,28].

2.3 Hearing assessment

The information on presence of hearing loss and use of hearing aids were extracted

from the NACC UDS Physical Evaluation form [29]. Information on hearing loss was

collected via self-report using a single hearing screening question: "Without a

hearing aid(s), is the subject's hearing functionally normal?", which provided possible

responses of 'yes' or 'no' [29]. Participants that responded 'yes' were defined as

individuals with hearing loss and those that responded 'no' were excluded from the

analysis. The determination of severity of hearing loss was not a part of a

standardized clinical evaluation given to NACC participants. Although participants

were allowed to wear hearing aids during the cognitive assessments, it was not

reported to NACC whether a participant was wearing a hearing aid during cognitive

testing. Missing codes were however entered when ADRCs had a reason to believe

that the test was invalid, including if they were aware that the participant was unable

to hear properly. All participants with missing codes were excluded from the analysis.

Information on hearing aid use was collected via self-report using a single question:

"Does the subject usually wear a hearing aid(s)?", which provided possible

responses of 'yes' or 'no' [29]. Participants that responded 'yes' were classified as

hearing aid users and those that responded 'no' were classified as non-users of

hearing aids. The consistent use (or non-use) of hearing aids was established if the

participant answered "yes" or "no" to the above question at every consecutive visit.

There was no additional information in the NACC-UDS on the number of

hours of daily use of hearing aids, the type of hearing aids used, or the history of

hearing aid use before the enrolment in the NACC-UDS.

Accordingly, data from individuals identified with impaired hearing at every annual

clinical evaluation, that consistently reported non-use or use of hearing aids as well

as functionally normal hearing when wearing hearing aids served as the sample for

our study. The subject's hearing was characterised as functionally normal with a

hearing aid if there was no evidence of reduced ability to do everyday activities such

as listening to the radio/television or talking with family/friends [29]. This information

was based on self-report. On the other hand, the group of participants excluded from

the analysis consisted of subjects without hearing impairment or with inconsistent

hearing impairment labelling as indicated by the records from each follow-up visit.

Participants with inconsistent records relating to the use or non-use of hearing aids

or without functionally normal hearing when wearing a hearing aid were also

excluded. The baseline characteristics of participants included and excluded from

the study are shown in Table A.1. Among 2114 patients with hearing impairment

included in the study, 636 subjects in Group 1 and 710 in Group 2 were classified as

using hearing aids.

2.4 Statistical analysis

Baseline summary statistics are presented as proportions for categorical data and

means with standard deviations (SD) for continuous variables. Unadjusted analyses

for comparison of demographic and clinical features between individuals with hearing

impairment that used and did not use hearing aids were performed with the Fisher's

exact test and unpaired t test [30]. The average annual rate of change in CDRSB

score in individuals using and not using hearing aids was compared by applying

Mann Whitney U method [30]. The assumption of normality of CDRSB data was

tested using Shapiro-Wilk test [30].

A Cox model with censored data was used to study time to incident dementia

diagnosis for MCI patients (Group 1) and death for individuals diagnosed with

dementia (Group 2) [31]. Censoring was accounted for in the analysis to allow for

valid inferences. Ignoring censoring and equating the observed follow-up time of

censored subjects with the unobserved total survival time would likely lead to an

overestimate of the overall survival probability. The proportional hazards assumption

for the Cox regression model fit was tested with the Schoenfeld residuals method

and satisfied for Group 1 (P = 0.7). The presence of non-proportional hazards was

observed in Group 2 (P = 0.001), with the proportional hazard assumption violated

for age (P < 0.001) and CDRSB (P = 0.007) variables. Accordingly, we used the Cox

proportional hazards regression model to model time to incident dementia for MCI

patients (Group 1) and implemented weighted Cox regression accounting for time-

varying effects to determine the survival rate of individuals diagnosed with dementia

(Group 2) [32]. Weighted Cox regression allowed for providing unbiased estimates of

hazard ratios irrespective of proportionality of hazards [32]. Hazard ratios (HR) with

95% confidence intervals (95% CI) were calculated for each group by comparing the

hazard rates for individuals with hearing impairment who used and did not use

hearing aids. All comparisons were adjusted by age, gender, years of education

(measured as the number of years of education completed) and CDRSB score to

remove their possible confounding effect. The linearity assumption of the relationship

between continuous confounding variables (i.e. age, education, CDRSB) and the

log-hazard of the time-to-event outcome was tested using the Box-Tidwell approach

and satisfied in both groups (p > 0.05) [33].

For each MCI individual in Group 1 and dementia participant in Group 2, time 'zero'

was defined as the date of the baseline evaluation. The diagnosis of MCI in Group 1

and dementia in Group 2 referred to the initial event while the endpoint event was

considered the conversion to all-cause dementia in Group 1 and the occurrence of

death in Group 2 (0 – censored, 1 – uncensored). Survival time was determined by

the year. MCI-nc subjects and dementia participants who did not die during the study

follow-up were censored at the last clinical assessment.

To avoid the inflation of false-positive findings, the Benjamini-Hochberg false

discovery rate (FDR) procedure was used to adjust for multiple hypothesis-testing

[34]. False discovery-adjusted P values (FDR P) less than 0.05 were considered as

statistically significant.

2.5 Sensitivity analysis

To validate the robustness of the main findings, we performed three different

sensitivity analyses: a) propensity score matching to control for measured

confounding, b) the analysis for unmeasured confounding to assess the sensitivity of

our main conclusions with respect to confounders not included in our study, and c)

the inverse probability of treatment weighting method to reduce selection bias within

our study population [35,36,37].

Propensity scores were generated for hearing aid status using multivariable logistic

regression model and adjusting for baseline covariates, including age, gender,

education, and CDRSB. The standardized mean difference between two patient

groups i.e., patients with and without hearing aids, was then calculated for each

covariate and compared before and after the matching process to determine

covariate balance between the two groups [34]. A standardized difference of less

than 0.1 was considered negligible in the prevalence of a covariate [34].

Sensitivity analysis for unmeasured confounding was conducted to measure the

potential influence an unmeasured covariate might have on the HR estimates of the

association between hearing aid use and a) incident dementia in Group 1 and b)

death in Group 2 [36]. We considered prevalence rates for the confounder of 5%,

10%, and 20% in the group of hearing aid users and three different values of HR

(0.5,2.0,4.0) for the association between the confounder and exposure. We then

varied the prevalence of the unmeasured confounder in the group of subjects without

hearing aids, from 10% to 30% to determine the extent to which its distribution under

these conditions would need to be imbalanced to influence the statistical significance

of our findings, i.e., when the upper limit of the 95% CI of HR crosses 1.0.

The inverse probability of treatment weighting method was used to attenuate

potential selection bias in the sampling and recruitment of NACC participants.

Weights were derived from propensity score modelling of the probability of hearing

aid use as a function of measured covariates using the Generalized Boosted Model

[37]. A multivariable Cox proportional hazards regression model and weighted Cox

regression model were then fitted using derived weights to examine the risk of

incident dementia and death for hearing aid users and non-users in Group 1 and 2

respectively.

3   Results

3.1 Participant Characteristics

Baseline demographic and clinical characteristics of participants by the hearing aid

status are presented in Table 1. Statistically significant differences in the use of

hearing aids were found between men and women both in Group 1 (P = 0.01) and

Group 2 (P = 0.02), with higher rates of hearing aid use in males. Hearing aid users

with baseline MCI were significantly older than MCI individuals not using hearing

aids ($P < 0.001$). Age was comparable for users/non-users of hearing aids in

Group 2 (P = 0.32). Hearing aid users with dementia had more years of education

completed ($P < 0.001$). The CDRSB score was significantly lower in both individuals

with baseline MCI and dementia who used hearing aids (P = 0.01 and $P < 0.001$

respectively).

3.2 Hearing aid status and risk of incident all-cause dementia in MCI participants

During the study follow-up, 497 MCI subjects in Group 1 developed dementia. The

median time to incident dementia was 2 years for non-hearing aid users and 4 years

for hearing aid users. The 5-year overall survival rate, which is the percentage of

participants that did not convert to dementia 5 year after the baseline MCI diagnosis,

was 19% for non-hearing aid users and 33% for individuals using hearing aids.

In the multivariable Cox proportional hazards regression model, the major risk factor

for MCI-to-dementia conversion was the CDRSB score (HR 1.39, 95%CI, 1.30-1.48,

FDR *P* < 0.001; Table 2), while a significantly reduced risk of dementia was

associated with the use of hearing aids (HR 0.73, 95%CI, 0.61-0.89, FDR *P* =

0.004).

The observed mean annual (SD) rate of change in CDRSB for non-hearing aid users

with MCI was 1.7 (1.95) points per year and significantly higher than the average

rate of change for hearing aid users of 1.3 (1.45) CDRSB points per year (*P* = 0.02).

3.3 Hearing aid status and mortality risk in participants with dementia

Group 2 consisted of 1175 individuals diagnosed with dementia at baseline: 349 of

those died during the follow-up. The median survival time for dementia participants

who did not use hearing aids was 6 years and the 5-year overall survival rate was

58%. For hearing aid users, the median survival time was 7 years and the 5-year

overall survival rate was 67%.

In the weighted Cox regression model accounting for time-varying effects, the

relationship between the use of hearing aids and mortality risk was not statistically

significant (HR, 0.98; 95% CI, 0.78-1.24; FDR *P* = 0.89; Table 3). Higher CDRSB

scores were associated with the increased risk of death (HR, 1.08; 95% CI, 1.05-

1.11; FDR $P < 0.001$).

The average (SD) annual rate of change in CDRSB score of 0.96 (1.02) for hearing

aid users with dementia was not significantly different from the 0.94 (1.19) point

increase in CDRSB score per year for individuals with dementia not using hearing

aids ($P = 0.75$).

3.4   Sensitivity analysis

The distributions of potential confounders were similar between the hearing aid and

non-hearing aid user groups after propensity score matching (standardized

difference < 0.1). Again, we found that MCI individuals using hearing aids were at

lower risk of developing dementia when compared to non-users of hearing aids (HR

0.67, 95%CI, 0.50-0.80, FDR $P = 0.01$) (Table 4). No link between hearing aid use

and risk of death was found for individuals with dementia (Table 4).

Sensitivity analysis for unmeasured confounding performed for each of two studied

groups produced virtually unchanged findings (Table 5). Within Group 1, we

observed a lower risk of incident dementia for hearing aid users with the estimated

HR for incident dementia in the group of participants with hearing aids below 1 for all

considered values of the strength of the confounder-outcome association (HR 0.5,

1.5, 2.0, 4.0), and the prevalence of potential confounder in the group of hearing aid

users (5%, 10%, 20%) and non-users (10%, 20%, 30%). No association between

hearing aid use and risk of death was found in Group 2.

The results of the inverse weighted propensity showed the increased risk of

dementia for MCI subjects in Group 1 (HR 0.72, 95%CI, 0.60-0.88, FDR $P = 0.003$)

and no statistically significant association between hearing aid use and risk of death

for dementia participants in Group 2.

A lack of improvement in hearing, when a hearing aid is used, may be an indicator of

central auditory processing issues rather than a faulty device. As such, we

performed additional analysis on a group of hearing aid users that included subjects

who still experienced auditory difficulties when wearing a hearing aid (Table A.3). In

total, 129 participants in Group 1 and 174 participants in Group 2 were identified as

those without functionally normal hearing when wearing a hearing aid. A majority of

them, 63% in Group 1 and 68% in Group 2, were males. We observed a lower risk of

developing dementia in subjects using hearing aids in Group 1 (HR, 0.74; 95% CI,

0.61-0.89; FDR $P = 0.003$) and no statistically significant relationship between the

use of hearing aids and mortality risk in Group 2 (HR, 0.99; 95% CI, 0.80-1.23; FDR

$P = 0.92$).

4   Discussion

Despite the prevalence of auditory impairment in dementia, hearing loss is often not

diagnosed and not treated even though hearing loss has been shown to be an

independent risk factor for poorer cognitive function [38], depression and loneliness

[39], and diminished functional status [39]. Several longitudinal studies indicated that

individuals with hearing impairment experience substantially higher risk of incident

all-cause dementia [12,40-42]. For instance, the study of Lin et al [12] observed a

strong relationship between hearing loss and risk of developing dementia. While the

authors observed no association between use of hearing aids and reduced risk of

dementia, they found a strong link between degree of hearing loss and dementia

incidence [12]. In Ray et al [42], the association between cognitive impairment and

degree of hearing loss was also observed but only in individuals who did not use
hearing aids.

Hypothesized mechanisms explaining the association between hearing impairment
and cognitive function included the reallocation of cognitive resources to auditory
perceptual processing [43,44,45], cognitive deterioration due to long-term deprivation
of auditory input [43,46], a common neurodegenerative process in the aging brain
[43], and social isolation caused by both sensory and cognitive loss [45]. In addition,
recent findings have suggested that hearing impairment manifested as central
auditory dysfunction may be an early marker for dementia [47]. Previous studies
concluded that intervention in the form of hearing aids may have a positive effect on
cognition [19] and reduce the impact of behavioural and psychological symptoms of
dementia [48].

In this study, we investigated the relationship between the use of hearing aids with
incidence and progression of dementia. Our results clearly suggest that the use of
hearing aids is independently associated with a decreased risk of incident all-cause
dementia diagnosis for MCI patients. Statistically significant differences in cumulative
survival functions by hearing aid status were found in Group 1, with accelerated
cognitive decline, as indicated by change in the CDRSB score in the MCI group. The
use of hearing aids was not associated with reduced risk of death in people with
dementia. Three different sensitivity analyses confirmed the robustness of our
findings.

So far, hearing aid usage has been linked to improvements in cognition as well as
psychological, social, and emotional functioning [13]. Amieva *et al.* [49] showed that
non-use of hearing aids was associated with faster cognitive decline as indicated by

the accelerated rate of change in Mini Mental State Examination score. Yet, no significant difference in cognitive decline was observed between hearing-impaired subjects using hearing aids and healthy individuals [49]. The recent study of Maharani *et al.* [16] adopted a different approach in examining differences in cognitive outcomes of hearing aid use. To prevent potential residual confounding caused by demographic differences between hearing aid users and non-users, the authors analysed rates of cognitive change before and after hearing aid use in the same individuals. They reported a significantly slower decline in episodic memory scores after patients started to use hearing aids.

The potential mechanisms behind the association between the use of hearing aids and cognitive loss, in particular the decreased risk for incident dementia, remain to be determined. Possible explanations include optimized communication and increased social engagement, with resulting lower rates of depression and loneliness, caused by the use of hearing aids and/or changes to the brain associated with the reduced impact of sensory deprivation on brain function [50]. There is also the possibility that facilitated access to auditory information for individuals using hearing aids may result in a reduction in cognitive resources consumed by listening and, hence, lead to improved cognitive ability [50].

We acknowledge that biases in our analysis could arise from multiple sources. First, there is selection bias associated with different recruitment strategies implemented by each ADRC. Those enrolling in ADRC cohorts are not random volunteers and therefore, are not representative of a wider population. Their level of education and income is likely above the national average and approximately 50% of subjects have a family history of dementia. These factors may limit generalizability of our findings.

Another methodological limitation of our analysis is its reliance on retrospective data.

Even though all ADRCs use standard evaluation procedures, there might be some

variation in diagnostic criteria between Centers. The lack of use of consistent

diagnostic definitions due to the retrospective design of this study can lead to an

increased risk of bias due to potential misclassification of the outcome. Furthermore,

bias may arise from the degree of accuracy with which subjects have been classified

with respect to their exposure status, i.e. hearing status and hearing aid use status.

We minimized this bias by considering only participants with a consistent record of

hearing impairment and use/non-use of hearing aids. In this way, we could obtain the

true effect of hearing aids on incident and progression of dementia. The inclusion of

participants with noisy labels could likely result in an over or underestimation of the

effect between exposure (hearing aid use) and outcome (incident dementia or

death).

It is also worth highlighting that the proportion of participants wearing hearing aids in

the final groups was considerably higher than the prevalence of hearing aid use in

the general population [7]. This potential selection bias might have been introduced

into the study at the data-gathering stage (the education level and income of NACC

volunteers may not be reflective of the general population) or during the process of

identifying the study population. In fact, a large number of participants with dementia

were excluded from the analysis due to the lack of consistency in reporting hearing

difficulty.

Gender, socioeconomic factors and cultural influences all play a role in the use of

hearing aids. It may also be the case that unrecorded intrinsic factors that influence

use of hearing aids, or lack thereof, play a significant role in the findings presented

here. Indeed, severity of cognitive decline may influence acceptance, compliance

and correct usage of hearing aids. With incremental changes and decline in

cognition, the capacity to comply with the use of hearing aids is likely to significantly

diminish [8]. Despite the fact that age, education, gender, and cognitive assessment

score were included in the analysis as potential confounders, other unmeasured

factors may have impact on the incident and progression of dementia. For instance,

hearing aids use in the U.S. is dependent on financial resources as hearing aids are

expensive and generally not covered by medical insurance [8]. Other potentially

important characteristics not considered in this study due to unavailability of data

include type of hearing aid used, hours of daily use, and use of other communicative

strategies. Consequently, whether these factors may have a significant effect on time

to incident dementia for MCI patients remains unknown and will require further study.

Nevertheless, we implemented measures to account for the potential impact of

confounding and selection bias in our study. Propensity score matching was applied

to control for measured confounding; the analysis for residual confounding was

implemented to assess the sensitivity of our main conclusions with respect to

confounders not included in our study; and potential selection bias was addressed

via the inverse probability of treatment weighting method. Our results remained

robust under different assumptions.

Finally, this study relies on self-reported hearing loss which is far less reliable than

audiometric screening. The use of information on hearing impairment via self-report

prevents any adjustment for the effect of degree of hearing loss when investigating

the impact of hearing aid usage on incidence and progression of dementia. It is also

worth noting that even though ADRCs are anticipated to enter missing codes when

they have reason to believe that the cognitive test is invalid, including if they are aware that the participant is unable to hear properly, we cannot exclude the possibility that hearing impairment, rather than cognitive function, impacted the ability to complete tasks on cognitive tests in participants with hearing loss who did not have hearing aids. Since verbal instructions that are used during cognitive testing depend significantly on hearing, hearing loss might have contributed to the overestimation of the level of cognitive impairment in some hearing-impaired individuals.

Irrespective of the limitations associated with the present analyses, the fact that significant benefit appears to be derived in delaying conversion of MCI to dementia with hearing aid use warrants further exploration. Properly designed clinical trials will definitively measure the potential benefit of hearing correction in those experiencing hearing loss.

5 Conclusion

Slower conversion from MCI to dementia in individuals using hearing aids suggests that effective identification and treatment of hearing loss may reduce the cumulative incidence of dementia. The competing risk of all-cause mortality and dementia among those with MCI should be examined in future work. One of the findings reviewed above suggested that higher-level cognitive processing involving memory in hearing-impaired individuals might be compromised because of mental resources being reallocated to perception and away from storing information. We believe this hypothesis should be further tested to see if the use of hearing aids can make word identification less effortful and thus, allow for freeing resources for higher-level processing that can in turn result in improvement in cognitive function. Furthermore,

more research is needed to better understand the relationship between hearing

impairment, changes in cognitive ability, and the role of hearing aids in preventing

milder forms of cognitive impairment. Such knowledge may provide new and novel

insights into prevention of cognitive decline. Most importantly, the magnitude and

causality of the effect of hearing loss treatment on cognitive decline and incident

dementia can only be established by conducting a well-designed clinical trial.

Public health campaigns are needed to demonstrate the scale and impact of hearing

loss and increase awareness regarding effective prevention strategies,

consequences of inaction and potential benefits of timely audiological intervention.

Acknowledgements

The NACC database is funded by NIA/NIH Grant U01 AG016976. NACC data are

contributed by the NIA-funded ADCs: P30 AG019610 (PI Eric Reiman, MD), P30

AG013846 (PI Neil Kowall, MD), P30 AG062428-01 (PI James Leverenz, MD) P50

AG008702 (PI Scott Small, MD), P50 AG025688 (PI Allan Levey, MD, PhD), P50

AG047266 (PI Todd Golde, MD, PhD), P30 AG010133 (PI Andrew Saykin, PsyD),

P50 AG005146 (PI Marilyn Albert, PhD), P30 AG062421-01 (PI Bradley Hyman, MD,

PhD), P30 AG062422-01 (PI Ronald Petersen, MD, PhD), P50 AG005138 (PI Mary

Sano, PhD), P30 AG008051 (PI Thomas Wisniewski, MD), P30 AG013854 (PI

Robert Vassar, PhD), P30 AG008017 (PI Jeffrey Kaye, MD), P30 AG010161 (PI

David Bennett, MD), P50 AG047366 (PI Victor Henderson, MD, MS), P30 AG010129

(PI Charles DeCarli, MD), P50 AG016573 (PI Frank LaFerla, PhD), P30 AG062429-

01(PI James Brewer, MD, PhD), P50 AG023501 (PI Bruce Miller, MD), P30

AG035982 (PI Russell Swerdlow, MD), P30 AG028383 (PI Linda Van Eldik, PhD),

P30 AG053760 (PI Henry Paulson, MD, PhD), P30 AG010124 (PI John Trojanowski,

MD, PhD), P50 AG005133 (PI Oscar Lopez, MD), P50 AG005142 (PI Helena Chui,

MD), P30 AG012300 (PI Roger Rosenberg, MD), P30 AG049638 (PI Suzanne Craft,

PhD), P50 AG005136 (PI Thomas Grabowski, MD), P30 AG062715-01 (PI Sanjay

Asthana, MD, FRCP), P50 AG005681 (PI John Morris, MD), P50 AG047270 (PI

Stephen Strittmatter, MD, PhD)

Funding

This work was supported by the Dr George Moore Endowment for Data Science at

Ulster University (MB); European Union INTERREG VA Programme (MB, PLM, ST,

LPM); Global Challenge Research Fund (MB, XD, QY, DW, WH), Alzheimer's

Research UK (MB, SB); European Union Regional Development Fund (PLM);

Northern Ireland Public Health Agency (PLM); Dementias Platform UK (DPUK) (SB);

Nutricia (ST); Bohringer-Ingelheim (ST); Genomics Medicine Ireland (ST); Vifor

Pharma (ST); and Putnam Associates (ST).

Conflict of Interest

The authors declare that they have no competing interests.

Ethics approval and consent to participate

The National Alzheimer's Coordinating Center Uniform Data Set (NACC-UDS) is

approved by the University of Washington Institutional Review Board and

participants provided informed consent at the ADRC where they completed their

study visits. The approval was received to use the NACC-UDS data in the present

work   (application   reference: 1026)

References

1. Alzheimer's Association. *Changing the trajectory of Alzheimer's disease: a national imperative*. Alzheimer's Association, USA, 2010.

2. Sabia S, Dugravot A, Dartigues JF, et al. Physical activity, cognitive decline, and risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ*. 2017;357:j2709.

3. Yaffe K. Modifiable Risk Factors and Prevention of Dementia: What Is the Latest Evidence? *JAMA Intern Med*. 2018;178(2):281–282.

4. Lourida I, Hannon E, Littlejohns TJ, et al. Association of Lifestyle and Genetic Risk With Incidence of Dementia. *JAMA*. 2019;322(5):430–437.

5. Gottesman RF, Albert MS, Alonso A, Coker LH, Coresh J, Davis SM, Deal JA, McKhann GM, Mosley TH, Sharrett AR, Schneider AL. Associations between midlife vascular risk factors and 25-year incident dementia in the Atherosclerosis Risk in Communities (ARIC) cohort. *JAMA Neurology*. 2017;74(10):1246-54.

6. Goman AM, Lin FR. Prevalence of hearing loss by severity in the United States. American journal of public health. 2016 Oct;106(10):1820-2.

7. Chien W, Lin F.R. Prevalence of hearing-aid use among older adults in the United States. Arch Intern Med. 2012;172:292–293.

8. McCormack A, Fortnum H. Why do people fitted with hearing aids not wear them?. *International Journal of Audiology*. 2013;52(5):360-8.

9. Kochkin S, Rogin CM. Quantifying the obvious: The impact of hearing instruments on quality of life. *Hear Rev*. 2000 Jan 1;7(1):6-34.

10. Deal JA, Albert MS, Arnold M, Bangdiwala SI, Chisolm T, Davis S, Eddins A, Glynn NW, Goman AM, Minotti M, Mosley T. A randomized feasibility pilot trial of hearing treatment for reducing cognitive decline: results from the Aging and

Cognitive Health Evaluation in Elders Pilot Study. Alzheimer's & Dementia: Translational Research & Clinical Interventions. 2017;3(3):410-5.

11. Lin FR, Metter EJ, O'Brien RJ, Resnick SM, Zonderman AB, Ferrucci L. Hearing loss and incident dementia. Archives of neurology. 2011;68(2):214-20.

12. Gurgel RK, Ward PD, Schwartz S, Norton MC, Foster NL, Tschanz JT. Relationship of hearing loss and dementia: a prospective, population-based study. Otol Neurotol. 2014;35(5):775-81.

13. Deal JA, Betz J, Yaffe K, et al. Hearing Impairment and Incident Dementia and Cognitive Decline in Older Adults: The Health ABC Study. J Gerontol A Biol Sci Med Sci. 2017;72(5):703-709.

14. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *The Lancet.* 2017;390(10113):2673-734.109

15. Yong M, Willink A, McMahon C, McPherson B, Nieman CL, Reed NS, Lin FR. Access to adults' hearing aids: policies and technologies used in eight countries. Bull World Health Organ. 2019 Oct 1;97(10):699-710.

16. Maharani A, Dawes P, Nazroo J, et al. Longitudinal relationship between hearing aid use and cognitive function in older Americans. *Journal of the American Geriatrics Society.* 2018;66(6):1130-6.

17. Mulrow CD, Tuley MR, Aguilar C. Sustained benefits of hearing aids. *Journal of Speech, Language, and Hearing Research.* 1992;35(6):1402-5.

18. Dawes P, Cruickshanks KJ, Fischer ME, Klein BE, Klein R, Nondahl DM. Hearing-aid use and long-term health outcomes: Hearing handicap, mental health, social engagement, cognitive function, physical health, and mortality. International journal of audiology. 2015 Nov 2;54(11):838-44.

19. Dawes P., Emsley R., Moore D. , Cruickshanks K.J. , Fortnum H . et al. 2015. Hearing loss and cognition: The role of hearing aids, social isolation, and depression. PLOS One.

20. Beekly DL, Ramos EM, van Belle G, et al. The national Alzheimer's coordinating center (NACC) database: an Alzheimer disease database. *Alzheimer Disease & Associated Disorders.* 2004;18(4):270-7.

21. Beekly DL, Ramos EM, Lee WW, et al. The National Alzheimer's Coordinating Cetner (NACC) database: the Uniform Data Set. *Alzheimer Dis Assoc Disord.* 2007;21(3):249–258.

22. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 1994. Washington.

23. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM: Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA work group under the auspices of Department of Health and Human Services Task Force on Alzheimer's disease. Neurology 1984, 34(7):939–944

24. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimers Dement. 2011;7:263–269.

25. Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV, et al. Current concepts in mild cognitive impairment. Arch Neurol 2001; 58:1985–92.

26. Morris JC. The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology*; 1993;43:2414.

27. Bucholc M, Ding X, Wang H, et al. A practical computerized decision support system for predicting the severity of Alzheimer's disease of an individual. *Expert Systems with Applications*. 2019;130:157-71.

28. Ding X, Bucholc M, Wang H, et al. A hybrid computational approach for efficient Alzheimer's disease classification based on heterogeneous data. *Scientific Reports*. 2018;8(1):9774.

29. University of Washington, NACC Uniform Dataset Guidebook Version 3.0. 2015. Available at:

https://www.alz.washington.edu/NONMEMBER/UDS/DOCS/VER3/UDS3_ivp_guide book.pdf (accessed: 30.08.2020)

30. Riffenburgh R. Statistics in Medicine. 2006. Elsevier

31. Cox DR. Regression models and life‑tables. Journal of the Royal Statistical Society: Series B. 1972;34(2):187-202.

32. Schemper M, Wakounig S, Heinze G. The estimation of average hazard ratios by weighted Cox regression. Statistics in medicine. 2009;28(19):2473-89.

33. Box GE, Tidwell PW. Transformation of the independent variables. *Technometrics*. 1962 Nov 1;4(4):531-50.

34. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B*. 1995; 57(1): 289–300.

35. Rubin DB. Estimating causal effects from large data sets using propensity scores. *Ann Intern Med* 1997;127 (8, pt 2) 757- 763.

36. Lin DY, Psaty BM, Kronmal RA. Assessing the sensitivity of regression results to unmeasured confounders in observational studies. Biometrics. 1998 Sep 1:948-63.

37. Shen C, Li X, Li L, Were MC. Sensitivity analysis for causal inference using inverse probability weighting. Biometrical journal. 2011 Sep;53(5):822-37.

38. Alattar AA, Bergstrom J, Laughlin GA, et al. Hearing impairment and cognitive decline in older, community-dwelling adults. *The Journals of Gerontology: Series A*. 2019.

39. Strawbridge WJ, Wallhagen MI, Shema SJ, Kaplan GA. Negative consequences of hearing impairment in old age: a longitudinal analysis. *The Gerontologist*. 2000;40(3):320-6.

40. Michalowsky B, Hoffmann W, Kostev K. Association Between Hearing and Vision Impairment and Risk of Dementia: Results of a Case-Control Study Based on Secondary Data. *Front Aging Neurosci*. 2019; 11.

41. Fritze T, Teipel S, Ovari A, Kilimann I, Witt G, Doblhammer G. Hearing Impairment Affects Dementia Incidence. An Analysis Based on Longitudinal Health Claims Data in Germany. PloS One. 2016;11(7): e0156876.

42. Ray, J., Popli, G. and Fell, G., 2018. Association of cognition and age-related hearing impairment in the English Longitudinal Study of Ageing. JAMA Otolaryngology–Head & Neck Surgery, 144(10), pp.876-882.

43. Uchida Y, Sugiura S, Nishita Y, Saji N, Sone M, Ueda H. Age-related hearing loss and cognitive decline – The potential mechanisms linking the two. *Auris Nasus Larynx*. 2019;46(1):1-9.

44. Tun PA, McCoy S, Wingfield A. Aging, hearing acuity, and the attentional costs of effortful listening. *Psychology and Aging*. 2009;24(3):761.

45. Wayne RV, Johnsrude IS. A review of causal mechanisms underlying the link between age-related hearing loss and cognitive decline. *Ageing Research Reviews*. 2015;23:154-66.

46. Peelle JE, Troiani V, Grossman M, Wingfield A. Hearing loss in older adults affects neural systems supporting speech comprehension. *Journal of Neuroscience*. 2011;31(35):12638-43.

47. Gates GA, Anderson ML, McCurry SM, Feeney MP, Larson EB. Central auditory dysfunction as a harbinger of Alzheimer dementia. *Archives of Otolaryngology - Head & Neck Surgery*. 2011;137(4):390-5.

48. Kar N. Behavioral and psychological symptoms of dementia and their management. *Indian Journal of Psychiatry*. 2009;51:S77.

49. Amieva H, Ouvrard C, Giulioli C, Meillon C, Rullier L, Dartigues JF. Self-reported hearing loss, hearing aids, and cognitive decline in elderly adults: A 25-year study. *Journal of the American Geriatrics Society*. 2015;63(10):2099-104.

50. Lin FR, Yaffe K, Xia J, et al. Hearing loss and cognitive decline in older adults. *JAMA Internal Medicine*. 2013;173(4):293-9.

Figures

Fig. 1. Selection of participants for study inclusion.

Table 1. Baseline demographic and clinical characteristics of participants by hearing aid status.

Characteristic	Group 1			Group 2		
	Hearing Aid Status			Hearing Aid Status		
	Used	Not Used	*P*-value	Used	Not Used	*P*-value
Gender, No. (%)						
Male	411 (64.6)	167 (55.1)	0.01	477 (67.2)	280 (60.2)	0.02
Female	225 (35.4)	136 (44.9)		233 (32.8)	185 (39.8)	
Age, mean (SD), years	79.6 (8.9)	77.3 (8.9)	< 0.001	78.2 (7.8)	77.7 (9.5)	0.32
Education, mean (SD), years[a]	16.4 (7.2)	15.5 (7.6)	0.07	15.4 (3.2)	14.5 (3.8)	< 0.001
CDRSB score, mean (SD)	1.4 (1.1)	1.6 (1.3)	0.01	4.7 (3.0)	6.5 (4.3)	< 0.001

Abbreviation: SD, standard deviation; CDRSB, Clinical Dementia Rating Sum of Boxes

[a] measured as the number of years of education completed

Table 2. Risk of incident all-cause dementia by hearing aid status after adjustment for age, gender, race, education, and CDRSB score. Hazard ratios with the 95% confidence interval (95%CI) calculated using Cox proportional hazards regression model. FDR: false discovery rate.

Characteristic	Level	Total	Hazard Ratio (95% CI)	P value	FDR P value
Hearing Aid	No	303			
	Yes	636	0.73 (0.61, 0.89)	0.002	0.004
Gender	Male	578			
	Female	361	1.03 (0.86, 1.24)	0.76	0.93
Age		939	1.01 (0.99, 1.02)	0.31	0.51
Education		939	1.00 (0.99, 1.01)	0.93	0.93
CDRSB score		939	1.39 (1.30, 1.48)	< 0.001	< 0.001

Abbreviation: CDRSB: Clinical Dementia Rating Sum of Boxes; FDR: false discovery rate.

Table 3. Risk of death by hearing aid status for individuals with dementia after adjustment for age, gender, race, education, and CDRSB score. Hazard ratios with the 95% confidence interval (95% CI) calculated using weighted Cox regression model accounting for time-varying effects. FDR: false discovery rate.

Characteristic	Level	Total	Hazard Ratio (95% CI)	P value	FDR P value
Hearing Aid	No	465			
	Yes	710	0.98 (0.78, 1.24)	0.89	0.89
Gender	Male	757			
	Female	418	0.79 (0.63, 0.99)	0.04	0.06
Age		1175	1.02 (1.00, 1.03)	0.02	0.05
Education		1175	1.01 (0.98, 1.05)	0.46	0.57
CDRSB score		1175	1.08 (1.05, 1.11)	< 0.001	< 0.001

Abbreviation: CDRSB: Clinical Dementia Rating Sum of Boxes; FDR: false discovery rate.

Table 4. Risk of incident all-cause dementia (Group 1) and risk of death (Group 2) by hearing aid status in the propensity score matched sample.

Characteristic	Level	Total	Hazard Ratio (95% CI)	P value	FDR P value
Group 1					
Hearing Aid	No	303			
	Yes	303	0.67 (0.50, 0.80)	0.004	0.01
Gender	Male	427			
	Female	179	0.92 (0.71, 1.19)	0.52	0.65
Age		606	1.01 (0.99, 1.02)	0.48	0.65
Education		606	0.99 (0.98, 1.01)	0.9	0.91
CDRSB score		606	1.37 (1.27, 1.50)	< 0.001	< 0.001
Group 2					
Hearing Aid	No	465			
	Yes	465	0.93 (0.69, 1.27)	0.66	0.66
Gender	Male	631			
	Female	299	0.91 (0.70, 1.21)	0.54	0.66
Age		930	1.01 (0.99, 1.03)	0.14	0.35
Education		930	1.02 (0.98, 1.07)	0.28	0.47
CDRSB score		930	1.07 (1.03, 1.10)	< 0.001	0.002

Abbreviation: CDRSB: Clinical Dementia Rating Sum of Boxes; FDR: false discovery rate.

Table 5. Sensitivity analysis for unmeasured confounding

Prevalence of unmeasured confounder (%)		HR adjusted for unmeasured confounder (95% CI)[a]			
Hearing aid used	Hearing aid not used	Unmeasured confounder HR 0.5	Unmeasured confounder HR 1.5	Unmeasured confounder HR 2.0	Unmeasured confounder HR 3.0
Group 1					
5	10	0.75 (0.62,0.91)	0.72 (0.59,0.87)	0.70 (0.58,0.85)	0.67 (0.56,0.82)
	20	0.80 (0.66,0.96)	0.68 (0.57,0.83)	0.64 (0.53,0.78)	0.58 (0.48,0.70)
	30	0.84 (0.70,0.97)	0.65 (0.54,0.79)	0.59 (0.49,0.72)	0.51 (0.42,0.61)
10	10	0.73 (0.61,0.89)	0.73 (0.61,0.89)	0.73 (0.61,0.89)	0.73 (0.61,0.89)
	20	0.78 (0.64,0.94)	0.70 (0.58,0.85)	0.67 (0.56,0.82)	0.63 (0.52,0.76)
	30	0.82 (0.68,0.99)	0.67 (0.55,0.81)	0.62 (0.51,0.75)	0.55 (0.46,0.67)
20	10	0.70 (0.58,0.84)	0.77 (0.64,0.93)	0.80 (0.66,0.97)	0.84 (0.71,0.99)
	20	0.73 (0.61,0.89)	0.73 (0.61,0.89)	0.73 (0.61,0.89)	0.73 (0.61,0.89)
	30	0.78 (0.64,0.94)	0.70 (0.58,0.85)	0.68 (0.56,0.82)	0.64 (0.53,0.78)
Group 2					
5	10	1.03 (0.81,1.31)	0.98 (0.77,1.25)	0.96 (0.75,1.22)	0.92 (0.72,1.17)
	20	1.09 (0.85,1.38)	0.93 (0.73,1.19)	0.88 (0.69,1.12)	0.79 (0.62,1.00)
	30	1.15 (0.90,1.47)	0.89 (0.70,1.14)	0.81 (0.63,1.03)	0.69 (0.54,0.88)
10	10	1.00 (0.79,1.28)	1.00 (0.79,1.28)	1.00 (0.79,1.28)	1.00 (0.79,1.28)
	20	1.06 (0.83,1.35)	0.96 (0.75,1.22)	0.92 (0.72,1.17)	0.86 (0.67,1.10)
	30	1.12 (0.88,1.43)	0.91 (0.72,1.17)	0.85 (0.66,1.08)	0.75 (0.59,0.96)
20	10	0.95 (0.74,1.21)	1.05 (0.82,1.34)	1.09 (0.86,1.39)	1.17 (0.92,1.49)
	20	1.00 (0.79,1.28)	1.00 (0.79,1.28)	1.00 (0.79,1.28)	1.00 (0.79,1.28)
	30	1.06 (0.83,1.35)	0.96 (0.75,1.22)	0.92 (0.73,1.18)	0.88 (0.69,1.12)

Abbreviation: HR, hazard ratio

[a] All models were adjusted for sex, age, education, and CDRSB score

Supplementary material

Table A.1. Baseline demographic and clinical characteristics of participants included and excluded from the study.

	Group 1			Group 2		
	MCI at baseline			Dementia at baseline		
	Excluded	Included		Excluded	Included	
Total	4461	939		8951	1175	
Gender, No. (%)						
Male	2055 (46.1)	578 (61.6)	< 0.001	4259 (47.6)	757 (64.4)	< 0.001
Female	2406 (53.9)	361 (38.4)		4692 (52.4)	418 (35.6)	
Age, mean (SD), years	74.7 (9.0)	78.9 (8.0)	< 0.001	73.4 (9.6)	78.0 (8.5)	< 0.001
Education, mean (SD), years[a]	15.5 (5.9)	16.1 (7.4)	0.03	15.3 (8.3)	15.1 (3.5)	0.12
CDRSB score, mean (SD)	1.4 (1.1)	1.5 (1.2)	0.01	5.7 (3.9)	5.4 (3.7)	0.01

Abbreviation: SD, standard deviation; CDRSB, Clinical Dementia Rating Sum of Boxes

[a] measured as the number of years of education completed

Table A.2. Sensitivity analysis based on the inverse probability weighting

Characteristic	Level	Total	Hazard Ratio (95% CI)	*P* value	FDR *P* value
Group 1					
Hearing Aid	No	303			
	Yes	636	0.72 (0.60, 0.88)	<0.001	0.003
Gender	Male	578			
	Female	361	0.98 (0.81, 1.20)	0.87	0.87
Age		939	1.01 (0.99, 1.02)	0.31	0.52
Education		939	1.00 (0.98, 1.01)	0.69	0.86
CDRSB score		939	1.35 (1.25, 1.46)	< 0.001	< 0.001
Group 2					
Hearing Aid	No	465			
	Yes	710	0.99 (0.80, 1.23)	0.92	0.92
Gender	Male	757			
	Female	418	0.73 (0.60, 0.91)	0.004	0.01
Age		1175	1.02 (1.00, 1.03)	0.01	0.02
Education		1175	1.01 (0.98, 1.04)	0.49	0.62
CDRSB score		1175	1.08 (1.05, 1.10)	< 0.001	< 0.001

Abbreviation: CDRSB: Clinical Dementia Rating Sum of Boxes; FDR: false discovery rate.

Table A.3. Risk of incident all-cause dementia (Group 1) and risk of death (Group 2) by hearing aid status. The group of hearing aid users consisted of subjects with functionally normal hearing with a device and those that exhibited reduced ability to do everyday activities such as listening to the radio/television or talking with family/friends when wearing a hearing aid

Characteristic	Level	Total	Hazard Ratio (95% CI)	P value	FDR P value
Group 1					
Hearing Aid	No	303			
	Yes	765	0.74 (0.61, 0.89)	0.001	0.003
Gender	Male	659			
	Female	409	1.05 (0.89, 1.25)	0.55	0.69
Age		1068	1.01 (0.99, 1.02)	0.32	0.54
Education		1068	1.00 (0.99, 1.01)	0.82	0.82
CDRSB score		1068	1.40 (1.32, 1.49)	< 0.001	< 0.001
Group 2					
Hearing Aid	No	465			
	Yes	884	0.99 (0.80, 1.23)	0.92	0.89
Gender	Male	876			
	Female	473	0.73 (0.60, 0.91)	0.004	0.01
Age		1349	1.02 (1.00, 1.03)	0.01	0.02
Education		1349	1.01 (0.98, 1.04)	0.49	0.61
CDRSB score		1349	1.07 (1.05, 1.10)	< 0.001	< 0.001

Abbreviation: CDRSB: Clinical Dementia Rating Sum of Boxes; FDR: false discovery rate.