

Assessing the Tendency of 2019-nCoV (COVID-19) Outbreak in China

Qinghe Liu¹, Deqiang Li¹, Zhicheng Liu¹, Zefei Gao¹, Junkai Zhu¹, Junyan Yang², Qiao Wang^{1*}

¹ School of Information Science and Engineering, Southeast University, Nanjing, 210096, China

² School of Architecture, Southeast University, Nanjing, 210096, China

*Correspondence to Q. Wang (Email: qiaowang@seu.edu.cn)

ABSTRACT.

Since December 8, 2019, the spread of 2019-nCoV (COVID-19) is increasing day by day. It is particularly important to predict the trend of the epidemic for the timely adjustment of the economy and industries. A Flow-SEHIR model is proposed to perform empirical analysis for Hubei Province and those areas in China outside Hubei.

It is shown that the time when new confirmed cases number has reached a peak level, i.e. the inflection point, is verified to be on February 6 - 10. The maximum of temporal infected cases number is represented by peak value. The predicted peak value in China except Hubei was estimated as 31300 (95% CI: 27700 - 36800). The peak arrival time is on March 3 - 9. The number of real-time patients in most areas of China outside Hubei will peak from March 2 to March 24. The peak values of more than 60% provinces or regions in China will be controlled within 1000. Considering the false-negative rate of nucleic acid detection is 0.5, the real peak cumulative number of patients in Hubei is estimated between 214400 - 472500.

Abbrevians: 2019-nCoV; COVID-19, coronavirus disease 2019; SARS-Cov-2, SARS coronavirus 2;

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1. INTRODUCTION

Coronavirus [1] is a single stranded RNA virus, which is spherical in shape and has petal-like spines. It usually causes mild respiratory infections and can sometimes be fatal. Since its first discovery and identification in 1965, there have been three large-scale outbreaks of highly treatable infectious diseases caused by coronavirus, namely, SARS in mainland China in 2003, MERS in Saudi Arabia in 2012 and MERS in South Korea in 2015 [2].

At the beginning of December, the first patient infected by a new type of coronavirus was found in Wuhan, Hubei, which is located in the central part of China. Then similar symptoms were found in the same city. After being infected, the main symptoms of patients in the early stage include low fever and fatigue, which were similar to the symptoms caused by general influenza, so it spreads in a large area without people's awareness. On January 12, 2020, the World Health Organization officially named the new coronavirus causing pneumonia in Wuhan as "2019 new coronavirus[3] (SARS-CoV-19)". On January 22, 2020, researchers speculated that the new coronavirus might come from wild bats[4]. During the Spring Festival of 2018 (from February 1 to March 12, 2018), the total estimated number of passenger trips in mainland China is about 2.97 billion [5]. As of 9 am on February 7, 2020, the cumulative reports of 31 provinces (districts and municipalities) received by the National Health Commission of the People's Republic of China [6] are shown in Table 1. In the table, Hubei Province's data of confirmed cases is temporarily excluded since Hubei Province, as a severe epidemic area, has the possibility of a large backlog of

confirmed cases, combined with the shortage of medical resources and the uncertainty of the diagnosis method, which makes the data in Hubei more unreliable than other provinces. Among these reports, 22112 cases of pneumonia with new coronavirus infection were found in Hubei Province[7].

The serious epidemic has brought many urgent problems. When the epidemic can be stable? When the growth of the number of confirmed cases can stop? To answer these questions, we perform an empirical analysis of the epidemic trend. In section 2 of this paper, the most up-to-date literature by 7 Feb 2020 related to COVID-19 epidemic situation will be discussed and briefly analyzed. Section 3 and 4 presents Flow-SEHIR model in details. Section 5 presents the experiments settings and empirical analysis. Finally, section 6 and 7 are results and conclusions.

2 RELATED ISSUES

On January 27, 2020, B. Tang *et al* [10] established a transmission dynamics model based on the transmission mechanism of new coronavirus in Wuhan, isolation and other strategies. They thought the reproduction number of infection is 3.8 and the intervention measures can reduce the reproduction number of infection and the risk of transmission. Based on the epidemic data from Jan. 10 to Jan. 22, they predicted the basic reproduction number of infection of novel coronavirus in Wuhan is 6.47 (95% CI: 5.71 – 7.23).

On January 27, 2020, J. A. Backer *et al* [11] estimated that the average incubation period should be 5.8 days (95% CI: 4.6-7.9 days) and the range of variation be 1.3-11.3

days, according to the travel records and symptoms of 34 confirmed cases found outside Wuhan. From which they considered that 14 days of isolation will ensure that no symptoms will occur. In addition, they compared the COVID-19 with SARS and MERS.

On January 31, 2020, J. Wu *et al* [12] used data from Dec 31, 2019, to Jan 28, 2020 thought that the basic reproduction number of infection was estimated by using Markov Chain Monte Carlo methods and presented by the the resulting posterior mean and 95% CI. They estimated that the basic reproduction number of infection in Wuhan in 2019 was 2.68 (95% CI: 2.47-2.86). In addition, as of January 25, the number of infections was 75815 (95% CI: 37304-130330). The epidemic doubling time was 6.4 days (95% CI: 5.8-7.1).

On February 2, 2020, Adam J Kucharski *et al* [13] from London School of Hygiene & Tropical Medicine used a stochastic transmission model, based on confirmed cases in Wuhan and exported cases originating in Wuhan and estimated that the reproduction number of infection from mid-December to mid-January 2020 fluctuated between 1.6-2.9.

Epidemiological study is often an afterwards research, such as our previous work[14]. However, there are many uncertainties and risks in the development of COVID-19, so we need to provide some suggestions from the data analysis work in time.

On Feb.9, 2020, X. Li *et al* [15] simulated the dynamics of the COVID-19 and their results show that after the the traffic blockage in Jan.23,2020, the reproductive number in Wuhan will be 2.5, lower than before. The confirmed cases increased in linear not in

exponential, which can confirm the effectiveness of strict control measures.

On Feb.12, 2020, Qiushi Lin *et al* [16] provide estimates of the daily trend in the size of the epidemic in Wuhan and found the number of cases that should be reported in Wuhan by January 11, 2020 is 4090 (95% CI: 3975 – 4206) and 56833 (95% CI: 55242 – 58449) by February 9, 2020.

Since the outbreak of the epidemic, many scholars have concentrated in Wuhan and other areas, and studied the impact of basic reproduction number of infection and traffic interruption. However, the outbreak of the epidemic has a huge impact on the economy and people's livelihood. Predicting the peak arrival date and evaluating the effectiveness of the control measures will play a decisive role in ensuring the smooth operation of the social economy. In this paper, two main areas (Hubei and other regions outside Hubei in China) are focused on for larger scale analysis. Deqiang Li *et al* [17] in our team conducted an in-depth analysis of the effects of traffic blockage and medical quarantine. It is shown that the quarantine is very effective for epidemic control, which can reduce the number of infections by nearly 90%.

3 COVID-19 TRANSMISSION MODEL: FLOW-SEHIR

3.1. SEIR Model.

The SEIR model is introduced in 1984 by Aron, J. and I. Schwartz [18], where S, E, I and R referring to the Susceptibles, Exposed, Infective, and Recovered, respectively. Compared with the standard SIR model, the SEIR model emphasized the importance of the Exposed. In the epidemic transmission, when susceptible ones are infected, most

of them will neither immediately get sick nor display symptoms, but patients in the exposed period are also infectious, and during the recovery they will possibly be immune. For the details of SEIR model, please refer to as [19][20]. Finally, the incubation period and the time-varying propagation rate are investigated in [21] and [22] respectively.

3.2. Flow-SEHIR Model.

Before building the model, we make 2 main assumptions about practical problems:

- (1). The interaction of people flow between domestic and foreign areas is ignored.
- (2). During the transmission of virus, the probability of infection in contact with each person is equal.

Furtherly considering the special communication feature in Chinese Springer Festival and the effectiveness of nucleic acid testing is questionable, we add the factor of flow to reflect the mobility of the population and use H denote ‘Hidden’, which means patients not detected by nucleic acid test. This forms our Flow-SEHIR model:

$$\begin{cases} \frac{dS}{dt} = F - \beta(I + lH)S \\ \frac{dE}{dt} = \beta(I + lH)S - (\frac{\alpha}{1-\omega} + \gamma_1)E \\ \frac{dH}{dt} = \frac{\omega}{1-\omega} \alpha E - \mu H \\ \frac{dI}{dt} = \alpha E - \gamma_2 I + \mu H \\ \frac{dR}{dt} = \gamma_1 E + \gamma_2 I \end{cases} \quad (3.1)$$

The flow $F(t)$ rewrites the standard SEIR model by the first equation of (3.1). In practice, $F(t)$ is estimated by the interregional migration intensities from the Baidu Migration API. The explanations of all parameters in the proposed model is shown in table 2.

3.3. Flow and the Network of Cities.

The graph model is used to explain traffic flows across the country. Each province is described as an independent node, and a system of Flow-SEHIR model as well. The flow intensity of each edge of the network is F , which represents potential impact on population flow. To perform this approach, we collected population migration data from Baidu API which provides daily intra-province population migration strength. For the i -th city or province, this floating population is defined as follows:

$$F_i = \sum_{j=1,2,\dots,N_j} (1 - \tau) \delta_{ji} S_{ji} \quad (3.3)$$

in which $(1 - \tau) \delta_{ji} S_{ji}$ means the flow factor of the corresponding node, δ_{ji} is the natural flow factor of the corresponding node independent of control policy of traffic blockage. It represents the ratio of floating population to potential population. Traffic blockage factor τ is introduced into the model, which is between 0 and 1. The higher the degree of traffic blockage over traffic flow implemented, the greater the flow of restriction follows. When $\tau=0$, the traffic flow is in the uncontrolled state.

3.4. Practical Consideration.

In Flow-SEHIR model, the transmission process of the COVID-19 can be divided into the following four main stages:

- (1). The outbreak is potentially spreading, but was not been aware of by people;
- (2). Starting from the first case was officially reported since December 8, 2019;
- (3). Official confirmation (including transmission control in key cities, such as Wuhan, Hubei);
- (4). The epidemic is under control and most of patients have been recovered.

Return trend, school opens days are mostly concentrated in this stage.

The proposed model simulates stage 2, 3 and 4. Assume that in each stage, the recovery rates are relatively consistent, in terms of the natural recovery of self-recovered and medical treatment. But at each stage, the infection rate and the population can vary dramatically. For the first two stages, the spread of the virus satisfies the exponential distribution. For the third stage, it is found that there is a large number of population flows in and out because of the transportation during Spring Festival.

For simulating the impact of return flow, we gradually reduced the traffic blockage factor τ from February 10 to February 29. It should be noted that due to the serious epidemic situation in Hubei Province, the traffic control factors are very low, so we mainly adjust from 0.9 to 0.8 in Hubei Province, and from 0.9 to 0.5 in areas outside Hubei Province.

Assuming that the suspected and infected population have been isolated and do not have the ability to transmit to the outside, starting with the number of potential susceptible S , then the transmission might increase by:

$$\beta = kR_0(1 - \varepsilon) \quad (3.2)$$

in which β is the rate of those who are transmitted to the infected from the susceptible, and k is a constant which will be estimated by real patient data. R_0 is the basic reproduction number. For the third stage, there are a large number of population flows out from Wuhan, because of the Spring Festival as well as the panic. In this case, F will increase sharply. After then, traffic blockage will make F decrease. Due to self-protection and quarantine, the quarantine factor ε increases, then β decline. For the recovery rate parameters, we collected the empirical data related to pneumonia as the prior distributions.

3.5. Parameter Estimation.

During calculation, (3.1) is implemented by following iterations with prescribed steps:

$$\begin{cases} S \leftarrow S + F - \beta(I + lH)S \\ E \leftarrow E + \beta(I + lH)S - \left(\frac{\alpha}{1-\omega} + \gamma_1\right)E \\ H \leftarrow H + \frac{\omega}{1-\omega}\alpha E - \mu H \\ I \leftarrow I + \alpha E - \gamma_2 I + \mu H \\ R \leftarrow R + \gamma_1 E + \gamma_2 I \end{cases} \quad (3.4)$$

In the process of parameter iteration, in order to get convergence parameters faster, we use Gradient Descent method to build BP network. In this model, the loss function f is defined by the least square method:

$$f = \frac{1}{M} \sum_{m=1}^M (z_m - \hat{z}_m)^2 \quad (3.5)$$

in which M means the length of the real data, z_m is the real data on the m th time tag, $m=1,2,\dots,M$ and \hat{z}_m is the estimated value.

3.6. Data Source and Processing.

In this study, we used all data from official open data sets online, and main sources are: National Health Commission of the people's Republic of China, Xinhua net, Baidu migration API. The detailed data capture links are shown in appendix 1.

National Health Commission of the people's Republic of China. The national health commission of the People's Republic of China regularly releases information on the epidemic situation in different parts of the country, which is officially released as public data, including local real-time diagnosis data, suspected data, cure data, etc.

Xinhua net. Xinhua net is mainly used to get official news on the epidemic, and the data is public.

Baidu migration API. Baidu migration data is free and open traffic flow data obtained on baidu platform, including traffic intensity, migration ratio, etc.

On February 12, 2020, the number of newly diagnosed patients in Hubei increased by about 14000 due to the change of diagnosis specification. For expanding the treatment scale, the confirmation of the disease will not only depend on the nucleic acid test, which will help to greatly reduce the proportion of false negative patients. Considering that this is a policy mutation, it is assumed that only this day is abnormal in terms of increment, and the normal increment level is restored after February 12. Therefore, in the experiment, the false negative ratio ω was adjusted to decrease on February 12, and the detection rate of false-negative patients μ was adjusted from 0 to 1 to deal with the occurrence of abnormal data.

4. MAIN RESULTS OF FLOW-SEHIR MODEL

4.1. Estimate of the Peak of Net Confirmed Cases.

We define peak value as the peak of $I(t)$, which is the cumulative number of infected cases minus the cumulative number of cured cases, i.e. the number of net infected cases. After this peak point, the temporary number of cases gradually decrease and the epidemic subsides.

We first simulated the epidemic trend of 34 provinces of China, and predicted the peak number of confirmed cases and its arrival time in each area, as shown in figure1, 2. On average, the peak number of real-time confirmed cases of Hubei is predicted to reach 63800, and the peak arrival time is on March 3, 2020. Tibet has both the minimum

peak value and the earliest peak arrival time. According to the data up to February 12, 2020, the number of confirmed cases in Tibet is only 1, and there is no suspected patient. The predicted peak number of net confirmed cases of COVID-19 in China will reach 85500 (95% CI: 76700 - 97500). On the one hand, it is shown that the transmission speed is very high, on the other hand, the rapid development of traffic caused the rapid development of cross transmission. The predicted peak values of different areas are divided to different intervals, which is listed in table 3. Except for Hubei, there are no other provinces that the peak value is more than 4000.

Table 3 distinguishes Hubei Province as a single category to reflect the corresponding relationships and differences. In terms of peak values, Hubei is far higher than other regions. Zhejiang, Guangdong, Henan, Hunan and Chongqing are the places most likely to become the next worst-hit areas, and the relevant departments should pay much attention to them. The peak value of more than 60% provinces or regions of China is predicted to be stable within 1000.

It is estimated that the peak arrival time of Hubei will be at the beginning of March, which is earlier than other provinces in China. Therefore, it is very urgent to analyze the progress of the disease in Hubei. Then, comparative analysis and discussion on more details of Hubei are conducted.

4.2. Simulation of Net Confirmed Cases.

Figure 4 and 5 show the simulation trend of Hubei Province and other regions, respectively. According to the results, the potential population affected by the epidemic will be stable in March 2020. Cumulative confirmed cases number in China is estimated

to reach 140000. As of February 5, 2020, the confirmed cases number in Hubei is 16678, only 16.7% of the population base of the potential population, and there is still a large number of infected cases that have not been tracked or confirmed. The epidemic situation in Hubei will not be completely controlled and eliminated until June 2020. Except that the trend of except-Hubei is basically consistent with that of Hubei, the change of the epidemic situation of except-Hubei is higher than that in Hubei. The trend of the confirmed cases' number in Hubei is steeper, which is consistent with the characteristics of a diffusion center, the change of which time is earlier and drastic. While comparing with the center of epidemics, the periphery presents the characteristics of postponement and slowness. The trend is usually steeper and ahead of time in the center of the epidemic compared with other regions. On the one hand, it is related to the timely protection measures that have taken to other areas. On the other hand, it is due to the information lag of the central location. Similarly, according to the data, compared with other regions in Hubei Province, Wuhan also shows obvious prepositiveness and steepness. In addition, the return trend will not cause the next peak of illness, and is relatively safe, due to the improvement of people's awareness of isolation and protection, as well as the importance of epidemic management in various industries.

The errors between the predicted data and the real data in Hubei, Guangdong, Jiangsu and Anhui are shown in Figure 6. Here we define the inflection point as the peak value of new infections number per day, and after the inflection point, the number of infected people will grow slowly and the epidemic will be gradually controlled. We can

get the inflection point after first-order difference calculation of the cumulative cases trend, which is expected to be on February 7 - February 9.

In figure 7 and 8, we took the upper quantile of 0.9 as the cure rate, and took the upper quantile of 0.1 as the prevalence rate, to represent a better case for simulation; and we took the upper quantile of 0.1 as the cure rate, took the upper quantile of 95% as the prevalence rate, to represent a worse case for simulation, and the results are shown in Table 4 and 5. The peak time fluctuations of Hubei and other regions are respectively about February 3, 2020 (95% CI: February 27 – March 18) and March 9, 2020 (95% CI: March 2 – March 24).

4.3. Estimation of the real number of patients in Hubei

Considering that there are a large number of false negative cases. Because these people have not received medical treatment, the transmission is stronger than the confirmed cases. Therefore, we assume that the potential transmission rate of false negative is 1 times of that of confirmed cases.

In figure 9, assuming l is 2, the peak number of patients is between 63800 and 268000. If l is 3, the maximum number of patients is 380000. Assuming l is 4, the peak value reaches 490000. If l reaches 5, the peak value will exceed 600000.

In figure 10, the results show that if the false negative rate decreased by 5% in one week, the peak number of patients would slow down by 11.62%. If the proportion decreased by 10% in one week, the peak number of patients would decrease by 21.91%. It can be seen that timely detection of false negative population is more effective for epidemic control.

5. CONCLUSION

We performed simulation analysis of Flow-SEHIR model, which is proposed for assessing the tendency of COVID-19 in China. From analysis, we draw the following conclusions:

- (1). The peak of areas except Hubei in China will come at March 9, 2020 (95% CI: March 2 –March 24). More than 60% of provinces and regions in China will be controlled within 1000. Rather than larger outbreak, the estimated number of infective patients in Hubei(in which Wuhan is situated) presents a front and steep feature, manifesting the overall trend that the epidemic tends to be consistent.
- (2). The predicted peak value in China except Hubei is 31300 (95% CI: 27700 - 36800). The predicted peak value in Hubei is estimated as 63800 (95% CI: 59300 - 76500), and its arrival time is on March 3, 2020 (95% CI: February 27 – March 18). The predicted peak value of China is 85500 (95% CI: 76700 - 97500) and cumulative confirmed cases number in China will reach 140000. It will take about 1.5-2 months from the peak to the end of the epidemic. Due to the people's sense of isolation and safety protection, the return trend will not have a lot of impact on the epidemic, which will continue to show a downward trend after February 10.
- (3). Assuming that the false-negative rate is 50% and the potential transmission rate is 1-4 times higher than that of the confirmed patients, the real peak number of patients in Hubei is likely to be between 214400-472500. Suppose that the transmission rate of the false negative patient is twice that of the confirmed patient, if the false negative rate decreased by 5% in one week, the peak number of patients

will slowed down by 11.62%. If the proportion decreased by 10% in one week, the peak number of patients will decreased by 21.91%, which shows timely detection of false negative population is more effective for epidemic control.

Reference

1. Chen Y, Liu Q, Guo D. Coronaviruses: genome structure, replication, and pathogenesis[J]. *Journal of Medical Virology*, 2020:doi:10.1002/jmv.25681
2. Bogoch II, Watts A, Thomas-Bachli A, Huber C, Kraemer MU, Khan K. Pneumonia of unknown etiology in Wuhan, China: potential for international spread via commercial air travel[J]. *Journal of Travel Medicine* 2020:doi:10.1093/jtm/taaa1008
3. 'Definition of suspected cases of unexplained pneumonia', the National Health Commission of the People's Republic of China (in Chinese). [<http://www.nhc.gov.cn/>]
4. Li W, Shi Z, Yu M, *et al.* Bats are natural reservoirs of SARS-like coronaviruses[J]. *Science*, 2005, 310(5748): 676-679.
5. Hu M. Visualizing the largest annual human migration during the Spring Festival travel season in China[J]. *Environment and Planning A: Economy and Space*, 2019, 51(8): 1618-1621.
6. National Health Commission of People's Republic of China: <http://www.nhc.gov.cn/> [cited 7 Feb, 2020]
7. Health Commission of Hubei Province: <http://wjw.hubei.gov.cn/> [cited 7 Feb, 2020]

8. Read JM, Bridgen JRE, Cummings DAT, *et al.* Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions[J]. medRxiv, 2020.
9. Zhao S, Ran J, MUSA SS, *et al.* Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak[J]. bioRxiv, 2020:doi:10.1016/j.ijid.2020.01.050
10. Jung S, Kinoshita R, Hayashi K, *et al.* Epidemiological identification of a novel infectious disease in real time: Analysis of the atypical pneumonia outbreak in Wuhan, China, 2019-20 [J]. medRxiv, 2020.doi:10.1101/2020.01.26.20018887
11. Tang B, Wang X, Li Q *et al.* Estimation of the transmission risk of 2019-nCov and its implication for public health interventions. Available at SSRN 3525558, 2020
12. Backer JA, Klinkenberg D, Wallinga J. The incubation period of 2019-nCoV infections among travellers from Wuhan, China[J]. medRxiv, 2020:doi: 10.1101/2020.01.27.20018986
13. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: A modelling study[J]. the Lancet, 2020: doi:10.1016/S0140-6736(20)30260-9
14. Kucharski AJ, Russell TW, Diamond C *et al.* . Early dynamics of transmission and control of 2019-nCoV: a mathematical modelling study[J]. medRxiv, 2020:doi:10.1101/2020. 01.3120019901
15. Lu J, Liu W, Xia R, *et al.* Effects of closing and reopening live poultry markets on

- the epidemic of human infection with avian influenza A virus[J]. Journal of biomedical research, 2016, 30(2): 112.
16. Ai, Siqi,Zhu, Guanghu,Tian, Fei et al. Population movement, city closure and spatial transmission of the 2019-nCoV infection in China[J]. medRxiv, 2020: 2020.02.04.20020339.
 17. Qiushi Lin, Taojun Hu, Xiao-Hua Zhou, Estimating the daily trend in the size of COVID-19 infected population in Wuhan[J]. medRxiv, 2020: 2020.02.12.20022277
 18. Deqiang Li, Qinghe Liu, Zhicheng Liu, Zefei Gao, Junkai Zhu,Junyan Yang,Qiao Wang, Estimating the Efficacy of Traffic Blockage and Quarantine for the Epidemic Caused by 2019-nCoV (COVID-19), medRxiv, 2020:022913.
 19. Aron JL, Schwartz IB. Seasonality and period-doubling bifurcations in an epidemic model[J].Journal of theoretical biology, 1984, 110(4): 665-679.
 20. Li MY, Muldowney JS. Global stability for the SEIR model in epidemiology[J]. Mathematical biosciences, 1995, 125(2): 155-164.
 21. Li MY, Smith HL, Wang L. Global dynamics of an SEIR epidemic model with vertical transmission[J]. SIAM Journal on Applied Mathematics, 2001, 62(1): 58-69.
 22. Kim S, Byun JH, Jung IH. An SEIR epidemic model where empirical distribution of incubation period has approximated by coxian distribution [J]. arXiv preprint arXiv:1906.05556, 2019.
 23. Smirnova A, deCamp L, Chowell G. Forecasting epidemics through nonparametric

estimation of time-dependent transmission rates using the SEIR model[J]. Bulletin of mathematical biology, 2019, 81(11): 4343-4365.

Appendix1 Data Source (All data used in paper is public.)

Provincial-Level Migration Data of China:

<http://qianxi.baidu.com/>

Reported Cases Data:

<http://www.nhc.gov.cn/>

<http://wjw.beijing.gov.cn/>

<http://wjw.fujian.gov.cn/>

<http://wsjk.tj.gov.cn/>

<http://wsjkw.hebei.gov.cn/>

<http://www.xjhfdc.gov.cn/>

<http://wsjkw.nx.gov.cn/>

<http://ynswsjkw.yn.gov.cn/wjwWebsite/web/index>

<http://www.gzhfdc.gov.cn/>

<http://wsjkw.sc.gov.cn/>

<http://wsjkw.cq.gov.cn/>

<http://wst.hainan.gov.cn/swjw/index.html>

<http://wsjkw.gxzf.gov.cn/>

<http://wsjkw.gd.gov.cn/>

<http://wjw.hunan.gov.cn/>

<http://wjw.hubei.gov.cn/>

<http://www.hnwsjsw.gov.cn/>

<http://wsjkw.shandong.gov.cn/>

<http://hc.jiangxi.gov.cn/>

<http://wjw.ah.gov.cn/>

<http://www.zjwjw.gov.cn/>

<http://wjw.jiangsu.gov.cn/>

<http://wsjkw.sh.gov.cn/>

<http://wsjkw.jl.gov.cn/>

<http://wjw.nmg.gov.cn/>

<http://wjw.beijing.gov.cn/>

<http://sxwjw.shaanxi.gov.cn/>

<http://wjw.shanxi.gov.cn/>

<http://wsjk.ln.gov.cn/>

<http://wsjk.gansu.gov.cn/>

<http://wsjkw.hlj.gov.cn/>

<https://wsjkw.qinghai.gov.cn/>

<http://wjw.xizang.gov.cn/>

Figures and Tables




Figure 1. Flow-SEHIR Model Diagram




Figure 2. Predicted peak number of patients in each province of China, the x-axis corresponds to different geographical areas and the y-axis is the logarithm calculation of the predicted number of people.




Figure 3. Predicted arrival time of peak number of confirmed cases in each province of China, the x-axis corresponds to different geographical areas and the y-axis is the prediction of peak arrival time at average level.




Figure 4. Simulated epidemic trend of Hubei with $\tau = 0.9$ and $\varepsilon = 0.25$






Figure 5. Simulated epidemic trend of China except Hubei with $\tau = 0.9$ and $\varepsilon = 0.25$





(g) Cumulants of Anhui

(h) Increment of Anhui

Figure 6. Comparisons between the predicted infective data and the real infective data




Figure 7. Iteration results of the number of confirmed cases in Hubei Province. Worse condition uses recovery rate of 0.05 CI, better condition uses recovery rate of 95% CI.




Figure 8. Iteration results of the number of confirmed cases in China except Hubei.

Worse condition uses recovery rate of 0.05 CI, better condition uses recovery rate of 95% CI.




Figure 9. Estimation of the real number of patients with different l (2, 3, 4, 5)




Figure 10. Estimation of the real number of patients with different decrease on false negative ratio (0%, 5%, 10%) within a week.

Table 1. Real time epidemic situation in China by 9am February 7 (*Data source: data released by the National Health Commission of the People's Republic of China and the Health Commissions of all provinces and autonomous regions*)

Cumulative total	Confirmed cases (except Hubei)	Death cases	Cured cases	Suspected cases
31 provinces of China	9049	637	1543	27657

Table 2. List of Symbols

S	Number of susceptible people
E	Number of exposed people
H	Number of false-negative people
I	Number of infective people
R	Number of recovered people
β	Rate of those susceptible and being infected.
α	Rate of those exposed and being infectious.
γ_1	Transmission rate to the recovered from the exposed.
γ_2	Transmission rate to the recovered from the infectious.
γ_3	Transmission rate to the recovered from the false-negative.
μ	Daily detection rate of false-negative patients.
ω	The proportion of false-negative in the actual population.
l	The ratio of transmission of the false-negative to that of the confirmed.

Table 3 Predicted peak number of confirmed cases of each province with $\tau = 0.9$, $\varepsilon = 0.25$ and the mean values of recovery rate and infection rate.

Provinces (autonomous region, or special administrative region)	Peak value interval
Hubei	>60000
Zhejiang, Guangdong, Henan, Hunan, Chongqing	[2000, 4000]
Anhui, Sichuan, Jiangxi, Shandong, Jiangsu,	[1000, 2000]

Beijing	
Shanghai, Fujian, Guangxi, Hainan	[500, 1000]
Shaanxi, Hebei, Heilongjiang, Liaoning, Yunnan, Tianjin, Shanxi, Gansu	[200, 500]
Inner Mongolia, Ningxia, Xinjiang, Guizhou, Hong Kong, Jilin, Macau, Qinghai, Taiwan, Tibet	<200

Table 4 Simulation results of epidemic situation in Hubei

	Peak value time	Peak value
Worse case	2020. 03.18	76500
Average case	2020. 03.03	63800
Better case	2020. 02.27	59300

Table 5 Simulation results of epidemic situation in China except Hubei

	Peak value time	Peak value
Worse case	2020. 03.24	36800
Average case	2020. 03.09	31300
Better case	2020. 03.02	27700