Application of n-of-1 clinical trials in personalized nutrition research: a trial protocol for Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR)

Yunyi Tian, MSc[1]; Yue Ma, PhD[1,2]; Yuanqing Fu, PhD[1,2], Ju-Sheng Zheng, PhD[1,2*]

[1] Zhejiang Provincial Laboratory of Life Sciences and Biomedicine, Key Laboratory of Growth Regulation and Translation Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, China

[2] Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China

*Corresponding author:

Ju-Sheng Zheng, PhD

Zhejiang Provincial Laboratory of Life Sciences and Biomedicine, Key Laboratory of Growth Regulation and Translation Research of Zhejiang Province, School of Life Sciences, Westlake University,18 Shilongshan Rd, Cloud Town, Hangzhou, China.

Tel: +86 (0)57186915303

Email: zhengjusheng@westlake.edu.cn

Yunyi Tian, Yue Ma, Yuanqing Fu and Ju-Sheng Zheng report no conflicts of interest.

This study was funded by National Natural Science Foundation of China (81903316),

Zhejiang Ten-thousand Talents Program (2019R52039) and Westlake University

(101396021801).

Trial registration: Clinicaltrials.gov (NCT04125602)

Abbreviations used: AUC24, area under the curve; CGM, continuous glucose

monitoring; HF-LC, high-fat, low-carbohydrate; LF-HC, low-fat, high-carbohydrate;

MAGE, mean amplitude of glycemic excursions; PMG, postprandial maximum

glucose; WE-MACNUTR, Westlake N-of-1 Trials for Macronutrient Intake.

Abstract

Personalized dietary recommendations can help with more effective disease

prevention. This study aims to investigate the individual postprandial glucose

response to diets with diverse macronutrient proportions at both individual level and

population level and explore the potential of the novel single-patient (n-of-1) trial for

the personalization of diet. Secondary outcomes include individual phenotypic

response and the effects of dietary ingredients on the composition and structure of gut

microbiota. Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) is a

multiple crossover feeding trial consisting of three successive 12-day dietary

intervention pairs including a 6-day wash-out period before each 6-day isocaloric

dietary intervention (a 6-day high-fat, low-carbohydrate (HF-LC) diet and a 6-day

low-fat, high-carbohydrate (LF-HC) diet). The results will help provide personalized

dietary recommendation on macronutrients in terms of postprandial blood glucose

response. Well-designed n-of-1 trial is likely to become an effective method of

optimizing individual health and advancing health care. This trial is registered at

clinicaltrials.gov (NCT04125602).

Keywords: personalized nutrition, dietary intervention, n-of-1, single patient trial,

high-fat diet, low carbohydrate diet, postprandial blood glucose, gut microbiome

Introduction

Diet and nutrition are key to maintain human health. Previous studies have showed

much interest in the metabolic effects of different ratios of dietary fat to carbohydrate

intake(1-5). Some evidence suggests that a high-fat, low-carbohydrate (HF-LC) diet

can improve glycemic control by reducing glycated hemoglobin and fasting glucose,

while others support the beneficial effect of a low-fat, high-carbohydrate (LF-HC)

diet with particular focus on the quality of carbohydrate(1, 6-10). One important

interpretation for the inconsistent results is the individualized or personalized

response to the dietary macronutrient intake, which is also called "personalized

nutrition"(11).

Personalized nutrition focuses on an individual's potentially unique dietary needs

instead of assuming a "one-size-fits-all" approach where everyone is thought to

benefit from the same diet (12). The general aim of personalized nutrition is to

improve health using nutritional, genetic, phenotypic and other information of the

individuals to develop targeted nutritional advice, services or other products(13-15).

Although specific dietary recommendation has been made for pregnant women,

infants, children, adults or elderly, they are still subgroup recommendation, which is

far from the stage of "personalization" or "precision".

Application of 'n-of-1' clinical trial or 'single patient' study represents a new

direction in personalized nutrition research (**Figure 1**). It can capture intra-individual

variability in health behaviors over time, aiming to identify individual response to a

given intervention in a controlled trial, which provides great opportunity to answer the

personalization potential of different diets, nutrients or nutrition supplements(16-18).

The idea of n-of-1 has been applied in special education, psychotherapy, psychology

and pharmaceutical studies for decades to test the individual response to specific

drugs or treatments(19-23). Stunnenberg et al. reported the efficacy of mexiletine on

reducing muscle stiffness in patients with nondystrophic myotonia (NDM) using a

series of n-of-1 trials(24). However, there is no published 'n-of-1' study in nutrition

field so far. Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) is a

novel series of clinical trials that use the macronutrient intake as an exemplar for the

new progress in the field.

We will describe the trial protocol of the WE-MACNUTR: a series of n-of-1 feeding

clinical trials among healthy adults. The feeding trial is composed of two dietary

interventions, a HF-LC diet and a LF-HC diet. The primary objective is to investigate

the response of postprandial blood glucose to the dietary interventions and the

primary outcomes include postprandial maximum glucose (PMG), the area under the

curve (AUC24) of postprandial glucose from 0.00 to 24.00 and the mean amplitude of

glycemic excursions (MAGE) obtained by measuring the arithmetic mean of the

differences between consecutive peaks and nadirs. Secondary objectives of this study

include evaluating different phenotypic responses, such as circulating lipid profile

changes to a specific diet among individuals and evaluating the impact of different

dietary components on the composition and structure of gut microbiota.

Methods

Study Design

The present n-of-1 trial is a multiple crossover feeding trial conducted in a single

participant, comparing her/his response to different interventions and assessing the

variability in these responses(25). In the WE-MACNUTR study, a series of n-of-1

trials are employed simultaneously and a common regimen of interventions is applied

to all participants (**Figure 2**). Participants experience three successive 12-day

intervention pairs including a 6-day wash-out period between each intervention. The

diets are isocaloric and all provided by the researchers, and the primary distinguishing

feature is their fat and carbohydrate contents, including a HF-LC diet and a LF-HC

diet. Prior studies have shown that dietary pattern may have a rapid influence on

glycemic control. Parr and colleagues performed an intervention study and

successfully observed a significant effect of high-carbohydrate diet (compared with a

high-fat diet) on various blood glucose measurements after a 5-day intervention(26).

In addition, gut microbial community was also reported to change substantially in

response to the dietary change within four days(27). Therefore, we set a wash-out

period of six days between each intervention arm (HF-LC and LF-HC), after

balancing the outcomes of the study (postprandial glycemic response, gut microbiota

change), and feasibility of the present feeding trials. Each 12-day intervention pair

comprises 6 days of HF-LC (three meals daily) and 6 days of LF-HC (three meals

daily) diets in a random order. The order of the diets in each pair is determined using

a block randomization. Major investigators and laboratory personnel responsible for

the measurements are masked to group allocation. Meal providers are aware of group

and diet allocation, but they are not involved in the rest of the trial.

Participants are required to complete a basic questionnaire on a daily basis after

dinner to summarize their eating behaviors and multiple lifestyle factors including

physical activity, mood and sleep patterns throughout the day. Biological samples

including blood, saliva, urine and fecal samples will be collected every 6 days for

metabolomics profiling using untargeted platforms. Saliva will also be collected for

oral microbiota investigation (**Figure 3**). The fasting venous blood samples will be

collected every 6 days only during the first set of the study as to reduce the burden

and increase the compliance of the participants. The participants will be asked to wear

a continuous glucose monitoring (CGM, Freestyle Libre Pro System), which measures

interstitial glucose every 15 minutes. The same CGM system was used by a recent

personalized nutrition study (PREDICT 1) to monitor individual postprandial

glycemic responses(28). As most individuals do not resemble "the average" in the

context of precision nutrition, using data sets generated from digital wearable devices,

such as CGM, will potentially help inform individualized food choices(29). In the

present study, participants will be asked to wear the sensor two days before the start

of each intervention period, and remove it by the end of each 6-day intervention

period. CGM will not be used during the washout periods. In addition, participants are

asked to continue their regular daily activities and exercise throughout the study

period, and wear a wrist-based triaxial accelerometer (AX3, Axivity) on the

non-dominant wrist to monitor the physical activity and exercise intensity during the

intervention period.

The study was approved by the Westlake University Internal Ethical Review Board in

Hangzhou, China, and registered with ClinicalTrials.gov (identifier: NCT04125602).

Written informed consent are obtained from all study participants.

Dietary Intervention

A dietitian designs the diet for the intervention and the washout period based on the

Chinese Dietary Guidelines (2016) and Chinese Dietary Reference Intakes (2013) as

well as the participants' demographic information, eating habits and physical activity

levels(30, 31). Besides, factors such as local food availability, the cooking methods of

the Westlake University canteen kitchen, and the current recipes are important factors

in the meal planning. Participants are instructed to consume only the provided foods

or beverages.

Wash-out Diet (WD)

Prior to each dietary intervention, all participants are provided with the same

standardized diets for 6 days as the 'wash-out diet' to reduce potential sources of bias

and eliminate any carry-over effect of previous intervention. The wash-out diet

consists of 30% of total energy (%E) from fat, 15%E from protein and 55%E from

carbohydrate based on the acceptable macronutrient distribution range (AMDR)(32).

HF-LC Diet Intervention

Throughout the 6-day HF-LC intervention, participants are provided with a HF-LC

diet, including a three-day diet consisting of 60%E from fat, 15%E from protein and

25%E from carbohydrate while the other three-day diet consisting of 70%E from fat,

15%E from protein and 15%E from carbohydrate.

LF-HC Diet Intervention

Throughout the 6-day LF-HC intervention, participants are provided with a LF-HC

diet, including a 3-day diet consisting of 20%E from fat, 15%E from protein and

65%E from carbohydrate while the other 3-day diet consisting of 10%E from fat,

15%E from protein and 75%E from carbohydrate.

Participants

Participants are recruited among students and staffs from Westlake University,

Hangzhou, China. Inclusion criteria are: adults aged between 18 and 65 years;

participants are able to provide written informed consent, and have access to smart

phones or computers. Exclusion criteria include: long-term gastrointestinal diseases;

neurological conditions and cognitive impairment; other clinically diagnosed medical

conditions including type 2 diabetes, hypertension, cardiovascular diseases, liver,

kidney diseases and/or other systemic diseases; taking antibiotics within the last two

weeks; hospitalization or surgery planned within the next three months; pregnant or

lactating women; tobacco, alcohol, or illicit drug abuse; vegan or food allergies; with

no access to a smart phone or computer with an internet connection; enrolled in

concurrent intervention study; and non-Chinese speaking participants. This study

emphasizes more on assessing individual responses to the intervention of interest

rather than drawing a general conclusion at population level, and therefore, study

participants cover a broad age range and will not be balanced by sex.

Measures

The primary outcomes are (1) postprandial maximum glucose (PMG). PMG is the

peak value of CGM within 3 hours after the first bite of a meal or the maximum value

of CGM between two meals when the interval is less than 3 hours. (2) area under the

curve (AUC24). AUC24 refers to total area under the CGM curve from 0.00 to 24.00.

(3) mean amplitude of glycemic excursions (MAGE). MAGE is obtained by

measuring the arithmetic mean of the differences between consecutive peaks and

nadirs, provided that the differences are greater than 1 SD around the mean glucose

values. Secondary outcomes include different phenotypic responses, such as

circulating lipid profile and gut microbiome profile, to a specific diet among

individuals (**Table 1**).

Sample Size Calculation

In the present study, a total of 30 participants will be enrolled. Bayesian hierarchical

model meta-analysis will be applied to combine the results from each n-of-1 trial to

generate pooled effect estimate at population level. However, no formula-based

methodology exists for sample size calculation for such design at population level(33).

Referring to the method reported by Stunnenberg et al., we performed a

simulation-based statistical power calculation at population level with

modifications(24). In brief, a prior distribution for the mean intervention effect

(HF-LC vs. LF-HC) on the PMG was prespecified based on results of a previous RCT

reported by Parr et al(26). In their study, 1.6 mmol/L (6.2 mmol/L vs. 7.8 mmol/L)

difference in the PMG was established between participants receiving high-fat versus

high-carbohydrate meals. We drew a random realization and simulated an

individual-specific virtual mean intervention effect for each participant. At the next

level of simulation, longitudinal measurements for each participant (3 sets, 2

intervention periods per set and 18 observations per intervention period) were

simulated assuming a normal distribution of the measurements centered around the

individual-specific virtual mean effect size, with a common within-participant

residual variance. Thereafter, all the simulated data from these n-of-1 trials were used

to perform a multilevel Bayesian meta-analysis specifying a linear mixed model with

flat non-informative priors at population level.

A previous study found that the difference in PMG between healthy participants aged

25-45 years and those older than 45 years was 3mg/dL (0.167mmol/L)(34, 35).

Therefore, we considered this magnitude of difference to be clinically meaningful and

determined the posterior probability on both intervention effect of at least 0.167

mmol/L from the simulation-based Bayesian meta-analysis based on the simulation

data. In order to keep a balance between the power of detecting meaningful

differences and false positive rate, a relatively strict 90% posterior probability on an

intervention effect of 0.167 was treated as a clinically positive result at population

level, while a stricter 99% posterior probability on an intervention effect of not being

equal to zero was treated as a statistically positive result. We repeated the above

procedures for 1000 times (corresponding with data from 1000 aggregated N-of-1

trials) and the fraction of meta-analyses that returned a positive conclusion was

treated as measure for the power. Consequently, the power was estimated to be 100%

for detecting the intervention effect of at 0.167 mmol/L difference in PMG.

On the other hand, in order to determine the false positive rate, we performed another

set of simulation under the assumption of no intervention effect. For each virtual

participant, we simulated data that allowed the participant to have an

individual-specific virtual mean intervention effect which centered around the true

population-level mean effect size (zero) with a normal distribution. At the next level

of simulation, longitudinal measurements for each participant (3 sets, 2 intervention

periods per set and 18 observations per intervention period) were simulated assuming

a normal distribution of the measurements centered around the individual-specific

virtual mean effect size, with a common within-participant residual variance. All data

from these 30 virtual individual N-of-1 trials were used to perform the multilevel

Bayesian meta-analysis at population level. A posterior probability of more than 90%

on an intervention effect of 0.167 mmol/L and more than 99% on an intervention

effect of not being equal to zero were treated as clinically and statistically false

positive results, respectively. After repeating the procedures for 1000 times, the

fraction of meta-analyses that returned a clinically or statistically false positive

conclusion was determined as measure for the type I error rate at population level,

which was found to be 0.023 and 0.048, respectively. Thus, this simulation-based

sample calculation indicates that with 30 participants completing the trial (3 sets, 2

intervention periods per set and 18 observations per intervention period), we will have

satisfactory type I error rate and enough power to detect the prespecified intervention

effect.

Statistical Analysis Plan

Data Management

Even though all participants are students/staffs who routinely have meals in campus,

we anticipate that some participants will skip some meals that we provide at dining

room and eat other foods instead, which adds uncertainty to the effects on

postprandial blood glucose. On this occasion, we will ask the volunteers to record all

the other foods they have eaten not provided by the researchers. Other major

violations, such as failure to complete at least one set of intervention, will prevent

statistical analysis at individual level and lead to exclusion of the participants from

meta-analysis at population level.

Primary Analysis

The primary analysis of intervention effect is the comparison of the effect of HF-LC

diet with that of LF-HC diet on postprandial blood glucose level. At individual level,

we use individual intervention effects to guide dietary decisions for each participant.

On the other hand, we generate estimates of intervention effect at population level by

combing the n-of-1 results with meta-analysis.

Analysis of Baseline Data

Descriptive statistics with demographics and baseline characteristics will be presented

for each participant.

Analysis of Individual N-of-1 Trials

Statistical analysis will be performed separately for each n-of-1 trial to estimate the

intervention effect at individual level. Bayesian models will be applied to estimate the

intervention effects (**Figure 4**). Posterior probabilities of outcomes will be calculated

using an interface that incorporates open-source R software (3.6.1) and open-source

OpenBUGS (version 3.2.3). The results will be reported for primary variables (e.g.,

peak concentration of postprandial blood glucose and AUC of postprandial glucose

concentration over 24 hours), and participants will be provided with an estimate of

differences in the variables and the probabilities that the differences are induced by

the different interventions (HF-LC vs. LF-HC).

Meta-analysis of N-of-1 Trials

A Bayesian multi-level model will be used to combine results of the multiple n-of-1

trials(24, 36, 37). Participant will be treated as a random effect and a common

within-participant residual variance will be assumed. Non-informative priors will be

applied for all model parameters, with mean parameters using normal prior

distributions with very large standard deviations and variation parameters using

inverse gamma distributions with both shape and scale parameters equal to 0.01.

Using the interface that incorporates R software (3.6.1) and OpenBUGS (version

3.2.3), combining the data from the individual n-of-1 trials will obtain posterior

distributions for the mean intervention effect at the population level. Secondary and

exploratory outcomes will be analyzed similarly.

Discussion

To advance the field of personalized nutrition, n-of-1 clinical trial appears to be a

promising study design to advocate, although the real-world example is rare. We will

use the WE-MACNUTR trial as an exemplar to showcase the study design of n-of-1

trial as to test the individualized response to macronutrient intake among healthy

adults. The study, if successful, will provide insight into the feasibility of n-of-1

approach to personalize or tailor dietary intervention to individuals.

Previous study suggested that the magnitude of postprandial responses to mixed meals

depended largely on the total amount of fat and carbohydrates intakes(1, 5, 6). The

American Diabetes Association (ADA) recommends monitoring carbohydrate intake

to achieve better glycemic control in patients with type 2 diabetes, which is based on

studies showing reduced postprandial glucose and triglyceride responses in

individuals consuming HF-LC diets(38-40). However, previous systematic reviews

discussed the effects of HF-LC and LF-HC diets on metabolic risk factors and showed

inconsistent results(5, 7, 41). Hsu et al suggested LF-HC diet consisting of high fiber

contents showed beneficial effects on glucose and insulin sensitivity among both

Asian Americans and Caucasian Americans(42). Several studies have reported that

both HF-LC and LF-HC diets reduced HbA1c and fasting glucose in obese adults

with T2D, while HF-LC diet achieved better improvements in glycemic control(1, 2).

Kamada et al suggested that changes in HbA1c and fasting plasma glucose did not

differ significantly between HF-LC and LF-HC diets among Japanese diabetic

patients(43). The benefits and drawbacks of different dietary patterns on health are

under intensive study these days despite the lack of a standardized definition

regarding the macronutrient contents(44-46). Therefore, n-of-1 trial has a huge

potential to help explore the main effects of a specific dietary intervention, and

identify the factors that influence individual response to nutritional factors. In the

present study, it is expected that the trial will provide information on the response of

postprandial blood glucose level among individuals to different dietary interventions,

namely HF-LC and LF-HC diets, enabling better understanding of intra-individual

differences in absorption, distribution and metabolism of macronutrients.

Individual human beings are not only unique in the aspect of host genome, but also in

the aspect of the gut microbiome, which represents the combined influence of the diet

and lifestyles, as well as host genetics(47, 48). Both animal and human studies have

demonstrated that the composition of the gut microbiome can be rapidly affected by a

specific dietary component exposure within four days(27, 49). Furthermore,

integration of machine learning algorithms with gut microbiome features have shown

powerful potentials to predict one's response to different dietary patterns in term of

postprandial glucose response(50). Zeevi *et al.* monitored the postprandial glucose

response in a cohort of 800 participants in Israel in response to identical meals. A

multidimensional data including gut microbiome features, anthropometrics, blood

parameters, and physical activity were integrated into a machine learning algorithm

that was capable of predicting personalized postprandial glucose response with the gut

microbiota(51). These new progresses have stimulated more research on the

application and integration of gut microbiome into the personalized nutrition field.

Results from this study may provide further evidence suggesting that n-of-1 trial is

feasible in characterizing individual microbiome profile.

With the aggregated data from isocaloric meal but different carbohydrate to fat ratio,

our study will make deeper investigations of the underlying interactions between

specific food components and microbiota species. Therefore, future methodological

studies on developing and implementing effective evaluation of personalized dietary

intervention could assist individuals in promoting healthy gut microbiota profile and

preventing cardiometabolic diseases. Another strength of the present study with

n-of-1 method is its flexibility which enables the study design to be personalized to

individuals' interests and requirements, and its high level of evidence for making

clinical decisions for individuals alongside systematic reviews of RCTs.

Limitations

The n-of-1 study does have limitations. Participation in feeding trial such as

WE-MACNUTR requires time and effort so the trial cannot be conducted in an

ideally controlled setting. Compliance of the participants to the intervention over time

is challenging, as they are all required to eat the provided foods with no extra food

intake throughout the feeding trial. Any extra snack or beverage intake could affect

individual blood glucose level. Besides, slight changes in cooking method, food

groups or food ingredients under inevitable circumstances would also jeopardize the

final results.

Conclusions

In summary, WE-MACNUTR trial, as an exemplar of nutritional n-of-1 trial, will

address the call for new method to advance the field of personalized nutrition.

WE-MACNUTR will potentially help clarify the individual postprandial glucose

response to diets with diverse macronutrient proportions, and help design and

optimize macronutrient composition in long-term dietary intervention studies. The

results of WE-MACNUTR will also be helpful in terms of understanding the

individual response of gut microbiome to macronutrients.

Acknowledgements

We thank all the volunteers participating in the Westlake N-of-1 Trials for

Macronutrient Intake (WE-MACNUTR). The authors' contributions were as follows:

JSZ designed the study and was the principal investigator of WE-MACNUTR; YT,

YM, YF and JSZ drafted the manuscript; and all authors read and approved the final

manuscript.

References

1. Tay J, Luscombe-Marsh ND, Thompson CH, Noakes M, Buckley JD, Wittert GA, Yancy WS, Jr., Brinkworth GD. Comparison of low- and high-carbohydrate diets for type 2 diabetes management: a randomized trial. Am J Clin Nutr. 2015;102(4):780-90. Epub 2015/08/01. doi: 10.3945/ajcn.115.112581. PubMed PMID: 26224300.

2. Tay J, Thompson CH, Luscombe-Marsh ND, Wycherley TP, Noakes M, Buckley JD, Wittert GA, Yancy WS, Jr., Brinkworth GD. Effects of an energy-restricted low-carbohydrate, high unsaturated fat/low saturated fat diet versus a high-carbohydrate, low-fat diet in type 2 diabetes: A 2-year randomized clinical trial. Diabetes Obes Metab. 2018;20(4):858-71. Epub 2017/11/28. doi: 10.1111/dom.13164. PubMed PMID: 29178536.

3. Kdekian A, Alssema M, Van Der Beek EM, Greyling A, Vermeer MA, Mela DJ, Trautwein EA. Impact of isocaloric exchanges of carbohydrate for fat on postprandial glucose, insulin, triglycerides, and free fatty acid responses-a systematic review and meta-analysis. Eur J Clin Nutr. 2020;74(1):1-8. Epub 2019/11/27. doi: 10.1038/s41430-019-0534-6. PubMed PMID: 31767988.

4. Accurso A, Bernstein RK, Dahlqvist A, Draznin B, Feinman RD, Fine EJ, Gleed A, Jacobs DB, Larson G, Lustig RH, et al. Dietary carbohydrate restriction in type 2 diabetes mellitus and metabolic syndrome: time for a critical appraisal. Nutr Metab (Lond). 2008;5:9. Epub 2008/04/10. doi: 10.1186/1743-7075-5-9. PubMed PMID: 18397522; PubMed Central PMCID: PMCPMC2359752.

5. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Sato M, Sugawara A, Totsuka K, Shimano H, Ohashi Y, et al. Influence of fat and carbohydrate proportions on the metabolic profile in patients with type 2 diabetes: a meta-analysis. Diabetes Care. 2009;32(5):959-65. Epub 2009/05/02. doi: 10.2337/dc08-1716. PubMed PMID: 19407076; PubMed Central PMCID: PMCPMC2671123.

6. Noakes TD, Windt J. Evidence that supports the prescription of low-carbohydrate high-fat diets: a narrative review. Br J Sports Med. 2017;51(2):133-9. Epub 2017/01/06. doi: 10.1136/bjsports-2016-096491. PubMed PMID: 28053201.

7. Brand-Miller J, Hayne S, Petocz P, Colagiuri S. Low–glycemic index diets in the management of diabetes: a meta-analysis of randomized controlled trials. Diabetes Care. 2003;26(8):2261-7.

8. Saslow LR, Mason AE, Kim S, Goldman V, Ploutz-Snyder R, Bayandorian H, Daubenmier J, Hecht FM, Moskowitz JT. An Online Intervention Comparing a Very Low-Carbohydrate Ketogenic Diet and Lifestyle Recommendations Versus a Plate Method Diet in Overweight Individuals With Type 2 Diabetes: A Randomized Controlled Trial. J Med Internet Res. 2017;19(2):e36. Epub 2017/02/15. doi: 10.2196/jmir.5806. PubMed PMID: 28193599; PubMed Central PMCID: PMCPMC5329646.

9. Bhanpuri NH, Hallberg SJ, Williams PT, McKenzie AL, Ballard KD, Campbell WW, McCarter JP, Phinney SD, Volek JS. Cardiovascular disease risk factor responses to a type 2 diabetes care model including nutritional ketosis induced by sustained carbohydrate restriction at 1 year: an open label, non-randomized, controlled study. Cardiovasc Diabetol. 2018;17(1):56. Epub 2018/05/02. doi: 10.1186/s12933-018-0698-8. PubMed PMID: 29712560; PubMed Central PMCID: PMCPMC5928595.

10. Wylie-Rosett J, Aebersold K, Conlon B, Isasi CR, Ostrovsky NW. Health effects of low-carbohydrate diets: where should new research go? Curr Diab Rep. 2013;13(2):271-8. Epub 2012/12/26. doi: 10.1007/s11892-012-0357-5. PubMed PMID: 23266565; PubMed Central PMCID: PMCPMC3595318.

11. Braconi D, Cicaloni V, Spiga O, Santucci A. An Introduction to Personalized Nutrition. Trends in Personalized Nutrition. 2019. p. 3-32.

12. Ordovas JM, Ferguson LR, Tai ES, Mathers JC. Personalised nutrition and health. BMJ. 2018;361:k2173. Epub 2018/06/15. doi: 10.1136/bmj.k2173. PubMed PMID: 29898881; PubMed Central PMCID: PMCPMC6081996 interests and declare the following interests: JMO is part of Habit advisory board.

13. German JB, Zivkovic AM, Dallas DC, Smilowitz JT. Nutrigenomics and personalized diets: What will they mean for food? Annu Rev Food Sci Technol. 2011;2:97-123. Epub 2011/12/02. doi: 10.1146/annurev.food.102308.124147. PubMed PMID: 22129377; PubMed Central PMCID: PMCPMC4414021.

14. Valdes AM, Walter J, Segal E, Spector TD. Role of the gut microbiota in nutrition and health. BMJ. 2018;361:36-44.

15. Fito M, Melander O, Martinez JA, Toledo E, Carpene C, Corella D. Advances in Integrating Traditional and Omic Biomarkers When Analyzing the Effects of the Mediterranean Diet Intervention in Cardiovascular Prevention. Int J Mol Sci. 2016;17(9). Epub 2016/09/07. doi: 10.3390/ijms17091469. PubMed PMID: 27598147; PubMed Central PMCID: PMCPMC5037747.

16. Schork NJ, Goetz LH. Single-Subject Studies in Translational Nutrition Research. Annu Rev Nutr. 2017;37:395-422. Epub 2017/07/19. doi: 10.1146/annurev-nutr-071816-064717. PubMed PMID: 28715990; PubMed Central PMCID: PMCPMC6383767.

17. McDonald S, Quinn F, Vieira R, O'Brien N, White M, Johnston DW, Sniehotta FF. The state of the art and future opportunities for using longitudinal n-of-1 methods in health behaviour research: a systematic literature overview. Health Psychol Rev. 2017;11(4):307-23. Epub 2017/04/14. doi: 10.1080/17437199.2017.1316672. PubMed PMID: 28406349.

18. Duan N, Kravitz RL, Schmid CH. Single-patient (n-of-1) trials: a pragmatic clinical decision methodology for patient-centered comparative effectiveness research. J Clin Epidemiol. 2013;66(8 Suppl):S21-8. Epub 2013/07/17. doi: 10.1016/j.jclinepi.2013.04.006. PubMed PMID: 23849149; PubMed Central PMCID: PMCPMC3972259.

19. Kravitz RL, Schmid CH, Marois M, Wilsey B, Ward D, Hays RD, Duan N, Wang Y, MacDonald S, Jerant A, et al. Effect of Mobile Device-Supported Single-Patient Multi-crossover Trials on Treatment of Chronic Musculoskeletal Pain: A Randomized Clinical Trial. JAMA Intern Med. 2018;178(10):1368-77. Epub 2018/09/08. doi: 10.1001/jamainternmed.2018.3981. PubMed PMID: 30193253; PubMed Central PMCID: PMCPMC6233756.

20. Lillie EO, Patay B, Diamant J, Issell B, Topol EJ, Schork NJ. The n-of-1 clinical trial: the ultimate strategy for individualizing medicine? Per Med. 2011;8(2):161-73. Epub 2011/06/23. doi: 10.2217/pme.11.7. PubMed PMID: 21695041; PubMed Central PMCID: PMCPMC3118090.

21. Senior HE, Mitchell GK, Nikles J, Carmont SA, Schluter PJ, Currow DC, Vora R, Yelland MJ, Agar M, Good PD. Using aggregated single patient (N-of-1) trials to determine the effectiveness of psychostimulants to reduce fatigue in advanced cancer patients: a rationale and protocol. Bmc Palliative Care. 2013;12(1):17.

22. Moeller JD, Dattilo J, Rusch F. Applying Quality Indicators to Single-Case Research Designs Used in Special Education: A Systematic Review. Psychology in the Schools. 2015;52(2):139-53. doi: 10.1002/pits.21801.

23. Norcross JC, Wampold BE. What works for whom: Tailoring psychotherapy to the person. J Clin Psychol. 2011;67(2):127-32. Epub 2010/11/26. doi: 10.1002/jclp.20764. PubMed PMID: 21108312.

24. Stunnenberg BC, Raaphorst J, Groenewoud HM, Statland JM, Griggs RC, Woertman W, Stegeman DF, Timmermans J, Trivedi J, Matthews E, et al. Effect of Mexiletine on Muscle Stiffness in Patients With Nondystrophic Myotonia Evaluated Using Aggregated N-of-1 Trials. JAMA. 2018;320(22):2344-53. Epub 2018/12/12. doi: 10.1001/jama.2018.18020. PubMed PMID: 30535218; PubMed Central PMCID: PMCPMC6583079.

25. Zucker DR, Ruthazer R, Schmid CH, Feuer JM, Fischer PA, Kieval RI, Mogavero N, Rapoport RJ, Selker HP, Stotsky SA, et al. Lessons learned combining N-of-1 trials to assess fibromyalgia therapies. The Journal of Rheumatology. 2006;33(10):2069-77.

26. Parr EB, Devlin BL, Callahan MJ, Radford BE, Blankenship JM, Dunstan DW, Hawley JA. Effects of providing high-fat versus high-carbohydrate meals on daily and postprandial physical activity and glucose patterns: a randomised controlled trial. Nutrients. 2018;10(5):557.

27. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, et al. Diet rapidly and reproducibly alters the human gut microbiome. Nature. 2014;505(7484):559-63. Epub 2013/12/18. doi: 10.1038/nature12820. PubMed PMID: 24336217; PubMed Central PMCID: PMCPMC3957428.

28. Berry SE, Valdes AM, Drew DA, Asnicar F, Mazidi M, Wolf J, Capdevila J, Hadjigeorgiou G, Davies R, Al Khatib H, et al. Human postprandial responses to

food and potential for precision nutrition. Nat Med. 2020;26(6):964-73. Epub 2020/06/13. doi: 10.1038/s41591-020-0934-0. PubMed PMID: 32528151.

29. Spector TD, Gardner CD. Challenges and opportunities for better nutrition science—an essay by Tim Spector and Christopher Gardner. BMJ. 2020;369:m2470. doi: 10.1136/bmj.m2470.

30. The Chinese Nutrition Society. Chinese Dietary Guidelines. Beijing: People's Medical Publishing House; 2016.

31. The Chinese Nutrition Society. Chinese Dietary Reference Intakes. Beijing: China Light Industry Press; 2013.

32. Trumbo P, Schlicker S, Yates AA, Poos M. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids. Journal of the American Dietetic Association. 2002;102(11):1621-30. doi: 10.1016/s0002-8223(02)90346-9.

33. Kravitz RL and the DEcIDE Methods Center N-of-1 Guidance Panel. Design and Implementation of N-of-1 Trials: A User's Guide. Rockville, MD: Agency for Healthcare Research and Quality; 2014 Feb.

34. Chan CL, Pyle L, Newnes L, Nadeau KJ, Zeitler PS, Kelsey MM. Continuous glucose monitoring and its relationship to hemoglobin A1c and oral glucose tolerance testing in obese and prediabetic youth. The Journal of Clinical Endocrinology & Metabolism. 2015;100(3):902-10.

35. Juvenile Diabetes Research Foundation Continuous Glucose Monitoring Study Group. Variation of interstitial glucose measurements assessed by continuous glucose monitors in healthy, nondiabetic individuals. Diabetes Care. 2010;33(6):1297-9.

36. Zucker DR, Schmid CH, McIntosh MW, D'Agostino RB, Selker HP, Lau J. Combining single patient (N-of-1) trials to estimate population treatment effects and to evaluate individual patient responses to treatment. J Clin Epidemiol. 1997;50(4):401-10.

37. Quintana M, Viele K, Lewis RJ. Bayesian Analysis: Using Prior Information to Interpret the Results of Clinical Trials. JAMA. 2017;318(16):1605-6. Epub 2017/10/27. doi: 10.1001/jama.2017.15574. PubMed PMID: 29067406.

38. Jung CH, Choi KM. Impact of High-Carbohydrate Diet on Metabolic Parameters in Patients with Type 2 Diabetes. Nutrients. 2017;9(4). Epub 2017/03/25. doi: 10.3390/nu9040322. PubMed PMID: 28338608; PubMed Central PMCID: PMCPMC5409661.

39. American Diabetes Association. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. Diabetes Care. 2008;31 Suppl 1:S61-78. Epub 2008/01/10. doi: 10.2337/dc08-S061. PubMed PMID: 18165339.

40. Feinman RD, Pogozelski WK, Astrup A, Bernstein RK, Fine EJ, Westman EC, Accurso A, Frassetto L, Gower BA, McFarlane SI, et al. Dietary carbohydrate restriction as the first approach in diabetes management: critical review and

evidence base. Nutrition. 2015;31(1):1-13. Epub 2014/10/08. doi: 10.1016/j.nut.2014.06.011. PubMed PMID: 25287761.

41. Santos FL, Esteves SS, da Costa Pereira A, Yancy WS, Jr., Nunes JP. Systematic review and meta-analysis of clinical trials of the effects of low carbohydrate diets on cardiovascular risk factors. Obes Rev. 2012;13(11):1048-66. Epub 2012/08/22. doi: 10.1111/j.1467-789X.2012.01021.x. PubMed PMID: 22905670.

42. Hsu WC, Lau KH, Matsumoto M, Moghazy D, Keenan H, King GL. Improvement of insulin sensitivity by isoenergy high carbohydrate traditional Asian diet: a randomized controlled pilot feasibility study. PLoS One. 2014;9(9):e106851. Epub 2014/09/17. doi: 10.1371/journal.pone.0106851. PubMed PMID: 25226279; PubMed Central PMCID: PMCPMC4167335.

43. Kamada C, Yoshimura H, Okumura R, Takahashi K, Iimuro S, Ohashi Y, Araki A, Umegaki H, Sakurai T, Yoshimura Y, et al. Optimal energy distribution of carbohydrate intake for Japanese elderly patients with type 2 diabetes: the Japanese Elderly Intervention Trial. Geriatr Gerontol Int. 2012;12 Suppl 1:41-9. Epub 2012/04/28. doi: 10.1111/j.1447-0594.2011.00811.x. PubMed PMID: 22435939.

44. Naude CE, Schoonees A, Senekal M, Young T, Garner P, Volmink J. Low carbohydrate versus isoenergetic balanced diets for reducing weight and cardiovascular risk: a systematic review and meta-analysis. PLoS One. 2014;9(7):e100652. Epub 2014/07/10. doi: 10.1371/journal.pone.0100652. PubMed PMID: 25007189; PubMed Central PMCID: PMCPMC4090010.

45. Hockaday TD, Hockaday JM, Mann JI, Turner RC. Prospective comparison of modified fat-high-carbohydrate with standard low-carbohydrate dietary advice in the treatment of diabetes: one year follow-up study. Br J Nutr. 1978;39(2):357-62. Epub 1978/03/01. doi: 10.1079/bjn19780045. PubMed PMID: 629924.

46. Schulze MB, Martinez-Gonzalez MA, Fung TT, Lichtenstein AH, Forouhi NG. Food based dietary patterns and chronic disease prevention. BMJ. 2018;361:k2396. Epub 2018/06/15. doi: 10.1136/bmj.k2396. PubMed PMID: 29898951; PubMed Central PMCID: PMCPMC5996879.

47. Zhernakova A, Kurilshikov A, Bonder MJ, Tigchelaar EF, Schirmer M, Vatanen T, Mujagic Z, Vila AV, Falony G, Vieira-Silva S. Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. Science. 2016;352(6285):565-9.

48. Celis-Morales C, Livingstone KM, Marsaux CF, Macready AL, Fallaize R, O'Donovan CB, Woolhead C, Forster H, Walsh MC, Navas-Carretero S, et al. Effect of personalized nutrition on health-related behaviour change: evidence from the Food4Me European randomized controlled trial. Int J Epidemiol. 2017;46(2):578-88. Epub 2016/08/16. doi: 10.1093/ije/dyw186. PubMed PMID: 27524815.

49. Tremaroli V, Backhed F. Functional interactions between the gut microbiota and host metabolism. Nature. 2012;489(7415):242-9. Epub 2012/09/14. doi: 10.1038/nature11552. PubMed PMID: 22972297.

50. Cani PD. Human gut microbiome: hopes, threats and promises. Gut. 2018;67(9):1716-25. Epub 2018/06/24. doi: 10.1136/gutjnl-2018-316723. PubMed PMID: 29934437; PubMed Central PMCID: PMCPMC6109275.

51. Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, Weinberger A, Ben-Yacov O, Lador D, Avnit-Sagi T, Lotan-Pompan M. Personalized nutrition by prediction of glycemic responses. Cell. 2015;163(5):1079-94.

Table 1. Outcome and Assessment Points

Variable	Measure	T_pS1 [a]	T_pS2 [b]	T_pS3 [c]
		Assessment point		
Primary outcomes				
Continuous glucose level				
PMG		●	●	●
AUC24	FreeStyle Libre Flash Glucose Monitoring System	●	●	●
MAGE		●	●	●
Secondary outcomes				
Lipid metabolism	Blood samples			
Inflammation	Blood samples			
Oral microbiota	Saliva	√	√	√
Gut microbiota profiling	Fecal samples	√	√	√
Fecal metabolites	Fecal samples	√	√	√
Metabolomics profiling	Blood samples			
	Fecal samples	√	√	√
	Urine samples	√	√	√
Physiological characteristics				
Weight	Kubei height scale			√
Blood pressure	YUWELL YE660D upper arm sphygmomanometer			√

Abbreviations: AUC24, total area under the continues glucose monitoring curve from 0.00 to 24.00; MAGE, mean amplitude of glycemic excursions; PMG, postprandial maximum glucose.

[a] T_pS1 represents set 1.

[b] T_pS2 represents set 2.

[c] T_pS3 represents set 3.

● per day during intervention periods

√ before and after intervention periods

Figure 1. The Development of Personalized Nutrition. Personalized nutrition was born in the context that a conventional 'one-size-fits-all' approach usually fail to meet individual's need for the nutritional requirements. "N-of-1" clinical trial is a novel study design for the research of personalized nutrition comparing with the traditional designs such as observational study and randomized controlled trial. Integration of multi-omics data including nutrigenomics, proteomics, metabolomics, microbiome and other phenotypes is key for the development of personalized nutrition.

Figure 2. Flow Diagram of the Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) Trial. The flowchart summarizes the preparation phase and the first set of the trial. Set 2 and set 3 share the same trial design as set 1 except for no blood sample collection before each wash-out or intervention period. The sequence of two types of six-day dietary interventions in each set is randomized using block randomization as LF-HC and HF-LC diets in set 1; HF-LC and LF-HC diets in set 2 and HF-LC and LF-HC diets in set 3, respectively. CGM: continuous glucose monitoring; GI diseases: gastrointestinal diseases; HF-LC: high-fat, low-carbohydrate; LF-HC: low-fat, high-carbohydrate; WE-MACNUTR: Westlake N-of-1 Trials for Macronutrient Intake.

Figure 3. The Timeline of the Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) Trial. The timeline illustrates a preparation period for participants recruitment, a baseline data collection period, and three feeding trial periods. The first set of the trial consists of two wash-out periods (highlighted in green) and two randomized dietary intervention periods (highlighted in grey). In all three sets of the trial, both wash-out and intervention periods last for 6 days respectively.

Figure 4. Representation of the Hierarchical Bayesian Estimation for the Primary Outcomes at Both Individual and Population Level (combination of single patient studies). The observed repeated measurements of the postprandial peak glucose on a given patient are combined into a sample mean and sample variance. The model assumes that the patients' measurements follow a normal distribution centered about that patient's true mean effect (μ_i) with variance σ_1^2. At the population level, the various patients' true mean (μ_i) are assumed to follow a normal distribution centered about an overall population mean (μ_0) with between-patient variance τ^2. For the Bayesian specification, prior distributions are assigned for β, μ_0, σ_1^2, and τ^2. In the present study, these prior distributions are standard non-informative prior distributions. X means the independent variable: dietary pattern (high-fat and low-carbohydrate vs. low-fat and high-carbohydrate). Secondary and exploratory outcomes will be analyzed similarly. PMG: postprandial maximum glucose.