Application of n-of-1 clinical trials in personalized nutrition research: a trial

protocol for Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR)

Yunyi Tian, MSc[1]; Yue Ma, PhD[1,2]; Yuanqing Fu, PhD[1,2], Ju-Sheng Zheng, PhD[1,2]

[1] School of Life Sciences, Westlake University, Hangzhou, China

[2] Institute of Basic Medical Sciences, Westlake Institute for Advanced Study,

Hangzhou, China

Corresponding author:

Ju-Sheng Zheng, PhD

School of Life Sciences, Westlake University,18 Shilongshan Rd, Cloud Town,

Hangzhou, China.

Tel: +86 (0)57186915303

Email: zhengjusheng@westlake.edu.cn

Abstract

BACKGROUND Diet and nutrition play essential roles in human health.

Personalized dietary recommendations or nutritional advice tailored to each individual

can help with more effective disease prevention. N-of-1 trials can provide a pragmatic

clinical means of addressing individual postprandial blood glucose variation in

response to different food ingredients or nutrients.

OBJECTIVE To investigate the individual postprandial glucose response to diets

with diverse macronutrient proportions at both individual level and population level

and the potential of the novel single-patient (n-of-1) trial for the personalization of

diet.

DESIGN Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) is a

multiple crossover feeding trial. Individual response to different dietary patterns in

terms of postprandial glucose response is the primary outcome. Secondary outcomes

include individual phenotypic response and the effects of dietary ingredients on the

composition and structure of gut microbiota.

SETTING Participants experience three successive 12-day dietary intervention pairs

including a 6-day wash-out period before each isocaloric dietary intervention. Two

different type of diets (a 6-day high-fat, low-carbohydrate (HF-LC) diet and a 6-day

low-fat, high-carbohydrate (LF-HC) diet) are assigned to an individual in a

randomized sequence using block randomization with a fixed block size of two. This

feeding trial takes place in Hangzhou, China.

PARTICIPANTS Target enrolment is 30 healthy individuals aged between 18 and 65

years. Exclusion criteria are inability or unwillingness to approved informed consent;

other serious medical conditions; food allergy; and no access to a smart phone or

computer with an internet connection.

DISCUSSION This trial addresses the feasibility of n-of-1 approach for personalizing

dietary intervention to individuals. The results will help provide personalized dietary

recommendation on macronutrients in terms of postprandial blood glucose response.

Well-designed n-of-1 trial is likely to become an effective method of optimizing

individual health and advancing health care.

TRIAL REGISTRATION ClinicalTrials.gov identifier: NCT04125602

Introduction

Diet and nutrition are key to maintain human health. Previous studies have showed

much interest in the metabolic effects of different ratios of dietary fat to carbohydrate

intake[1-5]. Some evidence suggests that a high-fat, low-carbohydrate (HF-LC) diet can

improve glycemic control by reducing glycated hemoglobin and fasting glucose,

while others support the beneficial effect of a low-fat, high-carbohydrate (LF-HC)

diet with particular focus on the quality of carbohydrate[1,6-10]. One important

interpretation for the inconsistent results is the individualized or personalized

response to the dietary macronutrient intake, which is also called 'personalized

nutrition'[11].

Personalized nutrition was born in the context that a conventional 'one-size-fits-all'

approach usually fail to meet individual's need for the nutritional requirements[12]. The

general aim of personalized nutrition is to improve health using nutritional, genetic,

phenotypic and other information about individuals to develop targeted nutritional

advice, services or other products[13-15]. Although specific dietary recommendation has

been made for pregnant women, infants, children, adults or elderly, they are still

subgroup recommendation, which is far from the stage of 'personalization' or

'precision'.

Application of 'n-of-1' clinical trial or 'single subject' study represents a new

direction in personalized nutrition research (**Figure 1**). It can capture intra-individual

variability in health behaviors over time, aiming to identify individual response to a

given intervention in a controlled trial, which provides great opportunity to answer the

personalization potential of different diets, nutrients or nutrition supplements[16-18]. The

idea of n-of-1 has been applied in special education, psychotherapy, psychology and

pharmaceutical studies for decades to test the individual response to specific drugs or

treatments[19-23]. Stunnenberg et al. reported the efficacy of mexiletine on reducing

muscle stiffness in patients with nondystrophic myotonia (NDM) using a series of

n-of-1 trials[24]. However, there is no published 'n-of-1' study in nutrition field so far.

Westlake N-of-1 Trials for Macronutrient Intake (WE-MACNUTR) is a novel clinical

trial that uses the macronutrient intake as an exemplar for the new progress in the

field.

We will describe the trial protocol of the WE-MACNUTR: a n-of-1 feeding clinical

trial among healthy adults. The feeding trial is composed of two dietary interventions,

a HF-LC diet and a LF-HC diet. The primary objective is to investigate the response

of postprandial blood glucose to the dietary interventions and the primary outcomes

include postprandial maximum glucose (PMG), the area under the curve (AUC24) of

postprandial glucose from 0.00 to 24.00 and the mean amplitude of glycemic

excursions (MAGE) obtained by measuring the arithmetic mean of the differences

between consecutive peaks and nadirs. Secondary objectives of this study include

evaluating different phenotypic responses, such as circulating lipid profile changes to

a specific diet among individuals and evaluating the impact of different dietary

components on the composition and structure of gut microbiota.

Methods

Study Design

The n-of-1 trial is a multiple crossover feeding trial conducted in a single participant,

comparing her/his response to different interventions and assesses the variability in

these responses[25]. In the WE-MACNUTR study, a series of n-of-1 trials are employed

simultaneously and a common regimen of interventions is applied to all participants

(**Figure 2**). Participants experience three successive 12-day intervention pairs

including a 6-day wash-out period between each intervention. The diets are isocaloric

and all provided by the researchers, and the primary distinguishing feature is their fat

and carbohydrate contents, including a HF-LC diet and a LF-HC diet. Each 12-day

intervention pair comprises 6 days of HF-LC (three meals daily) and 6 days of LF-HC

(three meals daily) diets in a random order. The order of the diets in each pair is

determined using a block randomization. Major investigators and laboratory personnel

responsible for the measurements are masked to group allocation. Meal providers are

aware of group and diet allocation, but they are not involved in the rest of the trial.

Participants are required to complete a basic questionnaire on a daily basis after

dinner to summarize their eating behaviors throughout the day. Biological samples

including blood, saliva, urine and fecal samples of each participant are collected at

baseline and every 6 days during the first 12-day intervention pairs (**Figure 3**).

Starting from the second 12-day intervention pair, only saliva, urine and fecal samples

are collected every six days. Prior to each intervention period, each participant is

required to wear a subcutaneous glucose sensor (on the back of the upper arm) for

continuous glucose monitoring (CGM) up to 14 days and an AX3 band that can

capture physical activity data from the wrist of each participant.

The study was approved by the Westlake University Internal Ethical Review Board in

Hangzhou, China, and registered with ClinicalTrials.gov (identifier: NCT04125602).

Written informed consent are obtained from all study participants.

Dietary Intervention

A dietitian designs the diet for the intervention and the washout period based on the

Chinese Dietary Guidelines (2016) and Chinese Dietary Reference Intakes (2013) as

well as the participants' demographic information, eating habits and physical activity

levels. Besides, factors such as local food availability, the cooking methods of the

Westlake University canteen kitchen, and the current recipes are important factors in

the meal planning. Participants are instructed to consume only the provided foods or

beverages.

Wash-out Diet (WD)

Prior to each dietary intervention, participants are provided with a 6-day diet as the

'wash-out diet' to reduce potential sources of bias and eliminate any carry-over effect

of previous intervention. The wash-out diet consists of 30% of total energy (%E) for

fat, 15%E for protein and 55%E for carbohydrate based on the acceptable

macronutrient distribution range (AMDR)[26].

HF-LC Diet Intervention

Throughout the 6-day intervention, participants are provided with a HF-LC diet,

including a three-day diet consisting of 60%E for fat, 15%E for protein and 25%E for

carbohydrate while the other three-day diet consisting of 70%E for fat, 15%E for

protein and 15%E for carbohydrate.

LF-HC Diet Intervention

Throughout the 6-day intervention, participants are provided with a LF-HC diet,

including a 3-day diet consisting of 20%E for fat, 15%E for protein and 65%E for

carbohydrate while the other 3-day diet consisting of 10%E for fat, 15%E for protein

and 75%E for carbohydrate.

Participants

Participants are recruited among students and staffs from Westlake University,

Hangzhou, China. Inclusion criteria are as follows: healthy Chinese adults aged

between 18 and 65 years; and ability to complete the study. Exclusion criteria are

inability or unwillingness to provide informed consent; neurological conditions that

might affect the assessment of the study measurement; hospitalization or surgery

planned within 3 months; gastrointestinal diseases; other serious medical conditions,

such as liver, kidney, or systemic disease; pregnant or lactating women; tobacco,

alcohol, or illicit drug abuse; antibiotics uses within two weeks prior to the trial;

participants on a vegan diet; food allergy; no access to a smart phone or computer

with an internet connection; concurrent intervention study.

Measures

The primary outcomes are (1) postprandial maximum glucose (PMG). PMG is the

peak value of CGM within 3 hours after the first bite of a meal or the maximum value

of CGM between two meals when the interval is less than 3 hours. (2) area under the

curve (AUC24). AUC24 refers to total area under the CGM curve from 0.00 to 24.00.

(3) mean amplitude of glycemic excursions (MAGE). MAGE is obtained by

measuring the arithmetic mean of the differences between consecutive peaks and

nadirs. Secondary outcomes include different phenotypic responses, such as

circulating lipid profile and gut microbiome profile, to a specific diet among

individuals (**Table 1**).

Sample Size Calculation

For an individual n-of-1 trial, sample size calculations are not recommended[27]. In the

present study, a total of 30 participants will be enrolled, and Bayesian hierarchical

model meta-analysis will be applied to combine the results from each n-of-1 trial

generating estimates of intervention effect at population level. However, no

formula-based methodology exists for sample size or power calculations for such

designs[28].

Referring to the method reported by Stunnenberg et al., we performed a

simulation-based statistical power calculation with modifications[24]. In brief, a prior

distribution for the mean intervention effect (HF-LC vs. LF-HC) on the PMG was

prespecified based on results of a previous RCT reported by Parr et al[29]. In their study,

1.6 mmol/L (0.62 mmol/L vs. 0.78 mmol/L) difference on the PMG was established

among overweight/obese participants. From this prior distribution, we drew a random

realization and simulated data (3 sets, 2 intervention periods per set and 18

observations per intervention period) accordingly for the 30 n-of-1 trials using R

software. Thereafter, all data from these n-of-1 trials were used to perform a Bayesian

meta-analysis with flat non-informative priors at population level. The Bayesian

meta-analysis resulted in a (marginal) posterior probability for the mean intervention

effect.

Chan et al. reported a difference of 19 mg/dL (1.05 mmol/L) in daytime PMG

between nondiabetic youth and prediabetic counterparts, whereas another study found

this difference between youth and middle-aged ones was 10mg/dL (0.56 mmol/L)[30,31].

Therefore, we determined the posterior probability on both intervention effect of at

least 1.05mmol/L and 0.56 mmol/L from the simulation-based Bayesian

meta-analysis, and a ≥80% posterior probability was treated as positive result. We

repeated the above procedures for 1000 times (corresponding with data from 1000

aggregated N-of-1 trials) and the fraction of meta-analyses that returned a positive

conclusion (i.e. a >80% posterior probability on a prespecified treatment effect) was

treated as measure for the power. Consequently, the power was estimated to be 99.7 %

and 100% for detecting the intervention effect of at least 1.05 mmol/L and 0.56

mmol/L respectively. Given the present study focus on the postprandial glucose

response to different diet patterns among healthy youths, we set the meaningful

difference as 0.56 mmol/L, which may be still of significance for reminding healthy

subjects to shift the structure of diet to improve postprandial glucose when this

meaningful difference is detected. On the other hand, in order to determine the type I

error, we performed another set of simulation under the assumption of no intervention

effect. For each virtual participant, we simulated data that allowed the participant to

have a fixed mean value and expected variability during both intervention periods. All

data from these 30 virtual individual N-of-1 trials was used to perform the Bayesian

meta-analysis at population level and a strict ≥20% posterior probability on a

meaningful intervention effect (i.e., 0.56 mmol/L) was treated as false positive result.

After repeating the procedures for 1000 times, the fraction of meta-analyses that

returned a false positive conclusion was determined as measure for the type I error

rate, which was found to be 0.001.

Thus, this simulation-based sample calculation indicates that with 30 participants

completing the trial (3 sets, 2 intervention periods per set and 18 observations per

intervention period), it will have satisfactory type I error rate and enough power to

detect the prespecified meaningful intervention effect.

Statistical Analysis Plan

Data Management

Even though all participants are students/staffs who routinely have meals in campus,

we anticipate that some participants will skip some meals that we provide at dining

room and eat other foods instead, which adds uncertainty to the effects on

postprandial blood glucose. On this occasion, we will ask the volunteers to record all

the other foods they have eaten not provided by the researchers, and we will then

perform a sensitivity analysis based on the data. Other major violations, such as

failure to complete at least one set of intervention, will prevent statistical analysis at

individual level and lead to exclusion of the participants from meta-analysis at

population level.

Primary Analysis

The primary analysis of intervention effect is the comparison of the effect of HF-LC

diet on postprandial blood glucose level compared with LF-HC diet. At individual

level, we use individual intervention effects to guide dietary decisions for each

participant. On the other hand, we generate estimates of intervention effect at

population level by combing the n-of-1 results with meta-analysis.

Analysis of Baseline Data

Descriptive statistics with demographics and baseline characteristics will be presented

for each participant.

Analysis of Individual N-of-1 Trials

Statistical analysis will be performed separately for each n-of-1 trial to estimate

intervention effect at individual level. Bayesian models will be applied for estimating

the intervention effects (**Figure 4**). Posterior probabilities of outcomes will be

calculated using an interface that incorporates open-source R software (3.6.1) and

open-source WinBUGS (1.4.3). The results will be reported for primary variables

(e.g., AUC of postprandial glucose concentration over 24 hours and peak

concentration of postprandial blood glucose), and participants will be provided with

an estimate of differences in the variables and the probabilities that the differences are

induced by the different interventions (HF-LC vs. LF-HC).

Meta-analysis of N-of-1 Trials

To combine results of the multiple n-of-1 trials, a Bayesian multi-level model will be

used[24,32,33]. Participant will be treated as a random effect and a common

within-participant residual variance will be assumed. Non-informative priors will be

applied for all model parameters, with mean parameters using normal prior

distributions with very large standard deviations and variation parameters using

inverse gamma distributions with both shape and scale parameters equal to 0.01.

Using the interface that incorporates R software (3.6.1) and WinBUGS (1.4.3),

combining the data from the individual n-of-1 trials will obtain posterior distributions

for the mean intervention effect at the population level and for the between-participant

variation. Secondary and exploratory outcomes will be analyzed similarly.

Discussion

To advance the field of personalized nutrition, n-of-1 clinical trial appears to be a

promising study design to advocate, although the real-world example is rare. We will

use the WE-MACNUTR trial as an exemplar to showcase the study design of n-of-1

trial as to test the individualized response to macronutrient intake among healthy

adults. The study, if successful, will provide insight into the feasibility of n-of-1

approach to personalize or tailor dietary intervention to individuals.

Previous study suggested that the magnitude of postprandial responses to mixed meals

depended largely on the total amount of fat and carbohydrates intakes[1,5,6].The

American Diabetes Association (ADA) recommends monitoring carbohydrate intake

to achieve better glycemic control in patients with type 2 diabetes, which is based on

studies showing reduced postprandial glucose and triglyceride responses in

individuals consuming HF-LC diets[34-36]. However, previous systematic reviews

discussed the effects of HF-LC and LF-HC diets on metabolic risk factors and showed

inconsistent results[5,7,37]. Hsu et al suggested LF-HC diet consisting of high fiber

contents showed beneficial effects on glucose and insulin sensitivity among both

Asian Americans and Caucasian Americans[38]. Several studies have reported that both

HF-LC and LF-HC diets reduced HbA1c and fasting glucose in obese adults with

T2D, while HF-LC diet achieved better improvements in glycemic control[1,2]. Kamada

et al suggested that changes in HbA1c and fasting plasma glucose did not differ

significantly between HF-LC and LF-HC diets among Japanese diabetic patients[39].

The benefits and drawbacks of different dietary patterns on health are under intensive

study these days despite the lack of a standardized definition regarding the

macronutrient contents[40-42]. Therefore, n-of-1 trial has a huge potential to explore the

main effects of a specific dietary intervention, and identify the factors that influence

individual response to nutritional factors. In the present study, it is expected that the

trial will provide information on the response of postprandial blood glucose level

among individuals to different dietary interventions, namely HF-LC and LF-HC diets,

enabling better understanding of intra-individual differences in absorption,

distribution and metabolism of macronutrients.

Individual human beings are not only unique in the aspect of host genome, but also in

the aspect of the gut microbiome, which represents the combined influence of the diet

and lifestyles, as well as host genetics[43,44]. Both animal and human studies have

demonstrated that the composition of the gut microbiome can be rapidly affected by a

specific dietary component exposure within four days[45,46]. Furthermore, integration of

machine learning algorithms with gut microbiome features have shown powerful

potentials to predict one's response to different dietary patterns in term of

postprandial glucose response[47]. Zeevi *et al.* monitored the postprandial glucose

response in a cohort of 800 participants in Israel in response to identical meals. A

multidimensional data including gut microbiome features, anthropometrics, blood

parameters, and physical activity were integrated into a machine learning algorithm

that was capable of predicting personalized postprandial glucose response with the gut

microbiota[48]. These new progresses have stimulated more research on the application

and integration of gut microbiome into the personalized nutrition field. Results from

this study may provide further evidence suggesting that n-of-1 trial is feasible in

characterizing individual microbiome profile.

With the aggregated data from isocaloric meal but different carbohydrate to fat ratio,

our study makes deeper investigations of the underlying interactions between specific

food components and microbiota species possible. Therefore, future methodological

studies on developing and implementing effective evaluation of personalized dietary

intervention could assist individuals in promoting healthy gut microbiota profile and

preventing cardiometabolic diseases. Another strength of the present study with

n-of-1 method is its flexibility which enables the study design to be personalized to

individuals' interests and requirements, and its high level of evidence for making

clinical decisions for individuals alongside systematic reviews of RCTs.

Limitations

The n-of-1 study does have limitations. Participation in feeding trial such as

WE-MACNUTR requires time and effort so the trial cannot be conducted in an

ideally controlled setting. Participants will face consciousness, reliability, and

persistence over time as they are required to follow restricted dietary patterns with no

extra food intake throughout the trial. Any extra snack or beverage intake could affect

individual blood glucose level. Besides, slight changes in cooking method, food

groups or food ingredients under inevitable circumstances would also jeopardize the

final results.

Conclusions

In summary, WE-MACNUTR trial, as an exemplar of nutritional n-of-1 trial, will

address the call for new method to advance the field of personalized nutrition.

WE-MACNUTR will potentially help clarify the individual postprandial glucose

response to diets with diverse macronutrient proportions, and help design and

optimize macronutrient composition in long-term dietary intervention studies. The

results of WE-MACNUTR will also be helpful in terms of understanding the

individual response in gut microbiome to macronutrients.

Acknowledgements

We thank all the volunteers participating in the Westlake N-of-1 Trials for

Macronutrient Intake (WE-MACNUTR).

Funding: This study was funded by National Natural Science Foundation of China

(81903316) and Westlake University (101396021801)

Contributors: JSZ designed the study and was the principal investigator of

WE-MACNUTR; YT, YM, YF and JSZ drafted the manuscript and approved the final

version.

References

1. Tay J, Luscombe-Marsh ND, Thompson CH, et al. Comparison of low- and
high-carbohydrate diets for type 2 diabetes management: a randomized trial.
Am J Clin Nutr. 2015;102(4):780-790.
2. Tay J, Thompson CH, Luscombe-Marsh ND, et al. Effects of an
energy-restricted low-carbohydrate, high unsaturated fat/low saturated fat diet
versus a high-carbohydrate, low-fat diet in type 2 diabetes: A 2-year
randomized clinical trial. *Diabetes Obes Metab.* 2018;20(4):858-871.
3. Kdekian A, Alssema M, Van Der Beek EM, et al. Impact of isocaloric
exchanges of carbohydrate for fat on postprandial glucose, insulin,
triglycerides, and free fatty acid responses-a systematic review and
meta-analysis. *Eur J Clin Nutr.* 2020;74(1):1-8.
4. Accurso A, Bernstein RK, Dahlqvist A, et al. Dietary carbohydrate restriction
in type 2 diabetes mellitus and metabolic syndrome: time for a critical
appraisal. *Nutr Metab (Lond).* 2008;5:9.
5. Kodama S, Saito K, Tanaka S, et al. Influence of fat and carbohydrate
proportions on the metabolic profile in patients with type 2 diabetes: a
meta-analysis. *Diabetes Care.* 2009;32(5):959-965.
6. Noakes TD, Windt J. Evidence that supports the prescription of
low-carbohydrate high-fat diets: a narrative review. *Br J Sports Med.*
2017;51(2):133-139.
7. Brand-Miller J, Hayne S, Petocz P, Colagiuri S. Low–glycemic index diets in
the management of diabetes: a meta-analysis of randomized controlled trials.
Diabetes Care. 2003;26(8):2261-2267.
8. Saslow LR, Mason AE, Kim S, et al. An Online Intervention Comparing a
Very Low-Carbohydrate Ketogenic Diet and Lifestyle Recommendations
Versus a Plate Method Diet in Overweight Individuals With Type 2 Diabetes:
A Randomized Controlled Trial. *J Med Internet Res.* 2017;19(2):e36.
9. Bhanpuri NH, Hallberg SJ, Williams PT, et al. Cardiovascular disease risk
factor responses to a type 2 diabetes care model including nutritional ketosis
induced by sustained carbohydrate restriction at 1 year: an open label,
non-randomized, controlled study. *Cardiovasc Diabetol.* 2018;17(1):56.
10. Wylie-Rosett J, Aebersold K, Conlon B, Isasi CR, Ostrovsky NW. Health
effects of low-carbohydrate diets: where should new research go? *Curr Diab*
Rep. 2013;13(2):271-278.
11. Braconi D, Cicaloni V, Spiga O, Santucci A. An Introduction to Personalized
Nutrition. In: *Trends in Personalized Nutrition.*2019:3-32.
12. Ordovas JM, Ferguson LR, Tai ES, Mathers JC. Personalised nutrition and
health. *BMJ.* 2018;361:k2173.

13. German JB, Zivkovic AM, Dallas DC, Smilowitz JT. Nutrigenomics and personalized diets: What will they mean for food? *Annu Rev Food Sci Technol.* 2011;2:97-123.

14. Valdes AM, Walter J, Segal E, Spector TD. Role of the gut microbiota in nutrition and health. *BMJ.* 2018;361:36-44.

15. Fito M, Melander O, Martinez JA, Toledo E, Carpene C, Corella D. Advances in Integrating Traditional and Omic Biomarkers When Analyzing the Effects of the Mediterranean Diet Intervention in Cardiovascular Prevention. *Int J Mol Sci.* 2016;17(9).

16. Schork NJ, Goetz LH. Single-Subject Studies in Translational Nutrition Research. *Annu Rev Nutr.* 2017;37:395-422.

17. McDonald S, Quinn F, Vieira R, et al. The state of the art and future opportunities for using longitudinal n-of-1 methods in health behaviour research: a systematic literature overview. *Health Psychol Rev.* 2017;11(4):307-323.

18. Duan N, Kravitz RL, Schmid CH. Single-patient (n-of-1) trials: a pragmatic clinical decision methodology for patient-centered comparative effectiveness research. *J Clin Epidemiol.* 2013;66(8 Suppl):S21-28.

19. Kravitz RL, Schmid CH, Marois M, et al. Effect of Mobile Device-Supported Single-Patient Multi-crossover Trials on Treatment of Chronic Musculoskeletal Pain: A Randomized Clinical Trial. *JAMA Intern Med.* 2018;178(10):1368-1377.

20. Lillie EO, Patay B, Diamant J, Issell B, Topol EJ, Schork NJ. The n-of-1 clinical trial: the ultimate strategy for individualizing medicine? *Per Med.* 2011;8(2):161-173.

21. Senior HE, Mitchell GK, Nikles J, et al. Using aggregated single patient (N-of-1) trials to determine the effectiveness of psychostimulants to reduce fatigue in advanced cancer patients: a rationale and protocol. *BMC Palliat Care.* 2013;12(1):17-17.

22. Moeller JD, Dattilo J, Rusch F. Applying Quality Indicators to Single-Case Research Designs Used in Special Education: A Systematic Review. *Psychol Sch.* 2015;52(2):139-153.

23. Norcross JC, Wampold BE. What works for whom: Tailoring psychotherapy to the person. *J Clin Psychol.* 2011;67(2):127-132.

24. Stunnenberg BC, Raaphorst J, Groenewoud HM, et al. Effect of Mexiletine on Muscle Stiffness in Patients With Nondystrophic Myotonia Evaluated Using Aggregated N-of-1 Trials. *JAMA.* 2018;320(22):2344-2353.

25. Zucker DR, Ruthazer R, Schmid CH, et al. Lessons learned combining N-of-1 trials to assess fibromyalgia therapies. *The Journal of Rheumatology.* 2006;33(10):2069-2077.

26. Trumbo P, Schlicker S, Yates AA, Poos M. Dietary Reference Intakes for
Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino
Acids. *J Am Diet Assoc.* 2002;102(11):1621-1630.
27. *Design and Impkementation of N-of-1 Trials: A User's Guide.* Rockville, MD:
Agency for Healthcare Research and Quality; 2014.
28. Stunnenberg BC, Woertman W, Raaphorst J, et al. Combined N-of-1 trials to
investigate mexiletine in non-dystrophic myotonia using a Bayesian approach;
study rationale and protocol. *BMC Neurol.* 2015;15:43.
29. Parr EB, Devlin BL, Callahan MJ, et al. Effects of providing high-fat versus
high-carbohydrate meals on daily and postprandial physical activity and
glucose patterns: a randomised controlled trial. *Nutrients.* 2018;10(5):557.
30. Chan CL, Pyle L, Newnes L, Nadeau KJ, Zeitler PS, Kelsey MM. Continuous
glucose monitoring and its relationship to hemoglobin A1c and oral glucose
tolerance testing in obese and prediabetic youth. *The Journal of Clinical
Endocrinology & Metabolism.* 2015;100(3):902-910.
31. Group JDRFCGMS. Variation of interstitial glucose measurements assessed
by continuous glucose monitors in healthy, nondiabetic individuals. *Diabetes
Care.* 2010;33(6):1297-1299.
32. Zucker DR, Schmid CH, McIntosh MW, D'Agostino RB, Selker HP, Lau J.
Combining single patient (N-of-1) trials to estimate population treatment
effects and to evaluate individual patient responses to treatment. *J Clin
Epidemiol.* 1997;50(4):401-410.
33. Quintana M, Viele K, Lewis RJ. Bayesian Analysis: Using Prior Information
to Interpret the Results of Clinical Trials. *JAMA.* 2017;318(16):1605-1606.
34. Jung CH, Choi KM. Impact of High-Carbohydrate Diet on Metabolic
Parameters in Patients with Type 2 Diabetes. *Nutrients.* 2017;9(4).
35. American Diabetes A, Bantle JP, Wylie-Rosett J, et al. Nutrition
recommendations and interventions for diabetes: a position statement of the
American Diabetes Association. *Diabetes Care.* 2008;31 Suppl 1:S61-78.
36. Feinman RD, Pogozelski WK, Astrup A, et al. Dietary carbohydrate restriction
as the first approach in diabetes management: critical review and evidence
base. *Nutrition.* 2015;31(1):1-13.
37. Santos FL, Esteves SS, da Costa Pereira A, Yancy WS, Jr., Nunes JP.
Systematic review and meta-analysis of clinical trials of the effects of low
carbohydrate diets on cardiovascular risk factors. *Obes Rev.*
2012;13(11):1048-1066.
38. Hsu WC, Lau KH, Matsumoto M, Moghazy D, Keenan H, King GL.
Improvement of insulin sensitivity by isoenergy high carbohydrate traditional
Asian diet: a randomized controlled pilot feasibility study. *PLoS One.*
2014;9(9):e106851.
39. Kamada C, Yoshimura H, Okumura R, et al. Optimal energy distribution of
carbohydrate intake for Japanese elderly patients with type 2 diabetes: the

Japanese Elderly Intervention Trial. *Geriatr Gerontol Int.* 2012;12 Suppl
1:41-49.

40. Naude CE, Schoonees A, Senekal M, Young T, Garner P, Volmink J. Low
carbohydrate versus isoenergetic balanced diets for reducing weight and
cardiovascular risk: a systematic review and meta-analysis. *PLoS One.*
2014;9(7):e100652.

41. Hockaday TD, Hockaday JM, Mann JI, Turner RC. Prospective comparison of
modified fat-high-carbohydrate with standard low-carbohydrate dietary advice
in the treatment of diabetes: one year follow-up study. *Br J Nutr.*
1978;39(2):357-362.

42. Schulze MB, Martinez-Gonzalez MA, Fung TT, Lichtenstein AH, Forouhi NG.
Food based dietary patterns and chronic disease prevention. *BMJ.*
2018;361:k2396.

43. Zhernakova A, Kurilshikov A, Bonder MJ, et al. Population-based
metagenomics analysis reveals markers for gut microbiome composition and
diversity. *Science.* 2016;352(6285):565-569.

44. Celis-Morales C, Livingstone KM, Marsaux CF, et al. Effect of personalized
nutrition on health-related behaviour change: evidence from the Food4Me
European randomized controlled trial. *Int J Epidemiol.* 2017;46(2):578-588.

45. David LA, Maurice CF, Carmody RN, et al. Diet rapidly and reproducibly
alters the human gut microbiome. *Nature.* 2014;505(7484):559-563.

46. Tremaroli V, Backhed F. Functional interactions between the gut microbiota
and host metabolism. *Nature.* 2012;489(7415):242-249.

47. Cani PD. Human gut microbiome: hopes, threats and promises. *Gut.*
2018;67(9):1716-1725.

48. Zeevi D, Korem T, Zmora N, et al. Personalized nutrition by prediction of
glycemic responses. *Cell.* 2015;163(5):1079-1094.

Table 1. Outcome and Assessment Points

Variable	Measure	Assessment point			
		T_p0^a	T_pS1^b	T_pS2^c	T_pS3^d
Primary outcomes					
Continuous glucose level					
PMG			●	●	●
AUC24	FreeStyle Libre Flash Glucose Monitoring System		●	●	●
MAGE			●	●	●
Secondary outcomes					
Lipid metabolism	Blood samples	√			
Inflammation	Blood samples	√			
Gut microbiota profiling	Fecal samples	√	√	√	√
Fecal metabolites	Fecal samples	√	√	√	√
Metabolomics profiling	Blood samples	√			
	Fecal samples	√	√	√	√
	Urine samples	√	√	√	√
Physiological characteristics					
Weight	Kubei height scale				√
Blood pressure	YUWELL YE660D upper arm sphygmomanometer				√

Abbreviations: PMG, postprandial maximum glucose; AUC24, total area under the CGM curve from

0.00 to 24.00; MAGE, mean amplitude of glycemic excursions.

[a]T_p0 represents baseline, prior to the first wash-out period of set 1.

[b]T_pS1 represents set 1.

[c]T_pS2 represents set 2.

[d]T_pS3 represents set 3.

● per day during intervention periods

√ before and after intervention periods

Figure 1. The Development of Personalized Nutrition. Personalized nutrition was born in
the context that a conventional 'one-size-fits-all' approach usually fail to meet individual's need for the
nutritional requirements. 'n-of-1' clinical trial is a novel study design for the research of personalized
nutrition comparing with the traditional designs such as observational study and randomized controlled
trial. Integration of multi-omics data including nutrigenomics, proteomics, metabolomics, microbiome
and other phenotypes is key for the development of personalized nutrition.

530 **Figure 2. Flow Diagram of the Westlake N-of-1 Trials for Macronutrient Intake**

531 **(WE-MACNUTR) Trial.** The flowchart summarizes the preparation phase and the first set of the

532 trial. Set 2 and set 3 share the same trial design as set 1 except for no blood sample collection before

533 each wash-out or intervention period. The sequence of two types of six-day dietary interventions in

534 each set is randomized using block randomization as LF-HC and HF-LC diets in set 1; HF-LC and

535 LF-HC diets in set 2 and HF-LC and LF-HC diets in set 3, respectively. HF-LC: high-fat,

536 low-carbohydrate; LF-HC: low-fat, high-carbohydrate.

537

538

539

540

Figure 3. The Timeline of the Westlake N-of-1 Trials for Macronutrient Intake
(WE-MACNUTR) Trial. The timeline illustrates a preparation period for participants
recruitment, a baseline data collection period, and three feeding trial periods. The first set of the trial
consists of two wash-out periods (highlighted in green) and two randomized dietary intervention
periods (highlighted in grey). In all three sets of the trial, both wash-out and intervention periods last
for 6 days respectively.

548
549

Figure 4. Representation of the Hierarchical Bayesian Estimation for the Primary Outcomes at Both Individual and Population Level (combination of single patient studies). The observed repeated measurements of the postprandial peak glucose on a given patient are combined into a sample mean and sample variance. The model assumes that the patients' measurements follow a normal distribution centered about that patient's true mean effect (μ_i) with variance σ_1^2. At the population level, the various patients' true mean (μ_i) are assumed to follow a normal distribution centered about an overall population mean (μ_0) with between-patient variance τ^2. For the Bayesian specification, prior distributions are assigned for β, μ_0, σ_1^2, and τ^2. In the present study, these prior distributions are standard non-informative prior distributions. X means the independent variable: dietary pattern (high-fat and low-carbohydrate vs. low-fat and high-carbohydrate); PMG: the peak value of CGM within 3 hours after the meal or the peak value of CGM between two meals when the interval is less than 3 hours. Secondary and exploratory outcomes will be analyzed similarly.