Estimation of the final size of the coronavirus epidemic by the logistic model

Milan Batista

University of Ljubljana, Slovenia

milan.batista@fpp.uni-lj.si

(Feb 2020)

Abstract

In the note, the logistic growth regression model is used for the estimation of the final size and its peak time of the coronavirus epidemic.

1 Introduction

One of the common questions in an epidemic is its final size and its peak time. For answering this question various models are used: analytical [1-4], stochastic [5], phenomenological [6, 7]

In this note, we will try to estimate a final epidemic size and its peak by the phenomenological model, i.e., with the logistic growth regression model [7, 8]. The estimation is made in two steps: first, data are sequentially fitted by logistic model and second, the estimated finale sizes are fitted by the Weibull function.

2 Logistic grow model

The logistic growth model originates in population dynamics [9]. The underlying assumption of the model is that the rate of change in the number of new cases per capita linearly decreases with the number of cases. So, if C is the number of cases, and t is the time, then the models read

$$\frac{1}{C}\frac{dC}{dt} = r\left(1 - \frac{C}{K}\right),\tag{1}$$

where r is infection rate, and K the final epidemic size. If $C(0) = C_0$ is the initial number of cases then the solution of (1) is

$$C = \frac{K}{1 + A\exp\left(-rt\right)} \tag{2}$$

where $A = \dfrac{K - C_0}{C_0}$ The growth rate $\dfrac{dC}{dt}$ reaches its maximum when $\dfrac{d^2C}{dt^2} = 0$. From this condition, we obtain that the growth rate peak occurs in time time

$$t_p = \frac{\ln A}{r} \tag{3}$$

At this time the number of cases and the growth rate are

$$C_p = \frac{K}{2}, \qquad \left(\frac{dC}{dt}\right)_p = \frac{rK}{4} \tag{4}$$

3 Logistic regression model

By the logistic regression model, we estimate the parameters of the logistic model. Using (2) we can write the regression model in the form

$$C = \frac{b_1}{1 + b_2\exp\left(-b_3t\right)} \tag{5}$$

where unknown parameters b_1, b_2, b_3 which are to be estimated from the data. In the logistic model, the parameters are constants while the regression model parameters depend on available data. So if $C_1, C_2, \ldots, C_n$ are the cases in times $t_1, t_2, \ldots, t_n$ then $b_i^{(n)} = b_i^{(n)}\left(C_1, C_2, \ldots, C_n\right)$ ($i = 1, 2, 3$), or in other words, parameters are time-dependent

$$b_i = b_i\left(t\right) \quad (\, i = 1, 2, 3\,) \tag{6}$$

Now, if we assume that the epidemy has a final size then $b_i = b_i\left(t\right)$ must converge to some finite values. By fitting the sequence $b_i^{(k)}$, $k = n_0, \ldots, n$ by Weibull function [5]

$$b_1 = K\left[1 - \exp\left(-k\left(t - t_0\right)^p\right)\right] \tag{7}$$

we obtain a final estimation of K, i.e., the epidemy size. k, p and t_0 are regression parameters.

4 Data

For regression analysis, we use two data sets.. For the period from 16 Jan to 21 Jan 2020 we use the data from the published graph[1] and for the period from 22 Jan 2020 to 16 Feb 2020 we use data from *worldmeters*[2]. The discrepancy between data set for a period of 22 Jan to 26 Jan is between 1% to 6%.

Results

For practical calculation, we use MATLAB's functions `lsqcurvefit` and `fitnlm` (see Apendix).

Based on available data (til 16 Feb 2020), the predicted final size of epidemy is about 100000 cases (see Table 1, see Fig 1, 2) and its peak was on 11 Feb 2020. With the new data, these estimates will change reliably within days.

Figure 1. Predicted evaluation of the coronavirus epidemic (data till 16.Feb 2020). Regression data are in Table 4.

---

[1] https://i.redd.it/f4nukz4ou9d41.png

[2] https://www.worldometers.info/coronavirus/

Table 1. Estimate logistic model parameters for (data till 16.Feb 2020).

	Estimate	SE	tStat	pValue
b1	**98028**	9403	10.425	3.8086e-11
b2	0.1943	0.01416	13.722	5.9161e-14
b3	137.73	25.611	5.3776	9.8884e-06

```
Number of observations: 31, Error degrees of freedom: 28
Root Mean Squared Error: 2.3e+03
R-Squared: 0.99,  Adjusted R-Squared 0.99
F-statistic vs. zero model: 1.82e+03, p-value = 3.55e-32
```

Figure 2. Predicted evaluation of the coronavirus epidemic rate (data till 16.Feb 2020)

The trend of model parameters is given in Table 2 while the graph of evaluation of the final size of the epidemy is shown as a bar graph in Fig 3. The parameters of the Weibull model (8) of evaluation of the size of epidemy, for data until 12. Feb 2020, are given in Table 3. We see that the final size of epidemy would be about 58000 infections and the epidemy would reach its peak on 4 Feb 2020. However, a new way of data collection distorts this prediction. By using the Weibull model we obtain the final size of the epidemy of about 1.2 million infections (see Table 4). We stress that this prediction should be taken with care because there is not yet enough data to use the Weibull model for the final prediction of epidemy size. This also applies to the final prediction of the epidemy peak time (Fig 5) where Weibull regression predicts paek time at day 218, i.e.at 22 Aug 2020.

Table 2. Results of logistic regression (see Eqs (2), (3), (4))

day	K (cases)	r (1/day)	A (day)	t_peak	dC_peak (cases/day)	date_peak
16	18165	0.45	452.557	13	2044	30.jan.20
17	21823	0.414	394.182	14	2260	31.jan.20
18	26068	0.383	350.317	15	2495	1.feb.20
19	31257	0.354	316.693	16	2766	2.feb.20
20	38821	0.324	290.899	17	3146	3.feb.20
21	44478	0.308	279.638	18	3420	4.feb.20
22	46179	0.303	275.404	18	3496	4.feb.20
23	48078	0.297	268.19	18	3570	4.feb.20
24	48596	0.295	265.423	18	3588	4.feb.20
25	49822	0.291	256.796	19	3621	5.feb.20
26	50903	0.286	247.804	19	3643	5.feb.20
27	51372	0.284	243.218	19	3650	5.feb.20
28	71332	0.226	152.945	22	4027	8.feb.20
29	93476	0.198	137.994	24	4627	10.feb.20
30	100472	0.192	137-028	25	4827	11.feb.20
31	98025	0.194	137.733	25	4751	11.feb.20

Figure 3. Evaluation of estimated final size of coronavirus epidemic (bars). Weibull regression data (red line) are given in Table 4.

Table 3. The estimated final size of epidemy from data till 11.Feb. 2020

	Estimate	SE	tStat	pValue
b1	**58121**	3746.8	15.512	8.424e-08
b2	0.18721	0.039055	4.7935	0.00098261
b3	12.085	7.1289	1.6952	0.12428

Number of observations: 12, Error degrees of freedom: 9
Root Mean Squared Error: 2.11e+03
R-Squared: 0.975, Adjusted R-Squared 0.97
F-statistic vs. zero model: 1.53e+03, p-value = 1.7e-12

Table 4. The estimated final size of the epidemy by Weibull model (7) from data till 16.Feb. 2020 (see Fig 3)

	Estimate	SE	tStat	pValue
b1	**1.1837e+06**	1.6115e-05	7.345e+10	1.6601e-145
b2	3.5997e-05	0.00028489	0.12635	0.90125
b3	1.0919	17.731	0.061583	0.95177
b4	2.3085	1.9105	1.2083	0.24693

Number of observations: 17, Error degrees of freedom: 14
Root Mean Squared Error: 8.14e+03
R-Squared: 0.917, Adjusted R-Squared 0.906
F-statistic vs. zero model: 268, p-value = 1.34e-12

Figure 5. Evaluation of estimated peak time of coronavirus epidemic (bars). Weibull regression data (red line) (Table 5).

Table 5. The estimated peak time size of the epidemy by Weibull model (7) from data till 16.Feb. 2020 (see Fig 5)

	Estimate	SE	tStat	pValue
b1	218.32	0.53019	411.79	5.4747e-30
b2	0.00047621	0.0040803	0.11671	0.90875
b3	-13.715	44.413	-0.30881	0.76201
b4	1.4657	1.8373	0.79771	0.43836

```
Number of observations: 17, Error degrees of freedom: 14
Root Mean Squared Error: 1.14
R-Squared: 0.923,  Adjusted R-Squared 0.912
F-statistic vs. zero model: 1.61e+03, p-value = 5.38e-18

Peak date 22-Aug-2020
```

5 Conclusion

On the base of available data, one can not yet predict the final size of coronavirus epidemy using the logistic model. If however, in the next three or four days the number of cases will be about say 72000 then we can predict that the final number of cases will be about 100 000, and that peak will be at about 32 days, i.e., at 18. Feb 2020.

Appendix. MATLAB program.

```
%FITVIRUS Estimation of coronavirus epidemy size by logistic model

close all

warning('off')

% obtain data
[sampleC,date0] = getData();

% form time
samplaTime = 0:1:length(sampleC)-1;
time = 0:1:2*length(samplaTime);

% form date
samplaDate = date0+samplaTime;
date = date0+time;

% initial guess
b0 = [max(sampleC) 1 max(sampleC)]';

% parameters evaluation
```

```matlab
    bb = NaN(3,length(sampleC));
    tpeak = NaN(length(sampleC),1);
    dCpeak = NaN(length(sampleC),1);

    %initial data set
    n0 = 15;

    % loop over periods
    opts = optimoptions('lsqcurvefit','Display','off');
    for n = n0:length(sampleC)
        [b,resnorm,~,exitflag,output] = lsqcurvefit(@fun,b0,...
            samplaTime(1:n),sampleC(1:n),[],[],opts);
        bb(:,n) = b;
        % peak time and growth rate
        tpeak(n) = log(b(3))/b(2);
        dCpeak(n) = dfun(b,tpeak(n));
    end

    % final fit
    fprintf('Regression parameters for complet data set\n')
    mdl = fitnlm(samplaTime(1:n),sampleC(1:n),@fun,b0)

    % print parameters evaluation
    fprintf('\nEvaluation of model parameters\n')
    fprintf('%4s %12s %10s %10s %10s %12s
%12s\n','day','K','r','A','t_peak','dC_peak','date_peak')
    fprintf('%4s %12s %10s %10s %10s %12s\n','
','(cases)','(1/day)','','(day)','(cases/day)')
    for n = n0:length(sampleC)
        fprintf('%4d %12d %10.3f %10.3f %10d %12d
%12s\n',n,fix(bb(1,n)),bb(2:3,n),...
            fix(tpeak(n)),fix(dCpeak(n)),datestr(ceil(tpeak(n))+date0))
    end

    % fit estimated final sizes
    fprintf('\nWeibull regression of predicted final sizes\n')
    % Initial guess
    f0 = [sampleC(end) 0.13 12 1]'
    n = length(sampleC);
    mdl2 = fitnlm(samplaTime(n0:n),bb(1,n0:n),@fun1,f0)
    f = mdl2.Coefficients.Estimate;

    % fit estimated peak time
    fprintf('\nWeibull regression of predicted peak time\n')
    % Initial guess
    ff0 = [tpeak(end) 0.13 12 1]'
    n = length(sampleC);
    mdl2 = fitnlm(samplaTime(n0:n),tpeak(n0:n),@fun1,ff0)
    ff = mdl2.Coefficients.Estimate;
    fprintf('Peak date %s\n',datestr(ceil(ff(1))+date0))

    warning('on')

    % plot evaluation of peak time
    figure
    hold on
    bar(samplaDate(n0:n),tpeak(n0:n)); %,'LineWidth',2)
    datetick('x',20,'keeplimits')
    plot(date(n0:n),fun1(ff,time(n0:n)),'r','LineWidth',2)
    datetick('x',20,'keeplimits')
    ylabel('Day')
```

```matlab
xlabel('Date')
title('Estimated peak time')
grid on
hold off

% plot evaluation of final size
figure
hold on
bar(samplaDate(n0:n),bb(1,n0:n)/1000); %,'LineWidth',2)
datetick('x',20,'keeplimits')
plot(date(n0:n),fun1(f,time(n0:n))/1000,'r','LineWidth',2)
datetick('x',20,'keeplimits')
ylabel('Infected (1000)')
xlabel('Date')
title('Estimated final size of epidemic')
grid on
hold off

% plot final prediction of epedimy evaluation
figure
hold on
plot(date,fun(b,time)/1000,'k','LineWidth',2)
scatter(samplaDate,sampleC/1000,50,'k','filled')
h = scatter(samplaDate,sampleC/1000,30,'w','filled');
h.Annotation.LegendInformation.IconDisplayStyle = 'off';
% add tick marks
datetick('x',20,'keeplimits')
% add axes labels
ylabel('Infected (1000)')
xlabel('Date')
% add legend
legend('Predicted','Actual','Location','best')
%add title
title('Coronavirus epidemic')
% add grid
grid on
hold off

% plot growth rate of epidemic
figure
hold on
plot(date,dfun(b,time),'LineWidth',2)
ylabel('Infected/day')
yyaxis right
plot(date,dfun(b,time)./fun(b,time),'LineWidth',2)
ylabel('Infected/per capita/day')
datetick('x',20,'keeplimits')
title('Coronavirus epidemic rate')
grid on
hold off

function y = fun(b,t)
% Logistic model
    y = b(1)./(1 + b(3)*exp(-b(2)*t));
end

function y = dfun(b,t)
% Grow rate
    y = b(1)*b(2)*b(3)*exp(-b(2)*t)./(1 + b(3)*exp(-b(2)*t)).^2;
end
```

```matlab
function y = fun1(b,t)
% Weibull model. Note b(3) !!!
    y = b(1)*(1 - exp(-b(2)*(t-b(3)).^b(4)));
end

function [C,date0] = getData()
%GETDATA Coronavirus data
%   data from 16 Jan to 21 Jan https://i.redd.it/f4nukz4ou9d41.png
%   data from from 22 Jan 2020 to 15 Feb 2020 are from
%   https://www.worldometers.info/coronavirus/
date0=datenum('2020/01/16'); % start date
C = [
11948
14551
17389
20628
24553
28276
31439
34876
37552
40553
43099
44919
60326
64437
67105
69270 %<------------- add new data here
]';
end
```

References

[1] D.A. Barbarossa MV, Kiss G, Nakata Y, Röst G, Vizi Z Transmission Dynamics and Final Epidemic Size of Ebola Virus Disease Outbreaks with Varying Interventions, PLoS ONE 10(7) (2015).

[2] F. Brauer, The Final Size of a Serious Epidemic, Bull Math Biol 81(3) (2019) 869-877.

[3] J.M.A. Danby, Computing applications to differential equations modelling in the physical and social sciences, Reston Publishing Company, Reston, Va., 1985.

[4] J.T. Wu, K. Leung, G.M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study, The Lancet (2020).

[5] J.C. Miller, A note on the derivation of epidemic final sizes, Bull Math Biol 74(9) (2012) 2125-2141.

[6] K.E. Fisman D, Tuite A. , Early Epidemic Dynamics of the West African 2014 Ebola Outbreak: Estimates Derived with a Simple Two-Parameter Model., PLOS Currents Outbreaks. (2014).

[7] B. Pell, Y. Kuang, C. Viboud, G. Chowell, Using phenomenological models for forecasting the 2015 Ebola challenge, Epidemics 22 (2018) 62-70.

[8] S.L. Chowell G, Viboud C, Kuang Y., West Africa Approaching a Catastrophic Phase or is the 2014 Ebola Epidemic Slowing Down? Different Models Yield Different Answers for Liberia. , PLOS Currents Outbreaks. (2014).

[9] R. Haberman, Mathematical models mechanical vibrations, population dynamics, and traffic flow an introduction to applied mathematics, Unabridged republication ed., SIAM, Philadelphia, 1998.