Estimation of the final size of the coronavirus epidemic by the logistic model

Milan Batista

University of Ljubljana, Slovenia

milan.batista@fpp.uni-lj.si

(Feb 2020)

Abstract

In the note, the logistic growth regression model is used for the estimation of the final size and its peak time of the coronavirus epidemic.

1 Introduction

One of the common questions in an epidemic is its final size and its peak time. For answering this question various models are used: analytical [1-4], stochastic [5], phenomenological [6, 7]

In Jan 2020 Read et al. estimate that on 4. Feb 2020 there will be from 132751 to 273649 infections.[8]. Xiong and Yan [9], using EIR model, for data til 7. Feb 2020 predicts that the final size of the epidemy size would be 115 022 infections. For data for the period between 16 Jan 2020 to 8 Feb 2020 Nesteruk [10], using the SIR model, the prediction was about 65 000. For data til 10 Feb 2020, Anastassopoulou et al. [11], using the SIRD model, forecast 140 000 (scenario I) to 1 million (Scenario 2) infections on 29 Feb 2020.

In this note, we will try to estimate a final epidemic size and its peak by the phenomenological model, i.e., with the logistic growth regression model [7, 12]. The estimation is made in two steps: first, daily data are by the logistic model - in this way, we obtain a series of estimates - and second, the finale size is estimated by Shanks transformation [13] and also fitted by the Weibull function.

2 Logistic grow model

The logistic growth model originates in population dynamics [14]. The underlying assumption of the model is that the rate of change in the number of new cases per

capita linearly decreases with the number of cases. So, if C is the number of cases, and t is the time, then the model reads

$$\frac{1}{C}\frac{dC}{dt} = r\left(1 - \frac{C}{K}\right), \tag{1}$$

where r is infection rate, and K the final epidemic size. If $C(0) = C_0$ is the initial number of cases then the solution of (1) is

$$C = \frac{K}{1 + A\exp(-rt)} \tag{2}$$

where $A = \dfrac{K - C_0}{C_0}$ When $t \ll 1$, assuming $K \gg C_0$, and terefore $A \gg 1$ we have the natural growth

$$C = \frac{Ke^{rt}}{e^{rt} + A} = \frac{C_0}{1 - C_0/K}\frac{e^{rt}}{1 + e^{rt}/A} \approx C_0 e^{rt} \tag{3}$$

When $t \to \infty$ the number of cases follows the Weibull function

$$C = K\left(1 - Ae^{-rt} + \cdots\right) \approx K\left[1 - e^{-r(t-t_0)}\right] \tag{4}$$

The growth rate $\dfrac{dC}{dt}$ reaches its maximum when $\dfrac{d^2C}{dt^2} = 0$. From this condition, we obtain that the growth rate peak occurs in time time

$$t_p = \frac{\ln A}{r} \tag{5}$$

At this time the number of cases and the growth rate are

$$C_p = \frac{K}{2}, \qquad \left(\frac{dC}{dt}\right)_p = \frac{rK}{4} \tag{6}$$

3 Logistic regression model

By the logistic regression model, we estimate the parameters of the logistic model. Using (2) we can write the regression model in the form

$$C = \frac{b_1}{1 + b_2 \exp\left(-b_3 t\right)} \tag{7}$$

where unknown parameters b_1, b_2, b_3 which are to be estimated from the data. In the logistic model, the parameters are constants, while the regression model parameters depend on available data. So if $C_1, C_2, \ldots, C_n$ are the cases in times $t_1, t_2, \ldots, t_n$ then $b_i^{(n)} = b_i^{(n)}\left(C_1, C_2, \ldots, C_n\right)$ ($i = 1, 2, 3$), or in other words, parameters are time-dependent

$$b_i = b_i\left(t\right) \quad (i = 1, 2, 3) \tag{8}$$

Now, if we assume that the epidemy has a final size, then $b_i = b_i\left(t\right)$ should converge to some finite values. In the early stage, we could try to estimate the final limit by the Shanks transformation [13]. For final epidemy size, the Shanks transformation reads

$$K\left(K_n\right) = \frac{K_{n+1}K_{n-1} - K_n^2}{K_{n+1} - 2K_n + K_{n-1}} \tag{9}$$

where K_n is the estimate in time t_n. By iterated use of (7) we may obtain a better estimate of the final value of K.

Alternatively, we can fit the sequence $b_i^{(k)}$, $k = n_0, \ldots, n$ by Weibull function (or any other suitable function)

$$b_1 = K\left[1 - \exp\left(-k\left(t - t_0\right)^p\right)\right] \tag{10}$$

we obtain a final estimation of K, i.e., the epidemy size. k, p and t_0 are regression parameters.

4 Data

For regression analysis, we use two data sets.. For the period from 16 Jan to 21 Jan 2020 we use the data from [10], and after that from *worldmeters*[1].

---

[1] https://www.worldometers.info/coronavirus/

5 Results

For practical calculation, we use MATLAB's functions *lsqcurvefit* and *fitnlm* (see Appendix).

Based on available data (til 25 Feb 2020), the predicted final size of epidemy is about 84000 cases (see Table 1, see Fig 1, 2), and its peak was on 9 Feb 2020. The regression has a high coefficient of determination 0.995, while p-value (< 0.000) indicates that all parameters are statistically significant. With the new data, these estimates will change reliably within days.

The trend of model parameters is given in Table 2 while the graph of evaluation of the final size of the epidemy is shown as a bar graph in Fig 3. The current estimate of epidemy size, using iterated Shanks transformation (7), is 83100 cases. (see Table 2a)

Table 1. Estimate logistic model parameters for.

	Estimate	SE	tStat	pValue
b1	**83756**	**1388.6**	60.315	1.4849e-38
b2	0.21841	0.0087237	25.036	8.5202e-25
b3	172.88	30.405	5.6859	1.6727e-06

```
Number of observations: 40, Error degrees of freedom: 37
Root Mean Squared Error: 2.18e+03
R-Squared: 0.995,  Adjusted R-Squared 0.995
F-statistic vs. zero model: 5.73e+03, p-value = 2.19e-49
```

Figure 1. Predicted evaluation of the coronavirus epidemic. The regression data are in Table 4.

Figure 2. Predicted evaluation of the coronavirus epidemic rate

Table 2. Data and results of logistic regression (see Eqs (2), (3), (4))

data			Regression			Peak		
date	day	C	K	r	A	day	dCdt	date
		(cases)	(cases)	(1/day)				
30.jan.20	15	9821	16419	0.469	482.885	13	1927	30.jan.20
31.jan.20	16	11948	18165	0.45	452.557	13	2044	30.jan.20
1.feb.20	17	14551	21823	0.414	394.184	14	2260	31.jan.20
2.feb.20	18	17389	26068	0.383	350.318	15	2495	1.feb.20
3.feb.20	19	20628	31257	0.354	316.693	16	2766	2.feb.20
4.feb.20	20	24553	38821	0.324	290.892	17	3146	3.feb.20
5.feb.20	21	28276	44479	0.308	279.637	18	3420	4.feb.20
6.feb.20	22	31439	46180	0.303	275.4	18	3496	4.feb.20
7.feb.20	23	34876	48078	0.297	268.19	18	3570	4.feb.20
8.feb.20	24	37552	48596	0.295	265.417	18	3588	4.feb.20
9.feb.20	25	40553	49822	0.291	256.799	19	3621	5.feb.20
10.feb.20	26	43099	50903	0.286	247.807	19	3643	5.feb.20
11.feb.20	27	44919	51372	0.284	243.22	19	3650	5.feb.20
12.feb.20	28	60326	71335	0.226	152.937	22	4027	8.feb.20
13.feb.20	29	64437	93484	0.198	137.983	24	4627	10.feb.20
14.feb.20	30	67100	100456	0.192	137.028	25	4826	11.feb.20
15.feb.20	31	69197	97849	0.194	137.789	25	4757	11.feb.20
16.feb.20	32	71329	94252	0.198	140.336	24	4672	10.feb.20
17.feb.20	33	73332	91467	0.202	143.931	24	4617	10.feb.20
18.feb.20	34	75198	89575	0.205	147.671	24	4589	10.feb.20
19.feb.20	35	75700	87525	0.208	153.464	24	4568	10.feb.20
20.feb.20	36	76676	86067	0.212	159.069	23	4560	9.feb.20
21.feb.20	37	77673	85090	0.214	163-862	23	4561	9.feb.20
22.feb.20	38	78651	84468	0.216	167.584	23	4564	9.feb.20
23.feb.20	39	79400	84039	0.217	170.598	23	4568	9.feb.20
24.feb.20	40	80088	**83756**	0.218	172.864	23	4573	9.feb.20

Table 2a. Iterated Shanks transform

Day	K	K(K)	K(K(K))	K(K(K(K)))	
32	94252				
33	91467	85566			
34	89575	113981	99045		
35	87524	82493	83090	83617	
36	86067	83101	83635	83405	83307
37	85090	83386	83236	83338	
38	84469	83071	83372		
39	84039	90072			
40	83576				

The parameters of the Weibull model (8) with $p = 1$, of evaluation of the size of epidemy, for data until 12. Feb 2020, are given in Table 3. We see that the final size of epidemy would be about 51000 infections and the epidemy would reach its peak on 5 Feb 2020. However, a new way of data collection distorts this prediction. By using the Weibull model, we obtain the final size of the epidemy of about 93000 infections (see Table 4). We stress that this prediction should be taken with some care because there is not yet enough data to use the Weibull model for the final prediction of epidemy size. Namely, from Table 4, we see that R^2 is relatively high (0.881) but regression coefficients for k have p-value > 0.2 so it is statistically insignificant. With current the epidemy peak time (Fig 4), using Weibull regression, was at day 25, i.e., on 11 Feb 2020. In this regression, all parameters are significant with p-value < 0.06

Figure 3. Evaluation of estimated final size of coronavirus epidemic (bars). Weibull regression data (red line) are given in Table 4.

Table 3. The estimated final size of epidemy from data till 12.Feb. 2020

	Estimate	SE	tStat	pValue
b1	**58121**	3746.8	15.512	8.424e-08
b2	0.18721	0.039055	4.7935	0.00098261
b3	12.085	7.1289	1.6952	0.12428

Number of observations: 12, Error degrees of freedom: 9
Root Mean Squared Error: 2.11e+03
R-Squared: 0.975, Adjusted R-Squared 0.97
F-statistic vs. zero model: 1.53e+03, p-value = 1.7e-12

Table 4. The estimated final size of the epidemy by Weibull model (8) (see Fig 3)

	Estimate	SE	tStat	pValue
b1	**92892**	4956.9	18.74	5.1676e-15
b2	0.23906	0.045772	5.2229	3.0763e-05
b3	165.65	150.51	1.1005	0.28299
b4	1	0	Inf	0

Number of observations: 25, Error degrees of freedom: 22
Root Mean Squared Error: 9.64e+03
R-Squared: 0.881, Adjusted R-Squared 0.87
F-statistic vs. constant model: 81.3, p-value = 6.9e-1

Figure 4. Evaluation of estimated peak time of coronavirus epidemic (bars). Weibull regression data (red line) (Table 5).

Table 5. The estimated peak time size of the epidemy by Weibull model (8) from data till 25.Feb. 2020 (see Fig 4)

```
          Estimate        SE        tStat       pValue
          ________     ________    ______     __________

b1         25.29        0.90744     27.87      1.179e-18
b2        0.15454       0.031128    4.9648     5.737e-05
b3        9.1989        4.6548      1.9762       0.0608
b4          1             0          Inf           0

Number of observations: 25, Error degrees of freedom: 22
Root Mean Squared Error: 1.31
R-Squared: 0.889,  Adjusted R-Squared 0.879
F-statistic vs. constant model: 87.9, p-value = 3.21e-11
Peak date 11-Feb-2020
```

In Table 6, we present the actual and predicted number of cases for a period from 20. Feb 2020. We see that the discrepancy between the actual and predicted number of cases is about 2%, and it is decreasing. The current model prediction for 1. March 2020 is 82983 cases.

Table 6. Actual and predicted number of cases

Day	date	Actual	Predicted	Discrepancy %
35	20.feb.20	75700	77354	2.19
36	21.feb.20	76676	78532	2.42
37	22.feb.20	77673	79505	2.36
38	23.feb.20	78651	80305	2.10
39	24.feb.20	79400	80959	1.96
40	25.feb.20	80088	81493	1.75
41	26.feb.20		81928	
42	27.feb.20		82280	
43	28.feb.20		82566	
44	29.feb.20		82797	
45	1.mar.20		82983	

5 Conclusion

On the base of available data, one can now predict that the final size of coronavirus epidemy using the logistic model will be about 85 000 cases. The peak of the epidemic

was on 9 Feb 2020. The more optimistic final size of about 83100 cases is obtained by iterated Shanks transformation.

Less reliable is the following prediction with Weibull function: the final number of infections will be about 93 000, and that peak was after 25 days, i.e., on 11. Feb 2020.

If data collection will not change, then based on the above results, we conclude that the epidemic is passing into the saturation phase.

Appendix. MATLAB program.

```matlab
%FITVIRUS Estimation of coronavirus epidemy size by logistic model

close all

warning('off')

% obtain data
[sampleC,date0] = getData();

% form time
samplaTime = 0:1:length(sampleC)-1;
time = 0:1:2*length(samplaTime);

% form date
samplaDate = date0+samplaTime;
date = date0+time;

% initial guess
b0 = [max(sampleC) 1 max(sampleC)]';

% parameters evaluation
bb = NaN(3,length(sampleC));
tpeak = NaN(length(sampleC),1);
dCpeak = NaN(length(sampleC),1);

%initial data set
n0 = 15;

% loop over periods
opts = optimoptions('lsqcurvefit','Display','off');
for n = n0:length(sampleC)
    [b,resnorm,~,exitflag,output] = lsqcurvefit(@fun,b0,...
        samplaTime(1:n),sampleC(1:n),[],[],opts);
    bb(:,n) = b;
    % peak time and growth rate
    tpeak(n) = log(b(3))/b(2);
    dCpeak(n) = dfun(b,tpeak(n));
end

% final fit
fprintf('Regression parameters for complet data set\n')
mdl = fitnlm(samplaTime(1:n),sampleC(1:n),@fun,b0)

% print parameters evaluation
```

```matlab
    fprintf('\nEvaluation of model parameters\n')
    fprintf('%4s %12s %10s %10s %10s %12s
%12s\n','day','K','r','A','t_peak','dC_peak','date_peak')
    fprintf('%4s %12s %10s %10s %10s %12s\n','
','(cases)','(1/day)','','(day)','(cases/day)')
    for n = n0:length(sampleC)
        fprintf('%4d %12d %10.3f %10.3f %10d %12d
%12s\n',n,fix(bb(1,n)),bb(2:3,n),...
            fix(tpeak(n)),fix(dCpeak(n)),datestr(ceil(tpeak(n))+date0))
    end

    % fit estimated final sizes
    fprintf('\nWeibull regression of predicted final sizes\n')
    % Initial guess
    f0 = [sampleC(end) 0.13 12 1]'
    n = length(sampleC);
    mdl2 = fitnlm(samplaTime(n0:n),bb(1,n0:n),@fun1,f0)
    f = mdl2.Coefficients.Estimate;

    % fit estimated peak time
    fprintf('\nWeibull regression of predicted peak time\n')
    % Initial guess
    ff0 = [tpeak(end) 0.13 12 1]'
    n = length(sampleC);
    mdl2 = fitnlm(samplaTime(n0:n),tpeak(n0:n),@fun1,ff0)
    ff = mdl2.Coefficients.Estimate;
    fprintf('Peak date %s\n',datestr(ceil(ff(1))+date0))

    warning('on')

    % plot evaluation of peak time
    figure
    hold on
    bar(samplaDate(n0:n),tpeak(n0:n)); %,'LineWidth',2)
    datetick('x',20,'keeplimits')
    plot(date(n0:n),fun1(ff,time(n0:n)),'r','LineWidth',2)
    datetick('x',20,'keeplimits')
    ylabel('Day')
    xlabel('Date')
    title('Estimated peak time')
    grid on
    hold off

    % plot evaluation of final size
    figure
    hold on
    bar(samplaDate(n0:n),bb(1,n0:n)/1000); %,'LineWidth',2)
    datetick('x',20,'keeplimits')
    plot(date(n0:n),fun1(f,time(n0:n))/1000,'r','LineWidth',2)
    datetick('x',20,'keeplimits')
    ylabel('Infected (1000)')
    xlabel('Date')
    title('Estimated final size of epidemic')
    grid on
    hold off

    % plot final prediction of epedimy evaluation
    figure
    hold on
    plot(date,fun(b,time)/1000,'k','LineWidth',2)
    scatter(samplaDate,sampleC/1000,50,'k','filled')
```

```matlab
h = scatter(samplaDate,sampleC/1000,30,'w','filled');
h.Annotation.LegendInformation.IconDisplayStyle = 'off';
% add tick marks
datetick('x',20,'keeplimits')
% add axes labels
ylabel('Infected (1000)')
xlabel('Date')
% add legend
legend('Predicted','Actual','Location','best')
%add title
title('Coronavirus epidemic')
% add grid
grid on
hold off

% plot growth rate of epidemic
figure
hold on
plot(date,dfun(b,time),'LineWidth',2)
ylabel('Infected/day')
yyaxis right
plot(date,dfun(b,time)./fun(b,time),'LineWidth',2)
ylabel('Infected/per capita/day')
datetick('x',20,'keeplimits')
title('Coronavirus epidemic rate')
grid on
hold off

function y = fun(b,t)
% Logistic model
    y = b(1)./(1 + b(3)*exp(-b(2)*t));
end

function y = dfun(b,t)
% Grow rate
    y = b(1)*b(2)*b(3)*exp(-b(2)*t)./(1 + b(3)*exp(-b(2)*t)).^2;
end

function y = fun1(b,t)
% Weibull model. Note b(3) !!!
    y = b(1)*(1 - exp(-b(2)*(t-b(3)).^b(4)));
end

function [C,date0] = getData()
%GETDATA Coronavirus data
%   data from from 22 Jan 2020 to 18 Feb 2020 are from
%   https://www.worldometers.info/coronavirus/
date0=datenum('2020/01/16'); % start date
C = [
```

```
    6058
    7813
    9821
    11948
    14551
    17389
    20628
    24553
    28276
    31439
    34876
    37552
    40553
    43099
    44919
    60326
    64437
    67100
    69197
    71329
    73332
    75198
    75700
    76676
    77673
    78651
    79400
    80088
%<-------------- add new data here
]';
end
```

References

[1] D.A. Barbarossa MV, Kiss G, Nakata Y, Röst G, Vizi Z Transmission Dynamics and Final Epidemic Size of Ebola Virus Disease Outbreaks with Varying Interventions, PLoS ONE, 10 (2015).

[2] F. Brauer, The Final Size of a Serious Epidemic, Bull Math Biol, 81 (2019) 869-877.

[3] J.M.A. Danby, Computing applications to differential equations modelling in the physical and social sciences, Reston Publishing Company, Reston, Va., 1985.

[4] J.T. Wu, K. Leung, G.M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study, The Lancet, (2020).

[5] J.C. Miller, A note on the derivation of epidemic final sizes, Bull Math Biol, 74 (2012) 2125-2141.

[6] K.E. Fisman D, Tuite A. , Early Epidemic Dynamics of the West African 2014 Ebola Outbreak: Estimates Derived with a Simple Two-Parameter Model., PLOS Currents Outbreaks. , (2014).

[7] B. Pell, Y. Kuang, C. Viboud, G. Chowell, Using phenomenological models for forecasting the 2015 Ebola challenge, Epidemics, 22 (2018) 62-70.

[8] J.M. Read, J.R.E. Bridgen, D.A.T. Cummings, A. Ho, C.P. Jewell, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions, medRxiv, (2020) 2020.2001.2023.20018549.

[9] H. Xiong, H. Yan, Simulating the infected population and spread trend of 2019-nCov under different policy by EIR model, medRxiv, (2020) 2020.2002.2010.20021519.

[10] I. Nesteruk, Statistics based predictions of coronavirus 2019-nCoV spreading in mainland China, medRxiv, (2020) 2020.2002.2012.20021931.

[11] C. Anastassopoulou, L. Russo, A. Tsakris, C. Siettos, Data-Based Analysis, Modelling and Forecasting of the novel Coronavirus (2019-nCoV) outbreak, medRxiv, (2020) 2020.2002.2011.20022186.

[12] S.L. Chowell G, Viboud C, Kuang Y., West Africa Approaching a Catastrophic Phase or is the 2014 Ebola Epidemic Slowing Down? Different Models Yield Different Answers for Liberia. , PLOS Currents Outbreaks., (2014).

[13] D. Shanks, Non-linear Transformations of Divergent and Slowly Convergent Sequences, Journal of Mathematics and Physics, 34 (1955) 1-42.

[14] R. Haberman, Mathematical models mechanical vibrations, population dynamics, and traffic flow an introduction to applied mathematics, Unabridged republication ed., SIAM, Philadelphia, 1998.