Title: Clinical features and progression of acute respiratory distress syndrome in

coronavirus disease 2019

Yanli Liu, MD[1,*]; Wenwu Sun, MD[1,*]; Jia Li, PhD[2,*]; Liangkai Chen, PhD[3,*,#]; Yujun

Wang, MD[1]; Lijuan Zhang, MD[1]; Li Yu, MD[1,#].

[1] Intensive Care Unit, the Central Hospital of Wuhan, Tongji Medical College,

Huazhong University of Science and Technology, Wuhan, China.

[2] Department of Health Technology and Informatics, the Hong Kong Polytechnic

University, Hung Hom, Kowloon, Hong Kong PRC.

[3] Department of Nutrition and Food Hygiene, Hubei Key Laboratory of Food

Nutrition and Safety, Ministry of Education Key Lab of Environment and Health,

School of Public Health, Tongji Medical College, Huazhong University of Science

and Technology, Wuhan, China.

* The first four authors contributed equally to this article.

 # Corresponding authors:

Li Yu, Email: <u>yuliwhzxyy@163.com</u>;

Liangkai Chen, Email: <u>clk@hust.edu.cn</u>.

Word count: 2,681 (including Research in context)

No. of Tables: 3

No. of Figures: 2

Summary

Background: The outbreak of severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2) results in a cluster of coronavirus disease 2019 (COVID-19). We

reported the clinical characteristics of COVID-19 patients with acute respiratory

distress syndrome (ARDS), and further investigated the treatment and progression of

ARDS in COVID-19.

Methods: This study enrolled 109 patients with COVID-19 admitted to the Central

Hospital of Wuhan, a designated hospital in Wuhan, China, from January 2 to

February 1, 2020. Patients were followed up to February 12, 2020. The clinical data

were collected from the electronic medical records. The differences in the treatment

and progression with the time and the severity of ARDS were determined.

Findings: Among 109 patients, mean age was 55 years, and 59 patients were male.

With a median 15 days (range, 4 to 30 days) follow-up period, 31 patients (28.4%)

died, while 78 (71.6%) survived and discharged. Of all patients, 53 (48.6%)

developed ARDS. Compared to non-ARDS patients, ARDS patients were elder (mean

age, 61 years *vs*. 49 years), and more likely to have the coexistent conditions,

including diabetes (20.8% *vs*. 1.8%), cerebrovascular disease (11.3% *vs*. 0%), and

chronic kidney disease (15.1% *vs*. 3.6%). Compared to mild ARDS patients, those

with moderate and severe ARDS had higher mortality rates. No significant effect of

antivirus, glucocorticoid, or immunoglobulin treatment on survival was observed in

patients with ARDS.

Interpretation: The mortality rate increased with the severity of ARDS in COVID-19,

and the effects of current therapies on the survival for these patients were not

satisfactory, which needs more attention from clinicians.

Funding: Health and Family Planning Commission of Wuhan Municipality.

Research in context

Evidence before this study

We searched PubMed and the China National Knowledge Infrastructure database for articles published up to Feb 24, 2020, using the keywords "novel coronavirus", "2019 novel coronavirus", "2019-nCoV", "SARS-CoV-2", "COVID-19", "pneumonia", "coronavirus", AND "clinical feature" ,"mortality", AND "acute respiratory distress syndrome", "ARDS", for articles published in both Chinese and English. We found several recent articles describing the clinical characteristics of COVID-19 patients. One autopsy report described pathological findings of a 50-year-old COVID-19 patient with ARDS. A recent research with 52 critically ill patients published in *The Lancet Respiratory Medicine* indicated that older patients concurrent ARDS are at increased risk of death. No published work about the comprehensive description of clinical features, treatment, and mortality according to the severity of ARDS in COVID-19 patients.

Added value of this study

We report the differences in the clinical manifestations between COVID-19 patients with and without ARDS. Among 109 patients, 53 (48.6%) of them developed ARDS. Compared with non-ARDS patients, patients with ARDS were elder and more likely to have coexistent diseases. The mortality rate in ARDS patients (49.1%) was significantly higher than that in non-ARDS patients (8.9%). The clinical characteristics of COVID-19 patients varied with the severity of ARDS. High mortality rates were found in patients with moderate and severe ARDS. The survival

of COVID-19 patients with ARDS was not significantly improved by the antivirus,

glucocorticoid, or immunoglobulin treatment.

Implications of all the available evidence

In the front-line epidemic area, COVID-19 patients with moderate-to-severe ARDS

had high mortality rates. The current medical treatments might not have a satisfactory

effect on the in-hospital survival of COVID-19 patients with ARDS. For clinicians, it

is necessary to pay close attention to COVID-19 patients who are at high risk for

ARDS. The risk stratification and therapeutic strategy for COVID-19 patients should

be tailored according the variations in ARDS severity. It is essential for intensive care

physicians to participate in treatment decision-making and management in the early

stages of the COVID-19 outbreak.

Introduction

The emergent outbreaks of coronavirus disease 2019 (COVID-19) caused by the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) remain a threat to the public health worldwide [1,2]. As of February 24, 2020, 77,262 cases were confirmed, and 2,595 death cases were recorded in China. The number of confirmed cases in other countries are ascending as well [3-5]. COVID-19 may impose a great socioeconomic, public health, and clinical burden, especially in the low-income and middle-income countries.

Growing studies identified the clinical features of COVID-19 [6-8]. Similar to SARS in 2003, this infectious disease results in the high possibility of the admission to the intensive care unit (ICU) and the mortality [6]. Notably, due to the cytokine cascade within a short period, the critically ill patients with COVID-19 were more likely to develop acute respiratory distress syndrome (ARDS) and receive oxygen treatment [6]. ARDS was the most common complication in COVID-19 patients, with a high mortality rate [6,7]. The latest publication in *The Lancet Respiratory Medicine*, Yang *et al.* [9] reported that 67% of critically ill patients develop ARDS. However, little is known regarding the clinical characteristics, treatment, and progression of COVID-19 patients according to the severity of ARDS.

In this study, we retrospectively reviewed the clinical data of patients with COVID-19 who were admitted to the Central Hospital of Wuhan, and determined the differences

in the clinical characteristics between COVID-19 patients with and without ARDS.

Additionally, to reveal the evolvement of ARDS in COVID-19, we also investigated

the clinical features and therapies of COVID-19 patients at each severity level of

ARDS. Our study might provide new insight into the risk stratification and

therapeutic strategy for COVID-19 patients.

Methods

Study Population

We retrospectively analyzed the data of confirmed COVID-19 patients admitted to the

Central Hospital of Wuhan between January 2 and February 1, 2020. The Central

Hospital of Wuhan is one of the first designated hospitals to receive patients with

COVID-19. According to the World Health Organization interim guidance [10], the

diagnostic criteria of COVID-19 was based on the virus RNA detection, the clinical

characteristics, the chest imaging, and the ruling out common pathogen. Patients with

malignant tumors, previous craniocerebral operation, or died on admission were

excluded. In this study, we also excluded patients who had been transferred to other

hospitals for advanced life support and patients with mild symptoms who had been

transferred to mobile cabin hospitals. The clinical outcomes of patients were followed

up to February 12, 2020. Patients who were still in hospital before February 12, 2020

were excluded. Finally, 109 confirmed COVID-19 patients were included in the

analyses. All data were anonymous, and the requirement for informed consent was

waived. The Ethics Committees of the Central Hospital of Wuhan approved this study.

Data Collection

Case report forms, nursing records, laboratory findings, and radiological features were

reviewed. All data were collected onto the standardized forms from the electronic

medical records in the hospital. Two senior physicians independently reviewed the

data. Information on demographic data, symptoms, underlying comorbidities, chest

computed tomographic images, and laboratory results were included. All treatment

measures were collected during the hospitalization, such as antiviral therapy,

antibacterial therapy, corticosteroid therapy, immune support therapy, and respiratory

support. The time of disease onset was defined as the day of symptom onset. The time

from the first symptom to the fever clinics, hospital admission, and clinical outcomes

were recorded. A confirmed respiratory tract specimen was defined as positive for

SARS-CoV-2. Repeated tests for SARS-CoV-2 were done in the confirmed patients to

show viral clearance before the hospital discharge. The Berlin definition was applied

to determine the presence and severity of ARDS [11]. Acute Physiology and Chronic

Health Evaluation II (APACHE II), Sequential Organ Failure Assessment (SOFA)

scores and CURB-65 criteria [12] were determined within 24 hours after admission.

Detection of Coronavirus by Real-Time Reverse Transcription Polymerase Chain

Reaction

Throat swab samples were collected from the suspected patients, and the method of

virus RNA detection was reported [7]. Briefly, the presence of SARS-CoV-2 in the

respiratory specimens was detected by real-time reverse transcription polymerase

chain reaction (RT-PCR). Two target genes were used, including open reading frame

lab (ORF1ab) and nucleocapsid protein (N), and the sequences were as follows:

ORF1ab: forward primer CCCTGTGGGTTTTACACTTAA; reverse primer

ACGATTGTGCATCAGCTGA. N: forward primer

GGGGAACTTCTCCTGCTAGAAT; reverse primer

CAGACATTTTGCTCTCAAGCTG. The real-time RT-PCR assay was conducted in

line with the manufacturer's protocol (Beijing Genomics Institution and Geneodx

biotechnology Co. Ltd). Positive test results for two target genes were considered as

laboratory-confirmed infection.

Statistical Analysis

Normally distributed continuous variables were presented as mean ± SD, compared

by Student's t-test. Skew distributed continuous variables were shown as median

(interquartile range [IQR]), analyzed by Mann–Whitney test or Kruskal–Wallis test.

Categorical variables were compared by Chi-square test or the Fisher's exact test.

Kaplan–Meier methods were used for survival plotting, and log-rank test for

comparison of survival curves. The dynamic trajectory in laboratory parameters was

plotted using GraphPad Prism 8 (GraphPad Software, Inc). Generalized linear mixed

models examined the differences in the laboratory data between ARDS and non-

ARDS groups over time. All statistical analysis were performed by the statistical

software packages R (http://www.R-project.org, The R Foundation) and the

EmpowerStats (http://www.empowerstats.com, X&Y Solutions, Inc., Boston, MA)

with a two-sided significance threshold of $P <0.05$.

Results

Clinical Characteristics

A total of 109 patients with COVID-19 were included in the current study. The

baseline clinical parameters of overall COVID-19 patients and COVID-19 patients

with and without ARDS were summarized in **Table 1**. Mean age of all subjects was

55 years (IQR, 43-66 years; range, 22-94 years), and 54.1% of all patients were male.

The most common symptom at the illness onset was fever (82.6%), followed by dry

cough (61.5%) and fatigue (56.9%). Seventy-six patients (69.7%) had underlying

comorbidities, including hypertension (33.9%), diabetes (11.0%), and chronic kidney

disease (9.2%). The first virus RNA detection rate was 24.8% in the fever clinics, and

all patients were reconfirmed by repeated virus RNA testing after admission. Among

109 patients, 53 (48.6%) of them developed ARDS, 100 (91.7%) of them had bilateral

involvement of the chest radiographs, and 28 (25.7%) patients received high-flow

nasal oxygen ventilation, while 31 (28.4%) died.

Compared with non-ARDS patients with COVID-19, ARDS patients were elder

(mean age, 61 years *vs*. 49 years), and more likely to have coexistent diabetes (20.8%

vs. 1.8%), cerebrovascular disease (11.3% *vs*. 0%), and chronic kidney disease (15.1%

vs. 3.6%). The bilateral involvement in the chest radiographs was identified in all

ARDS patients, but in 83.9% non-ARDS patients. The likelihood to receive high-flow

nasal ventilation for ARDS patients (49.1%) was significantly higher than that for

non-ARDS patients (3.6%). The mortality rate in ARDS patients (49.1%) was also

significantly higher than that in non-ARDS patients (8.9%).

Disease Severity Scores and Laboratory Findings in COVID-19 Patients With

and Without ARDS

The disease severity evaluation and laboratory examination of non-ARDS and ARDS

patients with COVID-19 on admission were shown in **Table 2**. Compared with non-

ARDS subjects, ARDS subjects had significantly higher disease severity on the day of

hospital admission (all *P* values <0.001 for CURB-65, SOFA, and APACHE II).

ARDS patients were more likely accompanied by lymphopenia on admission (*P* value

<0.001). Besides, the levels of lactate, neutrophil count, C-reactive protein (CRP), and

procalcitonin were significantly higher in ARDS patients than in non-ARDS patients

(median lactate level, 1.6 *vs.* 1.1 mmol/L; median neutrophil count level, 4.1 *vs.* 3.2

$\times 10^9$/L; median CRP level, 4.9 *vs.* 2.0 mg/dL; median procalcitonin level, 0.15 *vs.*

0.06 ng/mL). Significant increased levels of blood urea nitrogen, serum fibrinogen, D-

dimer, and lactate dehydrogenase in ARDS patients were observed, compared with

those in non-ARDS patients (all *P* values <0.05).

As shown in **Figure 1**, the dynamic trajectory in nine laboratory parameters were

tracked on Day 1, 3, 7 and 14, respectively. Significant differences in the levels of

neutrophil count, lymphocyte count, CRP, procalcitonin, and D-dimer were found

between ARDS and non-ARDS patients with COVID-19 (all P values <0.05). From

Day 1 to 14, ARDS patients had significantly lower lymphocyte count levels, while

the levels of other laboratory findings were higher in ARDS patients.

Clinical Profile and Progression of COVID-19 Patients With ARDS

According to the severity of ARDS, we further analyzed the clinical features,

treatment, and progression of 53 COVID-19 patients with ARDS. As presented in

Table 3, the increase in APACHE II, SOFA, and CURB criteria scores occurred

concomitantly with the severity of ARDS (all P values <0.001). The levels of lactate,

blood urea nitrogen, D-dimer, and lactate dehydrogenase ascended with the severity

of ARDS (all P values <0.05), whereas the lymphocyte count levels decreased (P

value = 0.045). Patients with moderate-to-severe ARDS were the most likely to

receive glucocorticoid therapy (P value = 0.02) and high-flow nasal oxygen

ventilation (P value <0.001). Compared to patients with mild ARDS, those with

moderate and severe ARDS had higher mortality rates (P value <0.001). Subsequent

survival analysis was shown in **Figure 2**. We found no significant effect of antivirus,

glucocorticoid, or immunoglobulin treatment on survival in COVID-19 patients with

ARDS (all log-rank tests P >0.05).

Discussion

In the present study of ARDS progression in COVID-19, we firstly identified the

differences in the clinical manifestations between COVID-19 patients with and

without ARDS. Second, the clinical characteristics of COVID-19 patients varied with

the severity of ARDS. High mortality rates were observed in patients with moderate-

to-severe ARDS. Third, we did not observe any apparent effect of the current

therapeutic interventions on the in-hospital survival of COVID-19 patients with

ARDS. Our findings might highlight the clinical significance in the enhanced

attention towards COVID-19 patients at high risks of ARDS.

To the best of our knowledge, this study is the first to investigate the clinical features

and progression of SARS-CoV-2 infected patients according to the severity of ARDS.

We revealed that elder patients, or complicated by diabetes, cerebrovascular disease,

and chronic kidney disease, were more likely to develop ARDS. This finding might

further explain the significant association of age and comorbidity conditions with the

poor clinical outcomes in a previous COVID-19 study [7]. In addition, the elevated

levels of neutrophil count, CRP, procalcitonin and D-dimer, but the decreased

lymphocyte count levels, were detected in patients with ARDS, rather than those

without ARDS. Similarly, the levels of lactate and D-dimer increased with the

progression of ARDS due to SARS-CoV-2, while severer lymphopenia occurred over

time. Our investigations into these laboratory indicators were comparable with other

COVID-19 studies [6,7,13], suggesting that cytokine cascade, excessive inflammatory

reaction, and coagulation dysfunction might evolve in the course of ARDS resulted

from SARS-CoV-2.

Our understanding of this ARDS progression could be furthered by a recent autopsy

report on a 50-year-old COVID-19 patient with ARDS [14], which was highlighted by

the inflammatory infiltration of lymphocytes in both lungs and the immune

hyperactivation in the patient. Interestingly, no obvious heart injury occurred in this

patient, while the liver injury might be caused by SARS-CoV-2 infection or

medication use [14]. These pathological results were concordant with our clinical

observation of a dysregulated inflammatory and immune state in COVID-19 patients

with ARDS, which might in turn help to guide our medical therapy.

The medical management of ARDS has advanced remarkably during the past decades

[15]. However, the general mortality rates of ARDS caused by all etiologies fluctuated

from 11% to 87% [16]. In our study, most patients with COVID-19 received antibiotics

and antivirus treatment, while over half of them were given by glucocorticoid and

intravenous immunoglobulin therapies. Moderate-to-severe ARDS patients were more

likely to use high-flow nasal oxygen ventilation. Unfortunately, the mortality rate was

extremely high in severe ARDS patients with COVID-19, and the survival of COVID-

19 patients with ARDS was not improved by the antivirus, glucocorticoid, or

immunoglobulin treatment. Some potential reasons might interpret the high mortality

we observed: First, due to the SARS-CoV-2 outbreak within a short period, the

designated hospitals were overloaded in short term, with inadequate medical facilities

and insufficient medical staff. Notably, intensive care resources are in short supply

and may fail to meet demand under the increasing number critically ill patients.

Second, the Central Hospital of Wuhan is close to Huanan Seafood Market, and the

inpatients were the first or the second generation infected patients. In addition, the

inadequate preparation and insufficient understanding of the disease might contribute

to increased mortality in the front-line epidemic area during the early outbreak period.

The epidemic spread knows no borders. We are deeply concerned about the current

outbreak of SARS-CoV-2. As the low-income and middle-income countries are more

vulnerable to the lack of medical resources, the mortality rate of infectious diseases

may be even higher. Therefore, from a policy perspective, the government should

provide capital, technology, and manpower to curb the epidemic. Meanwhile, in the

front-line epidemic area, limited medical resources should be allocated reasonably

and effectively. Based on our observation, some patients progress quickly to severe

pneumonia or ARDS without warning sign. Adequate critical care resources and the

involvement of intensive care physicians are essential in the early stages of the SARS-

CoV-2 outbreak.

The present study is subjected to several limitations. Due to the retrospective nature of

the present study, a systematic selection bias could be introduced. We could not

completely address the residual confounding factors as well. Although our laboratory

observations showed the significant changes with the time and the severity of ARDS,

the clinical predictive value remained to be determined. In addition, our results on the

effect of the current treatments should be considered with caution, and high-quality

clinical interventional studies are needed. Meanwhile, the benefits of the invasive

ventilation therapy on the disease prognosis should be further investigated. In the

future, a multi-center and follow-up study with a larger cohort is eagerly warranted.

Conclusions

In the current study from one of the designated hospitals in Wuhan, China, the

significant changes with the time and the severity of ARDS were observed in COVID-

19 patients. Moreover, COVID-19 patients with moderate and severe ARDS had high

mortality rates. The current medical treatments might not have a significant effect on

the in-hospital survival of COVID-19 patients with ARDS. Our observations might

help to guide the risk stratification and therapeutic strategy for COVID-19 patients.

Acknowledgement

We deeply regret and mourn all the lives lost in this disaster of SARS-CoV-2,

including our dearest colleague Dr. Wenliang Li. We would like to express our deepest

respect to all the people who are currently fighting against the outbreak of COVID-19.

Authors Contributions

LC and YL had full access to all the data in the study, and took responsibility for the

integrity of the data and the accuracy of the data analysis. YL, WS, LC, and LY made

substantial contributions to the study concept and design. YL and WS took

responsibility for obtaining ethical approval, collecting samples, and confirming data

accuracy. LC was in charge of the statistical analysis. YL, JL, and LC were in charge

of the manuscript draft. JL, LC, LZ, YW, and LY contributed to critical revision of the

report. All authors reviewed and approved the final version.

Conflict of Interest Disclosures

The authors declared no conflict of interest.

Role of the funding source

This study was supported by the Health and Family Planning Commission of Wuhan

Municipality, grant number WX18A02. The funders had no role in the design and

conduct of the study; collection, management, analysis, and interpretation of the data;

preparation, review, or approval of the manuscript; and decision to submit the

manuscript for publication.

Data sharing

The data that support the findings of this study are available from the corresponding

authors on reasonable request. We can provide participant data without names and

identifiers, but not the study protocol, or statistical analysis plan. After publication of

study findings, the data will be available for others to request. Once the data can be

made public, the research team will provide an email address for communication. The

corresponding authors have the right to decide whether to share the data or not based

on the research objectives and plan provided.

References

1.   Hui DS, E IA, Madani TA, et al. The continuing 2019-nCoV epidemic threat of novel
coronaviruses to global health - The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J*
Infect Dis 2020; **91**: 264-6.

2.   Velavan TP, Meyer CG. The COVID-19 epidemic. *Trop Med Int Health* 2020.

3.   Team C-NIRS. COVID-19, Australia: Epidemiology Report 3 (Reporting week ending 19:00
AEDT 15 February 2020). *Commun Dis Intell (2018)* 2020; **44**.

4.   Smith N, Fraser M. Straining the System: Novel Coronavirus (COVID-19) and Preparedness
for Concomitant Disasters. *Am J Public Health* 2020: e1-e2.

5.   Habibzadeh P, Stoneman EK. The Novel Coronavirus: A Bird's Eye View. *Int J Occup Environ*
Med 2020; **11**(2): 65-71.

6.   Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel
coronavirus in Wuhan, China. *Lancet* 2020; **395**(10223): 497-506.

7.   Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019
Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020.

8.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel
Coronavirus-Infected Pneumonia. *N Engl J Med* 2020.

9.   Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-
CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet*
Respiratory Medicine.

10.   World Health Organization. Clinical management of severe acute respiratory infection when
novel coronavirus (nCoV) infection is suspected: interim guidance. https://www.who.int/docs/default-
source/coronaviruse/clinical-management-of-novel-cov.pdf?sfvrsn=bc7da517_2. Published January 28,
2020. Accessed February 12, 2020.

11.   Force ADT, Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin
Definition. *JAMA* 2012; **307**(23): 2526-33.

12.   Lim WS, van der Eerden MM, Laing R, et al. Defining community acquired pneumonia
severity on presentation to hospital: an international derivation and validation study. *Thorax* 2003;
58(5): 377-82.

13.   Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019
novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;
395(10223): 514-23.

14.   Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute
respiratory distress syndrome. *The Lancet Respiratory Medicine*.

15.   Peck TJ, Hibbert KA. Recent advances in the understanding and management of ARDS.
F1000Res 2019; **8**.

16.   Maca J, Jor O, Holub M, et al. Past and Present ARDS Mortality Rates: A Systematic Review.
Respir Care 2017; **62**(1): 113-22.

Table 1. Baseline clinical features of subjects with COVID-19.

	No. (%)			
	Total (*n* = 109)	Non-ARDS (*n* = 56)	ARDS (*n* = 53)	*P*-value[a]
Age, median (IQR), y	55 (43-66)	49 (37-59)	61 (52-70)	**<0.001**
Male	59 (54.1)	31 (55.4)	28 (52.8)	0.79
Entering complaint				
Fever	90 (82.6)	48 (85.7)	42 (79.2)	0.37
Dry cough	67 (61.5)	36 (64.3)	31 (58.5)	0.53
Fatigue	62 (56.9)	26 (46.4)	36 (67.9)	**0.02**
Stethalgia	7 (6.4)	4 (7.1)	3 (5.7)	0.75
Diarrhea	12 (11.0)	6 (10.7)	6 (11.3)	0.92
Comorbidities				
COPD	4 (3.7)	2 (3.6)	2 (3.8)	>0.99
Hypertension	37 (33.9)	16 (28.6)	21 (39.6)	0.22
Diabetes	12 (11.0)	1 (1.8)	11 (20.8)	**0.002**
Cardiovascular disease	7 (6.4)	4 (7.1)	3 (5.7)	>0.99
Cerebrovascular disease	6 (5.5)	0 (0.0)	6 (11.3)	**0.01**
Chronic kidney disease	10 (9.2)	2 (3.6)	8 (15.1)	**0.049**
Onset of symptom to fever clinics, median (IQR), d	4 (2-6)	4 (2-7)	4 (2-6)	0.88
Onset of symptom to hospital admission, median (IQR), d	7 (5-9)	7 (5-9)	7 (5-9)	0.58
Hospital stays, median (IQR), d	15 (11-24)	15 (9-27)	16 (11-23)	0.41
Nucleic acid test (+) in fever clinics	27 (24.8)	12 (21.4)	15 (28.3)	0.41
Bilateral involvement of chest radiographs	100 (91.7)	47 (83.9)	53 (100)	**0.002**
Fever clinics treatment				
Antibiotics	105 (96.3)	55 (98.2)	50 (94.3)	0.35

Antivirus	105 (96.3)	55 (98.2)	50 (94.3)	0.35
Glucocorticoid therapy	43 (39.4)	20 (35.7)	23 (43.4)	0.41
Intravenous immunoglobulin	32 (29.4)	19 (33.9)	13 (24.5)	0.28
In-patient treatment				
Antibiotics	106 (97.2)	53 (94.6)	53 (100.0)	0.24
Antivirus	107 (98.2)	55 (98.2)	52 (98.1)	>0.99
Glucocorticoid therapy	72 (66.1)	35 (62.5)	37 (69.8)	0.42
Intravenous immunoglobulin	61 (56.0)	32 (57.1)	29 (54.7)	0.80
High-flow nasal ventilation	28 (25.7)	2 (3.6)	26 (49.1)	**<0.001**
Death	31 (28.4)	5 (8.9)	26 (49.1)	**<0.001**

390   Abbreviations: ARDS, acute respiratory distress syndrome; COPD, chronic obstructive pulmonary disease; IQR, interquartile range;

391   [a] *p* values indicate differences between non-ARDS and ARDS patients.

393 **Table 2. Severity of COVID-19 and laboratory tests in Non-ARDS and ARDS subjects.**

		Median (IQR)			
	Normal range	Total ($n = 109$)	Non-ARDS ($n = 56$)	ARDS ($n = 53$)	P-value[a]
Severity of illness scores and blood gas analysis on admission to hospital					
CURB-65	NA	0 (0-1)	0 (0-1)	1 (0-2)	**<0.001**
SOFA	NA	2 (1-4)	1 (0-1)	4 (2-5)	**<0.001**
APACHE II	NA	4 (2-8)	2 (1-3)	8 (5-10)	**<0.001**
Lactate, mmol/L	0.5-1.6	1.3 (0.8-2.0)	1.1 (0.8-1.6)	1.6 (1.1-2.2)	**<0.001**
$PaO_2:FiO_2$, mm Hg	400-500	296 (142-435)	475 (398-524)	145 (108-250)	**<0.001**
Laboratory findings on admission to hospital					
White blood cell count, $\times 10^9$/L	3.5-9.5	5.2 (4.0-7.0)	4.9 (3.8-6.2)	5.4 (4.5-8.0)	0.053
Neutrophil count, $\times 10^9$/L	1.8-6.3	3.6 (2.8-5.6)	3.2 (2.4-4.2)	4.1 (3.1-7.1)	**0.004**
Lymphocyte count, $\times 10^9$/L	1.1-3.2	0.9 (0.5-1.2)	1.0 (0.8-1.4)	0.7 (0.4-1.1)	**<0.001**
C-reactive protein, mg/dL	0-0.5	3.1 (1.1-5.4)	2.0 (0.9-3.2)	4.9 (2.7-6.5)	**<0.001**
Procalcitonin, ng/mL	<0.05	0.09 (0.06-0.20)	0.06 (0.05-0.09)	0.15 (0.08-0.37)	**<0.001**
Blood urea nitrogen, mmol/L	2.8-8	4.3 (3.3-6.5)	3.5 (2.9-4.8)	5.5 (4.0-7.1)	**<0.001**
Creatinine, μmol/L	57-111	65 (55-82)	64 (52-80)	67 (55-83)	0.42
Total bilirubin, mmol/L	0-21	9.0 (6.5-13.4)	8.3 (6.8-11.7)	10.5 (6.2-14.8)	0.23
Alanine aminotransferase, U/L	9-50	23 (15-36)	23 (14-41)	24 (16-31)	0.82
Aspartate aminotransferase, U/L	15-40	30 (21-40)	29 (19-38)	31 (25-44)	0.17
Fibrinogen, g/L	2-4	3.1 (2.7-3.6)	2.9 (2.7-3.3)	3.4 (2.9-3.9)	**0.001**
D-dimer, mg/L	0-500	570 (300-1178)	370 (250-650)	940 (470-1905)	**<0.001**
Lactate dehydrogenase, U/L	135-225	238 (185-341)	209 (183-267)	264 (190-448)	**0.011**
Creatine kinase, U/L	<190	91 (52-178)	100 (53-183)	83 (49-169)	0.29

| Creatine kinase–MB, U/L | <25 | 9.0 (6.1-14.8) | 8.5 (6.0-12.5) | 10.4 (7.0-15.1) | 0.28 |

Table 3. Clinical characteristics, laboratory findings, and treatment according to the severity of ARDS.

	Normal range	Median (IQR)			P-value[a]
		Mild ARDS ($n = 19$)	Moderate ARDS ($n = 24$)	Severe ARDS ($n = 10$)	
Age, y	NA	52 (46-63)	65 (57-70)	67 (55-71)	**0.04**
Severity of illness scores and blood gas analysis on admission to hospital					
CURB-65	NA	0 (0-1)	1 (1-2)	2 (1-3)	**<0.001**
SOFA	NA	2 (2-3)	4 (3-5)	6 (5-6)	**<0.001**
APACHE II	NA	3 (2-6)	8 (7-12)	10 (8-13)	**<0.001**
Lactate, mmol/L	0.5-1.6	1.2 (0.8-1.7)	1.8 (1.1-2.3)	5.2 (2.0-10.0)	**<0.001**
$PaO_2:FiO_2$, mm Hg	400-500	276 (247-280)	140 (117-153)	85 (57-88)	**<0.001**
Laboratory findings on admission to hospital					
White blood cell count, $\times 10^9$/L	3.5-9.5	4.6 (3.8-5.0)	7.3 (5.2-10.5)	7.0 (4.6-8.0)	**0.006**
Neutrophil count, $\times 10^9$/L	1.8-6.3	3.2 (2.3-3.8)	6.0 (4.2-9.0)	5.3 (3.7-7.3)	**0.002**
Lymphocyte count, $\times 10^9$/L	1.1-3.2	0.9 (0.7-1.3)	0.6 (0.4-0.9)	0.5 (0.4-0.8)	**0.045**
C-reactive protein, mg/dL	0-0.5	4.1 (2.2-5.9)	4.1 (1.9-9.7)	6.7 (5.5-12.2)	**0.027**
Procalcitonin, ng/mL	<0.05	0.10 (0.08-0.21)	0.16 (0.08-0.55)	0.28 (0.22-1.52)	0.09
Blood urea nitrogen, mmol/L	2.8-8	4.5 (3.5-6.1)	5.6 (4.5-7.7)	7.1 (6.0-8.0)	**0.03**
Creatinine, μmol/L	57-111	67 (58-74)	65 (54-99)	70 (55-80)	0.89
Total bilirubin, mmol/L	0-21	7.9 (5.8-10.4)	13.2 (9.3-18.6)	13.2 (6.2-13.5)	0.07
Alanine aminotransferase, U/L	9-50	24 (15-34)	24 (21-27)	21 (15-32)	0.90
Aspartate aminotransferase, U/L	15-40	29 (21-43)	34 (27-44)	29 (27-36)	0.45
Fibrinogen, g/L	2-4	3.3 (2.8-3.6)	3.4 (2.9-4.3)	3.6 (3.4-4.1)	0.41
D-dimer, mg/L	0-500	330 (220-798)	1220 (695-4035)	1385 (985-7550)	**<0.001**

Lactate dehydrogenase, U/L	135-225	190 (166-238)	347 (255-504)	458 (286-549)	**0.001**
Creatine kinase, U/L	<190	69 (44-119)	92 (58-185)	66 (53-188)	0.49
Creatine kinase–MB, U/L	<25	8.0 (6.5-11.6)	11.1 (6.8-16.1)	10.9 (8.9-12.0)	0.37
In-patient treatment, *n* (%)					
Ribavirin	NA	18 (94.7)	18 (75.0)	10 (100.0)	0.08
Oseltamivir	NA	5 (26.3)	8 (33.3)	1 (10.0)	0.43
Arbidol	NA	3 (15.8)	2 (8.3)	1 (10.0)	0.85
Glucocorticoid therapy	NA	9 (47.4)	21 (87.5)	7 (70.0)	**0.02**
Intravenous immunoglobulin	NA	13 (68.4)	9 (37.5)	7 (70.0)	0.08
High-flow nasal ventilation, *n* (%)	NA	1 (5.3)	19 (79.2)	6 (60.0)	**<0.001**
Hospital stays, d	NA	18 (13-25)	18 (15-25)	11 (10-13)	**0.03**
Death, *n* (%)	NA	1 (5.3)	15 (62.5)	10 (100.0)	**<0.001**

399 Abbreviations: APACHE II, Acute Physiology and Chronic Health Evaluation II; ARDS, acute respiratory distress syndrome; FiO$_2$, fraction of
400 inspired oxygen; IQR, interquartile range; NA, not available; PaO$_2$, partial pressure of oxygen; SOFA, Sequential Organ Failure Assessment
401 [a] *p* values indicate differences among mild, moderate, and severe ARDS patients.
402

Figure legends

Figure 1. Timeline charts illustrate the laboratory parameters in 109 patients with COVID-19 (56 non-ARDS and 53 ARDS) on day 1, day 3, day 7, and day 14 after admission. Data are represented as geometric mean and 95% confidence interval (one group only shows the upper or lower bar). The dash lines in black show the upper normal limit of each parameter, and the dash line in red shows the lower normal limit of lymphocyte count.

* *P* <0.05 for non-ARDS *vs*. ARDS.

Figure 2. Kaplan-Meier survival curves for 53 COVID-19 patients concurrent ARDS. (A) Ribavirin; (B) Oseltamivir; (C) Arbidol; (D) Glucocorticoid therapy; (E) Intravenous immunoglobulin.