TITLE

Novel *KITLG* regulatory variants are associated with lung function in African American children with asthma

AUTHORS

Angel CY Mak[1,*], Satria Sajuthi[2,^], Jaehyun Joo[3,^], Shujie Xiao[4,^], Patrick M Sleiman[5,6,^], Marquitta J White[1,^], Eunice Y Lee[1], Benjamin Saef[3], Donglei Hu[1], Hongsheng Gui[4], Kevin L Keys[1,7], Fred Lurmann[8], Deepti Jain[9], Gonçalo Abecasis[10], Hyun Min Kang[10], Deborah A. Nickerson[11,12,13], Soren Germer[14], Michael C Zody[14], Lara Winterkorn[14], Catherine Reeves[14], Scott Huntsman[1], Celeste Eng[1], Sandra Salazar[1], Sam S Oh[1], Frank D Gilliland[15], Zhanghua Chen[15], Rajesh Kumar[16], Fernando D Martínez[17], Ann Chen Wu[18], Elad Ziv[1], Hakon Hakonarson[5,6,#], Blanca E Himes[3,#], L Keoki Williams[4,#], Max A Seibold[3,#], Esteban G. Burchard[1,20]

[1] Department of Medicine, University of California San Francisco, San Francisco, CA, USA

[2] Center for Genes, Environment, and Health, National Jewish Health, Denver, CO, USA

[3] Department of Biostatistics, Epidemiology, and Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[4] Center for Individualized and Genomic Medicine Research, Department of Internal Medicine, Henry Ford Health System, Detroit, MI, USA

[5] Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, PA, USA

[6] Department of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA, USA

[7] Berkeley Institute for Data Science, University of California, Berkeley, CA, USA

[8] Sonoma Technology Inc, Petaluma, CA, USA

[9] Department of Biostatistics, University of Washington, Seattle, WA, USA

[10] Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA

[11] Department of Genome Sciences, University of Washington, Seattle, WA, USA

[12] Northwest Genomics Center, Seattle, WA, USA

[13] Brotman Baty Institute, Seattle, WA, USA

[14] New York Genome Center, New York, NY, USA

[15] Department of Preventive Medicine, Division of Environmental Health, Keck School of
Medicine, University of Southern California, Los Angeles, CA, USA

[16] Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago, IL, USA

[17] Asthma and Airway Disease Research Center, University of Arizona, Tucson, AZ, USA

[18] Precision Medicine Translational Research (PRoMoTeR) Center, Department of Population
Medicine, Harvard Medical School and Pilgrim Health Care Institute, Boston, MA, USA

[19] Department of Bioengineering and Therapeutic Sciences, University of California, San
Francisco, CA, USA

[^,#] These authors contributed equally to this work

[*] Corresponding author

Angel CY Mak
Address: Box 2911, 1550 4th Street, San Francisco, CA 94112
Tel: 415-514-9931
Email: angel.mak@ucsf.edu

AUTHOR CONTRIBUTIONS

ACYM, SSaj, MJW, SSO, FG, ZC, RK, FDM, ACW, EZ, and EGB designed and supervised the study.

ACYM, SSaj, JJ, SX, PS, MJW, EYL, BS, DH, SH, BEH, HH, LKW, and MAS supervised/performed

analyses and/or interpreted results. CE processed biospecimens for whole genome sequencing.

FL estimated air pollution exposure. DJ, GA, HMK, DAN, and SG contributed to TOPMed WGS

data generation and/or analysis. MZ, LW, and CR contributed to CCDG WGS data generation.

SSal coordinated study recruitment and processed phenotype data ACYM, SSaj, JJ, SX, PS, MJW,

EYL, BS, HG, KLK, SG, BEH, HH, LKW, MAS, and EGB wrote and/or critically reviewed the

manuscript.

SHORT TITLE

52 *KITLG* associated with FEV_1 in AA youths

KEYWORDS

54 GWAS, African American, FEV_1, gene-by-environment interaction, air pollution

ABSTRACT

Baseline lung function, quantified as forced expiratory volume in the first second of exhalation

(FEV_1), is a standard diagnostic criterion used by clinicians to identify and classify lung diseases.

Using whole genome sequencing data from the National Heart, Lung, and Blood Institute

TOPMed project, we identified a novel genetic association with FEV_1 on chromosome 12 in 867

African American children with asthma (p = 1.26 x 10^{-8}, β = 0.302). Conditional analysis within 1

Mb of the tag signal (rs73429450) yielded one major and two other weaker independent signals

within this peak. We explored statistical and functional evidence for all variants in linkage

disequilibrium with the three independent signals and yielded 9 variants as the most likely

candidates responsible for the association with FEV_1. Hi-C data and eQTL analysis demonstrated

that these variants physically interacted with *KITLG* and their minor alleles were associated with

increased expression of *KITLG* gene in nasal epithelial cells. Gene-by-air-pollution interaction

analysis found that the candidate variant rs58475486 interacted with past-year SO_2 exposure (p

= 0.003, β = 0.32). This study identified a novel protective genetic association with FEV_1,

possibly mediated through *KITLG*, in African American children with asthma.

INTRODUCTION

Asthma, a chronic pulmonary condition characterized by reversible airway obstruction, is one of the hallmark diseases of childhood in the United States (World Health Organization 2017). Asthma is also the most disparate common disease in the pediatric clinic, with significant variation in prevalence, morbidity, and mortality among U.S. racial/ethnic groups (Oh *et al.* 2016). Specifically, African American children carry a higher asthma disease burden compared to their European American counterparts (Akinbami *et al.* 2014; Akinbami 2015). Forced expiratory volume in the first second (FEV_1), a measurement of lung function, is a vital clinical trait used by physicians to assess overall lung health and diagnose pulmonary diseases such as asthma (Johnson and Theurer. 2014). We have previously shown that genetic ancestry plays an important role in FEV_1 variation and that African Americans have lower FEV_1 compared to European Americans regardless of asthma status (Kumar *et al.* 2010; Pino-Yanes *et al.* 2015). The disparity in lung function between populations may explain disparities in asthma disease burden. Understanding the factors that influence FEV_1 variation among individuals with asthma could lead to improved patient care and therapeutic interventions.

Twin and family-based studies estimate that the heritability of FEV_1 is as high as 81%, supporting a strong contribution by genetic factors in FEV_1 variation (Chatterjee and Das. 1995; Hukkinen *et al.* 2011; Palmer *et al.* 2001; Sillanpaa *et al.* 2017; Tian *et al.* 2017; Yamada *et al.* 2015). Genome-wide association studies (GWAS) of FEV_1, including among individuals with asthma, have identified several variants that contribute to lung function (Li *et al.* 2013; Liao *et al.* 2014; Repapi *et al.* 2010; Soler Artigas *et al.* 2011; Soler Artigas *et al.* 2015; Wain *et al.*

2017). Most of these previous GWAS, however, were performed in adult populations of

European descent, and their results may not generalize across populations or across the life

span of an individual (Carlson *et al.* 2013; Martin, A. R. *et al.* 2017; Wojcik *et al.* 2019). Previous

GWAS results are also limited due to their reliance on genotyping arrays. In particular, variation

in non-coding regions of the genome is not adequately covered by many genotyping arrays

because they were not designed while taking into account the population-specific genetic

variability of all populations (Kim, M. S. *et al.* 2018; Zhang and Lupski. 2015). Whole genome

sequencing (WGS) is a newer technology that captures nearly all common variation from coding

and non-coding regions of the genome and is unencumbered by genotype array design

constraints and differences in linkage disequilibrium patterns among populations. To date, no

large-scale WGS studies of lung function have been performed in African American children

with asthma (Martin *et al.* 2017).

In addition to genetics, FEV_1 is a complex trait that is significantly influenced by both genetic

variation and environmental factors, such as air pollution (Chatterjee and Das. 1995; Hukkinen

et al. 2011; Palmer *et al.* 2001; Sillanpaa *et al.* 2017; Tian *et al.* 2017; Yamada *et al.* 2015).

Exposure to ambient air pollution has been consistently associated with poor respiratory

outcomes, including reduced FEV_1 (Barraza-Villarreal *et al.* 2008; Brunekreef and Holgate. 2002;

Ierodiakonou *et al.* 2016; Wise 2019). We previously showed that exposure to sulfur dioxide

(SO_2), an air pollutant emitted by the burning of fossil fuels, is significantly associated with

reduced FEV_1 in African American children with asthma in the SAGE II study (Neophytou *et al.*

2016). Because the genetic variants associated with FEV_1 thus far do not account for the

majority of its estimated heritability, considering gene-environment interactions, specifically

gene-by-air-pollution, may improve our understanding of lung function genetics (Moore 2005;

Moore and Williams. 2009). Here, we performed a genome-wide association analysis using

WGS data to identify common genetic variants associated with FEV_1 in African American

children with asthma in SAGE II and investigated the effect of gene-by-air-pollution (SO_2)

interactions on FEV_1 associations.

METHODS

Study population

This study examined African American children between 8-21 years of age with physician-

diagnosed asthma from the Study of African Americans, Asthma, Genes & Environments (SAGE

II). All SAGE II participants were recruited from the San Francisco Bay Area. The inclusion and

exclusion are previously described in detailed (Oh *et al.* 2012; White *et al.* 2016) . Briefly,

participants were eligible if they were 8-21 years of age and self-identified as African American

and had four African American grandparents. Study exclusion criteria included the following: 1)

any smoking within one year of the recruitment date; 2) 10 or more pack-years of smoking; 3)

pregnancy in the third trimester; 4) history of lung diseases other than asthma (for cases) or

chronic illness (for cases and controls). Baseline lung function defined as forced expiratory

volume in the first second (FEV_1) was measured by spirometry prior to administering albuterol

as previously described (Oh *et al.* 2012).

TOPMed whole genome sequencing data

SAGE II DNA samples were sequenced as part of the Trans-Omics for Precision Medicine

(TOPMed) whole genome sequencing (WGS) program (Taliun *et al.* 2019). WGS was performed

at the New York Genome Center and Northwest Genomics Center on a HiSeq X system

(Illumina, San Diego, CA) using a paired-end read length of 150 base pairs (bp), with a minimum

of 30x mean genome coverage. DNA sample handling, quality control, library construction,

clustering and sequencing, read processing and sequence data quality control are described in

detail in the TOPMed website (TOPMed 2019). Variant calls were obtained from TOPMed data

freeze 8 VCF files corresponding to the GRCh38 assembly. Variants with a minimal read depth

of 10 (DP10) were used for analysis unless otherwise stated.

Genetic principal components, global ancestry, and kinship

estimation

Genetic principal components (PCs), global ancestry, and kinship estimation on genetic

relatedness were computed using biallelic single nucleotide polymorphisms (SNPs) with a PASS

flag from TOPMed freeze 8 DP10 data. PCs and kinship estimates were computed using the PC-

Relate function from the GENESIS R package (Conomos *et al.* 2015; Conomos *et al.* 2016) using

a workflow available from the Summer Institute in Statistical Genetics Module 17 course

website (Summer Institute in Statistical Genetics 2019). African global ancestry was computed

using the ADMIXTURE package (Alexander *et al.* 2009) in supervised mode using European

(CEU), African (YRI) and Native American (NAM) reference panels as previously described (Mak,

A. C. Y. *et al.* 2018).

FEV$_1$ GWAS

Non-normality of the distribution of FEV$_1$ values was tested with the Shapiro-Wilk test in R using

the shapiro.test function. Since FEV$_1$ was not normally distributed (p = 1.41 x 10^{-8} for FEV$_1$ and p

$= 1.05 \times 10^{-8}$ for $\log_{10}$ FEV_1), FEV_1 was regressed on all covariates (Age, sex, height, controller

medications, sequencing centers, and the first 5 genetic PCs) and the residuals were inverse-

normalized. These inverse-normalized residuals (FEV_1.res.rnorm) were the main outcome of the

discovery GWAS. The controller medication covariate included the use of inhaled

corticosteroids (ICS), long-acting beta-agonists (LABA), leukotriene inhibitors and/or an

ICS/LABA combo in the 2 weeks prior to the recruitment date.

Genome-wide single variant analysis was performed on the ENCORE server (https://github.com

/statgen/encore) using the linear Wald test (q.linear) originally implemented in EPACTS

([https://genome.sph.umich.edu /wiki/EPACTS](https://genome.sph.umich.edu/wiki/EPACTS)) and TOPMed freeze 8 data (DP0 PASS) with a

MAF filter of 0.1%. All pairwise relationships with degree 3 or more relatedness (kinship values

> 0.044) were identified, and one participant of the related pair was subsequently chosen at

random and removed prior to analysis. All covariates used to obtain FEV_1.res.rnorm were also

included as covariates in the GWAS as recommended in a recent publication (Sofer *et al.* 2019).

The association analysis was repeated using untransformed FEV_1 and FEV_1 percent predicted

(FEV_1.perc.predicted). FEV_1 percent predicted was defined as the percentage of measured FEV_1

relative to predicted FEV_1 estimated by the Hankinson lung function prediction equation for

African Americans (Hankinson *et al.* 1999). A secondary analysis that included smoking-related

covariates (smoking status and number of smokers in the family) was performed in PLINK 1.9

(version 1p9_2019_0304_dev) (Chang *et al.* 2015; Purcell and Chang. 2013). Regional

association results were plotted using LocusZoom 1.4 (Pruim *et al.* 2010) with a 500 kilobase

(Kb) flanking region. Linkage disequilibrium (R^2) was estimated in PLINK 1.9.

The function effectiveSize in the R package CODA was used to estimate the actual effective

number of independent tests and CODA-adjusted statistical and suggestive significance p-value

thresholds were defined as 0.05 and 1 divided by the effective number of tests, respectively

(Duggal *et al.* 2008). We compared the CODA-adjusted statistical significance threshold and the

widely used 5×10^{-8} GWAS genome-wide significance threshold (Pe'er *et al.* 2008) and selected

the more stringent threshold for genome-wide significance.

The following WGS quality control steps were applied to all reported variants from ENCORE to

ensure WGS variant quality: (1) The variant had VCF FILTER = PASS; (2) Variant quality was

confirmed via manual inspection on the BRAVO server based on TOPMed freeze 5 data

(University of Michigan and NHLBI TOPMed. 2018); (3) Variants were reanalyzed with linear

regression using PLINK 1.9 by applying the arguments --mac 5 --geno 0.1 --hwd 0.0001 using

TOPMed freeze 8 DP10 PASS data.

To determine if the rs73429450 association with FEV_1 was only identifiable using whole genome

sequencing data, we repeated the linear regression association analysis on signals that passed

the genome-wide significance threshold using PLINK 1.9 and genotype data generated with

Axiom Genome-Wide LAT 1 array (Affymetrix, Santa Clara, CA, dbGaP phs000921.v1.p1). These

array genotype data were imputed into the following reference panels: 1000 Genomes phase 3

version 5, Haplotype Reference Consortium (HRC) r1.1, the Consortium on Asthma among

African-ancestry Populations in the Americas (CAAPA) and the TOPMed phase 5 panels on the

Michigan Imputation Server (Das *et al.* 2016). It should be noted that 500 SAGE II subjects were

part of the TOPMed freeze 5 reference panel.

Ninety-eight GWAS FEV_1-associated loci were retrieved from GWAS catalog version 1.0.2-

associations_e93_r2019-01-31 (Buniello *et al.* 2019) using the trait name "Lung function

(FEV_1)". Overlap between these 98 loci with 20 Kb flanking regions and the discovery GWAS

FEV_1 associations (p < 0.001) were detected using the BEDTools intersect tool (Quinlan and Hall.

2010).

Conditional analysis

Conditional analysis was performed to identify all independent signals in a GWAS peak using

PLINK 1.9. All TOPMed freeze 8 DP10 variants within 1 megabase (Mb) of the tag association

signal and with association p-value of 1×10^{-4} or smaller in the discovery GWAS were included

in the analysis. Variants were first ordered by ascending p-value. A variant was considered to be

an independent signal if the association p-value after conditioning (conditional p-value) on the

tag signal was smaller than 0.05. Newly identified independent signals were included with the

tag signal for conditioning on the next variant.

Region-based association analysis

Region-based association analyses were performed in 1 Kb sliding windows with 500 bp

increments in a 1 Mb flanking region of the tag GWAS signal using the SKAT_CommonRare

function from the SKAT R package v1.3.2.1 (Ionita-Laza *et al.* 2013) . Default settings were used

with method = "C" and test.type = "Joint". A minor allele frequency (MAF) threshold of 0.01 was

used as the cutoff to distinguish rare and common variants. Variants were annotated in

TOPMed using the WGSA pipeline (Liu *et al.* 2016). Since SKAT imputes missing genotypes by

default by assigning mean genotype values (impute.method="fixed"), we chose to use low

coverage genotypes instead of SKAT imputation, and hence, TOPMed freeze 8 DP0 variants with

a VCF FILTER of PASS were included in the analysis. The function effectiveSize in the R package

CODA (Plummer *et al.* 2006) was used to estimate the effective number of independent

hypothesis tests for accurate Bonferroni multiple testing corrections. P-value thresholds for

statistical significance and suggestive significance were defined as 0.05 and 1 divided by the

effective number of tests, respectively (Duggal *et al.* 2008). If a region was suggestively

significant, region-based analyses were repeated with functional variants and/or rare variants

(MAF <= 0.01) to assess contribution of common, rare and/or functional variants. Region-based

analyses using rare variants only were performed using SKAT-O (Lee *et al.* 2012). The WGSA

annotation filters used to define functional variants are provided in File S1 (Supplementary Text

1). To study the contribution of individual variants to a region-based association p-value, drop-

one variant analysis was performed by repeating the region-based analysis multiple times and

dropping one variant only at a time.

Functional annotations and prioritization of genetic variants

The Hi-C Unifying Genomic Interrogator (HUGIN) (Ay *et al.* 2014; Martin, J. S. *et al.* 2017;

Schmitt *et al.* 2016)   was used to assign potential gene targets to each variant. ENCODE

annotations (ENCODE Project Consortium 2011; ENCODE Project Consortium 2012) were based

on overlap of the variants with functional data downloaded from the UCSC Table Browser

(Karolchik *et al.* 2004). These data included DNAase I hypersensitivity peak clusters (hg38

wgEncodeRegDnaseClustered table), transcription factor ChIP-Seq clusters (hg38

encRegTfbsClustered table) and histone modification ChIP-Seq peaks (hg19

wgEncodeBroadHistone<cell type><histone>StdPk tables). For DNase I hypersensitivity and

transcription factor binding sites, we focused on blood, bone marrow, lung and embryonic cells.

For histone modification ChIP-Seq, we focused on H3K27ac and H3K4me3 modifications in

human blood (GM12878), bone marrow (K562), lung fibroblast (NHLF), and embryonic stem

cells (H1-hESC). LiftOver tool (Hinrichs *et al.* 2006) was used to convert genomic coordinates

from hg19 to hg38. Candidate cis-regulatory elements (ccREs) were a subset of representative

DNase hypersensitivity sites with epigenetic activity further supported by histone modification

(H3K4me3 and H3K27ac) or CTCF-binding data from the ENCODE project. Overlap of variants

with ccREs were detected using the Search Candidate cis-Regulatory Elements by ENCODE

(SCREEN) web interface (ENCODE Project Consortium 2011; ENCODE Project Consortium 2012).

Prioritization of genetic variants was based on the presence of statistical, functional and/or

bioinformatic evidence as described in the Diverse Convergent Evidence (DiCE) prioritization

framework (Ciesielski *et al.* 2014). The priority score of each variant was obtained by summing

the number of evidence present for that variant.

Replication of GWAS associations

All replication analyses were performed in subjects with asthma. Replication of GWAS FEV_1

associations was attempted on TOPMed whole genome sequencing data generated from four

cohorts. These cohorts included Puerto Rican (n=1,109) and Mexican American (n=649) children

in the Genes-Environments and Admixture in Latino Americans (GALA II) study (Oh *et al.* 2012),

African American adults in the Study of Asthma Phenotypes and Pharmacogenomic Interactions

by Race-Ethnicity (SAPPHIRE, n=3,428) (Levin *et al.* 2014) and African American children in

Genetics of Complex Pediatric Disorders (GCPD-A, n=1,464) study (Ong *et al.* 2013). Age, sex,

height, controller medications and the first 5 PCs were used as covariates.

Additionally, replication of GWAS FEV_1 associations was attempted using data of black UK

Biobank subjects who had asthma (n=627) while adjusting for age, sex, height and the first 5

principal components. Asthma status was defined by ICD code or self-reported asthma. UK

Biobank genotype data was generated on Affymetrix UK BiLEVE axiom or UK Biobank Axiom

array and imputed into the Haplotype Reference Consortium, 1000G and UK 10K projects

(Bycroft *et al.* 2018; Canela-Xandri *et al.* 2018). Additional details on the UK Biobank study and

the replication procedures are available in File S1 (Supplementary Text 2).

RNA sequencing and expression quantitative trait loci (eQTL) analysis

Whole-transcriptome libraries of 370 nasal brushings from GALA II Puerto Rican children with

asthma were constructed by using the Beckman Coulter FX automation system (Beckman

Coulter, Fullerton, CA). Libraries were sequenced with the Illumina HiSeq 2500 system. Raw

RNA-Seq reads were trimmed using Skewer (Jiang *et al.* 2014) and mapped to human reference

genome hg38 using Hisat2 (Kim, D. *et al.* 2015). Reads mapped to genes were counted with

htseq-count and using the UCSC hg38 GTF file as reference (Anders *et al.* 2015). Cis-expression

quantitative trait locus (eQTL) analysis of *KITLG* was performed as described in the Genotype-

Tissue Expression (GTEx) project version 7 protocol (GTEx Consortium *et al.* 2017) using age,

sex, BMI, global African and European ancestries and 60 PEER factors as covariates.

Gene-by-air-pollution interaction analysis

We hypothesized that the effect of genetic variation on lung function in our study population

may differ by the levels of exposure to SO_2 (Neophytou *et al.* 2016). To test for an interaction

between a genetic variant and SO_2, an additional multiplicative interaction term (variant x SO_2

exposure) was included in the original GWAS model (see Method Section "FEV_1 GWAS"). The

SO_2 estimates used in the interaction analysis were first-year, past-year, and lifetime exposure

to ambient of SO_2, which were estimated as described previously (Neophytou *et al.* 2016).

Residuals of FEV_1 were plotted against exposure to SO_2 and stratified by the number of copies

of the minor allele. Residuals of FEV_1 were obtained as described in the Methods Section "FEV_1

GWAS".

Data availability

290   Local institutional review boards approved the studies (IRB# 10-02877). All subjects and legal

291   guardians provided written informed consent. TOPMed whole genome sequencing data are

292   available on dbGaP with the accession number: phs000921.v4.p1. The remaining sequence data

293   are available upon request. Supplemental materials available at figshare.

RESULTS

Novel lung function associations

Subject characteristics of the 867 African American children with asthma included in this study

are shown in Table 1, and the distribution of their FEV_1 values is in Figure S1. The CODA-

adjusted statistical significance threshold (2.10×10^{-8}) was selected as the genome-wide

significance level. According to this threshold, one SNP in chromosome 12 (chr12:88846435,

rs73429450, G>A) was associated with FEV_1.res.rnorm (Figure 1, p = 9.01 x 10^{-9}, β = 0.801). The

association between rs73429450 and lung function remained statistically significant when the

association was repeated using untransformed FEV_1 (p = 1.26 x 10^{-8}, β = 0.302) as the outcome

variable. The association between rs73429450 and lung function was suggestive using

FEV_1.perc.predicted (p = 1.69 x 10^{-7}, β = 0.100). Thirty-nine variants with association p < 0.001

overlapped with 14 out of 98 previously reported FEV_1-associated loci and their 20 Kb flanking

regions (Table S1). When including all variants with association p < 0.05, we found overlap with

all 98 previously reported loci. Our top FEV_1 association, rs73429450, did not overlap with any

previously reported loci in both scenarios and it is a novel association with FEV_1 in this study

population.

Secondary analysis that included covariates correcting for smoking status and number of

smokers in the family showed that smoking-related factors were not significantly associated

with FEV_1 in our pediatric SAGE cohort: using 657 out of 867 individuals with available smoking-

related covariates, the FEV_1.res.rnorm association p-values before and after including the

smoking-related covariates were 2.01 x 10^{-6} and 1.89 x 10^{-6}. Both p-values of the covariates

smoking status (p = 0.27) and number of smokers in the family (p = 0.54) were not significant.

Conditional analysis was performed on 45 variants with association p < 1 x 10^{-4} located within 1

Mb of the strongest association signal (rs73429450). Two weaker independent signals

(rs11312747, rs58475486) were identified (Table S2).

The minor allele frequency of rs73429450 in continental populations from the 1000 Genomes

Project (1000G) is 3% in Africans (AFR) and < 1% in Admixed Americans (AMR), Europeans (EUR)

and Asians (EAS and SAS) (1000 Genomes Project Consortium *et al.* 2015). Rs73429450 was not

included on the Affymetrix LAT1 genotyping array where SAGE participants were previously

genotyped. To determine if the rs73429450 association with FEV_1 was only identifiable using

whole genome sequencing data, we attempted to reproduce our results by imputing the

genotype of rs73429450 in 851 SAGE participants with available array data using 1000G phase 3

(n = 2,504), HRC r1.1 (n = 32,470), CAAPA (n = 883) and TOPMed freeze 5 (n = 62,784) reference

panels. Our results remained statistically significant when using the 1000G phase 3 (p = 4.97 x

10^{-8}, β = 0.79, imputation R^2 = 0.95) and TOPMed freeze 5 (p = 1.22 x 10^{-8}, β = 0.80, imputation

R^2 = 0.98) reference panels, but lost statistical significance when rs73429450 genotypes were

imputed using the HRC (p = 4.35 x 10^{-7}, β = 0.68, imputation R^2 = 0.94) and CAAPA (p = 1.95 x

10^{-7}, β = 0.80, imputation R^2 = 0.71) reference panels.

Region-based association analysis including all variants conditioned on the association signal

from rs73429450 was performed in its 1 Mb flanking region (chr12:87846435-89846435). No

windows were significantly associated after Bonferroni multiple testing correction (p < 2.81 x

10^{-4}, Figure S2), but 20 windows were suggestively associated with FEV_1.res.rnorm (p < 5.62 x

10^{-3}, Table S3). Two of 20 windows re-tested using only functional variants were suggestively

significant (region 4 and 16). Both of these windows were no longer suggestively significant

after removing the common variants, indicating that association signal from these regions was

mostly driven by common variants. Further investigation on region 16 using drop-one analysis

on the 2 rare and 1 common function variants confirmed the major contribution by the

common variant, rs1895710, as shown by the major increase in p-value (Table S4). The signal

was also slightly driven by the singleton, rs990979778. Drop-one analysis was not performed on

region 4 because there were only 1 common and 1 rare variants.

A Hi-C assay couples a chromosome conformation capture (3C) assay with next-generation

sequencing to capture long-range interactions in the genome. We identified a statistically

significant long-range chromatin interaction between the GWAS peak and the KIT ligand (*KITLG*,

also known as stem cell factor, *SCF*) gene in human fetal lung fibroblast cell line IMR90 (Table

S5). The long-range interaction detected in human primary lung tissue was not significant,

implying that the potential long-range interactions are specific to tissue type or developmental

stage.

Potential regulatory role of FEV_1-associated variants on KITLG

expression

To further elucidate potential regulatory relationships between the GWAS association peak and

KITLG, we analyzed whether variants in the peak were eQTL of *KITLG* in previously published

whole blood RNA-Seq data available from the same study participants (Mak, Angel CY *et al.*

2016) . The whole blood RNA-Seq data, however, did not yield evidence of expressed *KITLG*,

consistent with results in GTEx. We subsequently used RNA-Seq data from nasal epithelial cells

of 370 Puerto Rican children with asthma from the GALA II study, and found that five out of 45

variants were eQTL of *KITLG* (Table S6). While Puerto Ricans are a different population than

African Americans, they are both admixed populations with substantial African genetic

ancestry, and therefore could share eQTLs. All five eQTLs corresponded to one signal in a

region with strong linkage disequilibrium ($r^2 > 0.8$, markers 25, 26, 28, 29 and 32 in Figure 2).

Replication of genetic association with FEV$_1$

Subject Characteristics of our four replication cohorts (SAPPHIRE, GCPD-A, UK Biobank and

GALA II) are shown in Table S7. We attempted to replicate the association of the 45 SNPs in our

primary FEV$_1$ GWAS in each cohort. We used 0.05 as the suggestive p-value threshold and

0.0167 as the Bonferroni-corrected p-value threshold after correcting for 3 independent signals

(see conditional analysis in Results Section). A total of 20 variants were replicated at $p < 0.05$

with consistent direction of effect in black UK Biobank participants; 14 variants in SAPPHIRE and

2 variants in GCPD-A were significant but had an opposite direction of effect (Table S8).

We attempted to replicate the FEV$_1$.res.rnorm association in Mexican American (n = 649) and

Puerto Rican (n = 1,109) children with asthma from the GALA II study. In Mexican Americans,

we excluded 19 variants with MAF < 0.1% and associations for the remaining 26 variants did not

replicate (Table S9). In Puerto Ricans, we observed the same protective effect in 38 of the 45

variants in the locus, but the associations were not statistically significant (Table S9).

Incorporating statistical and functional evidence for candidate variant

prioritization

We combined and summarized all functional evidence for the top 45 variants, along with eQTL

findings from nasal epithelial RNA-Seq and replication results (Figure 2, Table 2). To facilitate

interpretation of the variant association with FEV$_1$, the effect sizes and p-values of both FEV$_1$ (β

and p) and FEV$_1$.res.rnorm (β_{norm} and p$_{norm}$) associations are reported in Table 2. CADD

functional prediction score and ENCODE histone modification ChIP-Seq peaks in embryonic,

blood, bone marrow, and lung-related tissues were also examined but not reported because

none of the variants had a CADD score greater than 10 and none overlapped with histone

modification sites. After summing up the number of evidence present for each variant to obtain

its priority score, rs73440122 received the highest functional score of 3 based on replication in

the UK Biobank, overlap with a DNase I hypersensitivity site in B-lymphoblastoid cells

(GM12865) and overlap with an SPI1 binding site in acute promyelocytic leukemia cells. Eight

other variants were prioritized with score > 2 or evidence of being an eQTL for *KITLG* in nasal

epithelial cells (Table 2, score marked with * or # respectively). These nine candidate variants

were selected for gene-by-air-pollution interaction analyses.

Gene-by-air-pollution interaction of rs58475486

We previously found that first year of life and lifetime exposure to SO$_2$
were associated with FEV$_1$ in African American children (Neophytou
et al. 2016). We investigated whether the effect of the nine prioritized
genetic variants associated with lung function varied by SO$_2$ exposure
(first year of life, past year, and lifetime exposure). Since the nine
variants represent three independent signals (see conditional
analysis in the Data availability

Local institutional review boards approved the studies (IRB# 10-02877). All subjects and legal

guardians provided written informed consent. TOPMed whole genome sequencing data are

available on dbGaP with the accession number: phs000921.v4.p1. The remaining sequence data

are available upon request. Supplemental materials available at figshare.

R Section), the Bonferroni-corrected p-value threshold was set to p = 0.0056 (correction for

nine tests; three signals and three exposure periods to SO_2). We observed a single statistically

significant interaction between the T allele of rs58475486 and past year exposure to SO_2 that

was positively associated with FEV_1 (p = 0.003, β = 0.32, Table 3, Figure 3A). Interestingly, six of

the remaining eight variants also displayed interaction effects with past year exposure to SO_2

that were suggestively associated (p < 0.05) with FEV_1 (Table 3). We also found a suggestive

interaction of the C allele of rs73440122 with first year exposure to SO_2 that was associated

with decreased FEV_1 (p = 0.045, β = -0.32, Figure 3B).

DISCUSSION

Variant rs73429450 (MAF = 0.030) was identified as the strongest association signal with FEV_1.

Each additional copy of the protective A allele of rs73429450 was associated with a 0.3 L

increase of FEV_1. We did not find any statistically significant contribution of rare variants to the

association signal from a 1 Kb sliding window analyses in the 1 MB flanking region centered on

rs73429450. We were surprised to identify a novel common variant (MAF = 0.030) associated

with lung function using whole genome sequence data in a population that was previously

analyzed for associations with lung function using genotype array data. Further investigation

revealed that our discovered variant, rs73429450, was not captured by the LAT 1 genotyping

array, and the association with lung function depended on the reference panel used to impute

the variant into our population. More surprisingly, our statistically significant finding was only

found to be suggestively significant using data imputed from the CAAPA reference panel (p =

1.95×10^{-7}, β = 0.80). Of the imputation reference panels that we assessed, CAAPA is one of the

more relevant reference panels for our study population because it is based on African

populations in the Americas. However, we note that the effect size estimated from CAAPA-

imputed data was comparable to that generated from WGS data. While whole genome

sequencing data is usually praised for enabling analysis of rare-variant contributions to

phenotype variability, our results show the utility of whole genome sequencing data for the

reliable analysis of common variants as well in the absence of relevant imputation panels.

Although rs73429450 had the lowest p-value from our whole genome sequencing association

analysis, we did not find the required amount of functional evidence to prioritize this marker

for inclusion in downstream gene-by-air-pollution analyses. Another variant, rs73440122, was

in moderate to strong linkage disequilibrium (r^2 = 0.76) with rs7349450 and had a similar MAF

(0.027) in our study population, but was only suggestively associated with FEV_1 in our

association analysis (p = 2.08×10^{-7}, Table2). In contrast to rs7349450, there were multiple lines

of evidence suggesting the functional relevance of rs73440122: rs73440122 received the

highest priority score, its association with FEV_1 replicated in black UK Biobank participants, and

it was one of the most likely drivers of FEV_1 variability among individuals, possibly mediated

through *KITLG*.

Bioinformatic interrogation of rs73440122 revealed that the variant overlapped with a ccRE

(SCREEN accession EH37E0279310), DNase I hypersensitivity site, and SPI1 ChIP-Seq clusters

that were indicative of a candidate open chromatin gene regulatory region. The transcription

factor SPI1, which is expressed in lung, spleen and whole blood, was previously linked to fibrotic

diseases and tissues (Wohlfahrt *et al.* 2019) and *KITLG* in erythroid progenitor cells (Quang *et al.*

1995), but there is no existing evidence of its role in lung function. Although rs73440122 itself is

not an eQTL of *KITLG*, it is located in a region that physically interacted with *KITLG* based on Hi-

C data in fetal lung fibroblast cells. Additionally, five neighboring FEV_1 associated variants were

identified as eQTLs of *KITLG*, although they also appeared to be an independent signal ($r^2 < 0.2$).

Overall, these results support regulatory interactions between this region and *KITLG*.

KITLG, a ligand of the KIT tyrosine kinase receptor, is expressed in lung and its decreased

expression was previously observed in patients with chronic obstructive pulmonary disease

(Bhattacharya *et al.* 2009). Inactivation of KIT signaling led to airspace enlargement and

contributed to declining lung function in mice (Lindsey *et al.* 2011). Polymorphisms of *KITLG*

have been previously associated with susceptibility to moderate-to-severe bronchopulmonary

dysplasia, a chronic inflammatory lung disease that affects preterm infants (Huusko *et al.* 2015).

These lines of evidence support the potential contribution of our novel locus to lung function

phenotype variation mediated through *KITLG*, especially in patients with inflammatory

conditions like asthma.

Gene-by-environment interactions likely account for a portion of the "missing" heritability of

many complex phenotypes (Moore and Williams. 2009). We hypothesized that a significant

portion of the heritability of lung function was due in part to gene-by-air-pollution (SO_2)

interaction effects. The interaction between rs58475486 and past year exposure to SO_2 that

was significantly associated with lung function supports our hypothesis. The T allele of

rs58475486 is common (8-14%) in African populations and showed a protective effect on lung

function in the presence of past year SO_2 exposure. SNP rs58475486 is located in a ccRE

(SCREEN accession EH37E0279296) and a *FOXA1* binding site in the A549 lung adenocarcinoma

cell line. *Foxa1* has a known compensatory role with *Foxa2* during lung morphogenesis in mice

(Wan *et al.* 2005). Deletion of both *Foxa1* and *Foxa2* inhibited cell proliferation, epithelial cell

differentiation, and branching morphogenesis in fetal lung tissue. Further functional validation

on the effect of rs58475486 on binding affinity of FOXA1 is necessary to confirm whether the

role of *FOXA1* in this ccRE is important for *KITLG* regulatory and lung function.

The higher frequency of the protective alleles of both rs73440122 and rs58465486 in African

populations appears to contradict previous findings that African ancestry was associated with

lower lung function (Kumar *et al.* 2010). One possible explanation for this seeming

inconsistency is that FEV_1 is a complex trait whose variation is influenced by many genetic

variants of small to moderate effect sizes whose influence on lung function may vary by

exposure to environmental factors. We found suggestive evidence that the interaction between

rs73440122 and first year exposure to SO_2 reverses the positive association of rs73440122 with

lung function to a negative one (Table 3). When assessed independently, our genetic

association analysis showed that the protective A allele of rs73440122 was associated with

higher lung function. However, with increasing levels of SO_2 exposure in the first year of life,

increasing copies of the A allele of rs73440122 were associated with decreased lung function.

Air pollution is known to negatively impact lung function, and we have previously shown that

the deleterious effects of air pollution on lung phenotypes may be significantly increased in

African American children compared to other populations experiencing the same amount of

exposure (Nishimura *et al.* 2013). It has also been reported that Latino and African American

populations often live in neighborhoods with high levels of air pollution (Mott 1995). The

increased susceptibility to negative pulmonary effects from air pollution exposure coupled with

the disproportionate exposure to air pollution experienced by the African American population

may also contribute to the lower lung function seen in this population despite the presence of

protective alleles.

One limitation of this study is that the FEV_1 genetic association and the eQTL analyses with

KITLG were performed in different populations due to data availability constraints. Although we

did not have RNA-Seq data from lung tissues from our study subjects, we previously

demonstrated that there is a high degree of overlap in gene expression profiles between nasal

and bronchial epithelial cells (Poole *et al.* 2014). The direction of effect of the association was

the same in GALA II Puerto Rican children with asthma but not statistically significant. This may

in part due to the significantly lower African Ancestry in Puerto Ricans compared to African

Americans.

We replicated 20 of 45 variants in black UK Biobank subjects and observed conflicting "flip-flop"

associations in African Americans from the SAPPHIRE and GCPD-A studies. In the past, flip-flop

associations were deemed as spurious results. Traditional association testing approach studies

the effect of each variant on phenotype independently and increases the chance of flip-flop

associations detected between studies. Differences in study design, sampling variation that

leads to variation in LD patterns, and lack of consideration of other disease influencing genetic

and/or environmental factors are all potential causes of flip-flop associations (Kraft *et al.* 2009;

Lin *et al.* 2007). Hence, it is not surprising to observe flip-flop associations when gene and

environment interactions were detected at our FEV_1 GWAS locus. It was previously shown that

flip-flop associations can occur between and within populations even in the presence of a

genuine genetic effect (Kraft *et al.* 2009; Lin *et al.* 2007). Further functional analysis is thus

required to validate the relationship between the candidate variants, *KITLG* and FEV_1. This may

include reporter assays to validate potential enhancer or repressor activity and CRISPR-based

editing assays to validate the regulatory role of the candidate variants on *KITLG*. Although

literature exists describing KIT signaling for lung function in mice (Lindsey *et al.* 2011),

additional knockout experiments in a model animal system may be necessary to confirm the

causal effect of *KITLG* on lung function.

The average concentration of ambient SO_2 exposure in our participants (Table 1) was lower

than the National Ambient Air Quality Standards. It is possible that SO_2 acted as a surrogate for

other unmeasured toxic pollutants emitted from local point sources. Major sources of SO_2 in

San Francisco Bay Area during the recruitment years of 2006 to 2011 include airports,

petroleum refineries, gas and oil plants, calcined petroleum coke plants, electric power plants,

cement manufacturing factories, chemical plants, and landfills (United States Environmental

Protection Agency 2008; United States Environmental Protection Agency 2011). The

Environmental Protection Agency's national emissions inventory data also showed that these

facilities emit Volatile Organic Compounds, heavy metals (lead, mercury, chromium, arsenic),

formaldehyde, ethyl benzene, acrolein, 1,3-butadiene, 1,4-dichlorobenzene, and

tetrachloroethylene into the air along with SO_2. These chemicals are highly toxic and inhaling

even a small amount may contribute to poor lung function. Another possibility is that exposure
to SO_2 captured unmeasured confounding socioeconomic factors.

This study identified a novel protective allele for lung function in African American children with
asthma. The protective association with lung function intensified with increased past year
exposure to SO_2. Our findings showcases the complexity of the relationship between genetic
and environmental factors impacting variation in FEV_1, highlights the utility of WGS data for
genetic research of complex phenotypes, and underscores the importance of including diverse
study populations in our exploration of the genetic architecture underlying lung function.

ACKNOWLEDGEMENTS

The Genes-Environments and Admixture in Latino Americans (GALA II) Study, the Study of
African Americans, Asthma, Genes and Environments (SAGE) Study and E.G.B. were supported
by the Sandler Family Foundation, the American Asthma Foundation, the RWJF Amos Medical
Faculty Development Program, the Harry Wm. and Diana V. Hind Distinguished Professor in
Pharmaceutical Sciences II, the National Heart, Lung, and Blood Institute (NHLBI)
[R01HL117004, R01HL128439, R01HL135156, X01HL134589]; the National Institute of
Environmental Health Sciences [R01ES015794]; the National Institute on Minority Health and
Health Disparities (NIMHD) [P60MD006902, R01MD010443], the National Human Genome
Research Institute [U01HG009080] and the Tobacco-Related Disease Research Program [24RT-
0025]. MJW was supported by the NHLBI [K01HL140218]. JJ and BEH were supported by the
NHLBI [R01HL133433, R01HL141992]. KLK was supported by the NHLBI [R01HL135156-S1], the
UCSF Bakar Institute, the Gordon and Betty Moore Foundation [GBMF3834], and the Alfred P.

Sloan Foundation [2013-10-27] grant to UC Berkeley through the Moore-Sloan Data Science

Environment Initiative. ACW was supported by the Eunice Kennedy Shriver National Institute of

Child Health and Human Development [1R01HD085993-01].

The SAPPHIRE study was supported by the Fund for Henry Ford Hospital, the American Asthma

Foundation, the NHLBI [R01HL118267, R01HL141485, X01HL134589], the National Institute of

Allergy and Infectious Diseases [R01AI079139], and the National Institute of Diabetes and

Digestive and Kidney Diseases [R01DK113003].

The GCPD-A study was supported by an Institutional award from the Children's Hospital of

Philadelphia and by the NHLBI [X01HL134589].

Part of this research was conducted using the UK Biobank Resource under Application Number

40375. We would like to thank UK Biobank participants and researchers who contributed or

collected data.

Whole genome sequencing (WGS) for the Trans-Omics in Precision Medicine (TOPMed)

program was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for

"NHLBI TOPMed: Gene-Environment, Admixture and Latino Asthmatics Study" (phs000920) and

"NHLBI TOPMed: Study of African Americans, Asthma, Genes and Environments" (phs000921)

was performed at the New York Genome Center (3R01HL117004-02S3) and the University of

Washington Northwest Genomics Center (HHSN268201600032I). WGS for "NHLBI TOPMed:

Study of Asthma Phenotypes & Pharmacogenomic Interactions by Race-Ethnicity" (phs001467)

and "Genetics of Complex Pediatric Disorders - Asthma" (phs001661) was performed at the

University of Washington Northwest Genomics Center (HHSN268201600032I). Centralized read

mapping and genotype calling, along with variant quality metrics and filtering were provided by

the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract

HHSN268201800002I). Phenotype harmonization, data management, sample-identity QC, and

general study coordination were provided by the TOPMed Data Coordinating Center (3R01HL-

120393-02S1; contract HHSN268201800001I). We gratefully acknowledge the studies and

participants who provided biological samples and data for TOPMed.

WGS of part of GALA II was performed by New York Genome Center under The Centers for

Common Disease Genomics of the Genome Sequencing Program (GSP) Grant (UM1 HG008901).

The GSP Coordinating Center (U24 HG008956) contributed to cross-program scientific initiatives

and provided logistical and general study coordination. GSP is funded by the National Human

Genome Research Institute, the National Heart, Lung, and Blood Institute, and the National Eye

Institute.

The TOPMed imputation panel was supported by the NHLBI and TOPMed study investigators

who contributed data to the reference panel. The panel was constructed and implemented by

the TOPMed Informatics Research Center at the University of Michigan (3R01HL-117626-02S1;

contract HHSN268201800002I). The TOPMed Data Coordinating Center (3R01HL-120393-02S1;

contract HHSN268201800001I) provided additional data management, sample identity checks,

and overall program coordination and support. We gratefully acknowledge the studies and

participants who provided biological samples and data for TOPMed.

The authors wish to acknowledge the following GALA II and SAGE study collaborators: Shannon

Thyne, UCSF; Harold J. Farber, Texas Children's Hospital; Denise Serebrisky, Jacobi Medical

Center; Rajesh Kumar, Lurie Children's Hospital of Chicago; Emerita Brigino-Buenaventura,

Kaiser Permanente; Michael A. LeNoir, Bay Area Pediatrics; Kelley Meade, UCSF Benioff

Children's Hospital, Oakland; William Rodríguez-Cintrón, VA Hospital, Puerto Rico; Pedro C.

Ávila, Northwestern University; Jose R. Rodríguez-Santana, Centro de Neumología Pediátrica;

Luisa N. Borrell, City University of New York; Adam Davis, UCSF Benioff Children's Hospital, Oakland;

Saunak Sen, University of Tennessee.

The authors acknowledge the families and patients for their participation and thank the

numerous health care providers and community clinics for their support and participation in

GALA II and SAGE. In particular, the authors thank the recruiters who obtained the data:

Duanny Alva, MD; Gaby Ayala-Rodríguez; Lisa Caine, RT; Elizabeth Castellanos; Jaime Colón;

Denise DeJesus; Blanca López; Brenda López, MD; Louis Martos; Vivian Medina; Juana Olivo;

Mario Peralta; Esther Pomares, MD; Jihan Quraishi; Johanna Rodríguez; Shahdad Saeedi; Dean

Soto; and Ana Taveras.

The authors thank María Pino-Yanes for providing feedback on this study and Thomas W

Blackwell for providing critical review on this manuscript.

The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health.

LITERATURE CITED

1000 Genomes Project Consortium, A. Auton, L. D. Brooks, R. M. Durbin, E. P. Garrison *et al*,

2015 A global reference for human genetic variation. Nature **526:** 68-74.

Akinbami, L. J., 2015 *Asthma Prevalence, Health Care use and Mortality: United States, 2003-05.*

[Online] Available at: http://www.cdc.gov/nchs/data/hestat/asthma03-05/asthma03-05.htm.

[Accessed 2020 Jan 8].

Akinbami, L. J., J. E. Moorman, A. E. Simon and K. C. Schoendorf, 2014 Trends in racial

disparities for asthma outcomes among children 0 to 17 years, 2001-2010. J. Allergy Clin.

Immunol. **134:** 547-553.e5.

Alexander, D. H., J. Novembre and K. Lange, 2009 Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. **19:** 1655-1664.

Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq--a python framework to work with high-

throughput sequencing data. Bioinformatics **31:** 166-169.

Ay, F., T. L. Bailey and W. S. Noble, 2014 Statistical confidence estimation for hi-C data reveals

regulatory chromatin contacts. Genome Res. **24:** 999-1011.

Barraza-Villarreal, A., J. Sunyer, L. Hernandez-Cadena, M. C. Escamilla-Nunez, J. J. Sienra-Monge

et al, 2008 Air pollution, airway inflammation, and lung function in a cohort study of mexico city

schoolchildren. Environ. Health Perspect. **116:** 832-838.

Bhattacharya, S., S. Srisuma, D. L. Demeo, S. D. Shapiro, R. Bueno *et al*, 2009 Molecular

biomarkers for quantitative and discrete COPD phenotypes. Am. J. Respir. Cell Mol. Biol. **40:**

359-367.

Brunekreef, B., and S. T. Holgate, 2002 Air pollution and health. Lancet **360:** 1233-1242.

Buniello, A., J. A. L. MacArthur, M. Cerezo, L. W. Harris, J. Hayhurst *et al*, 2019 The NHGRI-EBI

GWAS catalog of published genome-wide association studies, targeted arrays and summary

statistics 2019. Nucleic Acids Res. **47:** D1005-D1012.

Bycroft, C., C. Freeman, D. Petkova, G. Band, L. T. Elliott *et al*, 2018 The UK biobank resource

with deep phenotyping and genomic data. Nature **562:** 203-209.

Canela-Xandri, O., K. Rawlik and A. Tenesa, 2018 An atlas of genetic associations in UK biobank.

Nat. Genet. **50:** 1593-1599.

Carlson, C. S., T. C. Matise, K. E. North, C. A. Haiman, M. D. Fesinmeyer *et al*, 2013

Generalization and dilution of association results from european GWAS in populations of non-

european ancestry: The PAGE study. PLoS Biol. **11:** e1001661.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al*, 2015 Second-generation

PLINK: Rising to the challenge of larger and richer datasets. Gigascience **4:** 7-8. eCollection

2015.

Chatterjee, S., and N. Das, 1995 Lung function in indian twin children: Comparison of genetic

versus environmental influence. Ann. Hum. Biol. **22:** 289-303.

Ciesielski, T. H., S. A. Pendergrass, M. J. White, N. Kodaman, R. S. Sobota *et al*, 2014 Diverse

convergent evidence in the genetic analysis of complex disease: Coordinating omic, informatic,

and experimental evidence to better identify and validate risk factors. BioData Min. **7:** 10-10.

eCollection 2014.

Conomos, M. P., M. B. Miller and T. A. Thornton, 2015 Robust inference of population structure

for ancestry prediction and correction of stratification in the presence of relatedness. Genet.

Epidemiol. **39:** 276-293.

Conomos, M. P., A. P. Reiner, B. S. Weir and T. A. Thornton, 2016 Model-free estimation of

recent genetic relatedness. Am. J. Hum. Genet. **98:** 127-148.

Das, S., L. Forer, S. Schonherr, C. Sidore, A. E. Locke *et al*, 2016 Next-generation genotype

imputation service and methods. Nat. Genet. **48:** 1284-1287.

Duggal, P., E. M. Gillanders, T. N. Holmes and J. E. Bailey-Wilson, 2008 Establishing an adjusted

p-value threshold to control the family-wide type 1 error in genome wide association studies.

BMC Genomics **9:** 516-516.

ENCODE Project Consortium, 2012 An integrated encyclopedia of DNA elements in the human

genome. Nature **489:** 57-74.

ENCODE Project Consortium, 2011 A user's guide to the encyclopedia of DNA elements

(ENCODE). PLoS Biol. **9:** e1001046.

GTEx Consortium, Laboratory, Data Analysis &Coordinating Center (LDACC)-Analysis Working

Group, Statistical Methods groups-Analysis Working Group, Enhancing GTEx (eGTEx) groups,

NIH Common Fund *et al*, 2017 Genetic effects on gene expression across human tissues. Nature

550: 204-213.

Hankinson, J. L., J. R. Odencrantz and K. B. Fedan, 1999 Spirometric reference values from a

sample of the general U.S. population. Am. J. Respir. Crit. Care Med. **159:** 179-187.

Hinrichs, A. S., D. Karolchik, R. Baertsch, G. P. Barber, G. Bejerano *et al*, 2006 The UCSC genome

browser database: Update 2006. Nucleic Acids Res. **34:** 590.

Hukkinen, M., J. Kaprio, U. Broms, A. Viljanen, D. Kotz *et al*, 2011 Heritability of lung function: A

twin study among never-smoking elderly women. Twin Res. Hum. Genet. **14:** 401-407.

Huusko, J. M., M. Mahlman, M. K. Karjalainen, T. Kaukola, R. Haataja *et al*, 2015 Polymorphisms

of the gene encoding kit ligand are associated with bronchopulmonary dysplasia. Pediatr.

Pulmonol. **50:** 260-270.

Ierodiakonou, D., A. Zanobetti, B. A. Coull, S. Melly, D. S. Postma *et al*, 2016 Ambient air

pollution, lung function, and airway responsiveness in asthmatic children. J. Allergy Clin.

Immunol. **137:** 390-399.

Ionita-Laza, I., S. Lee, V. Makarov, J. D. Buxbaum and X. Lin, 2013 Sequence kernel association

tests for the combined effect of rare and common variants. Am. J. Hum. Genet. **92:** 841-853.

Jiang, H., R. Lei, S. W. Ding and S. Zhu, 2014 Skewer: A fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. BMC Bioinformatics **15:** 182-182.

Johnson, J. D., and W. M. Theurer, 2014 A stepwise approach to the interpretation of

pulmonary function tests. Am. Fam. Physician **89:** 359-366.

Karolchik, D., A. S. Hinrichs, T. S. Furey, K. M. Roskin, C. W. Sugnet *et al*, 2004 The UCSC table

browser data retrieval tool. Nucleic Acids Res. **32:** 493.

Kim, D., B. Langmead and S. L. Salzberg, 2015 HISAT: A fast spliced aligner with low memory

requirements. Nat. Methods **12:** 357-360.

Kim, M. S., K. P. Patel, A. K. Teng, A. J. Berens and J. Lachance, 2018 Genetic disease risks can be

misestimated across global populations. Genome Biol. **19:** 179-7.

Kraft, P., E. Zeggini and J. P. Ioannidis, 2009 Replication in genome-wide association studies.

Stat. Sci. **24:** 561-573.

Kumar, R., M. A. Seibold, M. C. Aldrich, L. K. Williams, A. P. Reiner *et al*, 2010 Genetic ancestry

in lung-function predictions. N. Engl. J. Med. **363:** 321-330.

Lee, S., M. J. Emond, M. J. Bamshad, K. C. Barnes, M. J. Rieder *et al*, 2012 Optimal unified

approach for rare-variant association testing with application to small-sample case-control

whole-exome sequencing studies. Am. J. Hum. Genet. **91:** 224-237.

Levin, A. M., Y. Wang, K. E. Wells, B. Padhukasahasram, J. J. Yang *et al*, 2014 Nocturnal asthma

and the importance of race/ethnicity and genetic ancestry. Am. J. Respir. Crit. Care Med. **190:**

266-273.

Li, X., G. A. Hawkins, E. J. Ampleford, W. C. Moore, H. Li *et al*, 2013 Genome-wide association

study identifies TH1 pathway genes associated with lung function in asthmatic patients. J.

Allergy Clin. Immunol. **132:** 313-20.e15.

Liao, S. Y., X. Lin and D. C. Christiani, 2014 Genome-wide association and network analysis of

lung function in the framingham heart study. Genet. Epidemiol. **38:** 572-578.

Lin, P. I., J. M. Vance, M. A. Pericak-Vance and E. R. Martin, 2007 No gene is an island: The flip-

flop phenomenon. Am. J. Hum. Genet. **80:** 531-538.

Lindsey, J. Y., K. Ganguly, D. M. Brass, Z. Li, E. N. Potts *et al*, 2011 C-kit is essential for alveolar

maintenance and protection from emphysema-like disease in mice. Am. J. Respir. Crit. Care

Med. **183:** 1644-1652.

Liu, X., S. White, B. Peng, A. D. Johnson, J. A. Brody *et al*, 2016 WGSA: An annotation pipeline

for human genome sequencing studies. J. Med. Genet. **53:** 111-112.

Mak, A. C. Y., M. J. White, W. L. Eckalbar, Z. A. Szpiech, S. S. Oh *et al*, 2018 Whole-genome

sequencing of pharmacogenetic drug response in racially diverse children with asthma. Am. J.

Respir. Crit. Care Med. **197:** 1552-1564.

Mak, A. C., M. J. White, C. Eng, D. Hu, S. Huntsman *et al*, 2016 *Whole Genome Sequencing to*

Identify Genetic Variation Associated with Bronchodilator Response in Minority Children with

Asthma.

Martin, A. R., C. R. Gignoux, R. K. Walters, G. L. Wojcik, B. M. Neale *et al*, 2017 Human

demographic history impacts genetic risk prediction across diverse populations. Am. J. Hum.

Genet. **100:** 635-649.

Martin, J. S., Z. Xu, A. P. Reiner, K. L. Mohlke, P. Sullivan *et al*, 2017 HUGIn: Hi-C unifying

genomic interrogator. Bioinformatics **33:** 3793-3795.

Moore, J. H., 2005 A global view of epistasis. Nat. Genet. **37:** 13-14.

Moore, J. H., and S. M. Williams, 2009 Epistasis and its implications for personal genetics. Am. J.

Hum. Genet. **85:** 309-320.

Mott, L., 1995 The disproportionate impact of environmental health threats on children of

color. Environ. Health Perspect. **103 Suppl 6:** 33-35.

Neophytou, A. M., M. J. White, S. S. Oh, N. Thakur, J. M. Galanter *et al*, 2016 Air pollution and

lung function in minority youth with asthma in the GALA II (genes-environments and admixture

in latino americans) and SAGE II (study of african americans, asthma, genes, and environments)

studies. Am. J. Respir. Crit. Care Med. **193:** 1271-1280.

Nishimura, K. K., J. M. Galanter, L. A. Roth, S. S. Oh, N. Thakur *et al*, 2013 Early-life air pollution

and asthma risk in minority children. the GALA II and SAGE II studies. Am. J. Respir. Crit. Care

Med. **188:** 309-318.

Oh, S. S., M. J. White, C. R. Gignoux and E. G. Burchard, 2016 Making precision medicine socially

precise. take a deep breath. Am. J. Respir. Crit. Care Med. **193:** 348-350.

Oh, S. S., H. Tcheurekdjian, L. A. Roth, E. A. Nguyen, S. Sen *et al*, 2012 Effect of secondhand

smoke on asthma control among black and latino children. J. Allergy Clin. Immunol. **129:** 1478-

83.e7.

Ong, B. A., J. Li, J. M. McDonough, Z. Wei, C. Kim *et al*, 2013 Gene network analysis in a

pediatric cohort identifies novel lung function genes. PLoS One **8:** e72899.

Palmer, L. J., M. W. Knuiman, M. L. Divitini, P. R. Burton, A. L. James *et al*, 2001 Familial

aggregation and heritability of adult lung function: Results from the busselton health study. Eur.

Respir. J. **17:** 696-702.

Pe'er, I., R. Yelensky, D. Altshuler and M. J. Daly, 2008 Estimation of the multiple testing burden

for genomewide association studies of nearly all common variants. Genet. Epidemiol. **32:** 381-

385.

Pino-Yanes, M., N. Thakur, C. R. Gignoux, J. M. Galanter, L. A. Roth *et al*, 2015 Genetic ancestry

influences asthma susceptibility and lung function among latinos. J. Allergy Clin. Immunol. **135:**

228-235.

Plummer, M., N. Best, K. Cowles and K. Vines, 2006 CODA: Convergence diagnosis and output

analysis for MCMC. R News **6:** 7-11.

Poole, A., C. Urbanek, C. Eng, J. Schageman, S. Jacobson *et al*, 2014 Dissecting childhood asthma

with nasal transcriptomics distinguishes subphenotypes of disease. J. Allergy Clin. Immunol.

133: 670-8.e12.

Pruim, R. J., R. P. Welch, S. Sanna, T. M. Teslovich, P. S. Chines *et al*, 2010 LocusZoom: Regional

visualization of genome-wide association scan results. Bioinformatics **26:** 2336-2337.

Purcell, S., and C. Chang, 2013 *Plink 1.9*. [Online] Available at: www.cog-

genomics.org/plink/1.9/. [Accessed 2019 Mar].

Quang, C. T., M. Pironin, M. von Lindern, H. Beug and J. Ghysdael, 1995 Spi-1 and mutant p53

regulate different aspects of the proliferation and differentiation control of primary erythroid

progenitors. Oncogene **11:** 1229-1239.

Quinlan, A. R., and I. M. Hall, 2010 BEDTools: A flexible suite of utilities for comparing genomic

features. Bioinformatics **26:** 841-842.

Repapi, E., I. Sayers, L. V. Wain, P. R. Burton, T. Johnson *et al*, 2010 Genome-wide association

study identifies five loci associated with lung function. Nat. Genet. **42:** 36-44.

Schmitt, A. D., M. Hu, I. Jung, Z. Xu, Y. Qiu *et al*, 2016 A compendium of chromatin contact maps

reveals spatially active regions in the human genome. Cell. Rep. **17:** 2042-2059.

Sillanpaa, E., S. Sipila, T. Tormakangas, J. Kaprio and T. Rantanen, 2017 Genetic and

environmental effects on telomere length and lung function: A twin study. J. Gerontol. A Biol.

Sci. Med. Sci. **72:** 1561-1568.

Sofer, T., X. Zheng, S. M. Gogarten, C. A. Laurie, K. Grinde *et al*, 2019 A fully adjusted two-stage

procedure for rank-normalization in genetic association studies. Genet. Epidemiol. **43:** 263-275.

Soler Artigas, M., L. V. Wain, S. Miller, A. K. Kheirallah, J. E. Huffman *et al*, 2015 Sixteen new

lung function signals identified through 1000 genomes project reference panel imputation. Nat.

Commun. **6:** 8658.

Soler Artigas, M., D. W. Loth, L. V. Wain, S. A. Gharib, M. Obeidat *et al*, 2011 Genome-wide

association and large-scale follow up identifies 16 new loci influencing lung function. Nat.

Genet. **43:** 1082-1090.

Summer Institute in Statistical Genetics, 2019 *PC-Relate*. [Online] Available at: https://uw-

gac.github.io/SISG_2019/pc-relate.html. [Accessed 2019 Jul 25].

Taliun, D., D. N. Harris, M. D. Kessler, J. Carlson, Z. A. Szpiech *et al*, 2019 Sequencing of 53,831

diverse genomes from the NHLBI TOPMed program. bioRxiv 563866.

Tian, X., C. Xu, Y. Wu, J. Sun, H. Duan *et al*, 2017 Genetic and environmental influences on

pulmonary function and muscle strength: The chinese twin study of aging. Twin Res. Hum.

Genet. **20:** 53-59.

TOPMed, 2019 *TOPMed Whole Gneome Sequencing Methods: Freeze 8*. [Online] Available at:

https://www.nhlbiwgs.org/topmed-whole-genome-sequencing-methods-freeze-8. [Accessed

2019 Dec 13].

United States Environmental Protection Agency, 2011 *National Emissions Inventory (NEI) 2011*

Data. [Online] Available at: https://www.epa.gov/air-emissions-inventories/2011-national-

emissions-inventory-nei-data. [Accessed 2020 Jan 8].

United States Environmental Protection Agency, 2008 *National Emissions Inventory (NEI) 2008*

Data. [Online] Available at: https://www.epa.gov/air-emissions-inventories/2008-national-

emissions-inventory-nei-data. [Accessed 2020 Jan 8].

University of Michigan, and NHLBI TOPMed, 2018 *BRAVO Variant Browser*. [Online] Available

at: https://bravo.sph.umich.edu/freeze5/hg38/. [Accessed 2019 Aug].

Wain, L. V., N. Shrine, M. S. Artigas, A. M. Erzurumluoglu, B. Noyvert *et al*, 2017 Genome-wide

association analyses for lung function and chronic obstructive pulmonary disease identify new

loci and potential druggable targets. Nat. Genet. **49:** 416-425.

Wan, H., S. Dingle, Y. Xu, V. Besnard, K. H. Kaestner *et al*, 2005 Compensatory roles of Foxa1

and Foxa2 during lung morphogenesis. J. Biol. Chem. **280:** 13809-13816.

White, M. J., O. Risse-Adams, P. Goddard, M. G. Contreras, J. Adams *et al*, 2016 Novel genetic

risk factors for asthma in african american children: Precision medicine and the SAGE II study.

Immunogenetics **68:** 391-400.

Wise, J., 2019 Air pollution is linked to infant deaths and reduced lung function in children. BMJ

366: l5772.

Wohlfahrt, T., S. Rauber, S. Uebe, M. Luber, A. Soare *et al*, 2019 PU.1 controls fibroblast

polarization and tissue fibrosis. Nature **566:** 344-349.

Wojcik, G. L., M. Graff, K. K. Nishimura, R. Tao, J. Haessler *et al*, 2019 Genetic analyses of

diverse populations improves discovery for complex traits. Nature **570:** 514-518.

World Health Organization, 2017 *Asthma*. [Online] Available at:

http://www.who.int/mediacentre/factsheets/fs307/en/. [Accessed 2020 Jan 8].

Yamada, H., Y. Yatagai, H. Masuko, T. Sakamoto, H. Iijima *et al*, 2015 Heritability of pulmonary

function estimated from genome-wide SNPs in healthy japanese adults. Respir. Investig. **53:** 60-

67.

Zhang, F., and J. R. Lupski, 2015 Non-coding genetic variants in human disease. Hum. Mol.

Genet. **24:** 102.

Figure 1.

A.

B.

Figure 1. Manhattan and LocusZoom plots from genome-wide association study of lung function*. (A) Manhattan plot from genome-wide association study of lung function* using linear regression in ENCORE. Red horizontal line: CODA-adjusted genome-wide significance p-value of 2.10×10^{-8}. Blue horizontal line: CODA-adjusted suggestive significance p-value of 4.19×10^{-7}. (B) LocusZoom plot of rs73429450 (chr12 : 88846435) and 500 Kb flanking region. Colors show linkage disequilibrium in the study population. * FEV_1.res.rnorm was used as the phenotype for the association testing.

Figure 2.

Figure 2. Integration of statistical and functional evidence for variant prioritization. Numbers and different shades of black in the LD plot represent LD in R^2. The three independent signals identified in the conditional analysis are marked with *. Nasal eQTL, variants eQTL of *KITLG* in nasal epithelial cells. ccREs, candidate cis-regulatory elements in SCREEN registry. ENCORE, DNase I hypersensitivity site and/or transcription factor ChIP-Seq overlapping with the variants.UK Biobank, SAPPHIRE, GCPD-A, replication results using Blacks in UK Biobank and African Americans in the SAPPHIRE and GCPD-A cohorts (R = replicated at $p < 0.05$; F = flip-flop association at $p < 0.05$). Candidate, candidate variants prioritized because of presence of two or more evidence or is nasal eQTL. + indicates presence of evidence.

Figure 3.

A

B

Figure 3. Gene-by-environment interaction analysis on FEV_1. FEV_1 residuals, residuals after FEV_1 was

regressed on the covariates age, sex, height, controller medications, sequencing centers and the first 5

genetic PCs. FEV_1 residuals was plotted against (A) past year exposure to SO_2 stratified by the number of copies of T allele of rs58475486, (B) first year of life exposure to SO_2 stratified by the number of copies of C allele of rs73440122.

Table 1. Descriptive characteristics of 867 African American children with asthma included in this study.

	African American (n=867)
Age	
Mean (SD)	14.1 (3.64)
Median [Min, Max]	13.8 [8.01, 21.8]
Sex	
Male	439 (50.6%)
Female	428 (49.4%)
Height (m)	
Mean (SD)	1.58 (0.145)
Median [Min, Max]	1.60 [1.14, 1.92]
Any control medications* in last 2 weeks	
No	543 (62.6%)
Yes	324 (37.4%)
ICS in last 2 weeks	
No	211 (24.3%)
Yes	306 (35.3%)
Missing	350 (40.4%)
LABA in last 2 weeks	
No	5 (0.6%)
Yes	94 (10.8%)
Missing	768 (88.6%)
Leukotriene inhibitor in last 2 weeks	
No	11 (1.3%)
Yes	68 (7.8%)
Missing	788 (90.9%)
African ancestry	
Mean (SD)	0.792 (0.129)
Median [Min, Max]	0.826 [0.118, 1.00]
Smoking status	
Never	793 (91.5%)
Past	72 (8.3%)
Current	0 (0%)
Missing	2 (0.2%)
Number of smokers in family	
0	469 (54.1%)
1	137 (15.8%)
2	42 (4.8%)
3+	10 (1.2%)
Missing	209 (24.1%)
SO2 first year exposure (ppb)	
Mean (SD)	1.59 (0.961)

Median [Min, Max]	1.50 [0.0300, 14.5]
Missing	227 (26.2%)
SO2 past year exposure (ppb)	
Mean (SD)	1.10 (0.302)
Median [Min, Max]	1.08 [0.120, 2.27]
Missing	206 (23.8%)
SO2 lifetime exposure (ppb)	
Mean (SD)	1.50 (0.371)
Median [Min, Max]	1.47 [0.964, 7.39]
Missing	206 (23.8%)
TOPMed sequencing center and phase	
phase 1, CAAPA	6 (0.7%)
phase 1, NYGC	460 (53.1%)
phase 3, NW	401 (46.3%)

Control medications include inhaled corticosteroid (ICS), long-acting beta-agonist (LABA), leukotriene inhibitor and/or ICS/LABA combo. SO_2 exposure are hourly exposure averaged over the specified time period before spirometry testing as previously described in Neophytou *et al.* 2016. ppb, parts per billion or $\mu g/m^3$.

Table 2. Integration of statistical and function evidence for variant prioritization.

rsID	Alt	β	p	β_{norm}	p_{norm}	MAF	1000 Genomes				ccRE	DNaseI	TFBS	Nasal eQTL	UKBB	SAPPHIRE	GCPD-A	Score
							ALL	AFR	AMR	EUR								
rs11835305	T	0.126	1.93E-05	0.320	3.69E-05	0.104	0.036	0.119	0.012	0.001	N	N	N	N	N	N	N	0
rs17015963	C	0.126	1.93E-05	0.320	3.69E-05	0.104	0.036	0.120	0.012	0.001	N	N	N	N	N	N	N	0
rs58475486	T	0.127	1.45E-05	0.323	2.81E-05	0.105	0.037	0.123	0.012	0.001	Y	N	FOXA1 (A549)	N	N	N	N	2*
rs17015979	T	0.127	1.45E-05	0.323	2.81E-05	0.105	0.037	0.123	0.012	0.001	N	N	N	N	N	N	N	0
rs57692452	C	0.245	1.63E-06	0.654	1.06E-06	0.033	0.011	0.030	0.006	0.001	Y	N	N	N	Y	FF	N	2*
rs112585732	T	0.235	4.35E-07	0.625	3.06E-07	0.041	0.016	0.050	0.006	0.001	N	N	N	N	N	N	N	0
rs113837356	T	0.270	3.44E-06	0.719	2.49E-06	0.025	0.008	0.027	0.006	0.001	N	N	N	-	Y	N	N	1
rs61441836	G	0.252	8.19E-07	0.671	5.46E-07	0.033	0.010	0.030	0.006	0.001	N	N	N	N	Y	FF	N	1
rs73438172	A	0.252	8.19E-07	0.671	5.46E-07	0.033	0.010	0.030	0.006	0.001	N	N	N	N	Y	FF	N	1
rs1044043958[i]	A	0.270	3.44E-06	0.719	2.49E-06	0.025	-	-	-	-	N	N	N	-	N	N	N	0
rs73438182	G	0.138	1.68E-04	0.378	9.18E-05	0.064	0.020	0.068	0.006	0.001	N	N	N	N	N	FF	FF	0
rs73438185	A	0.138	1.68E-04	0.378	9.18E-05	0.064	0.020	0.068	0.006	0.001	N	N	N	N	N	FF	FF	0
rs73438188	A	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	N	N	N	N	Y	N	N	1
rs73438190	C	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	N	N	N	N	N	FF	N	0
rs73438195	A	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	N	N	N	N	N	FF	N	0
rs111857459	T	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	N	N	N	N	N	FF	N	0
rs144369986[i]	T	0.285	1.21E-06	0.756	9.44E-07	0.025	0.008	0.026	0.006	0.001	N	N	N	-	N	N	-	0
rs73440106	G	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	N	N	N	N	N	FF	N	0
rs73440107	A	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	N	N	N	N	Y	N	N	1
rs111453514	C	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	N	N	N	N	Y	N	N	1
rs73440112	T	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	N	N	N	N	Y	N	N	1
rs73440115	G	0.297	6.97E-08	0.792	4.42E-08	0.028	0.011	0.028	0.006	0.001	N	N	N	N	Y	N	N	1
rs11312747	A	0.133	1.43E-05	0.357	8.51E-06	0.100	0.036	0.121	0.010	0.001	N	N	N	-	N	N	N	0
rs73440120	A	0.285	1.21E-06	0.756	9.44E-07	0.025	0.008	0.026	0.006	0.001	N	N	N	-	Y	N	N	1
rs111289668	G	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	Y	N	N	N	Y	N	N	2*
rs73440122	C	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	Y	GM12865	SPI1 (HL-60)	N	Y	N	N	3*
rs73440123	G	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	N	N	N	N	Y	N	N	1

rsID	alt	β	p	βnorm	pnorm	MAF	ALL	AFR	AMR	EUR	ccRE	DNaseI	TFBS	Nasal eQTL	UKBB	SAPPHIRE	GCPD-A	Score
rs17016065	G	0.112	3.19E-06	0.296	2.54E-06	0.177	0.075	0.217	0.017	0.009	N	N	N	Y	N	FF	N	1#
rs17016066	A	0.112	3.19E-06	0.296	2.54E-06	0.177	0.075	0.217	0.017	0.009	N	N	N	Y	N	FF	N	1#
rs147400083[i]	T	0.112	3.19E-06	0.296	2.54E-06	0.177	-	-	-	-	N	N	N	-	N	-	N	0
rs866852270	T	0.112	3.19E-06	0.296	2.54E-06	0.177	-	-	-	-	N	N	N	-	N	FF	N	0
rs141293300	C	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	Y	N	N	-	N	N	N	1
rs1398303	A	0.104	1.22E-05	0.274	1.12E-05	0.186	0.077	0.223	0.020	0.009	N	N	N	Y	N	FF	N	1#
rs61924868	T	0.104	1.24E-05	0.275	1.14E-05	0.185	0.078	0.223	0.020	0.009	N	N	N	Y	N	FF	N	1#
rs73440134	T	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	N	N	N	N	Y	N	N	1
rs73429413	G	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	N	N	N	N	Y	N	N	1
rs73429415	A	0.096	5.13E-05	0.253	4.84E-05	0.189	0.078	0.225	0.022	0.009	N	N	N	Y	N	N	N	1#
rs112449284	T	0.242	4.64E-06	0.640	3.87E-06	0.031	0.012	0.035	0.007	0.001	N	N	N	N	Y	N	N	1
rs111981782	C	0.296	5.78E-08	0.786	4.09E-08	0.029	0.012	0.033	0.006	0.001	N	N	N	N	Y	N	N	1
rs150942400	T	0.293	6.01E-08	0.780	4.01E-08	0.029	0.012	0.034	0.006	0.002	N	N	N	N	Y	N	N	1
rs147527487	C	0.086	8.49E-05	0.226	8.14E-05	0.205	0.095	0.249	0.016	0.005	N	N	N	N	N	N	N	0
rs111243672	A	0.258	6.41E-07	0.690	3.99E-07	0.032	0.014	0.037	0.007	0.004	N	N	N	N	Y	N	N	1
rs73429450	A	0.302	1.26E-08	0.801	9.01E-09	0.031	0.012	0.033	0.009	0.002	N	N	N	N	Y	N	N	1
rs758775577	C	0.217	2.22E-06	0.574	1.85E-06	0.041	-	-	-	-	N	N	N	N	N	N	N	0
rs142679473[i]	C	0.285	6.30E-08	0.756	4.62E-08	0.031	0.012	0.033	0.009	0.002	N	N	N	-	N	N	N	0

Candidate variants were prioritized if they had a functional score of greater than 2 (*) or if they are eQTL of KITLG in nasal epithelial cells (#). Indels were marked with the superscript i near the rsID. β (p) and β_{norm}(p_{norm}) are the effect sizes (p-values) of the genetic associations of the alternate allele (alt) with FEV_1 and FEV_1.res.rnorm respectively. MAF minor allele frequency. ALL/AFR/AMR/EUR, 1000 Genomes minor allele frequency from all/African/American/European populations. ccRE, candidate cis-regulatory elements derived from ENCODE data and stored in the SCREEN web interface. DNasel, ENCODE DNase I hypersensitivity clusters derived from assays in 95 cell (UCSC hg38 wgEncodeRegDnaseClustered table) in embryonic, blood, bone marrow or lung-related cell lines. TFBS, ENCODE transcription factor ChIP-Seq clusters representing 340 transcription factors and 129 cell types (UCSC hg38 encRegTfbsClustered table). Nasal eQTL, KITLG eQTLs in nasal epithelial cells. UKBB/SAPPHIRE/GCPD-A, replication association analysis of FEV_1 in UK Biobank Blacks and African Americans in SAPPHIRE and GCPD-A (details results in Supplementary Table 3). Score, priority score. GM12865, a B-lymphoblastoid cell line. HL-60, an acute promyelocytic

leukemia cell line. A549, a lung adenocarcinoma cell line. -, not available; FF, flip-flop association defined as replication association (p < 0.05) of the opposite

allele instead of the original allele identified in the discovery cohort.

Table 3. Gene-and-environment analysis on FEV_1

Variant	Exposure	β.variant	β.env	β.gxe	p.variant	p.env	p.gxe	
rs58475486_T	SO$_2$ first year	0.18	-0.04	-0.03	0.119	0.009	0.658	
rs57692452_C	SO$_2$ first year	0.61	-0.04	-0.24	0.006	0.007	0.091	
rs111289668_G	SO$_2$ first year	0.68	-0.04	-0.27	0.003	0.007	0.079	
rs73440122_C	SO$_2$ first year	0.73	-0.04	-0.32	0.002	0.008	0.045	*
rs17016065_G	SO$_2$ first year	0.23	-0.04	-0.08	0.004	0.027	0.108	
rs17016066_A	SO$_2$ first year	0.23	-0.04	-0.08	0.004	0.027	0.108	
rs1398303_A	SO$_2$ first year	0.23	-0.04	-0.09	0.004	0.030	0.082	
rs61924868_T	SO$_2$ first year	0.22	-0.04	-0.08	0.004	0.029	0.088	
rs73429415_A	SO$_2$ first year	0.23	-0.04	-0.09	0.004	0.031	0.069	
rs58475486_T	SO$_2$ past year	-0.21	-0.04	0.32	0.084	0.478	0.003	**
rs57692452_C	SO$_2$ past year	-0.09	0.01	0.29	0.680	0.839	0.100	
rs111289668_G	SO$_2$ past year	-0.18	0.01	0.41	0.444	0.869	0.037	*
rs73440122_C	SO$_2$ past year	-0.16	0.01	0.39	0.482	0.825	0.051	
rs17016065_G	SO$_2$ past year	-0.11	-0.03	0.20	0.299	0.583	0.026	*
rs17016066_A	SO$_2$ past year	-0.11	-0.03	0.20	0.299	0.583	0.026	*
rs1398303_A	SO$_2$ past year	-0.12	-0.04	0.21	0.231	0.521	0.021	*
rs61924868_T	SO$_2$ past year	-0.12	-0.04	0.21	0.248	0.534	0.023	*
rs73429415_A	SO$_2$ past year	-0.12	-0.04	0.20	0.246	0.539	0.026	*
rs58475486_T	SO$_2$ lifetime	-0.24	-0.13	0.26	0.390	0.001	0.173	
rs57692452_C	SO$_2$ lifetime	-0.44	-0.13	0.47	0.436	0.001	0.221	
rs111289668_G	SO$_2$ lifetime	-0.19	-0.13	0.32	0.764	0.001	0.444	
rs73440122_C	SO$_2$ lifetime	-0.15	-0.13	0.29	0.809	0.001	0.489	
rs17016065_G	SO$_2$ lifetime	0.40	-0.10	-0.19	0.043	0.013	0.143	
rs17016066_A	SO$_2$ lifetime	0.40	-0.10	-0.19	0.043	0.013	0.143	
rs1398303_A	SO$_2$ lifetime	0.36	-0.11	-0.17	0.064	0.012	0.184	
rs61924868_T	SO$_2$ lifetime	0.35	-0.11	-0.17	0.070	0.012	0.199	
rs73429415_A	SO$_2$ lifetime	0.34	-0.11	-0.16	0.078	0.012	0.207	

*, p < 0.05; **, p < bonferroni p value of 0.0056.