TITLE

Novel *KITLG/SCF* regulatory variants are associated with lung function in African American children with asthma

AUTHORS

Angel CY Mak[1,*], Satria Sajuthi[2,^], Jaehyun Joo[3,^], Shujie Xiao[4,^], Patrick M Sleiman[5,6,^], Marquitta J White[1,^], Eunice Y Lee[1], Benjamin Saef[3], Donglei Hu[1], Hongsheng Gui[4], Kevin L Keys[1,7], Fred Lurmann[8], Deepti Jain[9], Gonçalo Abecasis[10], Hyun Min Kang[10], Deborah A. Nickerson[11,12,13], Soren Germer[14], Michael C Zody[14], Lara Winterkorn[14], Catherine Reeves[14], Scott Huntsman[1], Celeste Eng[1], Sandra Salazar[1], Sam S Oh[1], Frank D Gilliland[15], Zhanghua Chen[15], Rajesh Kumar[16], Fernando D Martínez[17], Ann Chen Wu[18], Elad Ziv[1], Hakon Hakonarson[5,6,#], Blanca E Himes[3,#], L Keoki Williams[4,#], Max A Seibold[3,#], Esteban G. Burchard[1,19]

[1] Department of Medicine, University of California San Francisco, San Francisco, CA, USA

[2] Center for Genes, Environment, and Health, National Jewish Health, Denver, CO, USA

[3] Department of Biostatistics, Epidemiology, and Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[4] Center for Individualized and Genomic Medicine Research, Department of Internal Medicine, Henry Ford Health System, Detroit, MI, USA

[5] Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, PA, USA

[6] Department of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA, USA

[7] Berkeley Institute for Data Science, University of California, Berkeley, CA, USA

[8] Sonoma Technology Inc, Petaluma, CA, USA

[9] Department of Biostatistics, University of Washington, Seattle, WA, USA

[10] Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA

[11] Department of Genome Sciences, University of Washington, Seattle, WA, USA

[12] Northwest Genomics Center, Seattle, WA, USA

[13] Brotman Baty Institute, Seattle, WA, USA

[14] New York Genome Center, New York, NY, USA

[15] Department of Preventive Medicine, Division of Environmental Health, Keck School of Medicine,

University of Southern California, Los Angeles, CA, USA

[16] Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago, IL, USA

[17] Asthma and Airway Disease Research Center, University of Arizona, Tucson, AZ, USA

[18] Precision Medicine Translational Research (PRoMoTeR) Center, Department of Population

Medicine, Harvard Medical School and Pilgrim Health Care Institute, Boston, MA, USA

[19] Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco,

CA, USA

^,# These authors contributed equally to this work

* Corresponding author

Angel CY Mak

Address: Box 2911, 1550 4th Street, San Francisco, CA 94112

Tel: 415-514-9931

Email: angelcymak@gmail.com

AUTHOR CONTRIBUTIONS

ACYM, SSaj, MJW, SSO, FG, ZC, RK, FDM, ACW, EZ, and EGB designed and supervised the study.

ACYM, SSaj, JJ, SX, PS, MJW, EYL, BS, DH, SH, BEH, HH, LKW, and MAS supervised/performed

analyses and/or interpreted results. CE processed biospecimens for whole genome sequencing.

FL estimated air pollution exposure. DJ, GA, HMK, DAN, and SG contributed to TOPMed WGS data

generation and/or analysis. MZ, LW, and CR contributed to CCDG WGS data generation. SSal

coordinated study recruitment and processed phenotype data ACYM, SSaj, JJ, SX, PS, MJW, EYL,

BS, HG, KLK, SG, BEH, HH, LKW, MAS, and EGB wrote and/or critically reviewed the manuscript.

SHORT TITLE

51 *KITLG* associated with FEV_1 in AA youths

KEYWORDS

GWAS, African American, FEV_1, gene-by-environment interaction, GxE, air pollution, KITLG, SCF

ABSTRACT

Baseline lung function, quantified as forced expiratory volume in the first second of exhalation

(FEV_1), is a standard diagnostic criterion used by clinicians to identify and classify lung diseases.

Using whole genome sequencing data from the National Heart, Lung, and Blood Institute

TOPMed project, we identified a novel genetic association with FEV_1 on chromosome 12 in 867

African American children with asthma (p = 1.26 x 10^{-8}, β = 0.302). Conditional analysis within 1

Mb of the tag signal (rs73429450) yielded one major and two other weaker independent signals

within this peak. We explored statistical and functional evidence for all variants in linkage

disequilibrium with the three independent signals and yielded 9 variants as the most likely

candidates responsible for the association with FEV_1. Hi-C data and eQTL analysis demonstrated

that these variants physically interacted with *KITLG (aka SCF)* and their minor alleles were

associated with increased expression of *KITLG* gene in nasal epithelial cells. Gene-by-air-pollution

interaction analysis found that the candidate variant rs58475486 interacted with past-year SO_2

exposure (p = 0.003, β = 0.32). This study identified a novel protective genetic association with

FEV_1, possibly mediated through *KITLG*, in African American children with asthma.

INTRODUCTION

Asthma, a chronic pulmonary condition characterized by reversible airway obstruction, is one of

the hallmark diseases of childhood in the United States (World Health Organization 2017).

Asthma is also the most disparate common disease in the pediatric clinic, with significant

variation in prevalence, morbidity, and mortality among U.S. racial/ethnic groups (Oh *et al.* 2016).

Specifically, African American children carry a higher asthma disease burden compared to their

European American counterparts (Akinbami *et al.* 2014; Akinbami 2015). Forced expiratory

volume in the first second (FEV$_1$), a measurement of lung function, is a vital clinical trait used by

physicians to assess overall lung health and diagnose pulmonary diseases such as asthma

(Johnson and Theurer. 2014). We have previously shown that genetic ancestry plays an important

role in FEV$_1$ variation and that African Americans have lower FEV$_1$ compared to European

Americans regardless of asthma status (Kumar *et al.* 2010; Pino-Yanes *et al.* 2015). The disparity

in lung function between populations may explain disparities in asthma disease burden.

Understanding the factors that influence FEV$_1$ variation among individuals with asthma could

lead to improved patient care and therapeutic interventions.

Twin and family-based studies estimate that the heritability of FEV$_1$ ranged from 26% to 81%,

supporting the combined contribution by genetic and environmental factors in FEV$_1$ variation

(Chatterjee and Das. 1995; Chen *et al.* 1996; Hukkinen *et al.* 2011; Palmer *et al.* 2001; Sillanpaa

et al. 2017; Tian *et al.* 2017; Yamada *et al.* 2015). Genome-wide association studies (GWAS) of

FEV$_1$, including among individuals with asthma, have identified many variants that contribute to

lung function (Li *et al.* 2013; Liao *et al.* 2014; Repapi *et al.* 2010; Soler Artigas *et al.* 2011; Soler

Artigas *et al.* 2015; Wain *et al.* 2017). A search in NHGRI-EBI GWAS Catalog (version e98_r2020-

03-08) on baseline lung function (FEV$_1$) alone revealed 349 associations (Buniello *et al.* 2019).

Most of these previous GWAS, however, were performed in adult populations of European

descent, and their results may not generalize across populations or across the life span of an

individual (Carlson *et al.* 2013; Martin, A. R. *et al.* 2017; Wojcik *et al.* 2019). Previous GWAS results

are also limited due to their reliance on genotyping arrays. In particular, variation in non-coding

regions of the genome is not adequately covered by many genotyping arrays because they were

not designed to account for the population-specific genetic variability of all populations (Kim, M.

S. *et al.* 2018; Zhang and Lupski. 2015). Whole genome sequencing (WGS) is a newer technology

that captures nearly all common variation from coding and non-coding regions of the genome

and is unencumbered by genotype array design constraints and differences in linkage

disequilibrium patterns among populations. To date, no large-scale WGS studies of lung function

have been performed in African American children with asthma (Martin *et al.* 2017).

In addition to genetics, FEV_1 is a complex trait that is significantly influenced by both genetic

variation and environmental factors, such as air pollution (Chatterjee and Das. 1995; Hukkinen *et*

al. 2011; Palmer *et al.* 2001; Sillanpaa *et al.* 2017; Tian *et al.* 2017; Yamada *et al.* 2015). Exposure

to ambient air pollution has been consistently associated with poor respiratory outcomes,

including reduced FEV_1 (Barraza-Villarreal *et al.* 2008; Brunekreef and Holgate. 2002;

Ierodiakonou *et al.* 2016; Wise 2019). We previously showed that exposure to sulfur dioxide (SO_2),

an air pollutant emitted by the burning of fossil fuels, is significantly associated with reduced FEV_1

in African American children with asthma in the SAGE II study (Neophytou *et al.* 2016). Because

the genetic variants associated with FEV_1 thus far do not account for the majority of its estimated

heritability, considering gene-environment (GxE) interactions, specifically gene-by-air-pollution,

may improve our understanding of lung function genetics (Moore 2005; Moore and Williams.

2009). Here, we performed a genome-wide association analysis using WGS data to identify

common genetic variants associated with FEV_1 in African American children with asthma in SAGE

II and investigated the effect of GxE (SO_2) interactions on FEV_1 associations.

METHODS

Study population

This study examined African American children between 8-21 years of age with physician-

diagnosed asthma from the Study of African Americans, Asthma, Genes & Environments (SAGE

II). All SAGE II participants were recruited from the San Francisco Bay Area. The inclusion and

exclusion are previously described in detailed (Oh *et al.* 2012; White *et al.* 2016) . Briefly,

participants were eligible if they were 8-21 years of age and self-identified as African American

and had four African American grandparents. Study exclusion criteria included the following: 1)

any smoking within one year of the recruitment date; 2) 10 or more pack-years of smoking; 3)

pregnancy in the third trimester; 4) history of lung diseases other than asthma (for cases) or

chronic illness (for cases and controls). Baseline lung function defined as forced expiratory

volume in the first second (FEV_1) was measured by spirometry prior to administering albuterol as

previously described (Oh *et al.* 2012).

TOPMed whole genome sequencing data

SAGE II DNA samples were sequenced as part of the Trans-Omics for Precision Medicine (TOPMed)

whole genome sequencing (WGS) program (Taliun *et al.* 2019). WGS was performed at the New

York Genome Center and Northwest Genomics Center on a HiSeq X system (Illumina, San Diego,

CA) using a paired-end read length of 150 base pairs (bp), with a minimum of 30x mean genome

coverage. DNA sample handling, quality control, library construction, clustering and sequencing,

read processing and sequence data quality control are described in detail in the TOPMed website

(TOPMed 2019). Variant calls were obtained from TOPMed data freeze 8 VCF files corresponding

to the GRCh38 assembly. Variants with a minimal read depth of 10 (DP10) were used for analysis

unless otherwise stated.

Genetic principal components, global ancestry, and kinship estimation

Genetic principal components (PCs), global ancestry, and kinship estimation on genetic

relatedness were computed using biallelic single nucleotide polymorphisms (SNPs) with a PASS

flag from TOPMed freeze 8 DP10 data. PCs and kinship estimates were computed using the PC-

Relate function from the GENESIS R package (Conomos *et al.* 2015; Conomos *et al.* 2016) using a

workflow available from the Summer Institute in Statistical Genetics Module 17 course website

(Summer Institute in Statistical Genetics 2019). African global ancestry was computed using the

ADMIXTURE package (Alexander *et al.* 2009) in supervised mode using European (CEU), African

(YRI) and Native American (NAM) reference panels as previously described (Mak, A. C. Y. *et al.*

2018).

FEV_1 GWAS

Non-normality of the distribution of FEV_1 values was tested with the Shapiro-Wilk test in R using

the shapiro.test function. Since FEV_1 was not normally distributed (p = 1.41 x 10^{-8} for FEV_1 and p

= 1.05 x 10^{-8} for $\log_{10} FEV_1$), FEV_1 was regressed on all covariates (age, sex, height, controller

medications, sequencing centers, and the first 5 genetic PCs) and the residuals were inverse-

normalized. These inverse-normalized residuals (FEV$_1$.res.rnorm) were the main outcome of the

discovery GWAS. The controller medication covariate included the use of inhaled corticosteroids

(ICS), long-acting beta-agonists (LABA), leukotriene inhibitors and/or an ICS/LABA combo in the

2 weeks prior to the recruitment date.

Genome-wide single variant analysis was performed on the ENCORE server (https://github.com

/statgen/encore) using the linear Wald test (q.linear) originally implemented in EPACTS

(https://genome.sph.umich.edu /wiki/EPACTS) and TOPMed freeze 8 data (DP0 PASS) with a

MAF filter of 0.1%. All pairwise relationships with degree 3 or more relatedness (kinship values >

0.044) were identified, and one participant of the related pair was subsequently chosen at

random and removed prior to analysis. All covariates used to obtain FEV$_1$.res.rnorm were also

included as covariates in the GWAS as recommended in a recent publication (Sofer *et al.* 2019).

The association analysis was repeated using untransformed FEV$_1$ and FEV$_1$ percent predicted

(FEV$_1$.perc.predicted). FEV$_1$ percent predicted was defined as the percentage of measured FEV$_1$

relative to predicted FEV$_1$ estimated by the Hankinson lung function prediction equation for

African Americans (Hankinson *et al.* 1999). A secondary analysis that included smoking-related

covariates (smoking status and number of smokers in the family) was performed in PLINK 1.9

(version 1p9_2019_0304_dev) (Chang *et al.* 2015; Purcell and Chang. 2013). To study whether

association with FEV$_1$ is specific to SAGE II participants with asthma, we repeated the association

analysis adjusting for age, sex, height and the first 5 genetic PCs in SAGE II participants without

asthma on the ENCORE server. All of these participants were sequenced in the same center.

Regional association results were plotted using LocusZoom 1.4 (Pruim *et al.* 2010) with a 500

kilobase (Kb) flanking region. Linkage disequilibrium (R^2) was estimated in PLINK 1.9. LD plot was

generated using recoded genotype files (plink --recode 12) in Haploview (Barrett *et al.* 2005) .

The function effectiveSize in the R package CODA was used to estimate the actual effective

number of independent tests and CODA-adjusted statistical and suggestive significance p-value

thresholds were defined as 0.05 and 1 divided by the effective number of tests, respectively

(Duggal *et al.* 2008). We compared the CODA-adjusted statistical significance threshold and the

widely used 5×10^{-8} GWAS genome-wide significance threshold (Pe'er *et al.* 2008) and selected

the more stringent threshold for genome-wide significance.

The following WGS quality control steps were applied to all reported variants from ENCORE to

ensure WGS variant quality: (1) The variant had VCF FILTER = PASS; (2) Variant quality was

confirmed via manual inspection on the BRAVO server based on TOPMed freeze 5 data

(University of Michigan and NHLBI TOPMed. 2018); (3) Variants were reanalyzed with linear

regression using PLINK 1.9 by applying the arguments --mac 5 --geno 0.1 --hwe 0.0001 using

TOPMed freeze 8 DP10 PASS data.

To determine if the rs73429450 association with FEV_1 was only identifiable using whole genome

sequencing data, we repeated the linear regression association analysis on signals that passed

the genome-wide significance threshold using PLINK 1.9 and genotype data generated with

Axiom Genome-Wide LAT 1 array (Affymetrix, Santa Clara, CA, dbGaP phs000921.v1.p1). These

array genotype data were imputed into the following reference panels: 1000 Genomes phase 3

version 5, Haplotype Reference Consortium (HRC) r1.1, the Consortium on Asthma among

African-ancestry Populations in the Americas (CAAPA) and the TOPMed phase 5 panels on the

Michigan Imputation Server (Das *et al.* 2016). It should be noted that 500 SAGE II subjects were

part of the TOPMed freeze 5 reference panel.

A total of 349 GWAS FEV_1-associated entries were retrieved from the NHGRI-EBI GWAS Catalog

version 1.0.2-associations_e98_r2020-03-08 (Buniello *et al.* 2019) using the trait names "Lung

function (FEV_1)", "FEV1", "Lung function (forced expiratory volume in 1 second)" or

"Prebronchodilator FEV1". After adding 100 Kb flanking regions to each of the 349 entries, a total

of 230 non-overlapping region were obtained. To look up whether we replicated previously

GWAS loci while control for multiple testing penalties, we only used 279,495 common variants

(MAF >= 0.01) that overlapped with the 230 regions. The 279,495 common variants is equivalent

to 17,755 effective test based on CODA and 5.63×10^{-5} (1/17,755) was used as suggestive p-value

threshold for replication.

Conditional analysis

Conditional analysis was performed to identify all independent signals in a GWAS peak using

PLINK 1.9. All TOPMed freeze 8 DP10 variants within 1 megabase (Mb) of the tag association

signal and with association p-value of 1×10^{-4} or smaller in the discovery GWAS were included in

the analysis. Variants were first ordered by ascending p-value. A variant was considered to be an

independent signal if the association p-value after conditioning (conditional p-value) on the tag

signal was smaller than 0.05. Newly identified independent signals were included with the tag

signal for conditioning on the next variant.

Region-based association analysis

Region-based association analyses were performed in 1 Kb sliding windows with 500 bp increments in a 1 Mb flanking region of the tag GWAS signal using the SKAT_CommonRare function from the SKAT R package v1.3.2.1 (Ionita-Laza *et al.* 2013) . Default settings were used with method = "C" and test.type = "Joint". A minor allele frequency (MAF) threshold of 0.01 was used as the cutoff to distinguish rare and common variants. Variants were annotated in TOPMed using the WGSA pipeline (Liu *et al.* 2016). Since SKAT imputes missing genotypes by default by assigning mean genotype values (impute.method="fixed"), we chose to use low coverage genotypes instead of SKAT imputation, and hence, TOPMed freeze 8 DP0 variants with a VCF FILTER of PASS were included in the analysis. The function effectiveSize in the R package CODA (Plummer *et al.* 2006) was used to estimate the effective number of independent hypothesis tests for accurate Bonferroni multiple testing corrections. P-value thresholds for statistical significance and suggestive significance were defined as 0.05 and 1 divided by the effective number of tests, respectively (Duggal *et al.* 2008). If a region was suggestively significant, region-based analyses were repeated with functional variants and/or rare variants (MAF <= 0.01) to assess contribution of common, rare and/or functional variants. Region-based analyses using rare variants only were performed using SKAT-O (Lee *et al.* 2012). The WGSA annotation filters used to define functional variants are provided in File S1 (Supplementary Text 1). To study the contribution of individual variants to a region-based association p-value, drop-one variant analysis was performed by repeating the region-based analysis multiple times and dropping one variant only at a time.

Functional annotations and prioritization of genetic variants

The Hi-C Unifying Genomic Interrogator (HUGIN) (Ay *et al.* 2014; Martin, J. S. *et al.* 2017; Schmitt

et al. 2016) was used to assign potential gene targets to each variant. HUGIN uses the Hi-C data

generated from the primary human tissues from four donors used in the Roadmap Epigenomics

Project (Schmitt *et al.* 2016). ENCODE annotations (ENCODE Project Consortium 2011; ENCODE

Project Consortium 2012) were based on overlap of the variants with functional data downloaded

from the UCSC Table Browser (Karolchik *et al.* 2004). These data included DNAase I

hypersensitivity peak clusters (hg38 wgEncodeRegDnaseClustered table), transcription factor

ChIP-Seq clusters (hg38 encRegTfbsClustered table) and histone modification ChIP-Seq peaks

(hg19 wgEncodeBroadHistone<cell type><histone>StdPk tables). For DNase I hypersensitivity

and transcription factor binding sites, we focused on blood, bone marrow, lung and embryonic

cells. For histone modification ChIP-Seq, we focused on H3K27ac and H3K4me3 modifications in

human blood (GM12878), bone marrow (K562), lung fibroblast (NHLF), and embryonic stem cells

(H1-hESC). LiftOver tool (Hinrichs *et al.* 2006) was used to convert genomic coordinates from

hg19 to hg38. Candidate cis-regulatory elements (ccREs) were a subset of representative DNase

hypersensitivity sites with epigenetic activity further supported by histone modification

(H3K4me3 and H3K27ac) or CTCF-binding data from the ENCODE project. Overlap of variants with

ccREs were detected using the Search Candidate cis-Regulatory Elements by ENCODE (SCREEN)

web interface (ENCODE Project Consortium 2011; ENCODE Project Consortium 2012).

Prioritization of genetic variants was based on the presence of statistical, functional and/or

bioinformatic evidence as described in the Diverse Convergent Evidence (DiCE) prioritization

framework (Ciesielski *et al.* 2014). The priority score of each variant was obtained by counting

the number of statistical, functional, and/or bioinformatic evidences that support potential

biological function for that variant.

Replication of GWAS associations

All replication analyses were performed in subjects with asthma. Replication of GWAS FEV_1

associations was attempted on TOPMed whole genome sequencing data generated from four

cohorts. These cohorts included Puerto Rican (n=1,109) and Mexican American (n=649) children

in the Genes-Environments and Admixture in Latino Americans (GALA II) study (Oh *et al.* 2012),

African American adults in the Study of Asthma Phenotypes and Pharmacogenomic Interactions

by Race-Ethnicity (SAPPHIRE, n=3,428) (Levin *et al.* 2014) and African American children in

Genetics of Complex Pediatric Disorders (GCPD-A, n=1,464) study (Ong *et al.* 2013). Age, sex,

height, controller medications and the first 5 PCs were used as covariates.

Additionally, replication of GWAS FEV_1 associations was attempted using data of black UK

Biobank subjects who had asthma (n=627) while adjusting for age, sex, height and the first 5

principal components. Asthma status was defined by ICD code or self-reported asthma. UK

Biobank genotype data was generated on Affymetrix UK BiLEVE axiom or UK Biobank Axiom array

and imputed into the Haplotype Reference Consortium, 1000G and UK 10K projects (Bycroft *et*

al. 2018; Canela-Xandri *et al.* 2018). Additional details on the UK Biobank study and the

replication procedures are available in File S1 (Supplementary Text 2).

RNA sequencing and expression quantitative trait loci (eQTL) analysis

Whole-transcriptome libraries of 370 nasal brushings from GALA II Puerto Rican children with asthma were constructed by using the Beckman Coulter FX automation system (Beckman Coulter, Fullerton, CA). Libraries were sequenced with the Illumina HiSeq 2500 system. Raw RNA-Seq reads were trimmed using Skewer (Jiang *et al.* 2014) and mapped to human reference genome hg38 using Hisat2 (Kim, D. *et al.* 2015). Reads mapped to genes were counted with htseq-count and using the UCSC hg38 GTF file as reference (Anders *et al.* 2015). Cis-expression quantitative trait locus (eQTL) analysis of *KITLG* was performed as described in the Genotype-Tissue Expression (GTEx) project version 7 protocol (GTEx Consortium *et al.* 2017) using age, sex, BMI, global African and European ancestries and 60 PEER factors as covariates.

Gene-by-air-pollution interaction analysis

We hypothesized that the effect of genetic variation on lung function in our study population may differ by the levels of exposure to SO_2 (Neophytou *et al.* 2016). To test for an interaction between a genetic variant and SO_2, an additional multiplicative interaction term (variant x SO_2 exposure) was included in the original GWAS model (see Method Section "FEV_1 GWAS"). The SO_2 estimates used in the interaction analysis were first-year, past-year, and lifetime exposure to ambient of SO_2, which were estimated as described previously (Neophytou *et al.* 2016). Briefly, we obtained regional ambient daily air pollution data from the U.S. Environmental Protection Agency Air Quality System. SO_2 estimates for the participant's residential geographic coordinate were calculated as the inverse distance-squared weighted average from the four closest air pollution monitoring stations within 50 km of the participant's residence. We estimated yearly

exposure at the reported residential address by averaging all available daily measures (daily

average of 1-hour SO_2) in a given year. If the participant had a change of residential address in a

given year, we estimate yearly exposure as a time-weighted estimate based on the number of

months spent at each different address in that year. Average lifetime exposures were estimated

using all available yearly average estimates over the lifetime of the participant until the day of

spirometry testing. Since not all pollutants were measured daily, there are location- and

pollutant-dependent missing values. Residuals of FEV_1 were plotted against exposure to SO_2 and

stratified by the number of copies of the minor allele. Residuals of FEV_1 were obtained as

described in the Methods Section "FEV_1 GWAS".

Data availability

Local institutional review boards approved the studies (IRB# 10-02877). All subjects and legal

guardians provided written informed consent. TOPMed whole genome sequencing and

phenotype data from SAGE II are available on dbGaP under accession number phs000921.v4.p1.

Normalized gene count data for *KITLG* and supplemental materials are available at figshare.

RESULTS

Novel lung function associations

Subject characteristics of the 867 African American children with asthma included in this study

are shown in Table 1, and the distribution of their FEV_1 measurements (mean = 2.56 L, standard

deviation = 0.79 L) is in Figure S1. The CODA-adjusted statistical significance thresholds $2.10 \times 10^{-}$

[8] and 4.19 x 10^{-7} were used as the genome-wide and suggestive significance thresholds,

respectively. According to this threshold, one SNP in chromosome 12 (chr12:88846435,

rs73429450, G>A) was associated with FEV_1.res.rnorm (Figure 1, p = 9.01 x 10^{-9}, β = 0.801) at

genome-wide significance. The association between rs73429450 and lung function remained

statistically significant when the association was repeated using untransformed FEV_1 (p = 1.26 x

10^{-8}, β = 0.302) as the outcome variable. The association between rs73429450 and lung function

was suggestive using FEV_1.perc.predicted (p = 1.69 x 10^{-7}, β = 0.100). Twenty suggestive

associations corresponding to 4 tag signals are reported in Supplementary File S2. None of the

suggestive associations overlapped with any of the previously reported FEV_1-associated loci.

When considering only common variants and applying a p-value threshold of 5.63 x 10^{-5}, we

found replicated in 6 out of 230 previously reported FEV_1 associations (Table S1). Our top FEV_1

association, rs73429450, did not overlap with any previously reported loci and it is a novel

association with FEV_1 in this study population.

Secondary analysis that included covariates correcting for smoking status and number of smokers

in the family showed that smoking-related factors were not significantly associated with FEV_1 in

our pediatric SAGE cohort: using 657 out of 867 individuals with available smoking-related

covariates, the FEV_1.res.rnorm association p-values before and after including the smoking-

related covariates were 2.01 x 10^{-6} and 1.89 x 10^{-6}. Both p-values of the covariates smoking status

(p = 0.27) and number of smokers in the family (p = 0.54) were not significant.

Conditional analysis was performed on 45 variants with association p < 1 x 10^{-4} located within 1

Mb of the strongest association signal (rs73429450). Two weaker independent signals

(rs17016065, rs58475486) were identified (Table S2). None of the 45 variants showed association

with FEV_1.res.rnom in 251 SAGE II children without asthma (Table S3).

The minor allele frequency of rs73429450 in continental populations from the 1000 Genomes

Project (1000G) is 3% in Africans (AFR) and < 1% in Admixed Americans (AMR), Europeans (EUR)

and Asians (EAS and SAS) (1000 Genomes Project Consortium *et al.* 2015). Rs73429450 was not

included on the Affymetrix LAT1 genotyping array where SAGE participants were previously

genotyped. To determine if the rs73429450 association with FEV_1 was only identifiable using

whole genome sequencing data, we attempted to reproduce our results by imputing the

genotype of rs73429450 in 851 SAGE participants with available array data using 1000G phase 3

(n = 2,504), HRC r1.1 (n = 32,470), CAAPA (n = 883) and TOPMed freeze 5 (n = 62,784) reference

panels. Our results remained statistically significant when using the 1000G phase 3 ($p = 4.97 \times 10^{-8}$,

$\beta = 0.79$, imputation $R^2 = 0.95$) and TOPMed freeze 5 ($p = 1.22 \times 10^{-8}$, $\beta = 0.80$, imputation $R^2 = 0.98$)

0.98) reference panels, but lost statistical significance when rs73429450 genotypes were

imputed using the HRC ($p = 4.35 \times 10^{-7}$, $\beta = 0.68$, imputation $R^2 = 0.94$) and CAAPA ($p = 1.95 \times 10^{-7}$,

$\beta = 0.80$, imputation $R^2 = 0.71$) reference panels.

Region-based association analysis including all variants conditioned on the association signal

from rs73429450 was performed in its 1 Mb flanking region (chr12:87846435-89846435). No

windows were significantly associated after Bonferroni multiple testing correction ($p < 2.80 \times 10^{-4}$,

Figure S2), but 20 windows were suggestively associated with FEV_1.res.rnorm ($p < 5.60 \times 10^{-3}$,

Table S5). Two of 20 windows re-tested using only functional variants were suggestively

significant (region 4 and 16). Both of these windows were no longer suggestively significant after

removing the common variants, indicating that association signal from these regions was mostly

driven by common variants. Further investigation on region 16 using drop-one analysis on the 2

rare and 1 common function variants confirmed the major contribution by the common variant,

rs1895710, as shown by the major increase in p-value (Table S6). The signal was also slightly

driven by the singleton, rs990979778. Drop-one analysis was not performed on region 4 because

there were only 1 common and 1 rare variants.

A Hi-C assay couples a chromosome conformation capture (3C) assay with next-generation

sequencing to capture long-range interactions in the genome. We identified a statistically

significant long-range chromatin interaction between the GWAS peak and the KIT ligand (*KITLG*,

also known as stem cell factor, *SCF*) gene in human fetal lung fibroblast cell line IMR90 (Table S7).

The long-range interaction detected in human primary lung tissue was not significant, implying

that the potential long-range interactions are specific to tissue type or developmental stage.

Potential regulatory role of FEV$_1$-associated variants on KITLG expression

To further elucidate potential regulatory relationships between the GWAS association peak and

KITLG, we analyzed whether variants in the peak were eQTL of *KITLG* in previously published

whole blood RNA-Seq data available from the same study participants (Mak, Angel CY *et al.* 2016) .

The whole blood RNA-Seq data, however, did not yield evidence of expressed *KITLG*, consistent

with results in GTEx. We subsequently used RNA-Seq data from nasal epithelial cells of 370 Puerto

Rican children with asthma from the GALA II study, and found that five out of 45 variants were

eQTL of *KITLG* (Table S8). While Puerto Ricans are a different population than African Americans,

they are both admixed populations with substantial African genetic ancestry, and therefore could

share eQTLs. All five eQTLs corresponded to one signal in a region with strong linkage

disequilibrium ($r^2 > 0.8$, Figure S3).

Replication of genetic association with FEV_1

Subject characteristics of our four replication cohorts (SAPPHIRE, GCPD-A, UK Biobank and GALA

II) are shown in Table S9. We attempted to replicate the association of the 45 SNPs in our primary

FEV_1 GWAS in each cohort. We used 0.05 as the suggestive p-value threshold and 0.0167 as the

Bonferroni-corrected p-value threshold after correcting for 3 independent signals (see

conditional analysis in Results Section). A total of 20 variants were replicated at $p < 0.05$ with

consistent direction of effect in black UK Biobank participants; 14 variants in SAPPHIRE and 2

variants in GCPD-A were significant but had an opposite direction of effect (Table S10).

We attempted to replicate the FEV_1.res.rnorm association in Mexican American (n = 649) and

Puerto Rican (n = 1,109) children with asthma from the GALA II study. In Mexican Americans, we

excluded 19 variants with MAF < 0.1% and associations for the remaining 26 variants did not

replicate (Table S11). In Puerto Ricans, the associations were not replicated (Table S11).

Incorporating statistical and functional evidence for candidate variant prioritization

We combined and summarized all functional evidence for the top 45 variants, along with eQTL

findings from nasal epithelial RNA-Seq and replication results (Figure 2, Table 2 and S12). To

facilitate interpretation of the variant association with FEV_1, the effect sizes and p-values of both

FEV_1 (β and p) and FEV_1.res.rnorm (β_{norm} and p_{norm}) associations are also reported. CADD

functional prediction score and ENCODE histone modification ChIP-Seq peaks in embryonic,

blood, bone marrow, and lung-related tissues were also examined but not reported because

none of the variants had a CADD score greater than 10 and none overlapped with histone

modification sites. Rs73440122 received the highest priority score of 3 based on replication in

the UK Biobank, overlap with a DNase I hypersensitivity site in B-lymphoblastoid cells (GM12865)

and overlap with an SPI1 binding site in acute promyelocytic leukemia cells. Eight other variants

were prioritized with score > 2 or evidence of being an eQTL for *KITLG* in nasal epithelial cells

(Table 2, score marked with ^ or # respectively). These nine candidate variants were selected for

gene-by-air-pollution interaction analyses.

Gene-by-air-pollution interaction of rs58475486

We previously found that first year of life and lifetime exposure to SO_2 were associated with FEV_1

in African American children (Neophytou *et al.* 2016). We investigated whether the effect of the

nine prioritized genetic variants associated with lung function varied by SO_2 exposure (first year

of life, past year, and lifetime exposure). Since the nine variants represent three independent

signals (see conditional analysis in the Results Section), the Bonferroni-corrected p-value

threshold was set to p = 0.0056 (correction for nine tests; three signals and three exposure

periods to SO_2). We observed a single statistically significant interaction between the T allele of

rs58475486 and past year exposure to SO_2 that was positively associated with FEV_1 (p = 0.003, β

= 0.32, Table 3, Figure 3A). This interaction remains significant (p = 0.003, β = 0.32) in secondary

analyses adjusted for smoking status or a multiplicative interaction term of rs58475486 and

smoking status as additional covariates. Interestingly, six of the remaining eight variants also

displayed interaction effects with past year exposure to SO_2 that were suggestively associated (p

< 0.05) with FEV_1 (Table 3). We also found a suggestive interaction of the C allele of rs73440122

with first year exposure to SO_2 that was associated with decreased FEV_1 (p = 0.045, β = -0.32,

Figure 3B). The same allele also showed interaction with past year of exposure to SO_2 that was

suggestively associated with FEV_1 in the opposite direction (p = 0.051, β = 0.39).

DISCUSSION

Variant rs73429450 (MAF = 0.030) was identified as the strongest association signal with FEV_1.

Each additional copy of the protective A allele of rs73429450 was associated with a 0.3 L increase

of FEV_1. We did not find any statistically significant contribution of rare variants to the association

signal from a 1 Kb sliding window analyses in the 1 MB flanking region centered on rs73429450.

We were surprised to identify a novel common variant (MAF = 0.030) associated with lung

function using whole genome sequence data in a population that was previously analyzed for

associations with lung function using genotype array data. Further investigation revealed that

our discovered variant, rs73429450, was not captured by the LAT 1 genotyping array, and the

association with lung function depended on the reference panel used to impute the variant into

our population. More surprisingly, our statistically significant finding was only found to be

suggestively significant using data imputed from the CAAPA reference panel (p = 1.95 x 10^{-7}, β =

0.80). Of the imputation reference panels that we assessed, CAAPA is one of the more relevant

reference panels for our study population because it is based on African populations in the

Americas. However, we note that the effect size estimated from CAAPA-imputed data was

comparable to that generated from WGS data. While whole genome sequencing data is usually

praised for enabling analysis of rare-variant contributions to phenotype variability, our results

show the utility of whole genome sequencing data for the reliable analysis of common variants

as well in the absence of relevant imputation panels.

Although rs73429450 had the lowest p-value from our whole genome sequencing association

analysis, we did not find the required amount of functional evidence to prioritize this marker for

inclusion in downstream gene-by-air-pollution analyses. Another variant, rs73440122, was in

moderate to strong linkage disequilibrium (r^2 = 0.76) with rs7349450 and had a similar MAF

(0.027) in our study population, but was only suggestively associated with FEV_1 in our association

analysis (p = 2.08 x 10^{-7}, Table2). In contrast to rs73429450, there were multiple lines of evidence

suggesting the functional relevance of rs73440122: rs73440122 received the highest priority

score based on its replicated FEV_1 association in black UK Biobank participants and overlap with

ENCODE gene regulatory regions, making it one of the most likely drivers of FEV_1 variability

among individuals, possibly mediated through *KITLG*.

Bioinformatic interrogation of rs73440122 revealed that the variant overlapped with a ccRE

(SCREEN accession EH37E0279310), DNase I hypersensitivity site, and SPI1 ChIP-Seq clusters that

were indicative of a candidate open chromatin gene regulatory region (Table S12). The binding

evidence of SPI1 is highly relevant to the role of KITLG in type 2 inflammation (see below). Variant

rs73440122 is located in a region that physically interacted with *KITLG* based on Hi-C data in fetal

lung fibroblast cells. Additionally, five neighboring FEV_1 associated variants were identified as

eQTLs of *KITLG*, although they appeared to be an independent signal (r^2 < 0.2). Overall, these

results support regulatory interactions between our novel locus and *KITLG*.

Atopic or type 2 high asthma is the most common form of asthma in children (Comberiati *et al.*

2017). *KITLG,* more commonly known as stem cell factor (*SCF*)*,* is a ligand of the KIT tyrosine

kinase receptor. It plays an important role in type 2 inflammation in atopic asthma, especially in

inflammatory processes mediated through mast cells, IgE and group 2 innate lymphoid cells (Da

Silva and Frossard. 2005; Da Silva *et al.* 2006; Fonseca *et al.* 2019; Oliveira and Lukacs. 2003) . In

the airways, *KITLG* is expressed in bronchial epithelial cells, lung fibroblasts, bronchial smooth

muscle cells, endothelial cells, peripheral blood eosinophils, dendritic cells and mast cells (Hsieh

et al. 2005; Kassel *et al.* 1999; Oriss *et al.* 2014; Valent *et al.* 1992; Wen *et al.* 1996). *KITLG* is a

major growth factor of mast cells (Reviewed in Broudy 1997; Da Silva *et al.* 2006; Galli *et al.* 1994;

Galli *et al.* 1995). It promotes recruitment of mast cell progenitors into tissues (Reviewed in

Oliveira and Lukacs. 2003), prevents mast cell apoptosis (Iemura *et al.* 1994; Mekori *et al.* 1993)

and promotes release of inflammatory mediators such as proteases, histamine, chemotactic

factors, cytokines (Reviewed in Amin 2012; Borish and Joseph. 1992). While KITLG promotes the

production of cytokines like IL-13 upon IgE-receptor crosslinking on the surface of mast cells

(Kobayashi *et al.* 1998), IL-13 was also reported to up-regulate *KITLG* (Rochman *et al.* 2015).

Consistent with the critical role of KITLG for mast cells and type 2 inflammation, we found our

prioritized variant, rs73440122, overlapped with a SPI1 (aka PU.1) ChIP-Seq cluster. The

transcription factor SPI1 was demonstrated in SPI1 knockout mice to be necessary for the

development of B cells, T cells, neutrophils, macrophages, dendritic cells, and mast cells

(Anderson *et al.* 2000; Guerriero *et al.* 2000; McKercher *et al.* 1996; Scott *et al.* 1994; Scott *et al.*

1997; Walsh *et al.* 2002). It plays an essential role in macrophage differentiation in asthmatic and

other allergic inflammation (Qian *et al.* 2015; Yashiro *et al.* 2019). It was also shown to regulate

the cell fate between mast cells and monocytes (Ito *et al.* 2005; Ito *et al.* 2009; Nishiyama,

Nishiyama, Ito, Masaki, Maeda *et al.* 2004; Nishiyama, Nishiyama, Ito, Masaki, Masuoka *et al.*

2004). The presence of a SPI1 binding site in a candidate regulatory region of KITLG is therefore

highly relevant given the critical role of KITLG in mast cell survival and activation.

Higher levels of KITLG (Al-Muhsen *et al.* 2004; Da Silva *et al.* 2006; Tayel *et al.* 2017) and an

increased number of mast cells in the lung (Cruse and Bradding. 2016; Fajt and Wenzel. 2013;

Mendez-Enriquez and Hallgren. 2019) were detected in individuals with asthma. The percentage

of a subpopulation of circulating blood mast cell progenitors (Lin$^+$ CD34^{hi} CD117$^{int/hi}$ FcεRI$^+$) was

higher in individuals with a reduced lung function (Dahlin *et al.* 2016). These findings suggested

that higher *KITLG* expression and/or number of mast cells may be a contributing factor to lower

lung function. This notion was inconsistent with the association of our novel locus with higher

KITLG expression and increased lung function in SAGE II children with asthma. Interestingly, a

study of 20 subjects with severe asthma found that increased in the number of chymase-positive

mast cells in the small airway was associated with increased in lung function (Balzar *et al.* 2005).

Overall, while there is still controversy on the direction of effect, previous findings support the

association of our novel *KITLG* locus with lung function, especially in patients with allergic asthma.

Our novel locus likely represents part of a complex regulatory mechanism that modulates

immune cell differentiation, survival, and activation in highly cell-specific and context-dependent

manners. Further studies are required to study how this locus is regulated in different airway and

immune cells to affect lung function outcome in the context of asthma.

GxE interactions likely account for a portion of the "missing" heritability of many complex

phenotypes (Moore and Williams. 2009). We previously found that lung function in SAGE II

participants was associated with first year of life and lifetime exposures to SO_2 (1.66% decrease

[95% CI = -2.92 to -0.37] for first year of life and 5.30% decrease [95% CI = -8.43 to -2.06] for

lifetime exposures in FEV_1 per 1 ppb increases in SO_2) (Neophytou *et al.* 2016). We hypothesized

that a significant portion of the heritability of lung function was due in part to gene-by-air-

pollution (SO_2) interaction effects. The interaction between rs58475486 and past year exposure

to SO_2 that was significantly associated with lung function supports our hypothesis. The T allele

of rs58475486 is common (8-14%) in African populations and showed a protective effect on lung

function in the presence of past year SO_2 exposure. SNP rs58475486 is located in a ccRE (SCREEN

accession EH37E0279296) and a FOXA1 binding site in the A549 lung adenocarcinoma cell line.

FOXA1 has a known compensatory role with FOXA2 during lung morphogenesis in mice (Wan *et*

al. 2005). Deletion of both FOXA1 and FOXA2 inhibited cell proliferation, epithelial cell

differentiation, and branching morphogenesis in fetal lung tissue. Further functional validation

on the effect of rs58475486 on binding affinity of FOXA1 is necessary to confirm whether the role

of FOXA1 in this ccRE is important for *KITLG* regulatory and lung function.

The higher frequency of the protective alleles of both rs73440122 and rs58465486 in African

populations appears to contradict previous findings that African ancestry was associated with

lower lung function (Kumar *et al.* 2010). One possible explanation for this seeming inconsistency

is that FEV_1 is a complex trait whose variation is influenced by many genetic variants of small to

moderate effect sizes whose influence on lung function may vary by exposure to environmental

factors. We found suggestive evidence that the interaction between rs73440122 and first year

exposure to SO_2 reverses the positive association of rs73440122 with lung function to a negative

one (Table 3). When assessed independently, our genetic association analysis showed that the

protective A allele of rs73440122 was associated with higher lung function. However, with

increasing levels of SO_2 exposure in the first year of life, increasing copies of the A allele of

rs73440122 were associated with decreased lung function. Air pollution is known to negatively

impact lung function, and we have previously shown that the deleterious effects of air pollution

on lung phenotypes may be significantly increased in African American children compared to

other populations experiencing the same amount of exposure (Nishimura *et al.* 2013). It has also

been reported that Latino and African American populations often live in neighborhoods with

high levels of air pollution (Mott 1995). The increased susceptibility to negative pulmonary effects

from air pollution exposure coupled with the disproportionate exposure to air pollution

experienced by the African American population may also contribute to the lower lung function

seen in this population despite the presence of protective alleles. The overlap of the SPI1 binding

site with rs73440122 further supports gene-by-SO_2 interaction at this locus, since SPI1 played a

critical role in the development of type 2 inflammation in the airways through macrophage

polarization (Qian *et al.* 2015). We noted that the rs73440122 A allele also showed an interaction

approaching suggestive threshold with past year exposure to SO_2 that was positively associated

with FEV_1. The difference is not surprising because age of exposure may significantly impact the

effect of air pollution on lung function (Reviewed in Usemann *et al.* 2019) . Further studies are

required to better understand the effect of this suggestive interaction on lung function.

One strength of this study is the interrogation of independent lung function associated signals at

our novel locus. We identified evidence of three independent signals: the replicated signal that

showed evidence of regulatory functions (an open chromatin region with a SPI1/PU.1 binding

site), one signal that showed a statistically significant gene-by-SO_2 interaction on lung function,

and one signal that represents to *KITLG* eQTLs in the nasal epithelial cells together with suggestive

gene-by-SO_2 interaction. Our results demonstrated a glimpse of the complicated genetic

architecture behind complex traits.

One limitation of this study is that the FEV_1 genetic association and the eQTL analyses with *KITLG*

were performed in different populations due to data availability constraints. Although we did not

have RNA-Seq data from lung tissues from our study subjects, we previously demonstrated that

there is a high degree of overlap in gene expression profiles between nasal and bronchial

epithelial cells (Poole *et al.* 2014). The direction of effect of the association was the same in GALA

II Puerto Rican children with asthma but not statistically significant. This may in part due to the

significantly lower African Ancestry in Puerto Ricans compared to African Americans.

We replicated 20 of 45 variants in black UK Biobank subjects and observed conflicting "flip-flop"

associations in African Americans from the SAPPHIRE and GCPD-A studies. In the past, flip-flop

associations were deemed as spurious results. Traditional association testing approach studies

the effect of each variant on phenotype independently and increases the chance of flip-flop

associations detected between studies. Differences in study design, sampling variation that leads

to variation in LD patterns, and lack of consideration of other disease influencing genetic and/or

environmental factors are all potential causes of flip-flop associations (Kraft *et al.* 2009; Lin *et al.*

2007). Hence, it is not surprising to observe flip-flop associations when gene and environment

interactions were detected at our FEV_1 GWAS locus. It was previously shown that flip-flop

associations can occur between and within populations even in the presence of a genuine genetic

effect (Kraft *et al.* 2009; Lin *et al.* 2007). Further functional analysis is thus required to validate

the relationship between the candidate variants, *KITLG* and FEV_1. This may include reporter

assays to validate potential enhancer or repressor activity and CRISPR-based editing assays to

validate the regulatory role of the candidate variants on *KITLG*. Although literature exists

describing KIT signaling for lung function in mice (Lindsey *et al.* 2011), additional knockout

experiments in a model animal system may be necessary to study how *KITLG* contributed to

variation in lung function.

The average concentration of ambient SO_2 exposure in our participants (Table 1) was lower than

the National Ambient Air Quality Standards. It is possible that SO_2 acted as a surrogate for other

unmeasured toxic pollutants emitted from local point sources. Major sources of SO_2 in San

Francisco Bay Area during the recruitment years of 2006 to 2011 include airports, petroleum

refineries, gas and oil plants, calcined petroleum coke plants, electric power plants, cement

manufacturing factories, chemical plants, and landfills (United States Environmental Protection

Agency 2008; United States Environmental Protection Agency 2011). The Environmental

Protection Agency's national emissions inventory data also showed that these facilities emit

Volatile Organic Compounds, heavy metals (lead, mercury, chromium, arsenic), formaldehyde,

ethyl benzene, acrolein, 1,3-butadiene, 1,4-dichlorobenzene, and tetrachloroethylene into the

air along with SO_2. These chemicals are highly toxic and inhaling even a small amount may

contribute to poor lung function. Another possibility is that exposure to SO_2 captured

unmeasured confounding socioeconomic factors.

This study identified a novel protective allele for lung function in African American children with

asthma. The protective association with lung function intensified with increased past year

exposure to SO_2. Our findings showcase the complexity of the relationship between genetic and

environmental factors impacting variation in FEV_1, highlights the utility of WGS data for genetic

research of complex phenotypes, and underscores the importance of including diverse study

populations in our exploration of the genetic architecture underlying lung function.

ACKNOWLEDGEMENTS

The Genes-Environments and Admixture in Latino Americans (GALA II) Study, the Study of African

Americans, Asthma, Genes and Environments (SAGE) Study and E.G.B. were supported by the

Sandler Family Foundation, the American Asthma Foundation, the RWJF Amos Medical Faculty

Development Program, the Harry Wm. and Diana V. Hind Distinguished Professor in

Pharmaceutical Sciences II, the National Heart, Lung, and Blood Institute (NHLBI) [R01HL117004,

R01HL128439, R01HL135156, X01HL134589]; the National Institute of Environmental Health

Sciences [R01ES015794]; the National Institute on Minority Health and Health Disparities

(NIMHD) [P60MD006902, R01MD010443], the National Human Genome Research Institute

[U01HG009080] and the Tobacco-Related Disease Research Program [24RT-0025]. MJW was

supported by the NHLBI [K01HL140218]. JJ and BEH were supported by the NHLBI [R01HL133433,

R01HL141992]. KLK was supported by the NHLBI [R01HL135156-S1], the UCSF Bakar Institute, the

Gordon and Betty Moore Foundation [GBMF3834], and the Alfred P. Sloan Foundation [2013-10-

27] grant to UC Berkeley through the Moore-Sloan Data Science Environment Initiative. ACW was

supported by the Eunice Kennedy Shriver National Institute of Child Health and Human

Development [1R01HD085993-01].

The SAPPHIRE study was supported by the Fund for Henry Ford Hospital, the American Asthma

Foundation, the NHLBI [R01HL118267, R01HL141485, X01HL134589], the National Institute of

Allergy and Infectious Diseases [R01AI079139], and the National Institute of Diabetes and

Digestive and Kidney Diseases [R01DK113003].

The GCPD-A study was supported by an Institutional award from the Children's Hospital of

Philadelphia and by the NHLBI [X01HL134589].

Part of this research was conducted using the UK Biobank Resource under Application Number

40375. We would like to thank UK Biobank participants and researchers who contributed or

collected data.

Whole genome sequencing (WGS) for the Trans-Omics in Precision Medicine (TOPMed) program

was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for "NHLBI TOPMed:

Gene-Environment, Admixture and Latino Asthmatics Study" (phs000920) and "NHLBI TOPMed:

Study of African Americans, Asthma, Genes and Environments" (phs000921) was performed at

the New York Genome Center (3R01HL117004-02S3) and the University of Washington

Northwest Genomics Center (HHSN268201600032I). WGS for "NHLBI TOPMed: Study of Asthma

Phenotypes & Pharmacogenomic Interactions by Race-Ethnicity" (phs001467) and "Genetics of

Complex Pediatric Disorders - Asthma" (phs001661) was performed at the University of

Washington Northwest Genomics Center (HHSN268201600032I). Centralized read mapping and

genotype calling, along with variant quality metrics and filtering were provided by the TOPMed

Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Phenotype

harmonization, data management, sample-identity QC, and general study coordination were

provided by the TOPMed Data Coordinating Center (3R01HL-120393-02S1; contract

HHSN268201800001I). We gratefully acknowledge the studies and participants who provided

biological samples and data for TOPMed.

WGS of part of GALA II was performed by New York Genome Center under The Centers for

Common Disease Genomics of the Genome Sequencing Program (GSP) Grant (UM1 HG008901).

The GSP Coordinating Center (U24 HG008956) contributed to cross-program scientific initiatives

and provided logistical and general study coordination. GSP is funded by the National Human

Genome Research Institute, the National Heart, Lung, and Blood Institute, and the National Eye

Institute.

The TOPMed imputation panel was supported by the NHLBI and TOPMed study investigators who

contributed data to the reference panel. The panel was constructed and implemented by the

TOPMed Informatics Research Center at the University of Michigan (3R01HL-117626-02S1;

contract HHSN268201800002I). The TOPMed Data Coordinating Center (3R01HL-120393-02S1;

contract HHSN268201800001I) provided additional data management, sample identity checks,

and overall program coordination and support. We gratefully acknowledge the studies and

participants who provided biological samples and data for TOPMed.

The authors wish to acknowledge the following GALA II and SAGE study collaborators: Shannon

Thyne, UCSF; Harold J. Farber, Texas Children's Hospital; Denise Serebrisky, Jacobi Medical Center;

Rajesh Kumar, Lurie Children's Hospital of Chicago; Emerita Brigino-Buenaventura, Kaiser

Permanente; Michael A. LeNoir, Bay Area Pediatrics; Kelley Meade, UCSF Benioff Children's

Hospital, Oakland; William Rodríguez-Cintrón, VA Hospital, Puerto Rico; Pedro C. Ávila,

Northwestern University; Jose R. Rodríguez-Santana, Centro de Neumología Pediátrica; Luisa N.

Borrell, City University of New York; Adam Davis, UCSF Benioff Children's Hospital, Oakland;

Saunak Sen, University of Tennessee.

The authors acknowledge the families and patients for their participation and thank the

numerous health care providers and community clinics for their support and participation in

GALA II and SAGE. In particular, the authors thank the recruiters who obtained the data: Duanny

Alva, MD; Gaby Ayala-Rodríguez; Lisa Caine, RT; Elizabeth Castellanos; Jaime Colón; Denise

DeJesus; Blanca López; Brenda López, MD; Louis Martos; Vivian Medina; Juana Olivo; Mario

Peralta; Esther Pomares, MD; Jihan Quraishi; Johanna Rodríguez; Shahdad Saeedi; Dean Soto;

and Ana Taveras.

The authors thank María Pino-Yanes for providing feedback on this study and Thomas W

Blackwell for providing critical review on this manuscript.

The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health.

669   LITERATURE CITED

1000 Genomes Project Consortium, A. Auton, L. D. Brooks, R. M. Durbin, E. P. Garrison *et al*,

2015 A global reference for human genetic variation. Nature **526:** 68-74.

Akinbami, L. J., 2015 *Asthma Prevalence, Health Care use and Mortality: United States, 2003-05*.

[Online] Available at: http://www.cdc.gov/nchs/data/hestat/asthma03-05/asthma03-05.htm.

[Accessed 2020 Jan 8].

Akinbami, L. J., J. E. Moorman, A. E. Simon and K. C. Schoendorf, 2014 Trends in racial

disparities for asthma outcomes among children 0 to 17 years, 2001-2010. J. Allergy Clin.

Immunol. **134:** 547-553.e5.

Alexander, D. H., J. Novembre and K. Lange, 2009 Fast model-based estimation of ancestry in

unrelated individuals. Genome Res. **19:** 1655-1664.

Al-Muhsen, S. Z., G. Shablovsky, R. Olivenstein, B. Mazer and Q. Hamid, 2004 The expression of

stem cell factor and c-kit receptor in human asthmatic airways. Clin. Exp. Allergy **34:** 911-916.

Amin, K., 2012 The role of mast cells in allergic inflammation. Respir. Med. **106:** 9-14.

Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq--a python framework to work with high-

throughput sequencing data. Bioinformatics **31:** 166-169.

Anderson, K. L., H. Perkin, C. D. Surh, S. Venturini, R. A. Maki *et al*, 2000 Transcription factor

PU.1 is necessary for development of thymic and myeloid progenitor-derived dendritic cells. J.

Immunol. **164:** 1855-1861.

Ay, F., T. L. Bailey and W. S. Noble, 2014 Statistical confidence estimation for hi-C data reveals

regulatory chromatin contacts. Genome Res. **24:** 999-1011.

Balzar, S., H. W. Chu, M. Strand and S. Wenzel, 2005 Relationship of small airway chymase-

positive mast cells and lung function in severe asthma. Am. J. Respir. Crit. Care Med. **171:** 431-

439.

Barraza-Villarreal, A., J. Sunyer, L. Hernandez-Cadena, M. C. Escamilla-Nunez, J. J. Sienra-Monge

et al, 2008 Air pollution, airway inflammation, and lung function in a cohort study of mexico city

schoolchildren. Environ. Health Perspect. **116:** 832-838.

Barrett, J. C., B. Fry, J. Maller and M. J. Daly, 2005 Haploview: Analysis and visualization of LD

and haplotype maps. Bioinformatics **21:** 263-265.

Borish, L., and B. Z. Joseph, 1992 Inflammation and the allergic response. Med. Clin. North Am.

76: 765-787.

Broudy, V. C., 1997 Stem cell factor and hematopoiesis. Blood **90:** 1345-1364.

Brunekreef, B., and S. T. Holgate, 2002 Air pollution and health. Lancet **360:** 1233-1242.

Buniello, A., J. A. L. MacArthur, M. Cerezo, L. W. Harris, J. Hayhurst *et al*, 2019 The NHGRI-EBI

GWAS catalog of published genome-wide association studies, targeted arrays and summary

statistics 2019. Nucleic Acids Res. **47:** D1005-D1012.

Bycroft, C., C. Freeman, D. Petkova, G. Band, L. T. Elliott *et al*, 2018 The UK biobank resource

with deep phenotyping and genomic data. Nature **562:** 203-209.

Canela-Xandri, O., K. Rawlik and A. Tenesa, 2018 An atlas of genetic associations in UK biobank.

Nat. Genet. **50:** 1593-1599.

Carlson, C. S., T. C. Matise, K. E. North, C. A. Haiman, M. D. Fesinmeyer *et al*, 2013

Generalization and dilution of association results from european GWAS in populations of non-

european ancestry: The PAGE study. PLoS Biol. **11:** e1001661.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al*, 2015 Second-generation

PLINK: Rising to the challenge of larger and richer datasets. Gigascience **4:** 7-8. eCollection

2015.

Chatterjee, S., and N. Das, 1995 Lung function in indian twin children: Comparison of genetic

versus environmental influence. Ann. Hum. Biol. **22:** 289-303.

Chen, Y., S. L. Horne, D. C. Rennie and J. A. Dosman, 1996 Segregation analysis of two lung

function indices in a random sample of young families: The humboldt family study. Genet.

Epidemiol. **13:** 35-47.

Ciesielski, T. H., S. A. Pendergrass, M. J. White, N. Kodaman, R. S. Sobota *et al*, 2014 Diverse

convergent evidence in the genetic analysis of complex disease: Coordinating omic, informatic,

and experimental evidence to better identify and validate risk factors. BioData Min. **7:** 10-10.

eCollection 2014.

Comberiati, P., M. E. Di Cicco, S. D'Elios and D. G. Peroni, 2017 How much asthma is atopic in

children? Front. Pediatr. **5:** 122.

Conomos, M. P., M. B. Miller and T. A. Thornton, 2015 Robust inference of population structure

for ancestry prediction and correction of stratification in the presence of relatedness. Genet.

Epidemiol. **39:** 276-293.

Conomos, M. P., A. P. Reiner, B. S. Weir and T. A. Thornton, 2016 Model-free estimation of

recent genetic relatedness. Am. J. Hum. Genet. **98:** 127-148.

Cruse, G., and P. Bradding, 2016 Mast cells in airway diseases and interstitial lung disease. Eur.

J. Pharmacol. **778:** 125-138.

Da Silva, C. A., and N. Frossard, 2005 Regulation of stem cell factor expression in inflammation

and asthma. Mem. Inst. Oswaldo Cruz **100 Suppl 1:** 145-151.

Da Silva, C. A., L. Reber and N. Frossard, 2006 Stem cell factor expression, mast cells and

inflammation in asthma. Fundam. Clin. Pharmacol. **20:** 21-39.

Dahlin, J. S., A. Malinovschi, H. Ohrvik, M. Sandelin, C. Janson *et al*, 2016 Lin- CD34hi

CD117int/hi FcepsilonRI+ cells in human blood constitute a rare population of mast cell

progenitors. Blood **127:** 383-391.

Das, S., L. Forer, S. Schonherr, C. Sidore, A. E. Locke *et al*, 2016 Next-generation genotype

imputation service and methods. Nat. Genet. **48:** 1284-1287.

Duggal, P., E. M. Gillanders, T. N. Holmes and J. E. Bailey-Wilson, 2008 Establishing an adjusted

p-value threshold to control the family-wide type 1 error in genome wide association studies.

BMC Genomics **9:** 516-516.

ENCODE Project Consortium, 2012 An integrated encyclopedia of DNA elements in the human

genome. Nature **489:** 57-74.

ENCODE Project Consortium, 2011 A user's guide to the encyclopedia of DNA elements

(ENCODE). PLoS Biol. **9:** e1001046.

Fajt, M. L., and S. E. Wenzel, 2013 Mast cells, their subtypes, and relation to asthma

phenotypes. Ann. Am. Thorac. Soc. **10 Suppl:** 158.

Fonseca, W., A. J. Rasky, C. Ptaschinski, S. H. Morris, S. K. K. Best *et al*, 2019 Group 2 innate

lymphoid cells (ILC2) are regulated by stem cell factor during chronic asthmatic disease.

Mucosal Immunol. **12:** 445-456.

Galli, S. J., K. M. Zsebo and E. N. Geissler, 1994 The kit ligand, stem cell factor. Adv. Immunol.

55: 1-96.

Galli, S. J., M. Tsai, B. K. Wershil, S. Y. Tam and J. J. Costa, 1995 Regulation of mouse and human

mast cell development, survival and function by stem cell factor, the ligand for the c-kit

receptor. Int. Arch. Allergy Immunol. **107:** 51-53.

GTEx Consortium, Laboratory, Data Analysis &Coordinating Center (LDACC)-Analysis Working

Group, Statistical Methods groups-Analysis Working Group, Enhancing GTEx (eGTEx) groups,

NIH Common Fund *et al*, 2017 Genetic effects on gene expression across human tissues. Nature

550: 204-213.

Guerriero, A., P. B. Langmuir, L. M. Spain and E. W. Scott, 2000 PU.1 is required for myeloid-

derived but not lymphoid-derived dendritic cells. Blood **95:** 879-885.

Hankinson, J. L., J. R. Odencrantz and K. B. Fedan, 1999 Spirometric reference values from a

sample of the general U.S. population. Am. J. Respir. Crit. Care Med. **159:** 179-187.

Hinrichs, A. S., D. Karolchik, R. Baertsch, G. P. Barber, G. Bejerano *et al*, 2006 The UCSC genome

browser database: Update 2006. Nucleic Acids Res. **34:** 590.

Hsieh, F. H., P. Sharma, A. Gibbons, T. Goggans, S. C. Erzurum *et al*, 2005 Human airway

epithelial cell determinants of survival and functional phenotype for primary human mast cells.

Proc. Natl. Acad. Sci. U. S. A. **102:** 14380-14385.

Hukkinen, M., J. Kaprio, U. Broms, A. Viljanen, D. Kotz *et al*, 2011 Heritability of lung function: A

twin study among never-smoking elderly women. Twin Res. Hum. Genet. **14:** 401-407.

Iemura, A., M. Tsai, A. Ando, B. K. Wershil and S. J. Galli, 1994 The c-kit ligand, stem cell factor,

promotes mast cell survival by suppressing apoptosis. Am. J. Pathol. **144:** 321-328.

Ierodiakonou, D., A. Zanobetti, B. A. Coull, S. Melly, D. S. Postma *et al*, 2016 Ambient air

pollution, lung function, and airway responsiveness in asthmatic children. J. Allergy Clin.

Immunol. **137:** 390-399.

Ionita-Laza, I., S. Lee, V. Makarov, J. D. Buxbaum and X. Lin, 2013 Sequence kernel association

tests for the combined effect of rare and common variants. Am. J. Hum. Genet. **92:** 841-853.

Ito, T., C. Nishiyama, M. Nishiyama, H. Matsuda, K. Maeda *et al*, 2005 Mast cells acquire

monocyte-specific gene expression and monocyte-like morphology by overproduction of PU.1.

J. Immunol. **174:** 376-383.

Ito, T., C. Nishiyama, N. Nakano, M. Nishiyama, Y. Usui *et al*, 2009 Roles of PU.1 in monocyte-

and mast cell-specific gene regulation: PU.1 transactivates CIITA pIV in cooperation with IFN-

gamma. Int. Immunol. **21:** 803-816.

Jiang, H., R. Lei, S. W. Ding and S. Zhu, 2014 Skewer: A fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. BMC Bioinformatics **15:** 182-182.

Johnson, J. D., and W. M. Theurer, 2014 A stepwise approach to the interpretation of

pulmonary function tests. Am. Fam. Physician **89:** 359-366.

Karolchik, D., A. S. Hinrichs, T. S. Furey, K. M. Roskin, C. W. Sugnet *et al*, 2004 The UCSC table

browser data retrieval tool. Nucleic Acids Res. **32:** 493.

Kassel, O., F. Schmidlin, C. Duvernelle, B. Gasser, G. Massard *et al*, 1999 Human bronchial

smooth muscle cells in culture produce stem cell factor. Eur. Respir. J. **13:** 951-954.

Kim, D., B. Langmead and S. L. Salzberg, 2015 HISAT: A fast spliced aligner with low memory

requirements. Nat. Methods **12:** 357-360.

Kim, M. S., K. P. Patel, A. K. Teng, A. J. Berens and J. Lachance, 2018 Genetic disease risks can be

misestimated across global populations. Genome Biol. **19:** 179-7.

Kobayashi, H., Y. Okayama, T. Ishizuka, R. Pawankar, C. Ra *et al*, 1998 Production of IL-13 by

human lung mast cells in response to fcepsilon receptor cross-linkage. Clin. Exp. Allergy **28:**

1219-1227.

Kraft, P., E. Zeggini and J. P. Ioannidis, 2009 Replication in genome-wide association studies.

Stat. Sci. **24:** 561-573.

Kumar, R., M. A. Seibold, M. C. Aldrich, L. K. Williams, A. P. Reiner *et al*, 2010 Genetic ancestry

in lung-function predictions. N. Engl. J. Med. **363:** 321-330.

Lee, S., M. J. Emond, M. J. Bamshad, K. C. Barnes, M. J. Rieder *et al*, 2012 Optimal unified

approach for rare-variant association testing with application to small-sample case-control

whole-exome sequencing studies. Am. J. Hum. Genet. **91:** 224-237.

Levin, A. M., Y. Wang, K. E. Wells, B. Padhukasahasram, J. J. Yang *et al*, 2014 Nocturnal asthma

and the importance of race/ethnicity and genetic ancestry. Am. J. Respir. Crit. Care Med. **190:**

266-273.

Li, X., G. A. Hawkins, E. J. Ampleford, W. C. Moore, H. Li *et al*, 2013 Genome-wide association

study identifies TH1 pathway genes associated with lung function in asthmatic patients. J.

Allergy Clin. Immunol. **132:** 313-20.e15.

Liao, S. Y., X. Lin and D. C. Christiani, 2014 Genome-wide association and network analysis of

lung function in the framingham heart study. Genet. Epidemiol. **38:** 572-578.

Lin, P. I., J. M. Vance, M. A. Pericak-Vance and E. R. Martin, 2007 No gene is an island: The flip-

flop phenomenon. Am. J. Hum. Genet. **80:** 531-538.

Lindsey, J. Y., K. Ganguly, D. M. Brass, Z. Li, E. N. Potts *et al*, 2011 C-kit is essential for alveolar

maintenance and protection from emphysema-like disease in mice. Am. J. Respir. Crit. Care

Med. **183:** 1644-1652.

Liu, X., S. White, B. Peng, A. D. Johnson, J. A. Brody *et al*, 2016 WGSA: An annotation pipeline

for human genome sequencing studies. J. Med. Genet. **53:** 111-112.

Mak, A. C. Y., M. J. White, W. L. Eckalbar, Z. A. Szpiech, S. S. Oh *et al*, 2018 Whole-genome

sequencing of pharmacogenetic drug response in racially diverse children with asthma. Am. J.

Respir. Crit. Care Med. **197:** 1552-1564.

Mak, A. C., M. J. White, C. Eng, D. Hu, S. Huntsman *et al*, 2016 *Whole Genome Sequencing to*

Identify Genetic Variation Associated with Bronchodilator Response in Minority Children with

Asthma.

Martin, A. R., C. R. Gignoux, R. K. Walters, G. L. Wojcik, B. M. Neale *et al*, 2017 Human

demographic history impacts genetic risk prediction across diverse populations. Am. J. Hum.

Genet. **100:** 635-649.

Martin, J. S., Z. Xu, A. P. Reiner, K. L. Mohlke, P. Sullivan *et al*, 2017 HUGIn: Hi-C unifying

genomic interrogator. Bioinformatics **33:** 3793-3795.

McKercher, S. R., B. E. Torbett, K. L. Anderson, G. W. Henkel, D. J. Vestal *et al*, 1996 Targeted

disruption of the PU.1 gene results in multiple hematopoietic abnormalities. EMBO J. **15:** 5647-

5658.

Mekori, Y. A., C. K. Oh and D. D. Metcalfe, 1993 IL-3-dependent murine mast cells undergo

apoptosis on removal of IL-3. prevention of apoptosis by c-kit ligand. J. Immunol. **151:** 3775-

3784.

Mendez-Enriquez, E., and J. Hallgren, 2019 Mast cells and their progenitors in allergic asthma.

Front. Immunol. **10:** 821.

Moore, J. H., 2005 A global view of epistasis. Nat. Genet. **37:** 13-14.

Moore, J. H., and S. M. Williams, 2009 Epistasis and its implications for personal genetics. Am. J.

Hum. Genet. **85:** 309-320.

Mott, L., 1995 The disproportionate impact of environmental health threats on children of

color. Environ. Health Perspect. **103 Suppl 6:** 33-35.

Neophytou, A. M., M. J. White, S. S. Oh, N. Thakur, J. M. Galanter *et al*, 2016 Air pollution and

lung function in minority youth with asthma in the GALA II (genes-environments and admixture

in latino americans) and SAGE II (study of african americans, asthma, genes, and environments)

studies. Am. J. Respir. Crit. Care Med. **193:** 1271-1280.

Nishimura, K. K., J. M. Galanter, L. A. Roth, S. S. Oh, N. Thakur *et al*, 2013 Early-life air pollution

and asthma risk in minority children. the GALA II and SAGE II studies. Am. J. Respir. Crit. Care

Med. **188:** 309-318.

Nishiyama, C., M. Nishiyama, T. Ito, S. Masaki, N. Masuoka *et al*, 2004 Functional analysis of

PU.1 domains in monocyte-specific gene regulation. FEBS Lett. **561:** 63-68.

Nishiyama, C., M. Nishiyama, T. Ito, S. Masaki, K. Maeda *et al*, 2004 Overproduction of PU.1 in

mast cell progenitors: Its effect on monocyte- and mast cell-specific gene expression. Biochem.

Biophys. Res. Commun. **313:** 516-521.

Oh, S. S., M. J. White, C. R. Gignoux and E. G. Burchard, 2016 Making precision medicine socially

precise. take a deep breath. Am. J. Respir. Crit. Care Med. **193:** 348-350.

Oh, S. S., H. Tcheurekdjian, L. A. Roth, E. A. Nguyen, S. Sen *et al*, 2012 Effect of secondhand

smoke on asthma control among black and latino children. J. Allergy Clin. Immunol. **129:** 1478-

83.e7.

Oliveira, S. H., and N. W. Lukacs, 2003 Stem cell factor: A hemopoietic cytokine with important

targets in asthma. Curr. Drug Targets Inflamm. Allergy **2:** 313-318.

Ong, B. A., J. Li, J. M. McDonough, Z. Wei, C. Kim *et al*, 2013 Gene network analysis in a

pediatric cohort identifies novel lung function genes. PLoS One **8:** e72899.

Oriss, T. B., N. Krishnamoorthy, P. Ray and A. Ray, 2014 Dendritic cell c-kit signaling and

adaptive immunity: Implications for the upper airways. Curr. Opin. Allergy Clin. Immunol. **14:** 7-

12.

Palmer, L. J., M. W. Knuiman, M. L. Divitini, P. R. Burton, A. L. James *et al*, 2001 Familial

aggregation and heritability of adult lung function: Results from the busselton health study. Eur.

Respir. J. **17:** 696-702.

Pe'er, I., R. Yelensky, D. Altshuler and M. J. Daly, 2008 Estimation of the multiple testing burden

for genomewide association studies of nearly all common variants. Genet. Epidemiol. **32:** 381-

385.

Pino-Yanes, M., N. Thakur, C. R. Gignoux, J. M. Galanter, L. A. Roth *et al*, 2015 Genetic ancestry

influences asthma susceptibility and lung function among latinos. J. Allergy Clin. Immunol. **135:**

228-235.

Plummer, M., N. Best, K. Cowles and K. Vines, 2006 CODA: Convergence diagnosis and output

analysis for MCMC. R News **6:** 7-11.

Poole, A., C. Urbanek, C. Eng, J. Schageman, S. Jacobson *et al*, 2014 Dissecting childhood asthma

with nasal transcriptomics distinguishes subphenotypes of disease. J. Allergy Clin. Immunol.

133: 670-8.e12.

Pruim, R. J., R. P. Welch, S. Sanna, T. M. Teslovich, P. S. Chines *et al*, 2010 LocusZoom: Regional

visualization of genome-wide association scan results. Bioinformatics **26:** 2336-2337.

Purcell, S., and C. Chang, 2013 *Plink 1.9*. [Online] Available at: www.cog-

genomics.org/plink/1.9/. [Accessed 2019 Mar].

Qian, F., J. Deng, Y. G. Lee, J. Zhu, M. Karpurapu *et al*, 2015 The transcription factor PU.1

promotes alternative macrophage polarization and asthmatic airway inflammation. J. Mol. Cell.

Biol. **7:** 557-567.

Repapi, E., I. Sayers, L. V. Wain, P. R. Burton, T. Johnson *et al*, 2010 Genome-wide association

study identifies five loci associated with lung function. Nat. Genet. **42:** 36-44.

Rochman, M., A. V. Kartashov, J. M. Caldwell, M. H. Collins, E. M. Stucke *et al*, 2015

Neurotrophic tyrosine kinase receptor 1 is a direct transcriptional and epigenetic target of IL-13

involved in allergic inflammation. Mucosal Immunol. **8:** 785-798.

Schmitt, A. D., M. Hu, I. Jung, Z. Xu, Y. Qiu *et al*, 2016 A compendium of chromatin contact maps

reveals spatially active regions in the human genome. Cell. Rep. **17:** 2042-2059.

Scott, E. W., M. C. Simon, J. Anastasi and H. Singh, 1994 Requirement of transcription factor

PU.1 in the development of multiple hematopoietic lineages. Science **265:** 1573-1577.

Scott, E. W., R. C. Fisher, M. C. Olson, E. W. Kehrli, M. C. Simon *et al*, 1997 PU.1 functions in a

cell-autonomous manner to control the differentiation of multipotential lymphoid-myeloid

progenitors. Immunity **6:** 437-447.

Sillanpaa, E., S. Sipila, T. Tormakangas, J. Kaprio and T. Rantanen, 2017 Genetic and

environmental effects on telomere length and lung function: A twin study. J. Gerontol. A Biol.

Sci. Med. Sci. **72:** 1561-1568.

Sofer, T., X. Zheng, S. M. Gogarten, C. A. Laurie, K. Grinde *et al*, 2019 A fully adjusted two-stage

procedure for rank-normalization in genetic association studies. Genet. Epidemiol. **43:** 263-275.

Soler Artigas, M., L. V. Wain, S. Miller, A. K. Kheirallah, J. E. Huffman *et al*, 2015 Sixteen new

lung function signals identified through 1000 genomes project reference panel imputation. Nat.

Commun. **6:** 8658.

Soler Artigas, M., D. W. Loth, L. V. Wain, S. A. Gharib, M. Obeidat *et al*, 2011 Genome-wide

association and large-scale follow up identifies 16 new loci influencing lung function. Nat.

Genet. **43:** 1082-1090.

Summer Institute in Statistical Genetics, 2019 *PC-Relate*. [Online] Available at: https://uw-

gac.github.io/SISG_2019/pc-relate.html. [Accessed 2019 Jul 25].

Taliun, D., D. N. Harris, M. D. Kessler, J. Carlson, Z. A. Szpiech *et al*, 2019 Sequencing of 53,831

diverse genomes from the NHLBI TOPMed program. bioRxiv 563866.

Tayel, S. I., S. M. El-Hefnway, Abd El Gayed, E. M. and G. A. Abdelaal, 2017 Association of stem

cell factor gene expression with severity and atopic state in patients with bronchial asthma.

Respir. Res. **18:** 21-2.

Tian, X., C. Xu, Y. Wu, J. Sun, H. Duan *et al*, 2017 Genetic and environmental influences on

pulmonary function and muscle strength: The chinese twin study of aging. Twin Res. Hum.

Genet. **20:** 53-59.

TOPMed, 2019 *TOPMed Whole Gneome Sequencing Methods: Freeze 8*. [Online] Available at:

https://www.nhlbiwgs.org/topmed-whole-genome-sequencing-methods-freeze-8. [Accessed

2019 Dec 13].

United States Environmental Protection Agency, 2011 *National Emissions Inventory (NEI) 2011*

Data. [Online] Available at: https://www.epa.gov/air-emissions-inventories/2011-national-

emissions-inventory-nei-data. [Accessed 2020 Jan 8].

United States Environmental Protection Agency, 2008 *National Emissions Inventory (NEI) 2008*

Data. [Online] Available at: https://www.epa.gov/air-emissions-inventories/2008-national-

emissions-inventory-nei-data. [Accessed 2020 Jan 8].

University of Michigan, and NHLBI TOPMed, 2018 *BRAVO Variant Browser*. [Online] Available

at: https://bravo.sph.umich.edu/freeze5/hg38/. [Accessed 2019 Aug].

Usemann, J., F. Decrue, I. Korten, E. Proietti, O. Gorlanova *et al*, 2019 Exposure to moderate air

pollution and associations with lung function at school-age: A birth cohort study. Environ. Int.

126: 682-689.

Valent, P., E. Spanblochl, W. R. Sperr, C. Sillaber, K. M. Zsebo *et al*, 1992 Induction of

differentiation of human mast cells from bone marrow and peripheral blood mononuclear cells

by recombinant human stem cell factor/kit-ligand in long-term culture. Blood **80:** 2237-2245.

Wain, L. V., N. Shrine, M. S. Artigas, A. M. Erzurumluoglu, B. Noyvert *et al*, 2017 Genome-wide

association analyses for lung function and chronic obstructive pulmonary disease identify new

loci and potential druggable targets. Nat. Genet. **49:** 416-425.

Walsh, J. C., R. P. DeKoter, H. J. Lee, E. D. Smith, D. W. Lancki *et al*, 2002 Cooperative and

antagonistic interplay between PU.1 and GATA-2 in the specification of myeloid cell fates.

Immunity **17:** 665-676.

Wan, H., S. Dingle, Y. Xu, V. Besnard, K. H. Kaestner *et al*, 2005 Compensatory roles of Foxa1

and Foxa2 during lung morphogenesis. J. Biol. Chem. **280:** 13809-13816.

Wen, L. P., J. A. Fahrni, S. Matsui and G. D. Rosen, 1996 Airway epithelial cells produce stem cell

factor. Biochim. Biophys. Acta **1314:** 183-186.

White, M. J., O. Risse-Adams, P. Goddard, M. G. Contreras, J. Adams *et al*, 2016 Novel genetic

risk factors for asthma in african american children: Precision medicine and the SAGE II study.

Immunogenetics **68:** 391-400.

Wise, J., 2019 Air pollution is linked to infant deaths and reduced lung function in children. BMJ

366: l5772.

Wojcik, G. L., M. Graff, K. K. Nishimura, R. Tao, J. Haessler *et al*, 2019 Genetic analyses of

diverse populations improves discovery for complex traits. Nature **570:** 514-518.

World Health Organization, 2017 *Asthma*. [Online] Available at:

http://www.who.int/mediacentre/factsheets/fs307/en/. [Accessed 2020 Jan 8].

Yamada, H., Y. Yatagai, H. Masuko, T. Sakamoto, H. Iijima *et al*, 2015 Heritability of pulmonary

function estimated from genome-wide SNPs in healthy japanese adults. Respir. Investig. **53:** 60-

67.

Yashiro, T., S. Nakano, K. Nomura, Y. Uchida, K. Kasakura *et al*, 2019 A transcription factor PU.1

is critical for Ccl22 gene expression in dendritic cells and macrophages. Sci. Rep. **9:** 1161-9.

Zhang, F., and J. R. Lupski, 2015 Non-coding genetic variants in human disease. Hum. Mol. Genet.

24: 102.

Table 1. Descriptive characteristics of 867 African American children with asthma included in this study.

	African American (n=867)
Age	
Mean (SD)	14.1 (3.64)
Median [25%, 75%]	13.8 [10.98, 17.11]
Sex	
Male	439 (50.6%)
Female	428 (49.4%)
Height (m)	
Mean (SD)	1.58 (0.145)
Median [25%, 75%]	1.60 [1.47, 1.68]
Any control medications* in last 2 weeks	
No	543 (62.6%)
Yes	324 (37.4%)
ICS in last 2 weeks	
No	211 (24.3%)
Yes	306 (35.3%)
Missing	350 (40.4%)
LABA in last 2 weeks	
No	5 (0.6%)
Yes	94 (10.8%)
Missing	768 (88.6%)
Leukotriene inhibitor in last 2 weeks	
No	11 (1.3%)
Yes	68 (7.8%)
Missing	788 (90.9%)
African ancestry	
Mean (SD)	0.792 (0.129)
Median [25%, 75%]	0.826 [0.759, 0.869]
Smoking status	
Never	793 (91.5%)
Past	72 (8.3%)
Current	0 (0%)
Missing	2 (0.2%)
Number of smokers in family	
0	469 (54.1%)
1	137 (15.8%)
2	42 (4.8%)
3+	10 (1.2%)
Missing	209 (24.1%)

SO2 first year exposure (ppb)

Mean (SD)	1.59 (0.961)
Median [25%, 75%]	1.50 [1.24, 1.87]
Missing	227 (26.2%)

SO2 past year exposure (ppb)

Mean (SD)	1.10 (0.302)
Median [25%, 75%]	1.08 [0.910, 1.27]
Missing	206 (23.8%)

SO2 lifetime exposure (ppb)

Mean (SD)	1.50 (0.371)
Median [25%, 75%]	1.47 [1.40, 1.54]
Missing	206 (23.8%)

TOPMed sequencing center and phase

phase 1, CAAPA	6 (0.7%)
phase 1, NYGC	460 (53.1%)
phase 3, NW	401 (46.3%)

971 25% and 75%, 25 and 75 percentiles. Control medications include inhaled corticosteroid (ICS),

972 long-acting beta-agonist (LABA), leukotriene inhibitor and/or ICS/LABA combo. SO_2 exposure

973 are hourly exposure averaged over the specified time period before spirometry testing as

974 previously described in Neophytou *et al.* 2016. ppb, parts per billion or $\mu g/m^3$.

 Table 2. Genome-wide lung function association in SAGE II children with asthma.

M	rsID	Alt	β	p	β_{norm}	p_{norm}	MAF	1000 Genomes				Score
								ALL	AFR	AMR	EUR	
1	rs11835305	T	0.126	1.93E-05	0.320	3.69E-05	0.104	0.036	0.119	0.012	0.001	0
2	rs17015963	C	0.126	1.93E-05	0.320	3.69E-05	0.104	0.036	0.120	0.012	0.001	0
3	rs58475486*	T	0.127	1.45E-05	0.323	2.81E-05	0.105	0.037	0.123	0.012	0.001	2^
4	rs17015979	T	0.127	1.45E-05	0.323	2.81E-05	0.105	0.037	0.123	0.012	0.001	0
5	rs57692452	C	0.245	1.63E-06	0.654	1.06E-06	0.033	0.011	0.030	0.006	0.001	2^
6	rs112585732	T	0.235	4.35E-07	0.625	3.06E-07	0.041	0.016	0.050	0.006	0.001	0
7	rs113837356	T	0.270	3.44E-06	0.719	2.49E-06	0.025	0.008	0.027	0.006	0.001	1
8	rs61441836	G	0.252	8.19E-07	0.671	5.46E-07	0.033	0.010	0.030	0.006	0.001	1
9	rs73438172	A	0.252	8.19E-07	0.671	5.46E-07	0.033	0.010	0.030	0.006	0.001	1
10	rs1044043958&	A	0.270	3.44E-06	0.719	2.49E-06	0.025	-	-	-	-	0
11	rs73438182	G	0.138	1.68E-04	0.378	9.18E-05	0.064	0.020	0.068	0.006	0.001	0
12	rs73438185	A	0.138	1.68E-04	0.378	9.18E-05	0.064	0.020	0.068	0.006	0.001	0
13	rs73438188	A	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	1
14	rs73438190	C	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	0
15	rs73438195	A	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	0
16	rs111857459	T	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	0
17	rs144369986&	T	0.285	1.21E-06	0.756	9.44E-07	0.025	0.008	0.026	0.006	0.001	0
18	rs73440106	G	0.181	1.86E-05	0.486	1.22E-05	0.048	0.016	0.047	0.006	0.001	0
19	rs73440107	A	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	1
20	rs111453514	C	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	1
21	rs73440112	T	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	1
22	rs73440115	G	0.297	6.97E-08	0.792	4.42E-08	0.028	0.011	0.028	0.006	0.001	1
23	rs11312747&	A	0.133	1.43E-05	0.357	8.51E-06	0.100	0.036	0.121	0.010	0.001	0
24	rs73440120	A	0.285	1.21E-06	0.756	9.44E-07	0.025	0.008	0.026	0.006	0.001	1
25	rs111289668	G	0.297	6.97E-08	0.792	4.42E-08	0.028	0.010	0.027	0.006	0.001	2^
26	rs73440122	C	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	3^
27	rs73440123	G	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	1
28	rs17016065*	G	0.112	3.19E-06	0.296	2.54E-06	0.177	0.075	0.217	0.017	0.009	1#
29	rs17016066	A	0.112	3.19E-06	0.296	2.54E-06	0.177	0.075	0.217	0.017	0.009	1#
30	rs147400083&	T	0.112	3.19E-06	0.296	2.54E-06	0.177	-	-	-	-	0
31	rs866852270	T	0.112	3.19E-06	0.296	2.54E-06	0.177	-	-	-	-	0
32	rs141293300&	C	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	1
33	rs1398303	A	0.104	1.22E-05	0.274	1.12E-05	0.186	0.077	0.223	0.020	0.009	1#
34	rs61924868	T	0.104	1.24E-05	0.275	1.14E-05	0.185	0.078	0.223	0.020	0.009	1#

M	rsID											
35	rs73440134	T	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	1
36	rs73429413	G	0.292	2.08E-07	0.775	1.55E-07	0.027	0.011	0.030	0.006	0.001	1
37	rs73429415	A	0.096	5.13E-05	0.253	4.84E-05	0.189	0.078	0.225	0.022	0.009	1#
38	rs112449284	T	0.242	4.64E-06	0.640	3.87E-06	0.031	0.012	0.035	0.007	0.001	1
39	rs111981782	C	0.296	5.78E-08	0.786	4.09E-08	0.029	0.012	0.033	0.006	0.001	1
40	rs150942400	T	0.293	6.01E-08	0.780	4.01E-08	0.029	0.012	0.034	0.006	0.002	1
41	rs147527487	C	0.086	8.49E-05	0.226	8.14E-05	0.205	0.095	0.249	0.016	0.005	0
42	rs111243672	A	0.258	6.41E-07	0.690	3.99E-07	0.032	0.014	0.037	0.007	0.004	1
43	rs73429450*	A	0.302	1.26E-08	0.801	9.01E-09	0.031	0.012	0.033	0.009	0.002	1
44	rs758775577	C	0.217	2.22E-06	0.574	1.85E-06	0.041	-	-	-	-	0
45	rs142679473&	C	0.285	6.30E-08	0.756	4.62E-08	0.031	0.012	0.033	0.009	0.002	0

Score, priority score based on statistical and functional evidences which are reported in Table S12. M, marker number that corresponds to those in Figure 2 and Table S12. Candidate variants were prioritized if they had a priority score of greater than 2 (^) or if they are eQTL of KITLG in nasal epithelial cells (#). The three independent signals identified in the conditional analyses are marked with * near the rsID. Indels were marked with the superscript & near the rsID. β (p) and β_{norm}(p_{norm}) are the effect sizes (p-values) of the genetic associations of the alternate allele (alt) with FEV_1 and FEV_1.res.rnorm respectively. MAF minor allele frequency. ALL/AFR/AMR/EUR, 1000 Genomes minor allele frequency from all/African/American/European populations. -, not available.

986 Table 3. Gene-and-environment analysis on FEV_1

Variant	Exposure	n	Variant		Exposure		GxE		
			β	p	β	p	β	p	
rs58475486_T[1]	SO_2 first year	640	0.13	6.62E-06	-0.05	0.003	-0.03	0.658	
rs57692452_C[2]	SO_2 first year	640	0.25	1.16E-06	-0.05	0.003	-0.24	0.091	
rs111289668_G[2]	SO_2 first year	640	0.31	2.53E-08	-0.05	0.003	-0.27	0.079	
rs73440122_C[2]	SO_2 first year	640	0.31	7.78E-08	-0.05	0.003	-0.32	0.045	*
rs17016065_G[3]	SO_2 first year	640	0.11	8.82E-06	-0.05	0.003	-0.08	0.108	
rs17016066_A[3]	SO_2 first year	640	0.11	8.82E-06	-0.05	0.003	-0.08	0.108	
rs1398303_A[3]	SO_2 first year	640	0.10	3.15E-05	-0.05	0.003	-0.09	0.082	
rs61924868_T[3]	SO_2 first year	640	0.10	3.30E-05	-0.05	0.003	-0.08	0.088	
rs73429415_A[3]	SO_2 first year	640	0.09	1.09E-04	-0.05	0.003	-0.09	0.069	
rs58475486_T[1]	SO_2 past year	661	0.13	6.62E-06	0.05	0.362	0.32	0.003	**
rs57692452_C[2]	SO_2 past year	661	0.25	1.16E-06	0.05	0.362	0.29	0.100	
rs111289668_G[2]	SO_2 past year	661	0.31	2.53E-08	0.05	0.362	0.41	0.037	*
rs73440122_C[2]	SO_2 past year	661	0.31	7.78E-08	0.05	0.362	0.39	0.051	
rs17016065_G[3]	SO_2 past year	661	0.11	8.82E-06	0.05	0.362	0.20	0.026	*
rs17016066_A[3]	SO_2 past year	661	0.11	8.82E-06	0.05	0.362	0.20	0.026	*
rs1398303_A[3]	SO_2 past year	661	0.10	3.15E-05	0.05	0.362	0.21	0.021	*
rs61924868_T[3]	SO_2 past year	661	0.10	3.30E-05	0.05	0.362	0.21	0.023	*
rs73429415_A[3]	SO_2 past year	661	0.09	1.09E-04	0.05	0.362	0.20	0.026	*
rs58475486_T[1]	SO_2 lifetime	661	0.13	6.62E-06	-0.13	0.001	0.26	0.173	
rs57692452_C[2]	SO_2 lifetime	661	0.25	1.16E-06	-0.13	0.001	0.47	0.221	
rs111289668_G[2]	SO_2 lifetime	661	0.31	2.53E-08	-0.13	0.001	0.32	0.444	
rs73440122_C[2]	SO_2 lifetime	661	0.31	7.78E-08	-0.13	0.001	0.29	0.489	
rs17016065_G[3]	SO_2 lifetime	661	0.11	8.82E-06	-0.13	0.001	-0.19	0.143	
rs17016066_A[3]	SO_2 lifetime	661	0.11	8.82E-06	-0.13	0.001	-0.19	0.143	
rs1398303_A[3]	SO_2 lifetime	661	0.10	3.15E-05	-0.13	0.001	-0.17	0.184	
rs61924868_T[3]	SO_2 lifetime	661	0.10	3.30E-05	-0.13	0.001	-0.17	0.199	
rs73429415_A[3]	SO_2 lifetime	661	0.09	1.09E-04	-0.13	0.001	-0.16	0.207	

*, $p < 0.05$; **, p < bonferroni p value of 0.0056 for GxE analysis. n, sample sizes for the gene-

by-SO_2 interaction analysis. Superscript 1 to 3 in the variant column, variants that are in LD with

the 3 independent signals, rs58475486, rs73429450 and rs17016065, respectively. β, effect

sizes from the main effects of the variants, exposure and GxE interaction, respectively.

Figure 1.

A.

B.

Figure 1. Manhattan and LocusZoom plots from genome-wide association study of lung function*.

(A) Manhattan plot from genome-wide association study of lung function* using linear regression

in ENCORE. Red horizontal line: CODA-adjusted genome-wide significance p-value of 2.10×10^{-8}.

Blue horizontal line: CODA-adjusted suggestive significance p-value of 4.19×10^{-7}. (B) LocusZoom

plot of rs73429450 (chr12 : 88846435) and 500 Kb flanking region. Colors show linkage

disequilibrium in the study population. * FEV_1.res.rnorm was used as the phenotype for the

association testing.

1004

1005 Figure 2.

1006

Figure 2. Integration of statistical and functional evidence for variant prioritization. Numbers and

different shades of black in the LD plot represent LD in R^2. The three independent signals

identified in the conditional analysis are marked with *. Indels are marked with [&]. Nasal eQTL,

variants eQTL of *KITLG* in nasal epithelial cells. ccREs, candidate cis-regulatory elements in

SCREEN registry. ENCORE, DNase I hypersensitivity site and/or transcription factor ChIP-Seq

overlapping with the variants. UK Biobank, SAPPHIRE, GCPD-A, replication results using Blacks in

UK Biobank and African Americans in the SAPPHIRE and GCPD-A cohorts (R = replicated at p <

0.05; F = flip-flop association at p < 0.05). Candidate, candidate variants prioritized because of

presence of two or more evidence or is nasal eQTL. + indicates presence of evidence. Boxes in

the top panel were shaded grey if results were not available.

1017

1018   Figure 3.

1019   A

1020

1021   B

1022

1023   Figure 3. Gene-by-environment interaction analysis on FEV_1. FEV_1 residuals, residuals after FEV_1

1024   was regressed on the covariates age, sex, height, controller medications, sequencing centers and

1025   the first 5 genetic PCs. FEV_1 residuals was plotted against (A) past year exposure to SO_2 stratified

1026   by the number of copies of T allele of rs58475486, (B) first year of life exposure to SO_2 stratified

1027   by the number of copies of C allele of rs73440122.