Title: **Intrinsic growth rules of patients infected, dead and cured with 2019 novel**

coronavirus in mainland China

Chuanliang Han[1△], Yimeng Liu[2,3,4△], Saini Yang[2,3,4*]

[1]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing 100875, China

[2]Key Laboratory of Environmental Change and Natural Disaster, Ministry of

Education, Beijing Normal University, Beijing 100875, China

[3]State Key Laboratory of Earth Surface Processes and Resource Ecology, Beijing

Normal University, Beijing 100875, China

[4]Academy of Disaster Reduction and Emergency Management Ministry of

Emergency Management and Ministry of Education, Faculty of Geographical Science,

Beijing Normal University, Beijing 100875, China

[△]C.L.H and Y.M.L contributed equally to this work

*Correspondence authors:

Saini Yang,

Key Laboratory of Environmental Change and Natural Disaster, Ministry of

Education & State Key Laboratory of Earth Surface Processes and Resource Ecology

& Academy of Disaster Reduction and Emergency Management Ministry of

Emergency Management and Ministry of Education, Faculty of Geographical Science,

Beijing Normal University, Beijing 100875, China

Email: yangsaini@bnu.edu.cn

Conflict of Interest: The authors declare no competing conflicts of interest.

Acknowledgements: This study is sponsored by the National Key Research and

Development Program of China (2018YFC1508903), the National Natural Science

Foundation of China (41621061) and the support of International Center for

Collaborative Research on Disaster Risk Reduction (ICCR-DRR).

Abstract

Background

An outbreak of a novel coronavirus (SARS-CoV-2)-infected pneumonia (COVID-19)

was first occurred in Wuhan, China, in December 2019 and then spread rapidly to

other regions. It has been declared that at least one confirmed case infected by

SARS-CoV-2 was found in each province of China by late January 2020.

Methods

We collected the time series data of the cumulative number of confirmed infected,

dead, and cured cases from the health commissions in 31 provinces in mainland China.

A simple descriptive model in a logistic form was formulated to infer the intrinsic

epidemic rules of COVID-19. Furthermore, we compared the intrinsic epidemic rules

of COVID-19 in Hubei with that of severe acute respiratory syndrome (SARS) in

Beijing, which was obtained from the Ministry of Public Health of China in 2003.

Results

The feature of data of three population groups in all the provinces could be well

captured (R^2>0.95) by the proposed descriptive model with slight variation among

different provinces. By comparing fitted parameters, we found that the fitted curve of

infected case is the earliest to be saturated and has the lowest semi-saturation period

in both COVID-19 and SARS. While all three population groups of SARS is later to

saturate and has longer semi-saturation period than that of COVID-19.

Conclusions

Our model is robust and stable to depict the intrinsic growth rule for the cumulative

number of confirmed infected, dead, and cured cases in 30 provinces, which could

depict the epidemic situation in mainland China very well. Despite the virus caused

SARS (SARS-CoV) and the virus caused COVID-19 (SARS-CoV-2) are homologous,

their epidemiologic characteristics are quite different. The duration of the outbreak

would be shorter for COVID-19.

Introduction

In December 2019, a novel coronavirus-infected pneumonia causing human infection

was first identified in Wuhan, a city of Hubei province located in central China with a

population of 11 million people (Zhu et al., 2020; Hui et al., 2020). The coronavirus

was firstly named as 2019-nCoV on January 12, 2020 and may have a probable bat

origin (Zhou et al., 2020). Human to human transmission through 2019-nCoV has

been confirmed among closed contacts since the middle of December 2019 (Wang et

al. 2020; Li et al., 2020). As of February 12, 2020, infections of 2019-nCoV have been

identified in all Chinese provincial administrative regions and in dozens of countries

around the world (Munster et al., 2020; Holshue et al, 2020; Ki et al., 2020;

Shigemura et al., 2020; Nshiera et al., 2020). So far, the coronavirus has been

renamed as COVID-19 on February 12, 2020 officially. Different from a previous

famous beta coronavirus in 2003, severe acute respiratory syndrome coronavirus

(SARS-CoV) (Ksiazek et al., 2003; Kuiken et al., 2003; Drosten et al., 2003) that has

a mortality rates of 10% (WHO), that of COVID-19 is 3.25% in Hubei province, and

0.71% in rest of mainland China (from Chinese Center for Disease Control and

Prevention，CCDC) up to February 19, 2020.

Here, we used a descriptive model to fit data on the 2019-nCoV cases (infection, death,

cure) from January 20, 2020 to February 19, 2020, aiming to capture the

epidemiologic characteristics and dynamics of NCIP in 31 provinces in Mainland

China. Furthermore, we compared these characteristics with SARS in 2003 using the

same model structure in order to validate this model and understand the differences

among these two diseases.

Method

Sources of data

We obtained the time series data of 2019-nCoV cases from the National Health

Commission of China, and the provincial Health Commission of 31 provincial

administrative regions in Mainland China (January 20 to February 19, 2020). The data

used in this study include the number of confirmed 2019-nCoV cases, the cumulative

number of suspected, dead, and cured cases. All cases were laboratory confirmed

following the case definition by national health commission of China (from CCDC

website). The basic test procedure has been described in detail in previous work

(Huang et al., 2020; Li et al., 2020). We only chose the cases that are tested positive

by real-time RT-PCR with specific primers and probes to SARS-CoV-2 both at the

corresponding provincial CDC and the National Institute for Viral Disease Control at

CCDC.

The severe acute respiratory syndrome (SARS) in 2003 was another fatal coronavirus

epidemics over the last two decades, we collected the data (the cumulative number of

suspected, dead, and cured cases) of SARS during March 1 to August 16, 2003, in

Beijing from the National Health and Family Planning Commission of PRC.

Epidemic curve modeling

We assumed that the population of each region was susceptible to COVID-19, the

number of susceptible people per unit time was proportional to the number of

uninfected persons, and isolation or closure of the city could affect the spread effect of

infected to susceptible people to a certain extent. Previous studies have showed that

daily cumulative number of patients could be explained by logistic function (Huang et

al., 2003; Wang et al., 2003). Based on the above assumption and previous studies, we

modeled the epidemic information of virus as logistic form (Eq. (1))

112
$$\begin{cases} \frac{dI(t)}{dt} = \alpha I(t)(1 - \frac{I(t)}{N_1}) \\ I(0) = I_0 \end{cases} \quad (1)$$

113 where N_1 is maximum number of cumulative infections, $I(t)$ is cumulative number of

114 patients at time t, α is the incidence growth rate, and I_0 is the number of infected

115 cases at the initial time.

116 Analytical solution of Eq. (1) could be written in the form of Eq. (2)

117
$$I(t) = \frac{N_1}{1 + C_1 e^{-\alpha N_1 t}} \quad (2)$$

118
$$C_1 = (\frac{N_1}{I_0} - 1)e^{\alpha N_1 t_0} \quad (3)$$

119 Similarly, we could write the analytical solutions for the dead and cured cases by Eq.

120 (4) and Eq. (6).

121
$$D(t) = \frac{N_2}{1 + C_2 e^{-\beta N_2 t}} \quad (4)$$

122
$$C_2 = (\frac{N_2}{D_0} - 1)e^{\beta N_2 t_0} \quad (5)$$

123
$$R(t) = \frac{N_3}{1 + C_3 e^{-\gamma N_3 t}} \quad (6)$$

124
$$C_3 = (\frac{N_3}{R_0} - 1)e^{\gamma N_3 t_0} \quad (7)$$

125 Where β, γ is the growth rate of dead cases and the growth rate of cured cases,

126 respectively. N_2 and N_3 is maximum number of cumulative dead and cured cases. D_0

127 and R_0 is the number of dead and cured cases at initial time.

128 Here we simplified Eqs. (2),(4),(6) and unified them as Eq. (8)

129
$$N(t) = \frac{A}{1 + e^{-k(t-t_0)}} \quad (8)$$

where $N(t)$ is the general form of $I(t)$, $D(t)$ or $R(t)$, A denotes N_1, N_2, or N_3, and k

represents α, β, or γ, indicating the intrinsic rule for growth of cases. t_0 represents

the semi-saturation period, illustrating the inflection point of the sigmoid curve.

The data processing and modeling were performed on MATLAB (The MathWorks)

with custom scripts. The nonlinear least square (NLS) algorithm was adopted for data

fitting and parameter estimation. We used the MATLAB function "nlinfit" to

minimize the sum of squared differences between the data points and the fitted values.

Eq. (8) was used in this work as the descriptive model to depict the intrinsic growth

rule for infected, dead and cured cases.

Results

As of Feb 19, 2020, laboratory-conformed COVID-19 infections have been reported in each of the 31 Chinese provincial administrative region. So far, there have been a total of 75002 confirmed cases of 2019-nCoV infections in mainland China, including 2118 deaths and 16157 cases cured (From CCDC). Hubei province contributed a majority of cases to this statistical numbers (Fig 1A). Based on the limited infected case data in the early and middle phase of this new coronavirus outbreak, we firstly established a descriptive model with three parameters for each of the 31 provinces in mainland China by (Fig 1B & Fig 2B, See Methods). Fitting goodness of models was expressed by the R-square, which ranged from 0.95 to 0.999, indicating that the models we established fitted the data well. The panel B of Figure 1 illustrates examples of time series data of infected cases and corresponding fitting curves in four provinces (provinces of Xinjiang, Beijing, Guangdong, Zhejiang and Heilongjiang). It is clear that different province varied in patterns of increase of infected cases.

---------------------------- Insert Figure 1 near here -------------------------------------

In our model, k is an important parameter to depict the intrinsic growth rule of cumulative number of infected patients (Eq. (8)). Based on our assumption, the growth pattern was supposed to follow a sigmoid form, and the parameter k here denotes how rapid the number of infected cases get to saturate. If k is large, the S form of the curve would be sharp, and the curve will rise up very quickly to be saturated to the maximum number. While if k is only slightly larger than 0, the curve would show a slow and smooth increase from 0 to the maximum.

To further illustrate the spatial distribution of the infected cases' growth pattern, we exerted further analyses of the fitted infected case curves of 30 provinces. We excluded Tibet in this analysis because the number of confirmed infected case did not

169 increase since Jan 29, 2020. And the only one confirmed case in Tibet has been cured

170 on Feb 12. The spatial distribution of parameter k is shown in Fig 2A, illustrating

171 provinces varied in the increase patterns of cumulative infected cases. Fitted curves of

172 provinces such as Jilin and Qinghai has larger value of k, which means that the

173 number of infections peaked earlier in these regions than in other provinces. By

174 contrast, provinces like Hainan, Hebei, Ningxia, has smaller value of k than other

175 provinces, suggesting the infections of these provinces would continue increasing

176 when infections of other provinces reach saturation (Fig 2B). Fig 2C shows the spatial

177 distribution of predicted maximum number of infected cases (parameter A) in each

178 province of mainland China. Our model shows that the maximum number of infected

179 COVID-19 cases in mainland China would be 81498 (95% confidence interval:

180 [79099, 83896]). Among all the provinces in mainland China, the Hubei province is

181 predicted to have largest number of infected COVID-19 cases, which would be 68056

182 (95% confidence interval: [65679,70432]). Despite Tibet is predicted to have only one

183 infected case, the model predicted Qinghai province would have 19 cases (95%

184 confidence interval: [18, 19]), which is the fewest compared with other provinces in

185 mainland China.

187 ----------------------------- Insert Figure 2 near here -------------------------------------

189 Moreover, to compare the intrinsic growth rule of infected cases, dead cases and cured

190 cases between COVID-19 and SARS, we applied the descriptive model to fit the data

191 of SARS in Beijing 2003 (Fig 3A). Since the time series data of SARS was full phase

192 of the outbreak, the parameters estimated by our model was solid (R-square>0.95).

193 We found the k value of fitted curves of infected SARS cases in Beijing was 0.18,

194 which was much lower than that of Hubei province of COVID-19 ($k = 0.25$) (Fig 3

195 ABC). Furthermore, we found that all 30 provinces' k value were significantly larger

196 than 0.18 (Fig 3D). We also noted the t_0 of fitted curves of infected SARS cases in

Beijing was larger than that of fitted curves of infected COVID-19 cases in each

province in mainland China (Fig 3E). Moreover, the increasing rate (k) was negatively

correlated (Pearson, r=-0.47, p=0.0072) to semi-saturation period (Fig 3F). We

concluded that at the provincial level, the COVID-19 spread more quickly than the

SARS did. The infections of COVID-19 increased more rapid and would reach

saturation earlier than SARS according to our model.

---------------------------- Insert Figure 3 near here --------------------------------

The outbreak of SARS in Beijing was the most serious among all the provincial

administrative regions, and the outbreak of COVID-19 was most serious in Hubei

province. Here we used our descriptive model to fit the dead and cured cases' curves

for SARS in Beijing and COVID-19 in Hubei province (Fig 4 A-C, F-H, R-square≥

0.95), and compared the growth features of three types cases of SARS and COVID-19

by two parameters, k and t_0 (Fig 4 DEIJ). We found that k values of the three types

cases of SARS were lower than that of COVID-19 (Fig 4 DE), and semi-saturation

period for them in SARS was higher than COVID-19's, which suggested the duration

of COVID-19 would be shorter than that of SARS. For both SARS and COVID-19,

their infected curves have the lowest semi-saturation period (Fig 4 EJ), while cured

curves have the highest which is in keeping with the common sense that the cure is

always later than infection and death. For both SARS in Beijing and COVID-19 in

Hubei, the k values of death case curves were lowest compared with curves of

infected cases and cured cases. For both SARS and COVID-19, the k value of infected

case curve was larger than that of cured curve.

---------------------------- Insert Figure 4 near here --------------------------------

Discussion

We modeled the epidemiologic characteristics of 2019-nCoV for 31 provinces in

mainland China based on the data from Jan 20 to Feb 19 during the outbreak.

Furthermore, we compared these characteristics with SARS in 2003 using the same

model framework.

Compared with dynamical models such as SIR (Susceptible – Infectious - Recovered)

or SEIR (Susceptible – Exposed - Infectious - Recovered), our model is relatively

simple but robust. We used a differential equation in logistic form to describe the

dynamics of the infected, dead and healed cases caused by COVID-19 respectively

(See Methods). The increasing of the population no matter in which type of case

(infection, death or recovery) has multiple reasons. When the outbreak of COVID-19

originated in Wuhan, it did not cause too much concern at the beginning so that it

leads to many infections and starts to export potential exposed cases to other

provinces. But as the successive closure of the cities in Hubei province, and isolation

of the community in all the provinces in mainland China, the virus spread tendency

was weakened. In our model, we also considered these measures taken by the

government. We rewrote Eq. (1) to Eq. (9).

$$\frac{dI(t)}{dt} = \alpha I(t) - rI^2(t) \tag{9}$$

where α is growth rate of the number of infections; *I(t)* is cumulative number of

infections at time *t*; r could be understood as a parameter to reflect the effect of

measures aiming at controlling the virus spread. Here we have decomposed $I(t)$ into

two parts. The first part $\alpha I(t)$ could represent the rapid infections growth in the early

phase of the outbreak without any measures for epidemic control. The second part

$rI^2(t)$ illustrates the slower infections growth after taking effective isolation

measures for epidemic control. For death cases, this idea still holds. But things are

different for cured case. In contrast, at the early phase of the outbreak, the number of

cured cases increase slowly. But once there are effective treatments, the number of

cured cases will rise up quickly. The growth of cured cases would again be slow since

the number of patients is limited. All these facts were considered in our highly

summarized model (Eq. (8)) in a macroscopic scale. All data sets have a R-square

greater than 0.95 when fitted by our framework, so that we could further discuss some

intrinsic mechanisms for the growth rules of infection, death and recovery through the

parameters estimated from the model (See Methods).

The value of k from Eq. (8) is a critical parameter. k represents how rapid the number

of infected cases get to the saturation. We found the variability of k values among

fitted curves of infected COVID-19 cases in 30 provinces (Fig 2 AB) is large

(Ranging from 0.19 to 0.38). It is clear to show the real situation in each province and

this guides us to be more careful against the epidemic. Further we compared this

characteristics between COVID-19 and SARS in 2003 in typical provinces (Fig 3 AB).

We noted that k values of fitted COVID-19 infection curves in 30 provinces of

mainland China (exclude Tibet) are larger than that of SARS in Beijing 2003 (Fig 3

C), which suggests the time cost of this epidemic will be shorter compared with

SARS in 2003. This result may related to several aspects. Firstly, it may due to the

basic spread property of the virus. Besides, the isolation policy of all the provinces in

mainland China was more strictly enforced than it was in 2003. Moreover, the

improved clinical skills today may also helpful for shortening the duration of

epidemic.

Furthermore, we analyzed semi-saturation period (Fig 4 EK) (t_0 in Eq. (8)) and make

comparison in three types of cases (infection, death and recovery) (Fig 4 A-C G-I).

Our model could fit them very well (R-square $\geq (R-s)$. k values for the fitted

curves of the three types of cases caused by COVID-19 were larger than that of SARS,

while t_0 values were the opposite. This again suggests that the epidemic of COVID-19

will be quicker to pass away than SARS. For both SARS and COVID-19, fitted

curves of infections have lowest semi-saturation period (Fig 4 EK), while fitted

curves of cured cases have the highest semi-saturation period, which follows the

common sense that the cure was always later than infection and death since it costs

time to get on the research of effect plans to cure the patients. For both SARS and

COVID-19, the k value of infected case curve was larger than that of cured curve,

while that of death case is the lowest.

References

Drosten C, Günther S, Preiser W, et al. Identification of a novel coronavirus in

patients with severe acute respiratory syndrome. N Engl J Med 2003; 348: 1967–76.

Holshue ML et al. (2020) First Case of 2019 Novel Coronavirus in the United States.

N Engl J Med.

Huang C et al. (2020) Clinical features of patients infected with 2019 novel

coronavirus in Wuhan, China. Lancet.

HUANG De-sheng, GUAN Peng, ZHOU Bao-sen. Fitness of morbidity and

discussion of epidemic characteristics of SARS based on logistic models[J]. Chinese

Journal of Public Health, 2003, 19(6): 1-2.

Hui DS, E IA, Madani TA, Ntoumi F, Kock R, Dar O, Ippolito G, McHugh TD,

Memish ZA, Drosten C, Zumla A, Petersen E (2020) The continuing 2019-nCoV

epidemic threat of novel coronaviruses to global health - The latest 2019 novel

coronavirus outbreak in Wuhan, China. Int J Infect Dis 91:264-266.

Ki M, nCo VT (2020) Epidemiologic characteristics of early cases with 2019 novel

coronavirus (2019-nCoV) disease in Republic of Korea. Epidemiol Health:e2020007.

Ksiazek TG, Erdman D, Goldsmith CS, et al. A novel coronavirus associated with

severe acute respiratory syndrome. N Engl J Med 2003; 348: 1953–66.

Kuiken T, Fouchier RAM, Schutten M, et al. Newly discovered coronavirus as the

primary cause of severe acute respiratory syndrome. Lancet 2003; 362: 263–70.

Li Q et al. (2020) Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus-Infected Pneumonia. N Engl J Med.

Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM (2020) Public

responses to the novel 2019 coronavirus (2019-nCoV) in Japan: mental health

consequences and target populations. Psychiatry Clin Neurosci.

Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y,

Wang Y, Liu X, The compound Logistic model used to describe epidemic situation

dynamics of SARS in Beijing, JOURNAL OF CHINA JILIANG UNIVERSITY,

2005

WHO. Summary of probable SARS cases with onset of illness from 1 November

2002 to 31 July 2003. Dec 31, 2003.

https://www. who.int/csr/sars/country/table2004_04_21/en/ (accessedJan 19, 2020).

Zhao Y, Li Y, Wang X, Peng Z (2020) Clinical Characteristics of 138 Hospitalized

Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA.

Zhou P et al. (2020) A pneumonia outbreak associated with a new coronavirus of

probable bat origin. Nature.

Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R,

Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, China Novel

Coronavirus I, Research T (2020) A Novel Coronavirus from Patients with Pneumonia

in China, 2019. N Engl J Med.

Figure 1 Spatial distribution of the number of COVID-19 infected

cases in mainland China.

Panel A is the spatial distribution of the number of COVID-19 infected cases of

provinces in mainland China.

Panel B takes five provinces (Xinjiang, Beijing, Guangdong, Zhejiang and

Heilongjiang) as example to show the time series of infected cases and corresponding

fitted curve. The horizontal axis is the x_{th} day after Jan 1, 2020. And the vertical axis

denotes the number of infected cases in corresponding province. The black dots in

each panel are raw data of the cumulative infected cases. The red line is the fitted

curve by our descriptive model. The dashed red lines show the 95% confidence

interval of the fitted curves.

Figure 2 Spatial distribution of increasing rate (*k*) and the normalized fitted curves of each province of mainland China (exclude Tibet)

Panel A demonstrates the spatial distribution of parameter *k* of provinces in mainland China.

Panel B shows the fitted curves of infected cases in 30 provinces. The vertical axis is the number of infected cases (normalized by dividing the maximum number of infected cases in each province). The color of each province is consistent with Panel A.

Panel C is the spatial distribution of the predicted maximum number (parameter A in our model) of COVID-19 infected cases of provinces

Figure 3 Comparison of intrinsic epidemic rules between COVID-19 in Wuhan and SARS in Beijing (2003)

Panel A shows the raw data and fitted curve of SARS cases in Beijing. The horizontal axis is the x_{th} day after March 1, 2003. The vertical axis denotes the cumulative number of SARS cases in Beijing. The black dots are the raw data of the SARS cases. The red line is the fitted curve by our descriptive model. The dashed red lines are the 95% confidence interval of the fitted curves.

Panel B shows the raw data and fitted curve of COVID-19 cases in Hubei province. The horizontal axis is the x_{th} day after Jan 1, 2020. The vertical axis denotes the cumulative number of COVID-19 cases in Hubei. The black dots are raw data of the COVID-19 cases. The red line is the fitted curve by our descriptive model. The dashed red lines are the 95% confidence interval of the fitted curves.

Pannel C shows the fitted curves of infected SARS cases (gray curve) and COVID-19

(black curve), which are normalized by dividing the maximum number of infected

cases in each province.

Panel D illustrates the k value (increasing rate of infected cases) of fitted curves of

infected COVID-19 cases in each province in mainland China, where Hubei is

represented by the solid red square (the same in D and E). k value of fitted curves of

infected SARS cases in Beijing is represented by the solid red triangle (the same in D

and E).

Panel E illustrates the t_0 (semi-saturation period) of fitted curves of infected

COVID-19 cases in each province in mainland China, as well as t_0 of fitted curves of

infected SARS cases in Beijing. The scatter plot of k and t_0 is shown in Panel F,

indicating significant negative correlation between k and t_0.

Figure 4 Intrinsic rules of infection, death and recovery of COVID-19

and SARS2003

Panel A-C show the raw data and fitted curve of SARS cases in Beijing. Their

horizontal axises are the x_{th} day after Mar 1, 2003. The black dots in Panel A, B and C

are raw data of the SARS infected cases, dead cases and cured cases, respectively.

The red lines are the fitted curves by our descriptive model. The dashed red lines are

the 95% confidence interval of the fitted curves.

Panel D and Panel E respectively shows the k value and the t_0 of fitted curves of three

types cases caused by SARS (the circle, triangle and square denotes infected, dead

and cured cases, respectively, and the same in Panel I and J). The error bars show the

95% confidence interval of the parameters.

Panel F-H show the raw data and fitted curve of COVID-19 cases in Hubei. Their

horizontal axises are the x_{th} day after Jan 1, 2020. The black dots in Panel F, G and H

are raw data of the COVID-19 infected cases, dead cases and cured cases,

respectively. The red lines are the fitted curves by our descriptive model. The dashed

red lines are the 95% confidence interval of the fitted curves.

Panel I and Panel J respectively shows the k value and the t_0 of fitted curves of three

types cases caused by COVID-19, which is same with D and E.